
Article

Maxwell Equations in Homogeneous Spaces with Solvable Groups of Motions

V. V. Obukhov

Special Issue

Physics and Symmetry Section: Feature Papers 2022

Edited by

Prof. Dr. Tomohiro Inagaki and Dr. Olga Kodolova



Article

Maxwell Equations in Homogeneous Spaces with Solvable Groups of Motions

V. V. Obukhov ^{1,2}

¹ Institute of Scientific Research and Development, Tomsk State Pedagogical University (TSPU), 60 Kievskaya St., Tomsk 634041, Russia; obukhov@tspu.edu.ru

² Laboratory for Theoretical Cosmology, International Center of Gravity and Cosmos, Tomsk State University of Control Systems and Radio Electronics (TUSUR), 36, Lenin Avenue, Tomsk 634050, Russia

Abstract: The classification of exact solutions of Maxwell vacuum equations for the case where the electromagnetic fields and metrics of homogeneous spaces are invariant with respect to the motion group $G_3(VII)$ was completed. All non-equivalent exact solutions of Maxwell vacuum equations for electromagnetic fields and spaces with such symmetry were obtained. The vectors of the canonical frame of a homogeneous space of type VII according to the Bianchi classification and the electromagnetic field potentials were found.

Keywords: Maxwell equations; Klein–Gordon–Fock equation; algebra of symmetry operators; theory of symmetry; linear partial differential equations

1. Introduction



Citation: Obukhov, V.V. Maxwell Equations in Homogeneous Spaces with Solvable Groups of Motions. *Symmetry* **2022**, *14*, 2595. <https://doi.org/10.3390/sym14122595>

Academic Editors: Tomohiro Inagaki and Olga Kodolova

Received: 11 November 2022

Accepted: 2 December 2022

Published: 8 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

A special place in mathematical physics is occupied by the problem of the exact integration of the field equations for electromagnetic and gravitational fields. The problem can be successfully solved if the space and the electromagnetic fields possess some symmetry. Homogeneous spaces are one of the important examples of the space manifolds with symmetry. Stackel spaces are another example of such spaces. Both of these sets of spaces are applied in the theory of electromagnetism and gravitation due to the fact that, in these spaces, methods of commutative and noncommutative integration of equations of motion of single test particles can be applied.

The methods of commutative integration is based on the use of a commutative algebra of symmetry operators (integrals of motion) that form a complete set. The complete set includes first- and second-degree linear operators in momentum formed from complete sets of geometric objects consisting of vector and tensor Killing fields. The method is known as the method of the complete separation of variables. The theory of the complete separation of variables was mainly constructed in the works [1–7]. A description of the theory and detailed bibliography can be found in [8–10]. Examples of applications of the theory of complete separation of variables in the theory of gravitation can be found in the works [11–16]. The methods of non-commutative integration is based on the use of the algebra of symmetry operators, which are linear in momenta and constructed using noncommutative Killing vector fields forming noncommutative groups of motion of spacetime G_3 . Among these spacetime manifolds, the homogeneous spaces are of greatest interest for the theory of gravity (see, for example, [17–27]). The theory of the noncommutative integration method and development of the theory can be found in the works [28–33].

Thus, these two methods are essentially complementary and have similar classification problems (by solving a classification problem, we mean enumerating all metrics of the corresponding spaces that are not equivalent in terms of admissible transformations of privileged coordinate systems; likewise, all electromagnetic potentials of admissible elec-

tromagnetic fields that are not equivalent in terms of admissible gradient transformations). Among these classification problems, the most important are the following.

The classification of all metrics of the Stackel and homogeneous spaces in privileged coordinate systems. For Stackel spaces, this problem was solved in the papers cited above. For homogeneous spaces, this problem was solved in the work of Petrov (see [34]).

The classification of all (admissible) electromagnetic fields to which these methods can be applied. For the Hamilton–Jacobi and Klein–Gordon–Fock equations, this problem is completely solved in homogeneous spaces (see [30–32]). In Stackel spaces, it is completely solved for the Hamilton–Jacobi equation (see [8–10]) and partially solved for the Klein–Gordon–Fock equation.

The classification of all vacuum and electrovacuum solutions of the Einstein equations with metrics of Stackel and homogeneous spaces in admissible electromagnetic fields. This problem is completely solved for the Stackel metric (see, for example, [5,12,13] and bibliography in [8–10]). For homogeneous spaces, this classification problem has not yet been studied.

Thus, for the complete solution of the problem of uniform classification, it remains to integrate the Einstein–Maxwell vacuum equations using the previously found potentials of admissible electromagnetic fields and the known metrics of homogeneous spaces in privileged (canonical) coordinate systems. This problem can also be divided into two stages. In the first stage, all solutions of Maxwell vacuum equations for the potentials of admissible electromagnetic fields should be found.

In the paper [33], the first problem was decided for the case where there exist groups G_3 (II–VI) in the homogeneous spaces. The present work is devoted to the homogeneous spaces with groups of motion G_3 (VII). Thus, the classification problem for solvable groups of motions will be solved.

2. Maxwell Equations in the Homogeneous Spaces

Homogeneous Spaces

By definition, a space–time manifold V_4 is a homogeneous space if a three-parameter group of motions acts on it whose transitivity hypersurface V_3 is endowed with the Euclidean space signature. A semi-geodesic coordinate system $[u^i]$ is used. The metric V_4 has the form:

$$ds^2 = g_{ij}du^i du^j = -du^0{}^2 + g_{\alpha\beta}du^\alpha du^\beta, \quad \det|g_{\alpha\beta}| > 0. \quad (1)$$

Coordinate indices of the variables of the semi-geodesic coordinate system are denoted by lower-case Latin letters: $i, j, \dots = 0, 1, \dots, 3$. The coordinate indices of the variables of the local coordinate system on the hypersurface V_3 are denoted by lower-case Greek letters: $\alpha, \beta, \gamma, \dots = 1, \dots, 3$. The temporal variable is indexed by 0. Group indices and indices of a non-holonomic frame are denoted by $a, d, c, \dots = 1, \dots, 3$. The letters p, q denote the indices varying from 2 to 3. Summation is performed over repeated upper and lower indices within the index range.

Another definition of a homogeneous space exists, according to which, the spacetime V_4 is homogeneous if its subspace V_3 , endowed with the Euclidean space signature, admits a set of coordinate transformations (the group G_3 of motions spaces V_4) that allow us to connect any two points in V_3 (see, e.g., [35]). This definition directly implies that the metric tensor of the V_3 space can be represented as follows:

$$g_{\alpha\beta} = e_\alpha^a e_\beta^b \eta_{ab}(u^0), \quad e_{\alpha,0}^a = 0, \quad \eta_{ab} = \eta_{ab}(u^0). \quad (2)$$

while the form

$$\omega^a = e_\alpha^a du^\alpha$$

is invariant with respect to transformations of the group G_3 . The vectors of the frame e_a^α define a non-holonomic coordinate system in V_3 . The dual triplet of vectors e_a^α ($e_a^\alpha e_a^\beta = \delta_a^\beta$, $e_a^\alpha e_\beta^\alpha = \delta_\beta^\alpha$) constructs the operators of the G_3 algebra group:

$$\hat{Y}_a = e_a^\alpha \partial_a, \quad [\hat{Y}_a, \hat{Y}_b] = C_{ab}^c \hat{Y}_c. \quad (3)$$

In the following, this definition of homogeneous spaces is used. The electromagnetic field is invariant with respect to transformations of the group acting in the space. It has the form:

$$A_i = l_i^a \alpha_a \quad \alpha_a = \alpha_a(u^0). \quad (4)$$

3. Maxwell Equations

We consider the Maxwell equations with zero sources for electromagnetic potential (4):

$$\frac{1}{\sqrt{-g}} (\sqrt{-g} F^{ij})_{,j} = 0. \quad (5)$$

In the case where $i = 0$, from the system (5), it follows that: Answer: "Approved".

$$\frac{1}{\sqrt{-g}} (\sqrt{-g} g^{\alpha\beta} F_{0\beta})_{,\alpha} = \frac{1}{l} (l l_a^\alpha \eta^{ab} \dot{\alpha}_b)_{,\alpha} = (l_{a,\alpha}^\alpha + \frac{l_{|a}}{l}) \frac{\beta^a}{\eta} \quad (\beta^a = \eta^{ab} \eta \dot{\alpha}_b). \quad (6)$$

Notation used:

$$f_{|a} = l_a^\alpha f_{,\alpha}, \quad g = -\det |g_{\alpha\beta}| = -(\eta l)^2, \quad (\eta^2 = \det |\eta_{\alpha\beta}|, \quad l = \det |l_\alpha^a|).$$

The dots denote the time derivatives. Then, we have the first equation in the form:

$$(l_{a,\alpha}^\alpha + l_{|a}) \beta^a = 0. \quad (7)$$

If $i = \alpha$, from Equation (5), it follows that:

$$\frac{1}{\eta} (\eta g^{\alpha\beta} F_{0\beta})_{,\alpha} = \frac{1}{l} (l g^{\nu\beta} g^{\alpha\gamma} F_{\beta\gamma})_{,\nu} \Rightarrow \frac{1}{\eta} (\eta \eta^{ab} l_a^\alpha \dot{\alpha}_b)_{,\alpha} = \frac{1}{l} (l l_a^\nu l_b^\beta \eta^{ab} l_a^\alpha l_b^\gamma \eta^{\tilde{a}\tilde{b}} F_{\beta\gamma})_{,\nu} \Rightarrow \quad (8)$$

$$\frac{l_a^\alpha}{\eta} \dot{\beta}^a = \frac{1}{l} (l l_b^\beta l_a^\alpha l_b^\gamma F_{\beta\gamma})_{|a} \eta^{ab} \eta^{\tilde{a}\tilde{b}}. \quad (9)$$

$F_{\alpha\beta}$ can be found using the relations (2)–(4):

$$F_{\alpha\beta} = (l_{\beta,\alpha}^a - l_{\alpha,\beta}^a) \alpha_a = l_\beta^c l_c^\gamma l_\alpha^d l_d^\nu (l_{\gamma,\nu}^a - l_{\nu,\gamma}^a) \alpha_a = l_\beta^b l_\alpha^a l_\gamma^c (l_{a|b}^\gamma - l_{b|a}^\gamma) \alpha_c = l_\beta^b l_\alpha^a C_{ba}^c \alpha_c \Rightarrow \quad (10)$$

$$(l F^{\alpha\beta})_{,\beta} = \eta^{ab} \eta^{\tilde{a}\tilde{b}} C_{\tilde{b}\tilde{b}}^d \alpha_d ((l l_a^\alpha)_{|\tilde{a}} + l l_a^\alpha l_{\tilde{a},\gamma}^\gamma). \quad (11)$$

Structural constants of a group G_3 can be represent in the form:

$$C_{ab}^c = C_{12}^c \varepsilon_{\tilde{a}\tilde{b}}^{12} + C_{p3}^c \varepsilon_{\tilde{a}\tilde{b}}^{p3}, \quad (12)$$

where

$$\varepsilon_{ab}^{AB} = \delta_a^A \delta_b^B - \delta_b^A \delta_a^B.$$

From the relations:

$$(\varepsilon_{\tilde{a}\tilde{b}}^{AB} \eta^{a\tilde{a}} \eta^{b\tilde{b}}) = (\eta^{aA} \eta^{bB} - \eta^{aB} \eta^{bA}), \quad (13)$$

it follows that

$$\eta^2 \varepsilon_{cd}^c \eta^{ac} \eta^{bd} = (\eta_{33} \varepsilon_{12}^{ab} + \eta_{23} \varepsilon_{31}^{ab} + \eta_{13} \varepsilon_{23}^{ab}),$$

$$\begin{aligned}\eta^2 \varepsilon_{cd}^{31} \eta^{ac} \eta^{bd} &= (\eta_{22} \varepsilon_{31}^{ab} + \eta_{23} \varepsilon_{12}^{ab} + \eta_{12} \varepsilon_{23}^{ab}), \\ \eta^2 \varepsilon_{cd}^{23} \eta^{ac} \eta^{bd} &= (\eta_{13} \varepsilon_{12}^{ab} + \eta_{12} \varepsilon_{31}^{ab} + \eta_{11} \varepsilon_{23}^{ab}).\end{aligned}$$

Equations (5) take the form:

$$\eta \dot{\beta}^a = \delta_1^a (\gamma_1 C_{32}^1 - \gamma_2 (C_{31}^1 + \omega_3) + \gamma_3 (C_{21}^1 + \omega_2)) + \delta_2^a (\gamma_1 (C_{32}^2 + \omega_3) + \gamma_2 C_{13}^2 - \gamma_3 (C_{12}^2 + \omega_1)) + \delta_3^a (-\gamma_1 (C_{23}^3 + \omega_2) + \gamma_2 (C_{13}^3 + \omega_1) + \gamma_3 C_{21}^3), \quad (14)$$

$$\begin{aligned}\eta_{ab} \beta^b &= \eta \dot{\alpha}_a, \\ \omega_a \beta^a &= 0, \quad \omega_a = l_{a,\alpha}^\alpha + l_{|a}/l,\end{aligned} \quad (15)$$

where

$$\begin{aligned}\gamma_1 &= \sigma_1 \eta_{11} + \sigma_2 \eta_{12} + \sigma_3 \eta_{13}, \quad \gamma_2 = \sigma_1 \eta_{12} + \sigma_2 \eta_{22} + \sigma_3 \eta_{23}, \\ \gamma_3 &= \sigma_1 \eta_{13} + \sigma_2 \eta_{23} + \sigma_3 \eta_{33}, \quad \sigma_1 = C_{23}^a \alpha_a, \quad \sigma_2 = C_{31}^a \alpha_a, \quad \sigma_3 = C_{12}^a \alpha_a.\end{aligned}$$

Let us find sets of the Maxwell Equations (14)–(16) for all solvable groups.

Groups G_3 (I–VII)

The components of the metric tensor and structural constants C_{ab}^c were found by Petrov (see [28]). The components of the vector l_a^α were found in our work [35]:

$$e_a^\alpha = \delta_1^1 \delta_1^\alpha \exp(-ku^3) + \delta_2^2 (-\delta_1^\alpha \varepsilon u^3 \exp(-ku^3) + \delta_2^\alpha \exp(-nu^3)) + \delta_3^3 \delta_3^\alpha, \quad (17)$$

$$\begin{aligned}e_\alpha^a &= \delta_1^a \delta_1^1 \exp(ku^3) + \delta_2^2 (\delta_1^\alpha \varepsilon u^3 \exp nu^3 + \delta_2^\alpha \exp nu^3) + \delta_3^3 \delta_3^a, \\ C_{ab}^c &= k \delta_1^c \varepsilon_{ab}^{13} + (\varepsilon \delta_1^c + n \delta_2^c) \varepsilon_{ab}^{23}.\end{aligned} \quad (18)$$

Let us consider Maxwell Equations (14)–(16).

I. For the groups G(I–VI), the equations can be presented in the form:

(1) For the group $G_1(I)$ ($k = n = \varepsilon = 0$):

$$\dot{\beta}^a = 0, \quad \dot{\alpha}_a = \frac{1}{\eta} \eta_{ab} \beta^b \Rightarrow$$

Solution of the Maxwell Equations (14)–(16) has the form:

$$\beta^a = \text{const}, \quad \alpha_a = \beta^b \int \frac{1}{\eta} \eta_{ab} du^0; \quad (19)$$

(2) For the group $G_1(II)$ ($k = n = 0, \varepsilon = 1$):

$$\dot{\beta}^a = -\delta_1^a \alpha_1 \eta_{11}, \quad \dot{\alpha}_a = \frac{1}{\eta} \eta_{ab} \beta^b; \quad (20)$$

(3) For the group $G_1(III)$ ($k = 1, n = \varepsilon = 0$):

$$\dot{\beta}^a = -\delta_1^a \alpha_1 \eta_{22}, \quad \beta^3 = 0, \quad \dot{\alpha}_a = \frac{1}{\eta} \eta_{ab} \beta^b; \quad (21)$$

(4) For the group $G_1(IV)$ ($k = n = \varepsilon = 1$):

$$\dot{\beta}^a = -\delta_1^a ((\alpha_1 + \alpha_2) \eta_{11} + \alpha_2 \varepsilon \alpha_{12} - \alpha_1 \eta_{22}) + \delta_2^a ((\alpha_1 + \alpha_2) \eta_{11} - \alpha_1 \eta_{12}); \quad (22)$$

$$\beta^3 = 0, \quad \dot{\alpha}_a = \frac{1}{\eta} \eta_{ab} \beta^b;$$

(5) For the group $G_1(V)$ ($k = n = 1, \varepsilon = 0$):

$$\dot{\beta}^a = \delta_1^a(-\alpha_2\eta_{12} + \alpha_1\eta_{22}) + \delta_2^a(\alpha_1\eta_{12} - \alpha_2\eta_{11}), \quad \beta^3 = 0, \quad \dot{\alpha}_a = \frac{1}{\eta}\eta_{ab}\beta^b; \quad (23)$$

(6) For the group $G_1(VI)$ ($k = 1, n = 2, \varepsilon = 0$):

$$\dot{\beta}^a = -\delta_1^a(2\alpha_2\eta_{12} - \alpha_1\eta_{22}) + \delta_2^a(2\alpha_2\eta_{11} - \alpha_1\eta_{12}), \quad \beta^3 = 0, \quad \dot{\alpha}_a = \frac{1}{\eta}\eta_{ab}\beta^b. \quad (24)$$

Equations (20) and (24) were integrated into our work [35]. In the present paper, the solutions for the group $G(VII)$ were found.

II. Group $G(VII)$.

When obtaining the Maxwell equations for the groups $G_3(I-VI)$, the components of vector fields l_a^α could be constructed directly from the components of the metric tensor (see [35]). For the group $G(VII)$, this cannot be performed. Therefore, the vectors l_a^α must be found directly from the conditions (2). Consider these conditions for the structural constants of the group $G_3(VII)$:

$$C_{23}^a = -\delta_1^a + 2\delta_2^a \cos \alpha, \quad C_{13}^2 = 1, \quad \alpha = \text{const.}$$

By coordinate transformation of the form $\tilde{u}^\alpha = \tilde{u}^\alpha(u^\beta)$ the vector field l_3^α can be diagonalized:

$$l_3^\alpha = \delta_3^\alpha.$$

From the commutation relations, it follows that:

$$X_{1,3} = -X_2; \quad X_{2,3} = X_1 - 2X_2 \cos \alpha \Rightarrow l_2^\alpha = -l_{1,3}^\alpha, \quad l_{2,33}^\alpha + 2l_{1,3}^\alpha \cos \alpha + l_1^\alpha = 0. \quad (25)$$

Solution of the Equation (25) has the form:

$$l_1^\alpha = \exp(-q_3)(a_1^\alpha(u^p) \sin p_3 + b_1^\alpha(u^p) \cos p_3),$$

$$l_2^\alpha = -\exp(-q_3)(a_2^\alpha(u^p) \sin(p_3 - \alpha) + b_2^\alpha(u^p) \cos(p_3 - \alpha)),$$

where $p, q = 1, 2, q_3 = u^3 \cos \alpha, p_3 = u^3 \sin \alpha$. Since the operators X_p commute, the vectors a_q^p, a_q^p can be simultaneously diagonalized by coordinate transformations of the form $\tilde{u}^p = \tilde{u}^p(u^q)$:

$$a_q^p = \delta_q^p, \quad b_q^p = \delta_q^p,$$

From the commutation relations it follows that: $a_3^p = 0, b_3^p = 0$.

Thus, the vectors of the frame of the homogeneous space of type VII according to Bianchi can be represented in the form:

$$l_1^\alpha = \exp(-q_3)(\delta_1^\alpha \sin p_3 + \delta_2^\alpha \cos p_3), \quad (26)$$

$$l_2^\alpha = \exp(-q_3)(\delta_1^\alpha \sin(p_3 - \alpha) + \delta_2^\alpha \cos(p_3 - \alpha)), \quad l_3^\alpha = \delta_3^\alpha.$$

The Maxwell Equations will take the form:

$$\eta \dot{\beta}_a = \delta_1^a(\gamma_1 - 2\gamma_2 \cos \alpha) + \delta_2^a \gamma_2, \quad \Rightarrow \quad \gamma_2 = \eta \dot{\beta}_2, \quad \gamma_1 = \eta(\dot{\beta}_1 + 2\dot{\beta}_2 \cos \alpha). \quad (27)$$

The system of Maxwell's equations can be represented in the form:

$$\sigma \eta_{11} - \alpha_2 \eta_{12} = \gamma_1, \quad \sigma \eta_{12} - \alpha_2 \eta_{22} = \gamma_2 (\sigma = 2\alpha_2 \cos \alpha - \alpha_1); \quad (28)$$

$$\beta_1 \eta_{11} + \beta_2 \eta_{12} = \eta \dot{\alpha}_1, \quad \beta_1 \eta_{12} + \beta_2 \eta_{22} = \eta \dot{\alpha}_2, \quad \beta_3 = 0; \quad (29)$$

$$\eta\dot{\alpha}_3 = \beta_1\eta_{13} + \beta_2\eta_{23} \Rightarrow \alpha_3 = \int \frac{\beta_1\eta_{13} + \beta_2\eta_{23}}{\eta} du_0. \quad (30)$$

From Equations (28) and (29), it follows that:

$$\eta_{11}(\alpha_2\dot{\alpha}_2 - \sigma\dot{\alpha}_1)(\alpha_2\beta_1 + \sigma\beta_2) = \gamma_1\beta_2(\alpha_2\dot{\alpha}_2 - \sigma\dot{\alpha}_1) - \alpha_2\dot{\alpha}_2(\beta_1\gamma_1 + \beta_2\gamma_2). \quad (31)$$

$$\alpha_1\dot{\alpha}_2(\eta(\alpha_2\dot{\alpha}_2 - \sigma\dot{\alpha}_1) + \beta_1\gamma_1 + \beta_2\gamma_2) = 0. \quad (32)$$

When solving the system of Equations (31) and (32), the variants that need to be considered are:

A. $\alpha_2 \neq 0$. From the system of Equation (29), it follows:

$$\eta_{11}(\alpha_2\beta_1 + \sigma\beta_2) = \eta(\dot{\alpha}_1\alpha_2 + \dot{\beta}_1\beta_2), \quad \eta_{12} = \frac{1}{\alpha_2}(\sigma_1\eta_{11} - \eta\tilde{\beta}_1), \quad \eta_{22} = \frac{1}{\alpha_2^2}(\sigma_1^2\eta_{11} - \eta(\sigma_1\tilde{\beta}_1 + \alpha_2\dot{\beta}_2)). \quad (33)$$

When solving the set of Equations (31) and (33), the following variants must be considered:

1. $(\alpha_2\dot{\alpha}_2 - \sigma\dot{\alpha}_1) \neq 0 \Rightarrow \eta_{11} = \eta \frac{\dot{\alpha}_1\alpha_2 + \dot{\beta}_1\beta_2}{\alpha_2\beta_1 + \sigma\beta_2}$. We consider Equation (32):

We will use the following notations:

$$\alpha_1 = \sqrt{\rho} \sin(\omega/2), \quad \alpha_2 = \sqrt{\rho} \cos(\omega/2), \quad \Omega = (2\beta_2\dot{\beta}_1 \cos \alpha + \beta_1\dot{\beta}_1 + \beta_2\dot{\beta}_2), \quad \omega = \omega(u^0),$$

(1) Let $\alpha_1 \neq 0$. Then the Equation (34) can be reduced to the form:

$$2\alpha_2\dot{\alpha}_1 \cos \alpha - \alpha_1\dot{\alpha}_1 - \alpha_2\dot{\alpha}_2 = 2\beta_2\dot{\beta}_1 \cos \alpha + \beta_1\dot{\beta}_1 + \beta_2\dot{\beta}_2. \quad (34)$$

(a) $\dot{\omega} \neq 0$. In this case, we take the function ω as a new time variable and denote by the point the derivative on this variable. The functions β_p, ρ depend on ω . Then the Equation (34) can be reduced to the form:

$$\dot{\rho}(\cos \alpha \sin \omega - 1) + \cos \alpha(1 + \cos \omega)\rho = 2\Omega. \quad (35)$$

The function ρ can be represented in the form: $\rho = \mathfrak{R}(\omega)\tau(\omega)$, where

$$\mathfrak{R} = \int \frac{\cos \alpha(1 + \cos \omega)}{1 - \cos \alpha \sin \omega} d\omega,$$

The function τ has the form:

$$\tau = (c + 2 \int \frac{\Omega}{\mathfrak{R}(1 - \cos \alpha \sin \omega)} d\omega),$$

$$(b) \omega = a = const \Rightarrow \rho = (c - 2 \int \frac{\Omega}{1 - \cos \alpha \sin \omega} du^0),$$

$$2. \alpha_1 = 0, \quad \eta_{11} = \eta \frac{\beta^2\dot{\beta}}{\alpha_2}, \quad \eta_{12} = -\eta \frac{\beta^1\dot{\beta}}{\alpha_2}, \quad \eta_{22} = -\eta \frac{\beta^2 + 2\cos \alpha \beta^1\dot{\beta}}{\alpha_2}, \quad \beta = \ln(\beta^1 + 2\cos \alpha \beta^2)$$

The final solutions are represented in **Solutions**.

3. $\alpha_2\beta_1 + \sigma\beta_2 = 0, \eta_{11}$ is an arbitrary function of u^0 . In this case, there are two variants to consider:

(a) $\alpha_1 = 0 \Rightarrow$ function η_{pq} can be found from (33).

(b) $\alpha_1 \neq 0 \Rightarrow \dot{\alpha}_1\alpha_2 + \dot{\beta}_1\beta_2 = \dot{\alpha}_2\alpha_2 + \dot{\beta}_2\beta_2 = 0 \Rightarrow \alpha_2\dot{\alpha}_2 + \beta_2\dot{\beta}_2 = 0$ ($\beta = 2\beta_2 \cos \alpha + \beta_1$). From the last equation it follows that:

$$\alpha_2 = c \sin \omega \beta_2 = c \cos \omega.$$

B. $\alpha_2 = 0$. In this case, from the set of Equations (28) and (29), it follows that:

$$\alpha_1\dot{\alpha}_1 + \beta_1\dot{\beta}_1 + 2\cos \alpha \beta_1\dot{\beta}_2 = 0, \quad \alpha_1 = \sqrt{c - (\beta^1)^2} - 4\cos \alpha \int \beta_1\dot{\beta}_2 du^0.$$

The functions η_{ab} are determined from Equations (28) and (29). The results are given in the **Solutions**.

4. Solutions

In this section, all solutions of Maxwell's vacuum equations for homogeneous Bianchi type VII spaces and electromagnetic fields invariant with respect to the groups of motions $G_3(VII)$ are given. For all solutions, the functions α_3 and η_{33} have the form:

$$\alpha_3 = \int (\eta_{13}\beta^1 + \eta_{23}\beta^2)du^0, \quad \eta_{33} = \frac{\eta^2 - 2\eta_{12}\eta_{13}\eta_{23} + \eta_{11}\eta_{23}^2 + \eta_{22}\eta_{13}^2}{\eta_{11}\eta_{22} - \eta_{13}^2}.$$

Other functions that specify solutions are shown below.

4.1. $\alpha_2 \neq 0$

The functions η_{12}, η_{22} have the form:

$$\eta_{12} = \frac{1}{\alpha_2}(\sigma_1\eta_{11} - \eta\dot{\beta}), \quad \eta_{22} = \frac{1}{\alpha_2^2}(\sigma_1^2\eta_{11} - \eta(\sigma_1\dot{\beta}^1 + \alpha_2\dot{\beta})), \quad \sigma_1 = 2\alpha_2 \cos \alpha - \alpha_1, \quad \beta = 2\beta^2 \cos \alpha + \beta^1.$$

$$(1) \beta^1\alpha_2 + \beta_2\sigma_1 \neq 0, \eta_{11} = \eta \frac{\dot{\alpha}_1\alpha_2 + \beta^2\dot{\beta}}{\beta^1\alpha_2 + \beta_2\sigma_1}, \Omega = (\beta^1\dot{\beta}^1 + \beta^2\dot{\beta}^2) + 2\beta^2\dot{\beta}^1 \cos \alpha.$$

$$(a) \alpha_1 = \sqrt{\rho} \sin c, \quad \alpha_2 = \sqrt{\rho} \cos c, \quad \rho = \int \frac{2\Omega du^0}{\cos \alpha \sin c - 1}.$$

$$(b) \alpha_1 = \sqrt{\rho} \sin \frac{\omega}{2}, \quad \alpha_2 = \sqrt{\rho} \cos \frac{\omega}{2}, \quad \omega = \omega(u^0), \quad \beta^p = \beta^p(\omega), \quad \dot{\beta}^p = \partial \beta^p / \partial \omega,$$

$$\rho = \frac{\Re}{1 - \cos \alpha \sin \omega} (c - 2 \int \frac{\Omega(1 - \cos \alpha \sin \omega) d\omega}{\Re}), \Re = \exp \int \frac{\cos \alpha d\omega}{1 - \cos \alpha \sin \omega}$$

$$(c) \alpha_1 = 0, \eta_{11} = \eta \frac{\beta^2\tilde{\beta}}{\alpha_2}, \eta_{12} = -\eta \frac{\beta^1\tilde{\beta}}{\alpha_2}, \eta_{22} = -\eta \frac{\beta^2 + 2\cos \alpha \beta^1\tilde{\beta}}{\alpha_2} \tilde{\beta} = (\ln(\beta^1 + 2\cos \alpha \beta^2))_0.$$

(2) η_{11} is an arbitrary function of u^0 .

$$(a) \alpha_1 = 0, \eta_{12} = 2\eta_{11} \cos \alpha - \eta, \eta_{22} = 4\eta_{11} \cos^2 \alpha - \eta \frac{2\dot{\beta} \cos \alpha + \dot{\beta}}{\alpha_2}.$$

$$(b) \alpha_1 = ac \sin \omega, \alpha_2 = c \sin \omega, \beta_2 = c \cos \omega, \beta_1 = c(a - 2 \cos \alpha) \cos \omega, c, a = \text{const}$$

$$\eta_{12} = (2 \cos \alpha - a)\eta_{11} + a\eta, \eta_{22} = (2 \cos \alpha - a)^2\eta_{11} + \eta(a(2 \cos \alpha - a) + 1).$$

4.2. $\alpha_2 = 0$

$$1. \alpha_1 = \sqrt{c - (\beta^1)^2 - 4 \cos \alpha \int \beta^1 \dot{\beta}^2 du^0}, \eta_{11} = -\eta \frac{(2 \cos \alpha \dot{\beta}_2 + \dot{\beta}_1)}{\alpha_1}, \eta_{12} = -\eta \frac{\dot{\beta}_2}{\alpha_1}, \eta_{22} = \eta \frac{\dot{\beta}_2 \beta_1}{\beta_2 \alpha_1}.$$

2. $\beta_2 = 0, \alpha_1 = c \sin \omega, \beta_1 = c \cos \omega, \eta_{22}, \omega$ are arbitrary functions of u^0 .

$$\eta_{12} = 0, \quad \eta_{11} = -\eta \dot{\omega}, \quad \eta_{13} = \eta \frac{\dot{\alpha}_3}{\beta_1}.$$

All functions included in these expressions that are not additionally described (for example, η, η_{p3} , and so on) are arbitrary functions of u^0 .

5. Conclusions

In the paper, the classification of solutions of vacuum Maxwell equations for the case where the electromagnetic fields and the metrics of homogeneous spaces are invariant with respect to solvable groups of motions was completed (for the groups $G_3(I-VI)$, classification was carried out in the paper [35]). Since this classification was carried out in the canonical frame (2), it allows one to proceed with the classification of exact solutions of the vacuum Einstein–Maxwell equations for the found fields. This will be of interest for the study of the early stages of the evolution of the Universe.

Funding: The work is supported by Russian Science Foundation, project number N 23-21-00275.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data that support the findings of this study are available within the article.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Stackel, P. Über die integration der Hamiltonschen differentialechung mittels separation der variablen. *Math. Ann.* **1897**, *49*, 145–147. [\[CrossRef\]](#)
2. Eisenhart, L.P. Separable systems of stackel. *Ann. Math.* **1934**, *35*, 284–305. [\[CrossRef\]](#)
3. Levi-Civita, T. Sulla Integrazione Della Equazione Di Hamilton-Jacobi Per Separazione Di Variabili. *Math. Ann.* **1904**, *59*, 383–397. [\[CrossRef\]](#)
4. Jarov-Jrovoy, M.S. Integration of Hamilton-Jacobi equation by complete separation of variables method. *J. Appl. Math. Mech.* **1963**, *27*, 173–219.
5. Carter, B. New family of Einstein spaces. *Phys. Lett.* **1968**, *25*, 399–400. [\[CrossRef\]](#)
6. Shapovalov, V.N. Symmetry and separation of variables in the Hamilton-Jacobi equation. *Sov. Phys. J.* **1978**, *21*, 1124–1132. [\[CrossRef\]](#)
7. Shapovalov, V.N. Stackel’s spaces. *Sib. Math. J.* **1979**, *20*, 1117–1130. [\[CrossRef\]](#)
8. Obukhov, V.V. Hamilton-Jacobi equation for a charged test particle in the Stackel space of type (2.0). *Symmetry* **2020**, *12*, 1289. [\[CrossRef\]](#)
9. Obukhov, V.V. Hamilton-Jacobi equation for a charged test particle in the Stackel spaces of type (2.1). *Int. J. Geom. Meth. Mod. Phys.* **2020**, *14*, 2050186. [\[CrossRef\]](#)
10. Obukhov, V.V. Separation of variables in Hamilton-Jacobi and Klein-Gordon-Fock equations for a charged test particle in the stackel spaces of type (1.1). *Int. J. Geom. Meth. Mod. Phys.* **2021**, *18*, 2150036. [\[CrossRef\]](#)
11. McLenaghan, R.G.; Rastelli, G.; Valero, C. Complete separability of the Hamilton-Jacobi equation for the charged particle orbits in a Lienard-Wiehert field. *J. Math. Phys.* **2020**, *61*, 122903. [\[CrossRef\]](#)
12. Bagrov, V.G.; Obukhov, V.V. Classes of exact solutions of the Einstein-Maxwell equations. *Am. Phys.* **1983**, *40*, 181–188. [\[CrossRef\]](#)
13. Bagrov, V.G.; Obukhov, V.V. Separation of variables for the Klein-Gordon equation in special staekel space-times. *Class. Quant. Grav.* **1990**, *7*, 19–25. [\[CrossRef\]](#)
14. Osetrin, K.; Osetrin, E.; Osetrina, E. Geodesic deviation and tidal acceleration in the gravitational wave of the Bianchi type IV universe. *Eur. Phys. J. Plus* **2020**, *137*, 856. [\[CrossRef\]](#)
15. Osetrin, K.; Filippov, A.; Osetrin, E. Wave-like spatially homogeneous models of Stackel spacetimes (2.1) type in the scalar-tensor theory of gravity. *Mod. Phys. Lett. A* **2020**, *35*, 2050275. [\[CrossRef\]](#)
16. Kumaran, Y.; Ovgun, A. Deflection angle and shadow of the reissner-nordstrom black hole with higher-order magnetic correction in einstein-nonlinear-maxwell fields. *Symmetry* **2022**, *14*, 2054. [\[CrossRef\]](#)
17. Osetrin, K.; Kirnos, I.; Osetrin, E.; Filippov, A. Wave-like exact models with symmetry of spatial homogeneity in the quadratic theory of gravity with a scalar field. *Symmetry* **2021**, *13*, 1173. [\[CrossRef\]](#)
18. Breev, A.I.; Shapovalov, A.V. Non-commutative integration of the Dirac equation in homogeneous spaces. *Symmetry* **2020**, *12*, 1867. [\[CrossRef\]](#)
19. Mitsopoulos, A.; Mitsopoulos, A.; Tsamparlis, M.; Leon, G.A.; Paliathanasis, A. New conservation laws and exact cosmological solutions in Brans-Dicke cosmology with an extra scalar field. *Symmetry* **2021**, *13*, 1364. [\[CrossRef\]](#)
20. Breev, A.I.; Shapovalov, A.V. Vacuum quantum effects on Lie groups with bi-invariant metrics. *Int. J. Geom. Methods Mod. Phys. (IJGMMP)* **2019**, *16*, 1950122. [\[CrossRef\]](#)
21. Breev, A.I.; Shapovalov, A.V. Yang–Mills gauge fields conserving the symmetry algebra of the Dirac equation in a homogeneous space. *J. Phys. Conf. Ser.* **2014**, *563*, 012004. [\[CrossRef\]](#)
22. Epp, V.; Pervukhina, O.N. The Stormer problem for an aligned rotator. *Mon. Not. R. Astron. Soc.* **2018**, *474*, 5330–5339. [\[CrossRef\]](#)
23. Breev, A.I.; Shapovalov, A.V. Symmetry operators and separation of variables in the (2 + 1)-dimensional Dirac equation with external electromagnetic field. *Int. J. Geom. Methods Mod. Phys. (IJGMMP)* **2018**, *15*, 1850085.
24. Capozziello, S.; De Laurentis, M.; Odintsov, D. Hamiltonian dynamics and Noether symmetries in extended gravity cosmology. *Eur. Phys. J.* **2012**, *72*, 2068. [\[CrossRef\]](#)
25. Nojiri, S.; Odintsov, S.D.; Faraoni, V. Searching for dynamical black holes in various theories of gravity. *Phys. Rev. D* **2021**, *103*, 044055. [\[CrossRef\]](#)
26. Magazev, A.A.; Boldyreva, M.N. Schrodinger equations in electromagnetic fields: Symmetries and noncommutative integration. *Symmetry* **2021**, *13*, 1527. [\[CrossRef\]](#)
27. Breev, A.I.; Shirokov, I.V.; Magazev, A.A. Vacuum polarization of a scalar field on lie groups and homogeneous spaces. *Theor. Math. Phys.* **2011**, *167*, 468–483. [\[CrossRef\]](#)

28. Shapovalov, A.V.; Shirokov, I.V. Noncommutative integration method for linear partial differential equations. functional algebras and dimensional reduction. *Theor. Math. Phys.* **1996**, *106*, 1–10. [[CrossRef](#)]
29. Magazey, A.A. Integrating Klein-Gordon-Fock equations in an extremal electromagnetic field on Lie groups. *Theor. Math. Phys.* **2012**, *173*, 1654–1667. [[CrossRef](#)]
30. Obukhov, V.V. Algebra of symmetry operators for Klein-Gordon-Fock Equation. *Symmetry* **2021**, *13*, 727. [[CrossRef](#)]
31. Obukhov, V.V. Algebra of the symmetry operators of the Klein-Gordon-Fock equation for the case when groups of motions G_3 act transitively on null subsurfaces of spacetime. *Symmetry* **2022**, *14*, 346. [[CrossRef](#)]
32. Obukhov, V.V. Algebras of integrals of motion for the Hamilton-Jacobi and Klein-Gordon-Fock equations in spacetime with a four-parameter groups of motions in the presence of an external electromagnetic field. *J. Math. Phys.* **2022**, *63*, 023505. [[CrossRef](#)]
33. Obukhov, V.V. Maxwell Equations in Homogeneous Spaces for Admissible Electromagnetic Fields. *Universe* **2022**, *8*, 245. [[CrossRef](#)]
34. Petrov, A.Z. *Einstein Spaces*; Pergamon Press: Oxford, UK, 1969.
35. Landau, L.D.; Lifshits, E.M. *Theoretical Physics, Field Theory*, 7th ed.; Moskow Science Chief Editorial Board for Physical and Mathematical Literature; Nauka: Moscow, Russia, 1988; 512p, ISBN 5-02-014420-7.