

Recent developments in SHERPA

Jennifer Archibald¹, Tanju Gleisberg², Stefan Hoche¹, Frank Krauss¹,
Marek Schonherr³, Steen Schumann⁴, Frank Siegert¹[‡], Jan Winter⁵

1- IPPP Durham - Physics Department - Durham University - Durham DH1 3LE - United Kingdom
2- Stanford Linear Accelerator Center - Stanford University - Stanford CA 94309 - USA
3- Institute of Nuclear and Particle Physics - TU Dresden - D-01062 Dresden - Germany
4- The University of Edinburgh - Edinburgh EH9 3JZ - Scotland
5- Fermi National Accelerator Laboratory - Batavia - IL 60510 - USA

Some recent QCD-related developments in the SHERPA event generator are presented.

In the past decades, event generators such as PYTHIA [1, 2] and HERWIG [3, 4] have been central for nearly all physics analyses at particle physics experiments at the high-energy frontier. This will also hold true at the LHC, where a large number of interesting signals for new particles or new phenomena (the Higgs boson or any other manifestation of the mechanism behind electro-weak symmetry breaking, supersymmetry, extra dimensions etc.) is hampered by a plethora of severe, sometimes overwhelming backgrounds. Nearly all of them are largely influenced by QCD. Therefore it seems fair to say that the success of the LHC in finding new physics may very well depend on a deep and detailed understanding of old physics, like QCD. Examples for this include, among others, the central-jet veto for the vector boson fusion channel for Higgs production or topologies, where gauge bosons emerge in association with many jets, a background for many search channels.

In a reflection on increased needs by the experimental community, aiming at higher precision, incorporation of new physics models and so on, the work horses of old have undergone serious renovation efforts, resulting in new, improved versions of the respective codes, namely PYTHIA8 [5] and HERWIG++ [6]. In addition a completely new code, SHERPA [7], has been constructed and is in the process of maturing. The status of this code is the topic of this contribution.

SHERPA's hallmark property is the inclusion of higher-order tree-level QCD contributions, leading to an improved modelling of jet production. They are introduced through a full-edged matrix element generator, AMEGIC++ [8], which is capable of generating matrix elements and corresponding phase space mappings for processes with multi-particle final states in various models, including the Standard Model, anomalous gauge triple and quadruple couplings according to [9, 10], the Minimal Supersymmetric Standard Model with Feynman rules from [11], the ADD-model of extra dimensions [12, 13], and a model with an extra $U(1)$ singlet coupling to the Higgs boson only [14]. The code has been thoroughly tested and validated [15]. This code, however, is limited, especially in the treatment of many (6) external QCD particles. Therefore, in the near future, SHERPA will incorporate another, new matrix element generator, COMIX, which is based on Berends-Giele recursion relations [16] and colour-dressing [17] rather than colour-ordering. In Tabs. 1 and 2 some example cross sections for $gg \rightarrow ng$ at fixed energies and $pp \rightarrow bb + n$ jets obtained with this program are exhibited and compared to those from other programs.

supported by the HEPTOOLS Marie Curie RTN under contract number MRTN-CT-2006-035505

[†]Supported by STFC

[‡]supported by the MCnet Marie Curie RTN under contract number MRTN-CT-2006-035606

This manuscript has been authored by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics.

$gg \rightarrow ng$	Cross section [pb]				
n \sqrt{s} [GeV]	8 1500	9 2000	10 2500	11 3500	12 5000
Comix	0.755(3)	0.305(2)	0.101(7)	0.057(5)	0.019(2)
Maltoni	0.70(4)	0.30(2)	0.097(6)		
Alpgen	0.719(19)				

Table 1: Cross sections for multi-gluon scattering $gg \rightarrow ng$ at di erent center of mass energies \sqrt{s} , using the phase space cuts speci ed in [18] , compared to literature results. In parentheses the statistical error is stated in units of the last digit of the cross section.

$bb + n$ jets	[b]						
n	0	1	2	3	4	5	6
Comix	470.8(5)	8.83(2)	1.826(8)	0.459(2)	0.1500(8)	0.0544(6)	0.023(2)
ALPGEN	470.6(6)	8.83(1)	1.822(9)	0.459(2)	0.150(2)	0.053(1)	0.0215(8)
AMEGIC++	470.3(4)	8.84(2)	1.817(6)				

Table 2: Cross sections for $pp \rightarrow bb + n$ jets at the LHC, with cuts and de nitions according to those in the MC4LHC workshop, compared to literature results. In parentheses the statistical error is stated in units of the last digit of the cross section.

In addition, concerning the calculation of higher-order matrix elements and cross sections, there have been rst steps towards an automation of such caluclations at truly next-to leading order accuracy. They manifest themselves in the implementation of a procedure [19] to fully automatically construct and evaluate Catani-Seymour dipole subtraction terms [20] for the real part of such NLO calculations.

The results from the matrix element calculations are merged with the subsequent parton shower through the formalism of [21, 22]. The results of its implementation in SHERPA [23] has recently been compared with other algorithms [24]. Although there remains some dispute about the theoretical equivalence of the di erent approaches, the overall results show satisfying agreement with each other, such that they can be used with con dence for data analysis.

Furthermore, two new parton shower algorithms have been implemented. One is based on the Catani-Seymour subtraction kernels [25], following a proposal in [26]^a. It can be anticipated that this shower, together with the automated subtraction terms for the real part of the NLO calculations, will allow for a greatly alleviated construction of simulation of QCD radiation including NLO accuracy, in the spirit of MC@NLO or POWHEG. This new shower has been extensively compared with various data; as an illustrative example consider the azimuthal decorrelation of the two leading jets in jet production at the Tevatron, cf. Fig. 1, where the results of the simulation are confronted with data from [28]. The other new parton shower, presented in [29], constitutes a further development of the dipole-based shower formulation encoded in ARIADNE [30]. In the new model, initial state radiation is treated truly perturbatively, and is not replaced – as in the original proposal – by nal-state

^aThis algorithm has independently been also implemented in [27].

radiation involving the hadronic remnants and non-perturbative ideas to define their radiation properties. A first result underlining the power of this new formulation is shown in Fig. 2, where the p_{\perp} -distribution of the lepton pair in Drell-Yan processes at the Tevatron is confronted with data [31].




Figure 1: The azimuthal decorrelation of the two leading jets in pure QCD processes at the LHC, in different bins of the leading E_{\perp} . The new parton shower based on Catani-Seymour kernels is confronted with data from [28].

Figure 2: The p_{\perp} distribution of the Z boson in Drell-Yan processes at the Tevatron. The new dipole shower is confronted with data from [31].

Finally, the implementation of a new cluster hadronization model following [32] has been finished and it is now the default hadronization module of SHERPA. It has been supplemented with new modules for the simulation of hadron decays and of QED final state radiation in these decays.

In this contribution the current status of the SHERPA project has been summarised, with special emphasis on perturbative QCD issues, and some of its forthcoming attractions have been briefly discussed.

References

- [1] T. Sjostrand. High-energy physics event generation with pythia 5.7 and jetset 7.4. *Comput. Phys. Commun.*, 82:74, 1994.
- [2] Torbjorn Sjostrand, Stephen Mrenna, and Peter Skands. Pythia 6.4 physics and manual. *JHEP*, 05:026, 2006.
- [3] G. Marchesini et al. Herwig: A monte carlo event generator for simulating hadron emission reactions with interfering gluons. version 5.1 - april 1991. *Comput. Phys. Commun.*, 67:465–508, 1992.
- [4] G. Corcella et al. Herwig 6.5 release note. 2002.

- [5] Torbjorn Sjostrand, Stephen Mrenna, and Peter Skands. A brief introduction to pythia 8.1. *Comput. Phys. Commun.*, 178:852–867, 2008.
- [6] M. Bahr et al. Herwig++ physics and manual. 2008.
- [7] T. Gleisberg, S. Hoche, F. Krauss, A. Schalicke, S. Schumann, and J. Winter. Sherpa 1.alpha, a proof-of-concept version. *JHEP*, 0402:056, 2004.
- [8] F. Krauss, R. Kuhn, and G. So . Amegic++ 1.0: A matrix element generator in c++. *JHEP*, 0202:044, 2002.
- [9] K. Hagiwara, R. D. Peccei, D. Zeppenfeld, and K. Hikasa. Probing the weak boson sector in $e^+ e^- \rightarrow w^+ w^-$. *Nucl. Phys.*, B282:253, 1987.
- [10] G. J. Gounaris, J. Layssac, and F. M. Renard. O^- -shell structure of the anomalous z and gamma self-couplings. *Phys. Rev.*, D62:073012, 2000.
- [11] Janusz Rosiek. Complete set of feynman rules for the mssm – erratum. 1995.
- [12] Nima Arkani-Hamed, Savas Dimopoulos, and G. R. Dvali. The hierarchy problem and new dimensions at a millimeter. *Phys. Lett.*, B429:263–272, 1998.
- [13] T. Gleisberg, F. Krauss, K. T. Matchev, A. Schalicke, S. Schumann, and G. So . Helicity formalism for spin-2 particles. *JHEP*, 0309:001, 2003.
- [14] Athanasios Dedes, Terrance Figy, Stefan Hoche, Frank Krauss, and Thomas E. J. Underwood. Searching for nambu-goldstone bosons at the lhc. 2008.
- [15] T. Gleisberg, F. Krauss, C. G. Papadopoulos, A. Schalicke, and S. Schumann. Cross sections for multi-particle final states at a linear collider. *Eur. Phys. J. C*, 34:173, 2004.
- [16] Frits A. Berends and W. T. Giele. Recursive calculations for processes with n gluons. *Nucl. Phys.*, B306:759, 1988.
- [17] Claude Duhr, Stefan Hoche, and Fabio Maltoni. Color-dressed recursive relations for multi-parton amplitudes. *JHEP*, 08:062, 2006.
- [18] F. Maltoni, K. Paul, T. Stelzer, and S. Willenbrock. Color-flow decomposition of qcd amplitudes. *Phys. Rev.*, D67:014026, 2003.
- [19] Tanju Gleisberg and Frank Krauss. Automating dipole subtraction for qcd nlo calculations. *Eur. Phys. J.*, C53:501–523, 2008.
- [20] S. Catani and M. H. Seymour. A general algorithm for calculating jet cross sections in nlo qcd. *Nucl. Phys.*, B485:291–419, 1997.
- [21] S. Catani, F. Krauss, R. Kuhn, and B. R. Webber. Qcd matrix elements + parton showers. *JHEP*, 0111:063, 2001.
- [22] F. Krauss. Matrix elements and parton showers in hadronic interactions. *JHEP*, 0208:015, 2002.
- [23] A. Schalicke and F. Krauss. Implementing the me+ps merging algorithm. *JHEP*, 0507:018, 2005.
- [24] J. et al. Alwall. Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions. *Eur. Phys. J.*, C53:473–500, 2008.
- [25] Steen Schumann and Frank Krauss. A parton shower algorithm based on catani-seymour dipole factorisation. *JHEP*, 03:038, 2008.
- [26] Zoltan Nagy and Davison E. Soper. Matching parton showers to nlo computations. *JHEP*, 10:024, 2005.
- [27] Michael Dinsdale, Marko Ternick, and Stefan Weinzierl. Parton showers from the dipole formalism. *Phys. Rev.*, D76:094003, 2007.
- [28] V. M. Abazov et al. Measurement of dijet azimuthal decorrelations at central rapidities in pp collisions at $\sqrt{s} = 1.96$ tev. *Phys. Rev. Lett.*, 94:221801, 2005.
- [29] Jan-Christopher Winter and Frank Krauss. Initial-state showering based on colour dipoles connected to incoming parton lines. *JHEP*, 07:040, 2008.
- [30] Leif Lonnblad. Ariadne version 4: A program for simulation of qcd cascades implementing the color dipole model. *Comput. Phys. Commun.*, 71:15–31, 1992.
- [31] Anthony Allen A older et al. The transverse momentum and total cross section of e^+e^- pairs in the z boson region from pp collisions at $\sqrt{s} = 1.8$ tev. *Phys. Rev. Lett.*, 84:845–850, 2000.
- [32] J. Winter, F. Krauss, and G. So . A modified cluster-hadronization model. *Eur. Phys. J. C*, 36:381, 2004.