

Review

New Advancements in AdS/CFT in Lower Dimensions

Yolanda Lozano^{1,2,*} and Anayeli Ramirez^{1,2,*}¹ Department of Physics, University of Oviedo, Avda. Federico Garcia Lorca s/n, 33007 Oviedo, Spain² Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Calle de la Independencia 13, 33004 Oviedo, Spain.

* Correspondence: ylozano@uniovi.es (Y.L.); ramirezanayeli.uo@uniovi.es (A.R.)

† These authors contributed equally to this work.

Abstract: We review recent developments in the study of the AdS/CFT correspondence in lower dimensions. We start by summarising the classification of $\text{AdS}_3 \times \text{S}^2$ solutions in massive type IIA supergravity with $(0, 4)$ supersymmetries and the construction of their 2D dual quiver CFTs. These theories are the seed for further developments that we review next. First, we construct a new class of AdS_3 solutions in M-theory that describe M-strings in M5-brane intersections. Second, we generate a new class of $\text{AdS}_2 \times \text{S}^3$ solutions in massive IIA with four supercharges that we interpret as describing backreacted baryon vertices within the 5D $\mathcal{N} = 1$ QFT living in D4-D8 branes. Third, we construct two classes of AdS_2 solutions in Type IIB. The first are dual to discrete light-cone quantised quantum mechanics living in null cylinders. The second class is interpreted as dual to backreacted baryon vertices within 4D $\mathcal{N} = 2$ QFT living in D3-D7 branes. Explicit dual quiver field theories are given for all classes of solutions. These are used to compute the central charges of the CFTs that are shown to agree with the holographic expressions.

Keywords: string theory; AdS/CFT correspondence; supergravity

Citation: Lozano, Y.; Ramirez, A. New Advancements in AdS/CFT in Lower Dimensions. *Universe* **2021**, *7*, 250. <https://doi.org/10.3390/universe7070250>

Academic Editor: Norma G. Sanchez

Received: 28 June 2021

Accepted: 13 July 2021

Published: 19 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Lower dimensional CFTs¹ play a prominent role in the microscopic description of black holes and black strings. Since these exhibit, in the extremal case, AdS_2 and AdS_3 geometries close to their horizons, a deeper understanding of the AdS/CFT correspondence in lower dimensions is of key importance to their study.

The construction of AdS_2 and AdS_3 geometries and the identification of their dual superconformal field theories has been the focus of many interesting works. In general, the possible geometrical structures, supersymmetries preserved and topologies of the solutions increase as one increases the dimensions of the internal space, giving rise to a plethora of possible solutions (see for instance [1–45]). However, even if many classes of solutions with different amounts of supersymmetries have been constructed, this has only been paralleled with a detailed understanding of their dual CFTs for D1-D5 and D1-D5-D5' systems, and orbifolds thereof [16,46–51].

In this work, we review recent progress on the construction of $\text{AdS}_3/\text{CFT}_2$ and $\text{AdS}_2/\text{CFT}_1$ pairs with four supersymmetries where both sides of the correspondence are reasonably well-understood. These provide controlled settings where the AdS/CFT correspondence can be explicitly checked and where the black hole microstate counting programme can be carried out in detail.

Important progress in our understanding of the $\text{AdS}_3/\text{CFT}_2$ correspondence was provided by the recent constructions in [26]. These are solutions to massive Type IIA supergravity with $\mathcal{N} = (0, 4)$ supersymmetries (and $\text{SU}(2)$ R-symmetry) that are realised as fibrations of $\text{AdS}_3 \times \text{S}^2 \times M_4$ on an interval, with M_4 either a CY_2 or a 4d Kähler manifold. These solutions are dual to interesting classes of 2D CFTs admitting a quiver description in the UV that can be used to compute their degrees of freedom [52–54].

Besides providing explicit $\text{AdS}_3/\text{CFT}_2$ pairs, the constructions in [26,52–54] have been the seed of many other interesting developments. In a nutshell, new classes of AdS_3 solutions in M-theory with the same number of supersymmetries have been constructed in [34], which provide the holographic duals of the configurations of M-strings described in [55,56]. From the latter, new classes of solutions in massless Type IIA have been generated [36], which allow for an explicit defect interpretation as surface quivers embedded in a 6D CFT. Perhaps more interestingly, new examples of explicit $\text{AdS}_2/\text{CFT}_1$ duals have been derived from these $\text{AdS}_3/\text{CFT}_2$ pairs in both Type II supergravities [38,40,43].

The construction of new $\text{AdS}_2/\text{CFT}_1$ pairs is of special relevance. AdS_2 geometries arise as near horizon geometries of extremal black holes and therefore play a very important role in their microscopical studies. However, it is well-known that the precise realisation of an $\text{AdS}_2/\text{CFT}_1$ correspondence presents important technical and conceptual problems [57–60] that mainly originate from the fact that AdS_2 possesses two disjoint boundaries [61]. As a result, this correspondence is in need of a much deeper understanding.

A successful approach taken in [38,40,43] was to exploit its connection with the much better understood $\text{AdS}_3/\text{CFT}_2$ correspondence. At the geometrical level, AdS_3 and AdS_2 spaces are related by T-duality. This allows one to construct explicit $\text{AdS}_2/\text{CFT}_1$ pairs in which the CFT_1 arises as a discrete light-cone compactification of the 2D CFT dual to the AdS_3 solution, thus providing an explicit realisation of the constructions in [62–65]. Moreover, if two spheres are present in the internal space of an AdS_3 solution, such solutions are amenable to double analytical continuation techniques, which turn $\text{AdS}_3 \times S^2$ spaces into $\text{AdS}_2 \times S^3$ geometries. This latter class of solutions can then be dual to more general superconformal quantum mechanics (SCQM) than those arising from discrete light-cone compactification.

The aim of this review article is to summarise the main features of these recent developments. The paper is organised as follows. We start in Section 2 by reviewing the $\text{AdS}_3/\text{CFT}_2$ pairs constructed in [26,52–54], seeds of the forthcoming constructions. In Section 3, we review their uplift to M-theory, following [34], and briefly address their interpretation as duals to the M-strings in [55,56]. In Section 4, we turn our attention to the construction of new $\text{AdS}_2/\text{CFT}_1$ pairs in massive Type IIA, following [40]. We describe in some detail the associated dual SCQM, which allows one to interpret the solutions as backreacted D4-D0 baryon vertices in the 5D CFT living in D4'-D8 brane intersections. In Section 5, we discuss two more $\text{AdS}_2/\text{CFT}_1$ pairs, in this case in Type IIB, following [38,43]. A first class of solutions is constructed from the seed AdS_3 solutions reviewed in Section 2 using T-duality. These solutions are dual to explicit 1D CFTs that arise as discrete light-cone compactifications of the 2D CFTs reviewed in Section 2. The second class of solutions is constructed by T-dualising the AdS_2 class reviewed in Section 4. These solutions allow for an interesting interpretation as backreacted D5-D1 baryon vertices in the 4D $\mathcal{N} = 2$ QFT living in D3-D7 brane intersections, that we briefly discuss. Finally, in Section 6, we discuss open lines for further investigation.

2. $\text{AdS}_3/\text{CFT}_2$ in Massive IIA

In [26], a thorough classification of $\text{AdS}_3 \times S^2$ solutions to massive IIA supergravity consistent with a nontrivial Romans mass, with small (0, 4) supersymmetry and $\text{SU}(2)$ -structure, was obtained. The solutions are warped products of the form $\text{AdS}_3 \times S^2 \times M_4 \times I$, where M_4 is either a CY_2 or a 4D Kähler manifold. In this review, we restrict ourselves to the particular case when $M_4 = \text{CY}_2$, and there is no dependence on the coordinates of the CY_2 . These solutions provide the “seed” from which all supergravity backgrounds summarised in this work are derived. Furthermore, we review the proposal in [52–54] for their two-dimensional dual CFTs.

The Neveu–Schwarz (NS) sector of this class of solutions is as follows:

$$\begin{aligned}
 ds^2 &= \frac{u}{\sqrt{h_4 h_8}} \left(ds_{\text{AdS}_3}^2 + \frac{h_8 h_4}{4h_8 h_4 + u'^2} ds_{S^2}^2 \right) + \sqrt{\frac{h_4}{h_8}} ds_{\text{CY}_2}^2 + \frac{\sqrt{h_4 h_8}}{u} d\rho^2, \\
 e^{-\Phi} &= \frac{h_8^{\frac{3}{4}}}{2h_4^{\frac{1}{4}} \sqrt{u}} \sqrt{4h_8 h_4 + u'^2}, \quad H_3 = \frac{1}{2} d \left(-\rho + \frac{uu'}{4h_4 h_8 + u'^2} \right) \wedge \text{vol}_{S^2},
 \end{aligned}
 \tag{1}$$

where Φ is the dilaton, $H_3 = dB_2$ is the NS-NS three-form and the metric is written in string frame. The warping functions h_4, h_8 and u are functions of ρ , which parameterises an interval. We denote $' = \partial_\rho$. The RR fluxes are

$$\begin{aligned}
 F_0 &= h'_8, \quad F_2 = -\frac{1}{2} \left(h_8 - \frac{h'_8 u' u}{4h_8 h_4 + u'^2} \right) \text{vol}_{S^2}, \\
 F_4 &= - \left(d \left(\frac{uu'}{2h_4} \right) + 2h_8 d\rho \right) \wedge \text{vol}_{\text{AdS}_3} - h'_4 \text{vol}_{\text{CY}_2},
 \end{aligned}
 \tag{2}$$

with the higher fluxes related to them as $F_p = (-1)^{\lfloor p/2 \rfloor} \star_{10} F_{10-p}$. The background in (1) and (2) is a solution of the equations of motion if the functions h_4, h_8 and u satisfy

$$h''_4(\rho) = 0, \quad h''_8(\rho) = 0, \quad u''(\rho) = 0,
 \tag{3}$$

away from localised sources, which makes them linear functions of ρ . The first two equations are Bianchi identities, whereas $u'' = 0$ is a BPS equation.

The Page fluxes, defined as $\hat{F} = e^{-B_2} \wedge F$, are given by

$$\begin{aligned}
 \hat{F}_0 &= h'_8, \quad \hat{F}_2 = -\frac{1}{2} \left(h_8 - h'_8(\rho - 2\pi k) \right) \text{vol}_{S^2}, \\
 \hat{F}_4 &= - \left(\partial_\rho \left(\frac{uu'}{2h_4} \right) + 2h_8 \right) d\rho \wedge \text{vol}_{\text{AdS}_3} - h'_4 \text{vol}_{\text{CY}_2}.
 \end{aligned}
 \tag{4}$$

Here, we included large gauge transformations of B_2 of parameter k , $B_2 \rightarrow B_2 + \pi k \text{vol}_{S^2}$, for $k = 0, 1, \dots, P$. The transformations are performed every time a ρ -interval $\rho \in [2\pi k, 2\pi(k + 1)]$ is crossed. They ensure that B_2 satisfies the condition coming from String Theory:

$$\frac{1}{4\pi^2} \left| \int_{S^2} B_2 \right| \in [0, 1].
 \tag{5}$$

The most general solution to (3) is that h_4 and h_8 are piecewise linear functions. This allows for source branes in the geometry. In turn, u needs to be continuous for the preservation of supersymmetry. Here, we restrict it to the simpler case in which $u = \text{constant}$ ². In [52–54], piecewise linear solutions with the ρ direction bounded between 0 and $2\pi(P + 1)$, where h_4 and h_8 vanish, were proposed. These functions read

$$h_4(\rho) = \begin{cases} \frac{\beta_0}{2\pi} \rho & 0 \leq \rho \leq 2\pi, \\ \alpha_k + \frac{\beta_k}{2\pi} (\rho - 2\pi k) & 2\pi k \leq \rho \leq 2\pi(k + 1), \\ \alpha_P - \frac{\alpha_P}{2\pi} (\rho - 2\pi P) & 2\pi P \leq \rho \leq 2\pi(P + 1), \end{cases} \quad k = 1, \dots, P - 1
 \tag{6}$$

$$h_8(\rho) = \begin{cases} \frac{\nu_0}{2\pi} \rho & 0 \leq \rho \leq 2\pi, \\ \mu_k + \frac{\nu_k}{2\pi} (\rho - 2\pi k) & 2\pi k \leq \rho \leq 2\pi(k + 1), \\ \mu_P - \frac{\mu_P}{2\pi} (\rho - 2\pi P) & 2\pi P \leq \rho \leq 2\pi(P + 1). \end{cases} \quad k = 1, \dots, P - 1
 \tag{7}$$

The space begins at $\rho = 0$ in a smooth fashion. In turn, at $\rho = 2\pi(P + 1)$, the behaviour of the metric and dilaton is that of a superposition of D2-branes wrapped on AdS_3 and smeared on the $\text{CY}_2 \times S^2$, and of D6-branes wrapped on $\text{AdS}_3 \times \text{CY}_2$ ³.

The quantities $(\alpha_k, \beta_k, \mu_k, \nu_k)$ are integration constants and must satisfy

$$\alpha_k = \sum_{j=0}^{k-1} \beta_j, \quad \mu_k = \sum_{j=0}^{k-1} \nu_j. \tag{8}$$

in order for the metric and dilaton to be continuous. For the solutions defined by (6) and (7), the quantised charges associated with the Page fluxes given by (4) in the different $2\pi k \leq \rho \leq 2\pi(k + 1)$ intervals are

$$\begin{aligned} Q_{D2}^{(k)} &= \alpha_k = \sum_{j=0}^{k-1} \beta_j, & Q_{D6}^{(k)} &= \mu_k = \sum_{j=0}^{k-1} \nu_j \\ Q_{D4}^{(k)} &= \beta_k, & Q_{D8}^{(k)} &= \nu_k, & Q_{NS5}^{(k)} &= 1. \end{aligned} \tag{9}$$

The field theory duals to this class of solutions were studied in [52–54]⁴. We summarise them in the following subsection.

2.1. Two-Dimensional Dual CFTs

The backgrounds defined by Equations (1)–(3) are associated with the brane intersections depicted in Table 1. In these brane set-ups the D2- and D6-branes play the role of colour branes, while the D4- and D8-branes play the role of flavour branes. This is supported by the analysis of the Bianchi identities, which yields

$$\begin{aligned} dF_0 &= \sum_{k=1}^P \left(\frac{\nu_{k-1} - \nu_k}{2\pi} \right) \delta(\rho - 2\pi k) d\rho \\ d\hat{F}_4 &= \sum_{k=1}^P \left(\frac{\beta_{k-1} - \beta_k}{2\pi} \right) \delta(\rho - 2\pi k) d\rho \wedge \text{vol}_{CY_2}, \end{aligned} \tag{10}$$

indicating that, at the point $\rho = 2\pi k$, there is the possibility of having localised D8- and D4-branes. Indeed, explicit D8- and D4-branes need to be present at $\rho = 2\pi k$ if the slopes of h_8 and h_4 are different at both sides. Their numbers are given by

$$\Delta Q_{D8}^{(k)} = \nu_{k-1} - \nu_k, \quad \Delta Q_{D4}^{(k)} = \beta_{k-1} - \beta_k. \tag{11}$$

The associated Hanany–Witten brane set-up is then the one depicted in Figure 1.

Table 1. BPS brane intersection underlying the geometry given by (1)–(3). (x^0, x^1) are the directions where the 2D CFT lives. The directions (x^2, \dots, x^5) span the CY_2 , on which the D6- and the D8-branes are wrapped. The coordinate x^6 is the direction associated with ρ . Finally, (x^7, x^8, x^9) are the transverse directions realising the $SO(3)$ R-symmetry.

	0	1	2	3	4	5	6	7	8	9
D2	x	x					x			
D4	x	x						x	x	x
D6	x	x	x	x	x	x	x			
D8	x	x	x	x	x	x		x	x	x
NS5	x	x	x	x	x	x				




Figure 1. Generic Hanany–Witten brane set-up associated with the holographic background defined by the functions in (6) and (7). The vertical lines are NS5-branes, the horizontal lines represent D2- and D6-branes, and the crosses indicate D4- and D8-branes.

It was shown in [52–54] that these brane set-ups define 2D CFTs with $\mathcal{N} = (0, 4)$ SUSY. These CFTs describe the strongly coupled IR fixed points of the two-dimensional quantum field theories living in them. These field theories are encoded in the quivers depicted in Figure 2. Since the extension of the D2 and D6 branes is finite in the ρ direction, the field theory that lives in their intersection is effectively two dimensional at low energies. The quivers become non-anomalous when adequate flavour groups are attached at each node, coming from D4- and D8-branes. Their dynamics is described in terms of $(0, 4)$ vector multiplets and hypermultiplets, coming from the open strings that connect the different types of branes. This analysis was presented in [69]. It differs slightly from the one originally considered in [52–54], which also led to anomaly free quivers with the same central charge to leading order but was not based directly on open string quantisation. We next summarise the main ingredients of the quivers based on open string quantisation, following [69]:

- To each gauge node corresponds to a $(0, 4)$ vector multiplet, represented by a circle, plus a $(0, 4)$ hypermultiplet in the adjoint representation of the gauge group, represented by a grey line starting and ending on the same gauge group. In terms of $(0, 2)$ multiplets, the first consists of a vector multiplet and a Fermi multiplet in the adjoint, and the second consists of two chiral multiplets forming a $(0, 4)$ hypermultiplet.
- Between each pair of horizontal nodes there are two $(0, 2)$ Fermi multiplets, forming a $(0, 4)$ Fermi multiplet, and two $(0, 2)$ chiral multiplets, forming a $(0, 4)$ twisted hypermultiplet, each in the bifundamental representation of the gauge groups. The $(0, 4)$ Fermi multiplet and the $(0, 4)$ twisted hypermultiplet combine into a $(4, 4)$ twisted hypermultiplet. They are represented by horizontal black solid lines.
- Between each pair of vertical nodes, there are two $(0, 2)$ chiral multiplets forming a $(0, 4)$ hypermultiplet in the bifundamental representation of the gauge groups. They are represented by grey lines.
- Between each gauge node and any successive or preceding node, there is one $(0, 2)$ Fermi multiplet in the bifundamental representation. It is represented by dashed lines.
- Between each gauge node and its adjacent global symmetry node, there is one $(0, 2)$ Fermi multiplet in the fundamental representation of the gauge group. It is again represented by dashed lines.
- Between each gauge node and its opposing global symmetry node, there are two $(0, 2)$ Fermi multiplets, forming a $(0, 4)$ Fermi multiplet, and two $(0, 2)$ chiral multiplets, forming a $(0, 4)$ twisted hypermultiplet, each in the fundamental representation of the gauge groups. The $(0, 4)$ Fermi multiplet and the $(0, 4)$ twisted hypermultiplet combine into a $(4, 4)$ twisted hypermultiplet. They are represented by curvy black solid lines.




Figure 2. A generic quiver field theory in which the IR is dual to the holographic background defined by the functions in (6) and (7).

The previous fields contribute to the anomaly of a generic $SU(N_i)$ gauge group as follows:

- A (0, 2) vector multiplet contributes a factor of $-N_i$.
- A (0, 2) chiral multiplet in the adjoint representation contributes with a factor of N_i .
- A (0, 2) chiral multiplet in the bifundamental representation contributes with a factor of $\frac{1}{2}$.
- A (0, 2) Fermi multiplet in the adjoint representation contributes with a factor of $-N_i$.
- A (0, 2) Fermi multiplet in the fundamental of bifundamental representation contributes with a factor of $-\frac{1}{2}$.

Putting these together, we have that, for generic $SU(\alpha_k)$ and $SU(\mu_k)$ colour groups, the gauge anomaly cancellation conditions are, respectively,

$$F_k = \nu_{k-1} - \nu_k, \quad \tilde{F}_k = \beta_{k-1} - \beta_k, \tag{12}$$

for the F_k and \tilde{F}_k flavour groups. This is precisely the number of D8 and D4 flavour (source) branes in each interval, as shown by Equation (11).

In turn, as shown in [70], the right-handed central charge of the IR SCFT is calculated by associating it with the $U(1)_R$ current two-point function;

$$c_{cft} = 3 \text{Tr}[\gamma_3 Q_R^2], \tag{13}$$

where Q_R is the charge with respect to the $U(1)_R \subset SU(2)_R$ and the trace is over all Weyl fermions in the theory. The two left-handed fermions inside the (0, 4) vector multiplet have R-charge equal to 1. For hypermultiplets, we have that, for both right-handed fermions inside a (0, 4) hypermultiplet, the R-charge is -1 , while those in a (0, 4) twisted hypermultiplet have zero R-charge. Finally, the left-handed fermion inside the (0, 2) Fermi multiplet also has a vanishing R-charge. Putting this together, we find that

$$c_{cft} = 6(n_{hyp} - n_{vec}), \tag{14}$$

where n_{hyp} is the number of (untwisted) (0, 4) hypermultiplets and n_{vec} is the number of (0, 4) vector multiplets.

In [52–54], a number of dual holographic pairs were presented that provided stringent support for the validity of the proposed duality. In each example, the field theory central charge given by expression (14) was shown to coincide (for long quivers with large ranks,

when the background is a trustable dual description of the CFT) with the holographic central charge. This was computed from the Brown–Henneaux formula, giving

$$c_{hol} = \frac{3\pi}{2G_N} \text{Vol}_{\text{CY}_2} \int_0^{2\pi(P+1)} h_4 h_8 d\rho = \frac{3}{\pi} \int_0^{2\pi(P+1)} h_4 h_8 d\rho. \tag{15}$$

Here, we used that $G_N = 8\pi^6$, with $g_s = \alpha' = 1$, and that $\text{Vol}_{\text{CY}_2} = 16\pi^4$. For the functions h_4 and h_8 displayed above, this gives

$$c_{hol} = \sum_{k=1}^P \left(6\alpha_k \mu_k + 3(\alpha_k \nu_k + \beta_k \mu_k) + 2\beta_k \nu_k \right), \tag{16}$$

which can be shown to agree in the holographic limit with the expression coming from (14).

3. AdS₃ Solutions in M-Theory

In this section, we consider the uplift to eleven dimensions of the solutions discussed in the previous section, following [34]. In order to carry out this uplift, we need to restrict it to vanishing Romans’ mass, $F_0 = 0$, which imposes $h'_8 = 0$, and thus both the absence of D8-branes and the presence of a constant number of D6-branes between all pairs of NS5-branes. In the uplift to eleven dimensions, this number becomes a modding parameter of the geometry, associated with the KK-monopole charge.

The M-theory solutions are of the form $\text{AdS}_3 \times S^3 / \mathbf{Z}_k \times \text{CY}_2$ fibred over an interval. They read as follows:

$$\begin{aligned} ds_{11}^2 &= \Delta \left(\frac{u}{\sqrt{h_4 h_8}} ds_{\text{AdS}_3}^2 + \sqrt{\frac{h_4}{h_8}} ds_{\text{CY}_2}^2 + \frac{\sqrt{h_4 h_8}}{u} d\rho^2 \right) + \frac{h_8^2}{\Delta^2} ds_{S^3/\mathbf{Z}_k}^2, \\ G_4 &= -d \left(\frac{uu'}{2h_4} + 2\rho h_8 \right) \wedge \text{vol}_{\text{AdS}_3} + 2h_8 d \left(-\rho + \frac{uu'}{4h_4 h_8 + u'^2} \right) \wedge \text{vol}_{S^3/\mathbf{Z}_k} \\ &\quad - h'_4 \text{vol}_{\text{CY}_2}, \\ \Delta &= \frac{h_8^{1/2} (4h_4 h_8 + u'^2)^{1/3}}{2^{2/3} h_4^{1/6} u^{1/3}}, \end{aligned} \tag{17}$$

where $k = h_8$. The quotiented three-sphere is written as a Hopf fibration over S^2 ,

$$ds_{S^3/\mathbf{Z}_k}^2 = \frac{1}{4} \left[\left(\frac{dz}{k} + \omega \right)^2 + ds_{S^2}^2 \right] \quad \text{with} \quad d\omega = \text{vol}_{S^2}. \tag{18}$$

In these solutions, the symmetries $\text{SL}(2, \mathbf{R}) \times \text{SL}(2, \mathbf{R})$ and $\text{SU}(2)$ are geometrically realised by the AdS_3 and the quotiented three-sphere, respectively.

In the new class of solutions given by (17), the number of Type IIA D6-branes becomes the orbifold parameter in S^3 / \mathbf{Z}_k , $k = h_8$. This is associated with the KK-monopole charge. The D2- and D4-branes of the Type IIA solution become M2- and M5-branes, respectively. Their presence is captured by integrating the Page flux $\hat{G}_7 = G_7 - G_4 \wedge C_3$ and a nontrivial flux of G_4 through the CY_2 . In turn, the NS5 branes become M5'-branes. The uplifted brane set-up is the one depicted in Table 2. Recently, it was shown that the solutions emerge in the near horizon limit of this intersecting brane system [36]. The KK-monopoles (wrapped on the CY_2) and the M2 branes are stretched between parallel M5'-branes, with extra M5-branes providing for flavour groups.

Table 2. $\frac{1}{8}$ -BPS brane intersection underlying the $AdS_3 \times S^3/Z_k$ solutions in M-theory. The 2D CFT lives in the (x^0, x^1) directions, (x^2, \dots, x^5) span the CY_2 , x^6 is the “field theory” direction and (x^7, x^8, x^9) are the transverse directions on which the $SO(3)_R$ symmetry is realised. Finally, x^{10} is the extra eleventh direction, which spans the $S^1/Z_k \subset S^3/Z_k$ and plays the role of the Taub-NUT direction of the KK-monopoles.

	0	1	2	3	4	5	6	7	8	9	10
M2	x	x					x				
M5	x	x						x	x	x	x
KK	x	x	x	x	x	x	x				z
M5'	x	x	x	x	x	x					

In [34], it was argued that this brane intersection describes the M_A -strings introduced in [55,56], supplemented with extra M5-branes. The corresponding dual quivers are the ones depicted in Figure 3, with upper row nodes associated with M2-branes and lower row nodes associated with KK-monopoles. The M5-branes provide extra flavour groups that render the quivers non-anomalous (and the supergravity equations of motion satisfied).




Figure 3. Generic quiver field theory in which the IR is dual to the $AdS_3 \times S^3/Z_k \times CY_2 \times I$ M-theory solutions defined by the functions in (6) and (7).

Thus, the new solutions in M-theory provide for explicit AdS_3 geometries where M_A -strings can be studied holographically. In particular, the matching between the field theory and holographic computations of the central charge follows directly upon uplift from ten dimensions. The holographic central charge given by Equation (15) becomes, in the massless case,

$$c_{hol} = \frac{3}{\pi} h_8 \int_0^{2\pi(P+1)} d\phi h_4 = 6h_8 \sum_{k=1}^P \alpha_k = 6h_8 \sum_{k=1}^P Q_{M_2}^{(k)} = 6kQ_{M_2} = 6Q_{M_A}, \quad (19)$$

where Q_{M_A} stands for the total number of M_A -strings in the configuration, taking into account the orbifolding by Z_k . This result stresses out that the degrees of freedom of the strongly coupled conformal field theory that originate from the M_A -strings.

Furthermore, in [34], a second class of AdS_3 solutions to M-theory of the form $AdS_3/Z_k \times S^3 \times CY_2 \times I$ was obtained through a double analytic continuation from the previous solutions. In the background given in (17), the AdS_3 and S^3 factors can be swapped as follows:

$$AdS_3 \rightarrow -S^3, \quad S^3 \rightarrow -AdS_3, \quad (20)$$

to produce the new class of solutions, together with

$$u \rightarrow -iu, \quad h_4 \rightarrow ih_4, \quad h_8 \rightarrow ih_8, \quad \rho \rightarrow i\rho. \quad (21)$$

These solutions read

$$\begin{aligned}
 ds_{11}^2 &= \frac{h_8^2}{\Delta^2} ds_{\text{AdS}_3/\mathbf{Z}_k}^2 + \Delta \left(\frac{u}{\sqrt{h_4 h_8}} ds_{S^3}^2 + \sqrt{\frac{h_4}{h_8}} ds_{\text{CY}_2}^2 + \frac{\sqrt{h_4 h_8}}{u} d\rho^2 \right) \\
 G_4 &= -d \left(-\frac{uu'}{2h_4} + 2\rho h_8 \right) \wedge \text{vol}_{S^3} - 2h_8 d \left(\rho + \frac{uu'}{4h_4 h_8 - u'^2} \right) \wedge \text{vol}_{\text{AdS}_3/\mathbf{Z}_k} \\
 &\quad - h_4' \text{vol}_{\text{CY}_2}, \\
 \Delta &= \frac{h_8^{1/2} (4h_4 h_8 - u'^2)^{1/3}}{2^{2/3} h_4^{1/6} u^{1/3}},
 \end{aligned} \tag{22}$$

where $k = h_8$ and the quotiented AdS_3 subspace is written as a Hopf fibration over AdS_2 :

$$\begin{aligned}
 ds_{\text{AdS}_3/\mathbf{Z}_k}^2 &= \frac{1}{4} \left[\left(\frac{dz}{k} + \eta \right)^2 + ds_{\text{AdS}_2}^2 \right] \quad \text{with} \quad d\eta = \text{vol}_{\text{AdS}_2}, \\
 ds_{\text{AdS}_2}^2 &= -dt^2 \cosh^2 r + dr^2, \quad \eta = -\sinh r dt.
 \end{aligned} \tag{23}$$

Notice that the KK-monopoles become M0-branes, with the Taub-NUT direction of the KK-monopoles turned into the direction of propagation of the M0-branes, or waves. These solutions are associated with the M0-M2-M5-M5' brane intersections depicted in Table 3. They preserve the same number of supersymmetries as the original $\text{AdS}_3 \times S^3 / \mathbf{Z}_k \times \text{CY}_2 \times I$ solutions.

Table 3. $\frac{1}{8}$ -BPS brane intersection underlying the $\text{AdS}_3/\mathbf{Z}_k \times S^3$ solutions in M-theory. x^1 is the direction of propagation of the M0-branes, (x^2, \dots, x^5) span the CY_2 , x^6 is the direction along the ρ -interval and (x^7, x^8, x^9, x^{10}) are the transverse directions on which the $\text{SO}(4)$ R-symmetry is realised.

	0	1	2	3	4	5	6	7	8	9	10
M0	x	x									
M2	x	x					x				
M5	x	x						x	x	x	x
M5'	x	x	x	x	x	x					

4. $\text{AdS}_2/\text{CFT}_1$ in Massive IIA

A new class of $\text{AdS}_2 \times S^3 \times \text{CY}_2 \times I$ solutions to massless Type IIA supergravity can be obtained from (22) upon reduction along the Hopf fibre of the $\text{AdS}_3/\mathbf{Z}_k$ subspace given by (23). These backgrounds are associated with D0-F1-D4-D4' brane intersections that preserve $\mathcal{N} = 4$ supersymmetries in one dimension. These solutions can also be obtained through a double analytical continuation from the solutions reviewed in Section 2. In fact, this allows one to extend them to the massive case. The corresponding brane set-up is depicted in Table 4. In this manner, we find an $\text{AdS}_2 \times S^3 \times \text{CY}_2 \times I$ class of solutions to massive Type IIA supergravity with the NS-NS sector:

$$\begin{aligned}
 ds^2 &= \frac{u}{\sqrt{h_4 h_8}} \left(\frac{h_4 h_8}{4h_4 h_8 - u'^2} ds_{\text{AdS}_2}^2 + ds_{S^3}^2 \right) + \sqrt{\frac{h_4}{h_8}} ds_{\text{CY}_2}^2 + \frac{\sqrt{h_4 h_8}}{u} d\rho^2, \\
 e^{-2\Phi} &= \frac{h_8^{3/2} (4h_4 h_8 - u'^2)}{4h_4^{1/2} u}, \quad B_2 = -\frac{1}{2} \left(\rho + \frac{uu'}{4h_4 h_8 - u'^2} \right) \text{vol}_{\text{AdS}_2}.
 \end{aligned} \tag{24}$$

The background is further supported by the RR fluxes:

$$\begin{aligned}
 F_0 &= h'_8, & F_2 &= -\frac{1}{2} \left(h_8 + \frac{h'_8 u' u}{4h_8 h_4 - u'^2} \right) \text{vol}_{\text{AdS}_2}, \\
 F_4 &= \left(-d \left(\frac{u' u}{2h_4} \right) + 2h_8 d\rho \right) \wedge \text{vol}_{\text{S}^3} - h'_4 \text{vol}_{\text{CY}_2}.
 \end{aligned}
 \tag{25}$$

The warping functions h_8, h_4 and u depend on ρ , as in the seed solutions. Note that, in this case, $(4h_8 h_4 - u'^2) > 0$ in order to guarantee a real dilaton and a metric with the correct signature. Supersymmetry and the Bianchi identities of the fluxes (away from localised sources) are imposed by Equation (3), which make h_8, h_4 and u linear functions of ρ .

We quote the Page fluxes, $\hat{F} = e^{-B_2} \wedge F$, as follows:

$$\begin{aligned}
 \hat{F}_0 &= h'_8, & \hat{F}_2 &= -\frac{1}{2} \left(h_8 - h'_8 (\rho - 2\pi k) \right) \text{vol}_{\text{AdS}_2}, \\
 \hat{F}_4 &= \left(2h_8 d\rho - d \left(\frac{u' u}{2h_4} \right) \right) \wedge \text{vol}_{\text{S}^3} - h'_4 \text{vol}_{\text{CY}_2},
 \end{aligned}
 \tag{26}$$

where we included large gauge transformations of B_2 of parameter k , $B_2 \rightarrow B_2 + \pi k \text{vol}_{\text{AdS}_2}$ (see [40]).

Table 4. Brane set-up associated to the solutions (24) and (25). x^0 is the time direction of the ten dimensional spacetime; x^1, \dots, x^4 are the coordinates spanning CY_2 ; x^5 is the direction where the F1-strings are stretched; and x^6, x^7, x^8 , and x^9 are the coordinates where the $\text{SO}(4)$ R-symmetry is realised.

	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9
D0	x									
D4	x	x	x	x	x					
D4'	x						x	x	x	x
D8	x	x	x	x	x		x	x	x	x
F1	x					x				

An infinite family of backgrounds with $u = \text{constant}$ and h_4 and h_8 being piecewise continuous as in (6) and (7) were analysed in [40], and together with their dual SCQM description⁵. We summarise this description in the next subsection.

4.1. Dual Superconformal Quantum Mechanics

The superconformal quantum mechanics dual to the previous class of solutions was studied in [40]. The proposal therein is that a 1D $\mathcal{N} = 4$ quantum mechanics lives on the D0-D4-D4'-D8-F1 brane set-up depicted in Table 4 that describes the interactions between D0 and D4 brane instantons and F1 Wilson lines in the 5D gauge theory living in the intersection of D4' and D8 branes. This is a generalisation of the ADHM quantum mechanics described in [71] and of the quiver proposals discussed in [72,73].

In order to describe the quantum mechanics, the D0-D4-D4'-D8-F1 brane system was split into two subsystems, D4-D4'-F1 and D0-D8-F1, that were first studied independently. The first subsystem was interpreted as describing BPS F1 Wilson lines introduced in the 5D theory living on the D4'-branes by D4-branes [74]. Similarly, the D0-D8-F1 subsystem was interpreted as describing F1 Wilson lines introduced in the worldvolume of the D8-branes by D0-branes [75]. Indeed, the branes in the D4-D4'-F1 and D0-D8-F1 subsystems are displayed exactly as in the D3-D5-F1 brane configuration that describes Wilson lines in antisymmetric representations in 4D $\mathcal{N} = 4$ SYM, studied in [76,77].

For the solutions defined by (6) and (7), the quantised charges associated with the Page fluxes given by (26) in the different $2\pi k \leq \rho \leq 2\pi(k + 1)$ intervals are

$$\begin{aligned}
 Q_{D4}^{e(k)} = \alpha_k &= \sum_{j=0}^{k-1} \beta_j, & Q_{D0}^{e(k)} = \mu_k &= \sum_{j=0}^{k-1} \nu_j \\
 Q_{D4'}^{m(k)} = \beta_k, & & Q_{D8}^{m(k)} = \nu_k, & & Q_{F1}^{e(k)} = 1,
 \end{aligned}
 \tag{27}$$

where we use electric and magnetic charges, as explained in [40]. The Hanany–Witten brane set-up associated to these brane charges is the one depicted in Figure 4. In order to understand the quantum mechanics associated with this brane configuration, it is useful to go to Type IIB and S-dualise. Then, after performing Hanany–Witten moves, one can go back to Type IIA, where one can interpret the resulting brane set-up (depicted in Figure 5) as describing $U(\alpha_k)$ and $U(\mu_k)$ Wilson lines in the completely antisymmetric representations $(\beta_0, \beta_1, \dots, \beta_{k-1})$ of $U(\alpha_k)$ and $(\nu_0, \nu_1, \dots, \nu_{k-1})$ of $U(\mu_k)$, respectively. Given that the Wilson lines are in the completely antisymmetric representations, the D4–D4′–F1 and D0–D8–F1 subsystems describe baryon vertices [78].




Figure 4. Hanany–Witten brane set-up associated with the brane charges of the solutions.




Figure 5. Hanany–Witten brane set-up equivalent to the brane configuration in Table 4.

This is consistent with an interpretation of the AdS_2 solutions as describing backreacted baryon vertices within the 5D $\mathcal{N} = 1$ QFT living in D4′–D8 branes. In this interpretation, the SCQM arises in the very low energy limit of a system of D4′–D8 branes, dual to a 5D $\mathcal{N} = 1$ gauge theory, in which one-dimensional defects are introduced. The one-dimensional defects consist of D4-brane baryon vertices, connected to the D4′ with F1-strings, and D0-brane baryon vertices, connected to the D8 with F1-strings. In the IR, the gauge symmetry on both the D4′ and D8 branes becomes global, turning them from colour to flavour branes. In turn, the D4 and D0 defect branes become the new colour branes of the backreacted geometry. This is in agreement with the defect interpretation found for

this class of solutions in [36], where the AdS_2 geometries were shown to asymptote locally to the AdS_6 background of Brandhuber–Oz [79].

The previous SCQMs can be given a quiver-like description, which can be used to compute their central charge. From the brane set-up depicted in Figure 4, one can construct the quiver shown in Figure 6, where the gauge groups are associated with the colour D0- and D4-branes and the flavour groups are associated with the $D4'$ - and D8-branes. The quantised charges are the ones computed in Equation (27). The dynamics of the quiver is described in terms of (4, 4) vector multiplets, (4, 4) hypermultiplets in the adjoint representations and (4, 4) hypermultiplets in the bifundamental representations. The connection between colour and flavour branes is through twisted (4, 4) bifundamental hypermultiplets (bent black lines) and (0, 2) bifundamental Fermi multiplets (dashed lines). This follows directly from the analysis of Appendix B in [40].




Figure 6. One-dimensional quiver field theory in which the IR limit is dual to the AdS_2 backgrounds reviewed in Section 4.

As before, a check of the validity of the proposed quivers is given by matching the field theory and holographic central charges. In the case at hand, however, we are dealing with a one-dimensional field theory, for which the definition of central charge is subtle. We interpret the central charge as counting the ground states of the conformal quantum mechanics. In [40], the same expression used in Section 2.1 for the computation of the central charge of the 2D dual CFT was proposed to be valid for the 1D quiver mechanics. For these quivers, n_{hyp} and n_{vec} are, respectively, the numbers of $\mathcal{N} = 4$ (untwisted) hypermultiplets and vector multiplets. Using this expression, perfect agreement was found (in the large number of nodes with large ranks limit) between the quantum mechanics central charge and the holographic central charge, which in this case is obtained through the following integration:

$$c_{hol,1d} = \frac{3V_{int}}{4\pi G_N} = \frac{3}{4\pi} \frac{Vol_{CY_2}}{(2\pi)^4} \int_0^{2\pi(P+1)} (4h_4h_8 - u'^2) d\rho. \tag{28}$$

This is a striking result, since the superconformal quantum mechanics dual to our class of solutions does not seem to have a direct relation to 2D CFTs. Compared with the results in the literature for the dimension of the Higgs branch of $\mathcal{N} = 4$ quantum mechanics with gauge groups $\prod_v U(N_v)$ connected by bifundamentals [80], one sees that the expression $c_{cft} = 6(n_{hyp} - n_{vec})$ may be interpreted as an extension of the formulas therein to more general $\mathcal{N} = 4$ quivers including flavours. This is an interesting relation that deserves further investigation.

5. AdS_2 Solutions in Type IIB

In this section, we review two classes of AdS_2 solutions to Type IIB supergravity with 4 Poincaré supersymmetries. This is based on the works [38,43]. These solutions are obtained

from the backgrounds reviewed in Sections 2 and 4, respectively, upon T-duality. Moreover, both solutions are related to each other through a double analytical continuation.

The two classes of solutions consist of $AdS_2 \times S^2 \times CY_2 \times S^1$ geometries foliated over an interval. Despite this, they have substantially different dual field theory descriptions that are inherited from their respective T-dual origins. Although we do not review these results in this paper, it was shown in [43] that both types of solutions fit locally in the general class of $AdS_2 \times S^2 \times CY_2 \times \Sigma_2$ solutions of Type IIB supergravity found in [8,9], when there are no D3- and D7-branes present. In the presence of these branes, our solutions extend the previous classifications. In order to show this explicitly, it is necessary to perform a subtle zooming-in that was explained in [43].

5.1. Type A

Consider the class given by (1) and (2), where the AdS_3 subspace is written as a Hopf fibration over AdS_2 , as shown by expression (23) for $k = 1$ and $z = \psi$. By applying T-duality on the fibre direction ψ , new AdS_2 solutions preserving $\mathcal{N} = 4$ supersymmetry are obtained. These backgrounds have the structure of an $AdS_2 \times S^2 \times CY_2 \times S^1$ geometry warped over an interval. The NS-NS sector reads

$$\begin{aligned}
 ds^2 &= \frac{u}{\sqrt{h_4 h_8}} \left(\frac{1}{4} ds_{AdS_2}^2 + \frac{h_4 h_8}{4h_4 h_8 + u'^2} ds_{S^2}^2 \right) + \sqrt{\frac{h_4}{h_8}} ds_{CY_2}^2 + \frac{\sqrt{h_4 h_8}}{u} (d\rho^2 + d\psi^2), \\
 e^{-2\Phi} &= \frac{h_8}{4h_4} (4h_4 h_8 + u'^2), \\
 H_3 &= \frac{1}{2} d \left(-\rho + \frac{uu'}{4h_4 h_8 + u'^2} \right) \wedge vol_{S^2} + \frac{1}{2} vol_{AdS_2} \wedge d\psi,
 \end{aligned}
 \tag{29}$$

where the h_4, h_8 and u functions are inherited from the backgrounds (1) and (2) and thus have support on ρ .

The 10-dimensional RR fluxes are given by

$$\begin{aligned}
 F_1 &= h'_8 d\psi, \\
 F_3 &= -\frac{1}{2} \left(h_8 - \frac{h'_8 u' u}{4h_8 h_4 + u'^2} \right) vol_{S^2} \wedge d\psi + \frac{1}{4} \left(d \left(\frac{u' u}{2h_4} \right) + 2h_8 d\rho \right) \wedge vol_{AdS_2}, \\
 F_5 &= -(1 + \star) h'_4 vol_{CY_2} \wedge d\psi \\
 &= -h'_4 vol_{CY_2} \wedge d\psi + \frac{h'_4 h_8 u^2}{4h_4 (4h_4 h_8 + u'^2)} vol_{AdS_2} \wedge vol_{S^2} \wedge d\rho.
 \end{aligned}
 \tag{30}$$

The Type IIB equations of motion are satisfied by imposing the BPS equations and Bianchi identities given by (3). In turn, the Page forms $\hat{F} = e^{-B_2} \wedge F$ are given by

$$\begin{aligned}
 \hat{F}_1 &= h'_8 d\psi, \\
 \hat{F}_3 &= \frac{1}{2} (h'_8 (\rho - 2\pi k) - h_8) vol_{S^2} \wedge d\psi + \frac{1}{4} \left(\frac{u' (h_4 u' - u h'_4)}{2h_4^2} + 2h_8 \right) vol_{AdS_2} \wedge d\rho, \\
 \hat{F}_5 &= \frac{1}{16} \left(\frac{(u - (\rho - 2\pi k) u') (u h'_4 - h_4 u')}{h_4^2} + 4(\rho - 2\pi k) h_8 \right) vol_{AdS_2} \wedge vol_{S^2} \wedge d\rho \\
 &\quad - h'_4 vol_{CY_2} \wedge d\psi.
 \end{aligned}
 \tag{31}$$

The analysis of these fluxes suggests that the brane content that underlies this class of solutions is the one depicted in Table 5. Here, the D1- and D3-branes play the role of colour branes and the D7- and D3-branes play the role of flavour branes. As for the AdS_3 solutions reviewed in Section 2, an infinite family of AdS_2 backgrounds can be defined by the piecewise linear functions h_4 and h_8 given by Equations (6) and (7). We turn next to the description of the superconformal quantum mechanics dual to this class of solutions.

Table 5. Brane set-up underlying the geometry in (29) and (30). x^0 is the time direction, (x^1, \dots, x^4) span the CY_2 , x^5 is the direction associated with the ρ coordinate, (x^6, x^7, x^8) are the transverse directions realising the $SO(3)$ R-symmetry and x^9 is the ψ direction.

	0	1	2	3	4	5	6	7	8	9
D1	x					x				
D3	x						x	x	x	
D5	x	x	x	x	x	x				
D7	x	x	x	x	x		x	x	x	
NS5	x	x	x	x	x					x
F1	x									x

Dual Superconformal Quantum Mechanics

The $\mathcal{N} = 4$ superconformal quantum mechanics dual to the previous class of solutions was studied in [38]. At the geometrical level, these solutions are related to the AdS_3 solutions reviewed in Section 2 through T-duality. At the level of the dual CFTs, the superconformal quantum mechanics dual to the AdS_2 solutions should then arise from the 2D CFTs dual to the AdS_3 backgrounds upon dimensional reduction.

More concretely, in the coordinates used to obtain the AdS_2 geometry, defined in Equation (23), the boundaries of AdS_3 are two null cylinders [63]. For this reason, the 2D CFT that lives at these boundaries is effectively discrete light-cone quantised (DLCQ) because just one of the $SL(2, \mathbf{R}) \times SL(2, \mathbf{R})$ sectors of global AdS_3 survives the compactification. T-dualisation in these coordinates is then equivalent to starting with a given $\mathcal{N} = (0, 4)$ 2D CFT, such as those described in Section 2.1, and being discrete light-cone quantised, keeping $\mathcal{N} = 4$ as the SUSY right sector. This provides an explicit realisation of the constructions in [62–65]. Field theoretically, we start with the Lagrangian describing the 2D CFT dual to AdS_3 and dimensionally reduce it to a matrix model where only the time dependence and the zero modes in the T-dual direction are kept.

The concrete proposal in [38] is that the dynamics of the UV quantum mechanics is described by the dimensional reduction along the space-direction of the 2D QFTs discussed in Section 2.1. The quiver field theories are then the same ones depicted in Figure 2 but now associated with D1 and D5 colour branes and D3 and D7 flavour branes. In turn, the matter fields are $\mathcal{N} = 4$ multiplets, realised as dimensionally reduced 2D $(0, 4)$ multiplets. Note that these quivers inherit the anomaly cancellation condition of the 2D quivers, even if there is no anomaly cancellation condition in 1D.

As in previous AdS/CFT pairs, one can check the agreement between the field theory and holographic central charges to test the proposed duality. In this case, the usage of expression (14) for computing the quantum mechanics central charge is fully justified, since it arises from a 2D CFT upon compactification. As expected, perfect agreement is found with the holographic central charge, which is computed from the same expression (15) given its invariance under T-duality.

5.2. Type B

In this subsection, we review the AdS_2 solutions to Type IIB supergravity that arise from the backgrounds defined in (24) and (25) upon T-duality along the Hopf-fibre of the three sphere. The resulting class of solutions have an NS-NS sector

$$\begin{aligned}
 ds^2 &= \frac{u\sqrt{h_4 h_8}}{4h_4 h_8 - u'^2} ds_{AdS_2}^2 + \frac{u}{4\sqrt{h_4 h_8}} ds_{S^2}^2 + \sqrt{\frac{h_4}{h_8}} ds_{CY_2}^2 + \frac{\sqrt{h_4 h_8}}{u} (d\psi^2 + d\rho^2), \\
 e^{-2\Phi} &= \frac{h_8}{4h_4} (4h_4 h_8 - u'^2), \\
 H_3 &= -\frac{1}{2} d\left(\rho + \frac{uu'}{4h_4 h_8 - u'^2}\right) \wedge vol_{AdS_2} + \frac{1}{2} vol_{S^2} \wedge d\psi,
 \end{aligned}
 \tag{32}$$

and RR fluxes,

$$\begin{aligned}
 F_1 &= h'_8 d\psi, \\
 F_3 &= -\frac{1}{2} \left(h_8 + \frac{h'_8 u' u}{4h_8 h_4 - u'^2} \right) \text{vol}_{\text{AdS}_2} \wedge d\psi + \frac{1}{4} \left(-d \left(\frac{u' u}{2h_4} \right) + 2h_8 d\rho \right) \wedge \text{vol}_{\text{S}^2}, \\
 F_5 &= -(1 + \star_{10}) h'_4 \text{vol}_{\text{CY}_2} \wedge d\psi, \\
 &= -h'_4 \text{vol}_{\text{CY}_2} \wedge d\psi - \frac{h_8 h'_4 u'^2}{4h_4 (4h_8 h_4 - u'^2)} \text{vol}_{\text{AdS}_2} \wedge \text{vol}_{\text{S}^2} \wedge d\rho.
 \end{aligned}
 \tag{33}$$

Here, $\psi \in [0, 2\pi]$ is the T-dual coordinate. Note that $4h_4 h_8 - u'^2 \geq 0$ must be imposed to have well-defined supergravity fields. Supersymmetry holds whenever $u'' = 0$. In the same way, the Bianchi identities of the fluxes impose $h''_8 = 0$ and $h''_4 = 0$, away from localised sources. The ρ coordinate describes an interval that we take to be bounded between 0 and $2\pi(P + 1)$, as in the previous sections.

The Page fluxes read

$$\begin{aligned}
 \hat{F}_1 &= h'_8 d\psi, \\
 \hat{F}_3 &= \frac{1}{2} (h'_8 (\rho - 2\pi k) - h_8) \text{vol}_{\text{AdS}_2} \wedge d\psi + \frac{1}{4} \left(2h_8 + \frac{u' (u h'_4 - h_4 u')}{2h_4^2} \right) \text{vol}_{\text{S}^2} \wedge d\rho, \\
 \hat{F}_5 &= \frac{1}{4} \left(h_8 (\rho - 2\pi k) - \frac{(u - (\rho - 2\pi k) u') (u h'_4 - h_4 u')}{4h_4^2} \right) \text{vol}_{\text{AdS}_2} \wedge \text{vol}_{\text{S}^2} \wedge d\rho \\
 &\quad - h'_4 \text{vol}_{\text{CY}_2} \wedge d\psi.
 \end{aligned}
 \tag{34}$$

The analysis of these fluxes yields the brane set-up summarised in Table 6 as underlying this class of solutions. Here, the D1- and D5-branes play the role of colour branes and the D3- and D7-branes of flavour branes. Both the brane set-up and the warped $\text{AdS}_2 \times \text{S}^2 \times \text{CY}_2 \times \text{S}^1$ structure of this class of solutions are shared with those of the solutions reviewed in the previous section. The precise relation between the two types of backgrounds is through the double analytic continuation:

$$\begin{aligned}
 ds^2_{\text{AdS}_2} &\rightarrow -ds^2_{\text{S}^2}, & ds^2_{\text{S}^2} &\rightarrow -ds^2_{\text{AdS}_2}, & e^\Phi &\rightarrow i e^\Phi, & F_i &\rightarrow -i F_i, \\
 u &\rightarrow -iu, & h_4 &\rightarrow ih_4, & h_8 &\rightarrow ih_8, & \rho &\rightarrow i\rho, & \psi &\rightarrow -i\psi.
 \end{aligned}
 \tag{35}$$

Table 6. Brane set-up underlying the geometry in (32) and (33). x^0 is the time direction; x^1, \dots, x^4 are the coordinates spanned by the CY_2 ; x^5 is the direction associated with the ρ coordinate; (x^6, x^7, x^8) are the transverse coordinates realising the $\text{SO}(3)$ R-symmetry; and x^9 is the ψ direction.

	0	1	2	3	4	5	6	7	8	9
D1	x									x
D3	x						x	x	x	
D5	x	x	x	x	x					x
D7	x	x	x	x	x		x	x	x	
NS5	x	x	x	x	x	x				
F1	x					x				

5.2.1. Dual Superconformal Quantum Mechanics

The superconformal quantum mechanics dual to this new class of solutions was discussed in [43]. Given that they are obtained via T-duality from the class of AdS_2 solutions reviewed in Section 4.1, they should be dual to the same superconformal quantum mechanics. In this case, the Wilson lines arise from the massive fermionic strings that stretch between D1-branes (and D5-branes) in the k th interval and D7-branes (and D3-branes) in all other intervals. In turn, the Wilson lines are in the $(\nu_0, \dots, \nu_{k-1})$ and $(\beta_0, \dots, \beta_{k-1})$

completely antisymmetric representation of the $U(\mu_k)$ and $U(\alpha_k)$ gauge groups, respectively. As we indicated in Section 4.1, given that the Wilson lines are in the completely antisymmetric representation, the D1-D7-F1 and D5-D3-F1 subsystems describe baryon vertices [78].

This is consistent with an interpretation of our AdS_2 solutions as describing backreacted baryon vertices within the 4D $\mathcal{N} = 2$ QFT living in D3-D7 branes. In this interpretation, the SCQM arises in the very low energy limit of a system of D3-D7 branes, dual to a 4D $\mathcal{N} = 2$ QFT, in which one-dimensional defects are introduced. The one-dimensional defects consist of D5-brane baryon vertices, connected to the D3 with F1-strings, and D1-brane baryon vertices, connected to the D7 with F1-strings. In the IR, the gauge symmetry on both the D7 and D3 branes becomes global, turning them from colour to flavour branes. In turn, the D5 and D1 defect branes become the new colour branes of the backreacted geometry. It would be very interesting to explicitly realise this defect interpretation geometrically.

6. Discussion

In this review article, we summarised the salient features of the recent developments in [26,34,38,40,43,52–54], in relation to the construction of AdS_3/CFT_2 and AdS_2/CFT_1 pairs with four supercharges. For clarity, we summarised the connections between these new classes of solutions in Figure 7.




Figure 7. Connections between the different classes of solutions reviewed in this paper.

The construction of these new dual pairs extends existing classifications of AdS_3 and AdS_2 solutions to the case with four supersymmetries. Moreover, the analysis in these references complements the construction of the backgrounds with a comprehensive study of the 2D and 1D CFTs dual to them. The proposed CFTs are described in the UV by means of explicit quivers from which observables such as the central charge can be computed and checked against holographic calculations. The holographic central charge is interpreted as the entanglement entropy of black strings or black holes with AdS_3 or AdS_2 near horizon geometries. This can then be cross-checked against the field theory computation, within a well controlled setting where the microstate counting programme can be put to work. This line of research remains to be further exploited. See [81–83]. In particular, it would be interesting to apply exact calculational techniques (see [84]) to the new classes of solutions, since this would provide for a deeper understanding of the IR regime of the different theories.

A study that would be interesting to pursue further is the interpretation of the new classes of solutions as surface or line-defect CFTs within higher dimensional conformal field theories. Recent progress in this direction has shown that some subclasses of the AdS_3 and AdS_2 solutions to massive IIA supergravity (constructed in Sections 2 and 4) asymptote locally to the AdS_6 background of Brandhuber–Oz [79]. This means that they can be

interpreted as surface or line-defect CFTs, respectively, within the 5D $\text{Sp}(N)$ fixed point theory dual to the Brandhuber–Oz solution [36]. In the AdS_2 case, this is in nice agreement with our proposed interpretation of these solutions as backreacted D4-D0 baryon vertices in a system of D4'-D8 branes. In this description, the one-dimensional defects consist of D4-D0 baryon vertices connected to the D4'-D8 branes with (fermionic) F1-strings. In the IR, the gauge symmetry on the D4'-D8 branes becomes global, turning them from colour to flavour branes. In turn, the D4-D0 defect branes become the new colour branes of the backreacted geometry.

In order to complete the previous picture, we should keep in mind that D4-D8 brane set-ups must include O8 orientifold fixed planes in order to flow to a 5D fixed point theory in the UV [85]. It would be interesting to clarify to what extent the behaviour found in [40] at both ends of the space, compatible with the presence of O8 orientifold fixed planes, would provide for a fully consistent global picture. It is expected that, in this set-up, baryon vertices affected by the orbifold projection are removed from the spectrum, corresponding to the fact that USp baryons are unstable against their decay into mesons. A similar interpretation for the D2-D6-NS5 brane surface defects within the D4'-D8 brane intersection, put forward in [36] for the AdS_3 case, still remains to be elucidated. In both the AdS_2 and AdS_3 cases, an explicit realisation of the quiver CFTs as embedded in the 5D quiver CFT associated with the D4'-D8 brane system remains to be found.

This is in contrast with the interpretation of a subclass of the M-theory pairs reviewed in Section 3 as surface defects within the 6D (1, 0) CFT living in M5-branes on ALE singularities, found in [36]. In this case, it has been possible to explicitly realise the 2D quiver CFTs as embedded in the 6D quiver CFT associated with the M5-branes intersecting with KK-monopoles.

We argued that the Type B AdS_2 solutions reviewed in Section 5.2 describe backreacted D5-D1 baryon vertices in the 4D $\mathcal{N} = 2$ QFT living in D3-D7 intersections. In this case, there is no holographic analogue, and it would be interesting to see if these solutions asymptote locally to an AdS_5 background. This would provide further support to the proposed defect interpretation.

Finally, we stress that the full class of AdS_3 solutions discussed in [26,52–54], which constitute the basis for the developments reviewed in this paper, is much broader than the subset of solutions that have been the focus of our CFT analysis. In particular, there is evidence that interesting new CFTs arise when there is dependence on the internal structure of the CY_2 manifold. Work is in progress in this direction [69].

Funding: This research was partially funded by the Spanish government grant number PGC2018-096894-B-100. AR is supported by CONACyT-Mexico.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: We thank Niall Macpherson, Carlos Nunez and Stefano Speziali for their collaboration in some of the results reviewed in this article and especially Chris Couzens, Niall Macpherson and Carlos Nunez for careful reading of the manuscript. We are indebted to Prof. Norma Sanchez for inviting us to contribute this review article to the Open Access Special Issue “Women Physicists in Astrophysics, Cosmology and Particle Physics” to be published in *Universe* (ISSN 2218-1997, IF 1.752).

Conflicts of Interest: The authors declare no conflicts of interest.

Notes

¹ By which we mean one and two dimensional.

² The $u \neq \text{constant}$ case was studied in [37,54].

³ This is also compatible with a superposition of O2-O6 planes. The string theory interpretation of smeared orientifold fixed planes is however unclear.

⁴ See [66–68] for further developments.

⁵ A concrete example with $u' \neq 0$ was analysed in [45].

References

1. Argurio, R.; Giveon, A.; Shomer, A. Superstring theory on $AdS_3 \times G/H$ and boundary $N = 3$ superconformal symmetry. *J. High Energy Phys.* **2000**, *2000*, 010. [[CrossRef](#)]
2. Cvetič, M.; Lu, H.; Pope, C.; Vazquez-Poritz, J.F. AdS in warped space-times. *Phys. Rev. D* **2000**, *62*, 122003. [[CrossRef](#)]
3. Kim, N. AdS_3 solutions of IIB supergravity from D3-branes. *J. High Energy Phys.* **2006**, *2006*, 094. [[CrossRef](#)]
4. Gauntlett, J.P.; Kim, N.; Waldram, D. Supersymmetric AdS_3 , AdS_2 and Bubble Solutions. *J. High Energy Phys.* **2007**, *2007*, 005. [[CrossRef](#)]
5. Gauntlett, J.P.; Mac Conamhna, O.A.P.; Mateos, T.; Waldram, D. Supersymmetric AdS_3 solutions of type IIB supergravity. *Phys. Rev. Lett.* **2006**, *97*, 171601. [[CrossRef](#)]
6. D'Hoker, E.; Estes, J.; Gutperle, M. Gravity duals of half-BPS Wilson loops. *J. High Energy Phys.* **2007**, *2007*, 063. [[CrossRef](#)]
7. Donos, A.; Gauntlett, J.P.; Sparks, J. $AdS_3 \times (S^{*3} \times S^{*3} \times S^{*1})$ Solutions of Type IIB String Theory. *Class. Quant. Grav.* **2009**, *26*, 065009. [[CrossRef](#)]
8. Chiodaroli, M.; Gutperle, M.; Krym, D. Half-BPS Solutions locally asymptotic to $AdS_3 \times S^{*3}$ and interface conformal field theories. *J. High Energy Phys.* **2010**, *2010*, 066. [[CrossRef](#)]
9. Chiodaroli, M.; D'Hoker, E.; Gutperle, M. Open Worldsheets for Holographic Interfaces. *J. High Energy Phys.* **2010**, *2010*, 060. [[CrossRef](#)]
10. Kim, N. Comments on AdS_2 solutions from M2-branes on complex curves and the backreacted Kähler geometry. *Eur. Phys. J. C* **2014**, *74*, 2778. [[CrossRef](#)]
11. Lozano, Y.; Macpherson, N.T.; Montero, J.; Colgáin, E.O. New $AdS_3 \times S^2$ T-duals with $\mathcal{N} = (0, 4)$ supersymmetry. *J. High Energy Phys.* **2015**, *2015*, 121. [[CrossRef](#)]
12. Kelekci, O.; Lozano, Y.; Montero, J.; Colgáin, E.; Park, M. Large superconformal near-horizons from M-theory. *Phys. Rev. D* **2016**, *93*, 086010. [[CrossRef](#)]
13. Couzens, C.; Lawrie, C.; Martelli, D.; Schafer-Nameki, S.; Wong, J.M. F-theory and AdS_3/CFT_2 . *J. High Energy Phys.* **2017**, *2017*, 043. [[CrossRef](#)]
14. Dibitetto, G.; Petri, N. BPS objects in $D = 7$ supergravity and their M-theory origin. *J. High Energy Phys.* **2017**, *2017*, 041. [[CrossRef](#)]
15. Dibitetto, G.; Petri, N. 6d surface defects from massive type IIA. *J. High Energy Phys.* **2018**, *2018*, 039. [[CrossRef](#)]
16. Eberhardt, L. Supersymmetric AdS_3 supergravity backgrounds and holography. *J. High Energy Phys.* **2018**, *2018*, 087. [[CrossRef](#)]
17. Corbino, D.; D'Hoker, E.; Uhlemann, C.F. $AdS_2 \times S^6$ versus $AdS_6 \times S^2$ in Type IIB supergravity. *J. High Energy Phys.* **2018**, *2018*, 120. [[CrossRef](#)]
18. Couzens, C.; Martelli, D.; Schafer-Nameki, S. F-theory and $AdS_3/CFT_2(2, 0)$. *J. High Energy Phys.* **2018**, *2018*, 008. [[CrossRef](#)]
19. Dibitetto, G.; Passias, A. $AdS_2 \times S^7$ solutions from D0-F1-D8 intersections. *J. High Energy Phys.* **2018**, *2018*, 190. [[CrossRef](#)]
20. Dibitetto, G.; Lo Monaco, G.; Passias, A.; Petri, N.; Tomasiello, A. AdS_3 Solutions with Exceptional Supersymmetry. *Fortsch. Phys.* **2018**, *66*, 1800060. [[CrossRef](#)]
21. Dibitetto, G.; Petri, N. Surface defects in the D4 – D8 brane system. *J. High Energy Phys.* **2019**, *2019*, 193. [[CrossRef](#)]
22. Dibitetto, G.; Petri, N. AdS_2 solutions and their massive IIA origin. *J. High Energy Phys.* **2019**, *2019*, 107. [[CrossRef](#)]
23. Corbino, D.; D'Hoker, E.; Kaidi, J.; Uhlemann, C.F. Global half-BPS $AdS_2 \times S^6$ solutions in Type IIB. *J. High Energy Phys.* **2019**, *2019*, 039. [[CrossRef](#)]
24. Macpherson, N.T. Type II solutions on $AdS_3 \times S^3 \times S^3$ with large superconformal symmetry. *J. High Energy Phys.* **2019**, *2019*, 089. [[CrossRef](#)]
25. Hong, J.; Macpherson, N.T.; Pando Zayas, L.A. Aspects of AdS_2 classification in M-theory: solutions with mesonic and baryonic charges. *J. High Energy Phys.* **2019**, *2019*, 127. [[CrossRef](#)]
26. Lozano, Y.; Macpherson, N.T.; Nunez, C.; Ramirez, A. AdS_3 solutions in Massive IIA with small $\mathcal{N} = (4, 0)$ supersymmetry. *J. High Energy Phys.* **2020**, *2020*, 129. [[CrossRef](#)]
27. Passias, A.; Prins, D. On AdS_3 solutions of Type IIB. *J. High Energy Phys.* **2020**, *2020*, 048. [[CrossRef](#)]
28. Couzens, C. $\mathcal{N} = (0, 2)$ AdS_3 solutions of type IIB and F-theory with generic fluxes. *J. High Energy Phys.* **2021**, *2021*, 038. [[CrossRef](#)]
29. Couzens, C.; het Lam, H.; Mayer, K. Twisted $\mathcal{N} = 1$ SCFTs and their AdS_3 duals. *J. High Energy Phys.* **2020**, *2020*, 032. [[CrossRef](#)]
30. Legramandi, A.; Macpherson, N.T. AdS_3 solutions with from $\mathcal{N} = (3, 0)$ from $S^3 \times S^3$ fibrations. *Fortsch. Phys.* **2020**, *68*, 2000014. [[CrossRef](#)]
31. Dibitetto, G.; Lozano, Y.; Petri, N.; Ramirez, A. Holographic description of M-branes via AdS_2 . *J. High Energy Phys.* **2020**, *2020*, 037. [[CrossRef](#)]
32. Lüst, D.; Tsimpis, D. AdS_2 type-IIA solutions and scale separation. *J. High Energy Phys.* **2020**, *2020*, 060. [[CrossRef](#)]
33. Corbino, D. Warped AdS_2 and $SU(1, 1|4)$ symmetry in Type IIB. *J. High Energy Phys.* **2020**, *2021*, 60. [[CrossRef](#)]
34. Lozano, Y.; Nunez, C.; Ramirez, A.; Speziali, S. M-strings and AdS_3 solutions to M-theory with small $\mathcal{N} = (0, 4)$ supersymmetry. *J. High Energy Phys.* **2020**, *2020*, 118. [[CrossRef](#)]
35. Chen, K.; Gutperle, M.; Vicino, M. Holographic Line Defects in $D = 4, N = 2$ Gauged Supergravity. *Phys. Rev. D* **2020**, *102*, 026025. [[CrossRef](#)]
36. Faedo, F.; Lozano, Y.; Petri, N. Searching for surface defect CFTs within AdS_3 . *J. High Energy Phys.* **2020**, *2020*, 52. [[CrossRef](#)]
37. Dibitetto, G.; Petri, N. AdS_3 from M-branes at conical singularities. *J. High Energy Phys.* **2020**, *2020*, 129.
38. Lozano, Y.; Nunez, C.; Ramirez, A.; Speziali, S. New AdS_2 backgrounds and $\mathcal{N} = 4$ Conformal Quantum Mechanics. *J. High Energy Phys.* **2021**, *2021*, 277. [[CrossRef](#)]

39. Passias, A.; Prins, D. On supersymmetric AdS₃ solutions of Type II. *arXiv* **2020**, arXiv:2011.00008.
40. Lozano, Y.; Nunez, C.; Ramirez, A.; Speziali, S. AdS₂ duals to ADHM quivers with Wilson lines. *J. High Energy Phys.* **2021**, *2021*, 145. [[CrossRef](#)]
41. Faedo, F.; Lozano, Y.; Petri, N. New $\mathcal{N} = (0, 4)$ AdS₃ near-horizons in Type IIB. *arXiv* **2020**, arXiv:2012.07148.
42. Legramandi, A.; Lo Monaco, G.; Macpherson, N.T. All $\mathcal{N} = (8, 0)$ AdS₃ solutions in 10 and 11 dimensions. *J. High Energy Phys.* **2021**, *2021*, 263. [[CrossRef](#)]
43. Lozano, Y.; Nunez, C.; Ramirez, A. AdS₂ × S² × CY₂ solutions in Type IIB with 8 supersymmetries. *J. High Energy Phys.* **2021**, *2021*, 110. [[CrossRef](#)]
44. Balaguer, J.R.; Dibitetto, G.; Fernández-Melgarejo, J.J. New IIB intersecting brane solutions yielding supersymmetric AdS₃ vacua. *arXiv* **2021**, arXiv:2104.03970.
45. Ramirez, A. AdS₂ geometries and non-Abelian T-duality in non-compact spaces. *arXiv* **2021**, arXiv:2106.09735.
46. Eberhardt, L.; Gaberdiel, M.R.; Li, W. A holographic dual for string theory on AdS₃ × S³ × S³ × S¹. *J. High Energy Phys.* **2017**, *2017*, 111. [[CrossRef](#)]
47. Datta, S.; Eberhardt, L.; Gaberdiel, M.R. Stringy $\mathcal{N} = (2, 2)$ holography for AdS₃. *J. High Energy Phys.* **2018**, *2018*, 146. [[CrossRef](#)]
48. Gaberdiel, M.R.; Gopakumar, R. Tensionless string spectra on AdS₃. *J. High Energy Phys.* **2018**, *2018*, 085. [[CrossRef](#)]
49. Eberhardt, L.; Zadeh, I.G. $\mathcal{N} = (3, 3)$ holography on AdS₃ × (S³ × S³ × S¹)/Z₂. *J. High Energy Phys.* **2018**, *2018*, 143. [[CrossRef](#)]
50. Eberhardt, L.; Gaberdiel, M.R.; Gopakumar, R. The Worldsheet Dual of the Symmetric Product CFT. *J. High Energy Phys.* **2019**, *2019*, 103. [[CrossRef](#)]
51. Eberhardt, L.; Gaberdiel, M.R.; Gopakumar, R. Deriving the AdS₃/CFT₂ correspondence. *J. High Energy Phys.* **2020**, *2020*, 136. [[CrossRef](#)]
52. Lozano, Y.; Macpherson, N.T.; Nunez, C.; Ramirez, A. 1/4 BPS solutions and the AdS₃/CFT₂ correspondence. *Phys. Rev. D* **2020**, *101*, 026014. [[CrossRef](#)]
53. Lozano, Y.; Macpherson, N.T.; Nunez, C.; Ramirez, A. Two dimensional $\mathcal{N} = (0, 4)$ quivers dual to AdS₃ solutions in massive IIA. *J. High Energy Phys.* **2020**, *2020*, 140. [[CrossRef](#)]
54. Lozano, Y.; Macpherson, N.T.; Nunez, C.; Ramirez, A. AdS₃ solutions in massive IIA, defect CFTs and T-duality. *J. High Energy Phys.* **2019**, *2019*, 013. [[CrossRef](#)]
55. Haghighat, B.; Kozcaz, C.; Lockhart, G.; Vafa, C. Orbifolds of M-strings. *Phys. Rev. D* **2014**, *89*, 046003. [[CrossRef](#)]
56. Gadde, A.; Haghighat, B.; Kim, J.; Kim, S.; Lockhart, G.; Vafa, C. 6d String Chains. *J. High Energy Phys.* **2018**, *2018*, 143. [[CrossRef](#)]
57. Maldacena, J.M.; Michelson, J.; Strominger, A. Anti-de Sitter fragmentation. *J. High Energy Phys.* **1999**, *1999*, 011. [[CrossRef](#)]
58. Deneff, F.; Gaiotto, D.; Strominger, A.; Van den Bleeken, D.; Yin, X. Black Hole Deconstruction. *J. High Energy Phys.* **2012**, *2012*, 071. [[CrossRef](#)]
59. Maldacena, J.; Stanford, D. Remarks on the Sachdev-Ye-Kitaev model. *Phys. Rev. D* **2016**, *94*, 106002. [[CrossRef](#)]
60. Maldacena, J.; Stanford, D.; Yang, Z. Conformal symmetry and its breaking in two dimensional Nearly Anti-de-Sitter space. *Prog. Theor. Exp. Phys.* **2016**, *2016*, 12C104. [[CrossRef](#)]
61. Harlow, D.; Jafferis, D. The Factorization Problem in Jackiw-Teitelboim Gravity. *J. High Energy Phys.* **2020**, *2020*, 177. [[CrossRef](#)]
62. Strominger, A. AdS₂ quantum gravity and string theory. *J. High Energy Phys.* **1999**, *1999*, 007. [[CrossRef](#)]
63. Balasubramanian, V.; Naqvi, A.; Simon, J. A Multiboundary AdS orbifold and DLCQ holography: A Universal holographic description of extremal black hole horizons. *J. High Energy Phys.* **2004**, *2004*, 023. [[CrossRef](#)]
64. Balasubramanian, V.; de Boer, J.; Sheikh-Jabbari, M.; Simon, J. What is a chiral 2d CFT? And what does it have to do with extremal black holes? *J. High Energy Phys.* **2010**, *2010*, 017. [[CrossRef](#)]
65. Azeyanagi, T.; Nishioka, T.; Takayanagi, T. Near Extremal Black Hole Entropy as Entanglement Entropy via AdS₂/CFT₁. *Phys. Rev. D* **2008**, *77*, 064005. [[CrossRef](#)]
66. Filippas, K. Non-integrability on AdS₃ supergravity backgrounds. *J. High Energy Phys.* **2020**, *2020*, 027. [[CrossRef](#)]
67. Speziali, S. Spin 2 fluctuations in 1/4 BPS AdS₃/CFT₂. *J. High Energy Phys.* **2020**, *2020*, 079. [[CrossRef](#)]
68. Roychowdhury, D. Fragmentation and defragmentation of strings in type IIA and their holographic duals. *arXiv* **2021**, arXiv:2104.11953.
69. Couzens, C.; Lozano, Y.; Petri, N.; Vandoren, S. 2021. In preparation.
70. Putrov, P.; Song, J.; Yan, W. (0, 4) dualities. *J. High Energy Phys.* **2016**, *2016*, 185. [[CrossRef](#)]
71. Kim, H.C. Line defects and 5d instanton partition functions. *J. High Energy Phys.* **2016**, *2016*, 199. [[CrossRef](#)]
72. Assel, B.; Sciarappa, A. Wilson loops in 5d $\mathcal{N} = 1$ theories and S-duality. *J. High Energy Phys.* **2018**, *2018*, 082. [[CrossRef](#)]
73. Assel, B.; Sciarappa, A. On monopole bubbling contributions to 't Hooft loops. *J. High Energy Phys.* **2019**, *2019*, 180. [[CrossRef](#)]
74. Tong, D.; Wong, K. Instantons, Wilson lines, and D-branes. *Phys. Rev. D* **2015**, *91*, 026007. [[CrossRef](#)]
75. Chang, C.M.; Ganor, O.; Oh, J. An index for ray operators in 5d E_n SCFTs. *J. High Energy Phys.* **2017**, *2017*, 018. [[CrossRef](#)]
76. Yamaguchi, S. Wilson loops of anti-symmetric representation and D5-branes. *J. High Energy Phys.* **2006**, *2006*, 037. [[CrossRef](#)]
77. Gomis, J.; Passerini, F. Wilson Loops as D3-Branes. *J. High Energy Phys.* **2007**, *2007*, 097. [[CrossRef](#)]
78. Witten, E. Baryons and branes in anti-de Sitter space. *J. High Energy Phys.* **1998**, *1998*, 006. [[CrossRef](#)]
79. Brandhuber, A.; Oz, Y. The D-4 - D-8 brane system and five-dimensional fixed points. *Phys. Lett. B* **1999**, *460*, 307–312. [[CrossRef](#)]
80. Deneff, F. Quantum quivers and Hall/hole halos. *J. High Energy Phys.* **2002**, *2002*, 023. [[CrossRef](#)]

81. Haghighat, B.; Murthy, S.; Vafa, C.; Vandoren, S. F-Theory, Spinning Black Holes and Multi-string Branches. *J. High Energy Phys.* **2016**, *2016*, 009. [[CrossRef](#)]
82. Couzens, C.; het Lam, H.; Mayer, K.; Vandoren, S. Black Holes and (0, 4) SCFTs from Type IIB on K3. *J. High Energy Phys.* **2019**, *2019*, 043. [[CrossRef](#)]
83. Hull, C.; Marcus, E.; Stemerding, K.; Vandoren, S. Black holes in string theory with duality twists. *J. High Energy Phys.* **2020**, *2020*, 086. [[CrossRef](#)]
84. Benini, F.; Hristov, K.; Zaffaroni, A. Black hole microstates in AdS₄ from supersymmetric localization. *J. High Energy Phys.* **2016**, *2016*, 054. [[CrossRef](#)]
85. Seiberg, N. Five-dimensional SUSY field theories, nontrivial fixed points and string dynamics. *Phys. Lett. B* **1996**, *388*, 753–760. [[CrossRef](#)]