

B Physics at the Tevatron

Karen Gibson^{*}[†]

University of Pittsburgh

E-mail: krg20@pitt.edu

I present recent results in B physics from the CDF and D0 experiments at the Fermilab Tevatron. A number of lifetime results, spectroscopy results, and CP violation results are presented, all of which have been released in 2008. The breadth of the work in B physics that has been produced by the Tevatron experiments indicates the overall success of the program, as well as the new and improved results which can still be anticipated in the final years of Run II.

*2008 Physics at LHC
September 29 - 4 October 2008
Split, Croatia*

^{*}Speaker.

[†]On behalf of the CDF and D0 Collaborations

1. Introduction

Over the past five years, the CDF and D0 experiments have produced prolific B physics results during Run II of the Fermilab Tevatron. These results have been complementary in nature to the excellent B physics programs conducted by the BABAR and Belle experiments, both of which operate at electron-positron colliders tuned to the $\Upsilon(4S)$ threshold (called B factories), giving the experiments very clean, high-statistics samples of B^0 and B^+ mesons[1] to study.¹ In contrast, the $\sqrt{s} = 1.96 \text{ TeV}/c^2$ $p\bar{p}$ collisions at the Tevatron produce all flavors of B hadrons (in addition to a lot of other things), giving relatively large samples of B_s^0 mesons and Λ_b^0 baryons, as well as large samples of light B^0 and B^+ mesons. This has allowed the Tevatron to measure the frequency of $B_s^0 - \bar{B}_s^0$ oscillations [2] and discover new B baryons, such as the $\Sigma_b^{(*)\pm}$ [3] and the Ξ_b^- [4, 5]. With more than 4 fb^{-1} of data on tape, the samples of B hadrons which will be available for analysis will allow (and in some cases already are allowing) CDF and D0 to produce CP violation and lifetime measurements in B^0 and B^+ that compete with the B factories for the most precise results in the world.

In fact, the Tevatron experiments are in a special position; the LHC experiments do not yet have data for B physics, while the B factories have either ceased operation (at SLAC) or are transitioning from their first run (at KEK) to upgrades which will allow much more integrated luminosity in a few years time. I will highlight recent results from the Tevatron which were made public in 2008 and give some indication of results that can be anticipated when the full Tevatron dataset becomes available. The results presented are divided into three main categories: production, lifetimes, and CP violation.

When presenting results from the CDF and D0 experiments, it is worthwhile to note that the two detectors have different strengths in detecting B hadrons. The CDF detector features a strong tracking system that provides good mass and vertex resolution, and the ability to trigger on displaced tracks, which allows CDF to select a large sample of fully-reconstructed B events. The D0 detector has excellent calorimetry and muon identification that provides a large sample of semi-leptonic B decays. The D0 experiment also reverses the direction of the magnetic field in the detector regularly, which is beneficial in direct CP violation measurements by minimizing systematic uncertainties due to charge asymmetries in the data.

2. Production

In all aspects of particle physics, it is important both to look for things that we expect and also for things that we don't expect. Flavor physics is no exception to this rule and discoveries of the X , Y , and Z particles by the Belle and BABAR experiments [6] demonstrate that unexpected particles are still present in current datasets. Although the Y and Z particles are unobserved by the Tevatron experiments, CDF has made several important contributions to the study of the $X(3872)$ particle. The Tevatron experiments have also recently observed some b -baryons for the first time, including the Σ_b^\pm baryon, the Ξ_b^- baryon, and, recently, the Ω_b^- baryon. The existence of these particles has been predicted, but until Run II none had been observed.

¹The Belle experiment has collected some data at the $\Upsilon(5S)$ threshold, which decays to $B_s^0 - \bar{B}_s^0$ pairs, but the sample, which is presently small, has not yet produced significant physics results.

The $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ was first observed by the Belle collaboration in 2003 [6]. Although five years have passed since its observation, it is still not known whether it is a D^*D bound state, a tetra-quark state, or something else. The CDF collaboration has searched for a mass splitting in order to test the tetra-quark hypothesis. If no mass splitting is observed, a precision measurement of the absolute mass is interesting, as it checks whether the mass is within the threshold of a bound-state D^*D “molecule” [7].

In order to test the tetra-quark interpretation, the CDF collaboration fits the mass distribution of the $X(3872)$ with a Breit-Wigner convolved with detector resolution and generates pseudo-experiments to determine whether the two mass hypothesis is consistent with the distribution observed in data. The data shows no evidence of a mass splitting and an upper limit on the mass splitting is set. The absolute mass is then measured, finding a mass which is below the threshold of a D^*D bound state, but which is within 1σ of the threshold, leaving open the possibility of the bound-state explanation. The limits on the mass splitting and the absolute mass are

$$\begin{aligned}\Delta m(X(3872)) &< 3.2(3.6) \text{ MeV}/c^2 \text{ at 90\% (95\%) CL,} \\ m(X(3872)) &= 3871.61 \pm 0.16 \text{ (stat.)} \pm 0.19 \text{ (syst.) MeV}/c^2.\end{aligned}\quad (2.1)$$

One of the most exciting physics results from the Tevatron this year is D0’s recent observation of the Ω_b^- baryon [8] in the decay $\Omega_b^- \rightarrow J/\psi [\rightarrow \mu^+ \mu^-] \Omega^- [\rightarrow \Lambda^0 (\rightarrow p \pi^-) K^-]$. This observation was made with 1.3 fb^{-1} of data and builds on D0’s previous observation of the $\Xi_b^- \rightarrow J/\psi [\rightarrow \mu^+ \mu^-] \Xi^- [\rightarrow \Lambda^0 (\rightarrow p \pi^-) \pi^-]$ [4]. Special selection techniques, such as a boosted decision tree and re-processing the data with a higher impact parameter requirement on tracks to increase acceptance, improved the Ω^- reconstruction significantly. The invariant mass distribution of the Ω_b^- is shown in Fig. 1. D0 observes $17.8 \pm 4.9 \text{ (stat.)} \pm 0.8 \text{ (syst.)}$ events with 5.4σ significance and measures the mass $m(\Omega_b^-) = 6.165 \pm 0.010 \text{ (stat.)} \pm 0.013 \text{ (syst.) GeV}/c^2$. D0 calculates the production fraction times branching ratio of the observed Ω_b^- decays relative to the observed Ξ_b^- decay to be

$$\frac{f(b \rightarrow \Omega_b^-) Br(\Omega_b^- \rightarrow J/\psi / \Omega^-)}{f(b \rightarrow \Xi_b^-) Br(\Xi_b^- \rightarrow J/\psi / \Xi^-)} = 0.80 \pm 0.32 \text{ (stat.)} \begin{array}{l} 0.14 \\ 0.22 \end{array} \text{ (syst.)}. \quad (2.2)$$

3. Lifetimes

The proper decay times of B hadrons are fundamental quantities, the measurement of which, along with measurements of their masses and decay rates to different final states, helps to paint a complete picture of B hadrons. Precise measurements of B hadron lifetimes test the predictions of the Heavy Quark Effective Theory (HQET), which predicts $\tau(B^+) > \tau(B^0) \approx \tau(B_s^0) > \tau(\Lambda_b^0)$ ². Previously, measurements of $\tau(B_s^0)$ and $\tau(\Lambda_b^0)$ showed $1 - 2\sigma$ discrepancies with the HQET predictions, which were difficult to explain theoretically.

A recent CDF measurement of the B_s^0 lifetime in hadronic $B_s^0 \rightarrow D_s^- [\rightarrow \phi (\rightarrow K^+ K^-) \pi^-] \pi^+$ decays finds a lifetime of $c\tau(B_s^0) = 455 \pm 12 \text{ (stat.)} \pm 7 \text{ (syst.)} \mu\text{m}$ [9], which is quite compatible with the current average B^0 lifetime $c\tau(B_s^0) = 458.7 \pm 2.7 \mu\text{m}$ [10]. A similar measurement of

²All of these lifetimes are predicted to be significantly longer than the lifetime of the B_c^+ .




Figure 1: Ω_b^- invariant mass observed by the D0 collaboration.

the Λ_b^0 lifetime made by CDF in the hadronic decay channel $\Lambda_b^0 \rightarrow \Lambda_c^+ [\rightarrow pK^-\pi^+]\pi^+$ also finds much better compatibility with HQET than had previously been established, with CDF finding $c\tau(\Lambda_b^0) = 423 \pm 14$ (stat.) ± 9 (syst.) μm and $c\tau(\Lambda_b^0)/c\tau(B^0) = 0.92 \pm 0.04$ [11]. This measurement resolves 3σ discrepancy between recent Λ_b^0 lifetime measurements and provides the most precise measurement of the Λ_b^0 lifetime to-date. Both CDF and D0 collaborations have also measured the B_c^+ meson lifetime in semi-leptonic modes. D0 has measured the lifetime in the decay channel $B_c^+ \rightarrow J/\psi\mu^+X$, while CDF measures the lifetime in both $B_c^+ \rightarrow J/\psi\mu^+X$ and $B_c^+ \rightarrow J/\psi e^+X$ decays. The measured lifetimes are consistent between the two experiments, with D0 finding $c\tau(B_c^+) = 134 \pm 11$ (stat.) ± 10 (syst.) μm [12] and CDF measuring $c\tau(B_c^+) = 142 \pm 15$ (stat.) ± 6 (syst.) μm [13]. All of these recent lifetime measurements are in good agreement with theoretical predictions [14].

4. CP Violation

There are three known varieties of CP violation possible in hadron systems. CP violation in the decay of hadrons is called direct CP violation and is the only type of CP violation possible for charged mesons such as the B^+ meson. Indirect CP violation can occur in the mixing of neutral mesons; examples of which include CP asymmetries in semi-leptonic decays of neutral mesons. Finally, CP violation can occur in the interference between mixing and decay. A well-known example of this in B physics is the CP -violating phase β in the $B^0 \rightarrow J/\psi K_s^0$ decay channel, although the analogous CP -violating phase β_s , which can be measured in $B_s^0 \rightarrow J/\psi\phi$ decays, has recently generated considerable theoretical and experimental interest. The most powerful measurements of CP -violating phases generally involve the use of flavor-tagging, which determines the flavor of the B hadron at production. This task is straight-forward for charged particles like the B^+ , which are self-tagging, but is much more difficult for neutral mesons, such as the B^0 and B_s^0 . Powerful flavor tags include same-side kaon tags and opposite-side lepton and jet-charge tags.

Mixing in neutral meson systems is described by a Schrodinger equation

$$i\frac{d}{dt}(|B_s^0\rangle|\bar{B}_s^0\rangle) = (M)^s - \frac{i}{2}\Gamma^s(|B_s^0\rangle|\bar{B}_s^0\rangle), \quad (4.1)$$

where M^s is the mass matrix and Γ^s is the decay matrix that relates the flavor eigenstates $|B_s^0\rangle$ and $|\bar{B}_s^0\rangle$ to the mass eigenstates $|B_s^H\rangle$ and $|B_s^L\rangle$, such that

$$|B_s^H\rangle = p|B_s^0\rangle - q|\bar{B}_s^0\rangle, \quad (4.2)$$

$$|B_s^L\rangle = p|B_s^0\rangle + q|\bar{B}_s^0\rangle. \quad (4.3)$$

where $|q/p| = |(V_{tb}V_{ts}^*) / (V_{tb}^*V_{ts})| \equiv 1$ is related to the strength of the $t \rightarrow s$ and $t \rightarrow b$ weak transitions described by the CKM quark matrix [15]. The mass difference between the two mass eigenstates gives rise to the oscillation frequency $\Delta m_s \equiv m_H - m_L$ between the B_s^0 and \bar{B}_s^0 and is related to the off-diagonal mass matrix element $\Delta m_s \approx 2|M_{12}^s|$, while the width difference $\Delta\Gamma \equiv \Gamma_L^s - \Gamma_H^s$ is related to the off-diagonal decay matrix element gives rise to a lifetime difference between the two mass eigenstates $\Delta\Gamma \approx 2|\Gamma_{12}^s \cos(\phi_s)|$, where $\phi_s = \arg(-M_{12}/\Gamma_{12}) \sim 0.004$ in the standard model.

A primary reason for interest in the CP -violating phase β_s in $B_s^0 \rightarrow J/\psi\phi$ decays is that this phase is particularly sensitive to the presence of new physics, since it is theoretically well-predicted from elements of the CKM quark matrix to be nearly zero, $\beta_s = \arg\left(-\frac{V_{ts}V_{tb}^*}{V_{cs}V_{cb}^*}\right) \sim 0.02$ [16]. Consequently, any observation of a large CP -violating phase would be an indication of new physics. If present, new physics is likely to appear in the mixing of B_s^0 into \bar{B}_s^0 , which involves $t \rightarrow s$ quark transitions in the standard model but which could also involve new, heavy particles.

In late 2007 CDF announced the first flavor-tagged measurement of β_s in 1.3 fb^{-1} of data and observed a large deviation from the standard model prediction with a 1.5σ significance [17]. CDF updated this result for the summer of 2008, using data through 2.8 fb^{-1} , and finds that the discrepancy remains and its significance increases to 1.8σ [18]. Intriguingly, D0 finds a completely compatible result in 2.8 fb^{-1} of data, observing a discrepancy with the standard model of 1.7σ [19]. A recent combination of the original 1.3 fb^{-1} CDF result and the D0 result by the Heavy Flavor Averaging Group (HFAG), shown in Fig. 2, finds a discrepancy with the standard model of 2.2σ . While not statistically significant, this result is very intriguing, particularly when taken in conjunction with the asymmetry of direct CP asymmetries in B^0 and B^+ decays to $K\pi$ final states observed by the Belle and **B**_A**B**_R collaborations [20]. George Hou et al. have predicted that both sets of discrepancies could be due to the presence of a heavy t' quark, with a mass between $300 \text{ GeV}/c^2$ and $\sim 1 \text{ TeV}/c^2$ [21], as both effects involve $t \rightarrow s$ quark transitions.

The D0 collaboration has also recently announced a time-dependent, flavor-tagged measurement of the semi-leptonic B_s^0 asymmetry a_{sl}^s , measured in the decay $B_s^0 \rightarrow D_s^- [\rightarrow \phi\pi^-]\mu^+\nu_\mu X$ [22]. The time dependence of this decay depends on $\tau(B_s^0)$, $\Delta\Gamma$, Δm_s . D0 performs a maximum likelihood fit to extract $a_{sl}^s = -0.0024 \pm 0.0117 \text{ (stat.)}^{+0.0015}_{-0.0024} \text{ (syst.)}$, which is the most precise measurement of this quantity to date.

Another probe of the mixing-related quantities in the B_s^0 system is the measurement of the branching ratio of $B_s^0 \rightarrow D_s^{(*)+}D_s^{(*)-}$ to determine $\Delta\Gamma$. D0 has measured this branching fraction by fully reconstructing one D_s in the $\phi [\rightarrow K^+K^-]\pi$ decay mode and the other D_s semi-leptonically in the $\phi\mu\nu_\mu X$ decay mode [23]. Under certain theoretical assumptions [16], the $B_s^0 \rightarrow D_s^{(*)+}D_s^{(*)-}$ decay is nearly CP even, giving

$$2Br(B_s^0 \rightarrow D_s^{(*)+}D_s^{(*)-}) \approx \Delta\Gamma^{CP} \left[\frac{1 + \cos\phi_s}{2\Gamma_L} + \frac{1 - \cos\phi_s}{2\Gamma_H} \right].. \quad (4.4)$$




Figure 2: HFAG combination of CDF and D0 results in $\phi_s^{J/\psi\phi} - \Delta\Gamma$ plane. Note that $\phi_s^{J/\psi\phi} - \Delta\Gamma \equiv -2\beta_s$.

D0 finds a branching ratio of $Br(B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}) = 0.042 \pm 0.015 \text{ (stat.)} \pm 0.017 \text{ (syst.)}$. Under standard model assumptions $\phi_s = 0.004 \approx 0$ and $\Delta\Gamma^{CP} = \Delta\Gamma$ and D0 extracts $\Delta\Gamma/\Gamma = 0.088 \pm 0.030 \text{ (stat.)} \pm 0.017 \text{ (syst.)}$, which is consistent with the HFAG average $\Delta\Gamma/\Gamma = 0.096^{+0.048}_{-0.053}$ [24].

Finally, D0 also has a recent measurement of the direct CP asymmetry in $B^+ \rightarrow J/\psi K^+(\pi^+)$ decays [25]. The standard model predicts that $A_{CP}(B^+ \rightarrow J/\psi K^+) \sim 0.003$, although new physics may give rise to asymmetries as large ~ 0.01 . Clearly this is a measurement which requires high precision in order to test the standard model. The CP asymmetry measured is defined as

$$A_{CP}(B^+ \rightarrow J/\psi K^+(\pi^+)) = \frac{N(B^- \rightarrow J/\psi K^-(\pi^-)) - N(B^+ \rightarrow J/\psi K^+(\pi^+))}{N(B^- \rightarrow J/\psi K^-(\pi^-)) + N(B^+ \rightarrow J/\psi K^+(\pi^+))} \quad (4.5)$$

D0 obtains $A_{CP}(B^+ \rightarrow J/\psi K^+) = +0.007 \pm 0.0061 \text{ (stat.)} \pm 0.0027 \text{ (syst.)}$, which is approaching the sensitivity necessary for test of the standard model, and $A_{CP}(B^+ \rightarrow J/\psi K^+) = -0.09 \pm 0.08 \text{ (stat.)} \pm 0.03 \text{ (syst.)}$.

5. Outlook

Looking beyond the present set of B physics results from the Tevatron, a number of new and updated measurements can be anticipated. Most of the Tevatron CP violation measurements will be updated with two to four times the data presently used by the end of Run II. In addition, now that the flavor-tagging is well-established and statistics are increasing, more tagged CP violation measurements can be expected. The lifetime measurements can be updated with higher statistics; updated $B \rightarrow J/\psi X$ lifetime measurements are expected with twice the statistics of previous measurements, which should give the most precise B^+ and Λ_b^0 lifetimes to date. Finally, while never

something to be counted on, it is reasonable that the Tevatron may still hold a surprise or two concerning the observation of new states. This is an exciting period for Tevatron B physics, with many significant contributions made already and many more to come within the next couple of years.

References

- [1] Charge-conjugate states are implied throughout the paper unless otherwise specified.
- [2] A. Abulencia et al. (CDF Collaboration), Phys. Rev. Lett. **97**, 242003 (2006).
- [3] T. Aaltonen et al (CDF Collaboration), Phys. Rev. Lett. **99**, 202001 (2007).
- [4] V. M. Abazov et al. (D0 Collaboration), Phys. Rev. Lett. **99**, 052001 (2007).
- [5] T. Aaltonen et al. (CDF Collaboration), Phys. Rev. Lett. **99** 052002 (2007).
- [6] S. K. Choi et al. (Belle Collaboration), Phys. Rev. Lett. **91**, 262001 (2003);
P. Pakhlov et al. (Belle Collaboration), Phys. Rev. Lett. **100**, 202001 (2008);
K. Abe et al. (Belle Collaboration), Phys. Rev. Lett. **94**, 182002 (2005);
B. Aubert et al. (BABAR Collaboration), Phys. Rev. Lett. **95**, 142001 (2005).
- [7] <http://www-cdf.fnal.gov/physics/new/bottom/080724.blessed-X-Mass/cdfpubnote.ps>
- [8] V. M. Abazov et al. (D0 Collaboration) arXiv:0808.4142 [hep-ex] (2008).
- [9] http://www-cdf.fnal.gov/physics/new/bottom/080207.blessed-bs-lifetime/pubnote_v2.ps
- [10] C. Amsler et al., Physics Letters B **667**, 1 (2008).
- [11] http://www-cdf.fnal.gov/physics/new/bottom/080703.blessed-lblcpi-ct/lambdab_lifetime_public.pdf
- [12] V. M. Abazov et al. (D0 Collaboration), arXiv:0805.2614 [hep-ex] (2008).
- [13] http://www-cdf.fnal.gov/physics/new/bottom/080327.blessed-BC_LT_SemiLeptonic/blessed-bc-lifetime.pdf
- [14] V. V. Kiselev, hep-ph/0308214 (2003).
- [15] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963);
M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [16] I. Dunietz, R. Fleischer, and U. Nierste", Phys. Rev. D **63**, 114015 (2001).
- [17] T. Aaltonen et al. (CDF Collaboration), Phys. Rev. Lett. **100**, 161802 (2008).
- [18] http://www-cdf.fnal.gov/physics/new/bottom/080724.blessed-tagged_BsJPsiPhi_update_prelim/public_note.pdf
- [19] V. M. Abazov et al. (D0 Collaboration), arXiv:0802.2255 [hep-ex] (2008).
- [20] The Belle Collaboration, Nature, **452**, 332 (2008);
B. Aubert et al. (BABAR Collaboration), "Phys. Rev. Lett." **99**, 021603 (2007);
B. Aubert et al. (BABAR Collaboration), Phys. Rev. D **76**, 091102 (2007).
- [21] W.-S. Hou, M. Nagashima, and A. Soddu, Phys. Rev. Lett. **95**, 141601 (2005);
W.-S. Hou, M. Nagashima, and A. Soddu, Phys. Rev. D **76**, 016004 (2007).
- [22] <http://www-d0.fnal.gov/Run2Physics/WWW/results/prelim/B/B55/B55.pdf>
- [23] V. M. Abazov et al. (D0 Collaboration), arXiv:0811.2173 [hep-ex] (2008).
- [24] E. Barberio et al., arXiv:0808.1297 and online update at <http://www.slac.stanford.edu/xorg/hfag>.
- [25] V. M. Abazov et al. (D0 Collaboration), Phys. Rev. Lett. **100**, 211802 (2008).