



*symmetry*

IMPACT  
FACTOR  
2.2

CITESCORE  
5.4

Article

---

# On the Analytic Continuation of Appell's Hypergeometric Function $F_2$ to Some Symmetric Domains in the Space $\mathbb{C}^2$

---

Roman Dmytryshyn

Special Issue

Theory and Applications of Special Functions II

Edited by


Dr. Diego Caratelli



<https://doi.org/10.3390/sym16111480>

Article

# On the Analytic Continuation of Appell's Hypergeometric Function $F_2$ to Some Symmetric Domains in the Space $\mathbb{C}^2$

Roman Dmytryshyn 

Department of Mathematical and Functional Analysis, Vasyl Stefanyk Precarpathian National University, 57 Shevchenko Str., 76018 Ivano-Frankivsk, Ukraine; roman.dmytryshyn@pnu.edu.ua

**Abstract:** The paper considers the problem of representation and extension of Appell's hypergeometric functions by a special family of functions—branched continued fractions. Here, we establish new symmetric domains of the analytical continuation of Appell's hypergeometric function  $F_2$  with real and complex parameters, using their branched continued fraction expansions whose elements are polynomials in the space  $\mathbb{C}^2$ . To do this, we used a technique that extends the domain of convergence of the branched continued fraction, which is already known for a small domain, to a larger domain, as well as the PC method to prove that it is also the domain of analytical continuation. A few examples are provided at the end to illustrate this.

**Keywords:** Appell's hypergeometric function; branched continued fraction; analytic continuation; convergence; approximation by rational functions

**MSC:** 33C65; 32A17; 32D99; 40A99; 41A20

## 1. Introduction

This paper considers the Appell's hypergeometric function  $F_2$  defined as (see, [1,2])

$$F_2(\alpha, \beta, \beta'; \gamma, \gamma'; \mathbf{z}) = \sum_{p,q=0}^{+\infty} \frac{(\alpha)_{p+q} (\beta)_p (\beta')_q}{(\gamma)_p (\gamma')_q} \frac{z_1^p z_2^q}{p! q!}, \quad |z_1| + |z_2| < 1,$$

where  $\alpha, \beta, \beta', \gamma, \gamma' \in \mathbb{C}$ ,  $\gamma, \gamma' \notin \{0, -1, -2, \dots\}$ ,  $(\cdot)_k$  is the Pochhammer symbol,  $\mathbf{z} = (z_1, z_2) \in \mathbb{C}^2$ .

Appell's hypergeometric function  $F_2$  surprisingly appears in various applications, in particular, in materials science for the compute of the canonical partition function of the model of heteropolymer in the form of a freely jointed chain [3], in probability theory and statistics for the study of the compound gamma bivariate distribution [4], in the theory of the quantum Hall effect for the explicit evaluation of the matrix elements of the Coulomb interaction of two-body [5], in the quantum field theory for the evaluation of Feynman integrals [6] and a two-loop diagram of the propagator-type (the so-called propagator seagull) [7], in the spectral theory of atom, molecule and plasma for the compute of multipole matrix elements [8].

Many works are devoted to the study of the Appell's hypergeometric function  $F_2$  itself, in particular, to the establishment of recurrence relations [9,10], reduction and transformation formulas [11], to the construction of analytic continuations [12–14], integral representations [15,16] and asymptotic expansions [16–20]. We also note the work [21], which presents the *Mathematica* package AppellF2.wl, dedicated to the evaluation of the Appell's hypergeometric function  $F_2$ .

This paper discusses the representation and analytical extension of the Appell's hypergeometric function  $F_2$  due a special family of functions—branched continued fractions.



**Citation:** Dmytryshyn, R. On the Analytic Continuation of Appell's Hypergeometric Function  $F_2$  to Some Symmetric Domains in the Space  $\mathbb{C}^2$ . *Symmetry* **2024**, *16*, 1480. <https://doi.org/10.3390/sym16111480>

Academic Editor: Junesang Choi

Received: 1 October 2024

Revised: 18 October 2024

Accepted: 30 October 2024

Published: 6 November 2024



**Copyright:** © 2024 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Domains of analytical continuation will be symmetric domains of convergence of co-called confluent branched continued fractions.

Research in the direction mentioned above was started in [22], where a formal branched continued fraction expansion was constructed for the following ratio

$$\frac{F_2(\alpha, \beta, \beta'; \gamma, \gamma'; \mathbf{z})}{F_2(\alpha + 1, \beta, \beta'; \gamma + 1, \gamma'; \mathbf{z})}$$

In [23], the following formal branched continued fraction expansion

$$\frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; \mathbf{z})}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; \mathbf{z})} = 1 - z_1 - \frac{h_1 z_2}{1 - \frac{h_2 z_2}{1 - z_1 - \frac{h_3 z_2}{1 - \frac{h_4 z_2}{1 - z_1 - \frac{h_5 z_2}{1 - \dots}}}}}, \tag{1}$$

where

$$h_{2k-1} = \frac{(\beta' + k - 1)(\gamma' - \alpha + k - 1)}{(\gamma' + 2k - 2)(\gamma' + 2k - 1)} \quad \text{and} \quad h_{2k} = \frac{(\alpha + k)(\gamma' - \beta' + k)}{(\gamma' + 2k - 1)(\gamma' + 2k)}, \quad k \geq 1, \tag{2}$$

was considered, and it was shown that

$$\Psi_{d,h} = \left\{ \mathbf{z} \in \mathbb{C}^2 : z_1 \notin [1 - d, +\infty), z_2 \notin \left[ \frac{d}{4h}, +\infty \right) \right\}, \quad h > 0, 0 < d < 1,$$

is the domain of the analytical continuation of the function on the left side of (1) provided that  $0 < h_k \leq h$  for all  $k \geq 1$ .

The following theorem holds:

**Theorem 1.** Suppose that  $\alpha, \beta',$  and  $\gamma'$  are complex constants such that

$$|h_k| - \text{Re}(h_k) \leq pq, \quad k \geq 1, \tag{3}$$

where  $h_k, k \geq 1,$  are given in (2) herewith  $\gamma' \notin \{0, -1, -2, \dots\},$

$$p > 0 \quad \text{and} \quad 0 < q < 1. \tag{4}$$

Then, the following statements hold:

(A) The branched continued fraction

$$1 - z_1 + \frac{h_1 z_2}{1 + \frac{h_2 z_2}{1 - z_1 + \frac{h_3 z_2}{1 + \frac{h_4 z_2}{1 - z_1 + \frac{h_5 z_2}{1 + \dots}}}}}, \tag{5}$$

converges uniformly on every compact subset of the domain

$$\Omega_{p,q}^{d,h} = \Omega_{p,q} \cup \Omega^{d,h}, \tag{6}$$

where

$$\Omega_{p,q} = \left\{ \mathbf{z} \in \mathbb{C}^2 : \operatorname{Re}(z_1 e^{-i/2} \arg(z_2)) < (1-q) \cos\left(\frac{\arg(z_2)}{2}\right), |z_2| < \frac{1 + \cos(\arg(z_2))}{4p} \right\} \quad (7)$$

and

$$\Omega^{d,h} = \left\{ \mathbf{z} \in \mathbb{C}^2 : |z_1| < \frac{1-d}{2}, |z_2| < \frac{d(1-d)}{2h} \right\}, \quad (8)$$

where

$$h = \sup_{k \in \mathbb{N}} |h_k| \quad \text{and} \quad 0 < d < 1, \quad (9)$$

to the function  $f(\mathbf{z})$  holomorphic in the domain  $\Omega_{p,q}^{d,h}$ .

(B) The function  $f(\mathbf{z})$  is an analytic continuation of the function

$$\frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} \quad (10)$$

in the domain (6).

This paper is organized as follows. In Section 2, we give the necessary definitions and statements and prove Theorem 2, the result of which is a certain contribution to the theory of branched continued fractions. In the next section, we prove Theorem 1 and several important consequences from it. In Section 4, we give some examples. Finally, we collect our conclusions in Section 5.

## 2. Definitions and Auxiliary Results

Let us recall the necessary concepts of convergence in the theory of branched continued fractions (see, [24,25]).

We set  $i(0) = 0$ ,  $\mathfrak{J}_0 = \{0\}$ , and

$$\mathfrak{J}_k = \{i(k) : i(k) = (i_1, i_2, \dots, i_k), 1 \leq i_r \leq 2, 1 \leq r \leq k\}, \quad k \geq 1.$$

By Bodnar ([25], p. 15) for each  $r \geq 1$  the symbol  $\mathbf{u}^{(r)}$  denotes a vector in  $\mathbb{C}^{2^r}$  with components  $u_{j(r)}$ ,  $j(r) \in \mathfrak{J}_r$ ; for each  $r \geq 1$ ,  $k \geq 1$  and for each multiindex  $i(k) \in \mathfrak{J}_k$  the symbol  $\mathbf{u}_{i(k)}^{(r)}$  is a vector in  $\mathbb{C}^{2^r}$  with components  $u_{i(k),j(r)}$ ,  $i(k) \in \mathfrak{J}_k$ ,  $1 \leq j_s \leq 2$ ,  $1 \leq s \leq r$ ,  $j_0 = i_k$ , with the following order of components:

- (i)  $u_{n(r)} \prec u_{m(r)}$  ( $u_{i(k),n(r)} \prec u_{i(k),m(r)}$ ), if  $n(r) \prec m(r)$ ;
- (ii)  $n(r) \prec m(r)$ , if  $n_1 < m_1$  or there exists index  $s$ ,  $1 \leq s < r$ , such that  $n_p = m_p$ ,  $1 \leq p \leq s$ , and  $n_{s+1} < m_{s+1}$ .

Let the ordered pair of sequences

$$\langle \{a_{i(k)}\}_{i(k) \in \mathfrak{J}_k, k \in \mathbb{N}}, \{b_{i(k)}\}_{i(k) \in \mathfrak{J}_k, k \in \mathbb{N}_0} \rangle,$$

of complex numbers such that:

- (\*)  $a_{i(k)} \neq 0$  for all  $i(k) \in \mathfrak{J}_k, k \geq 1$ ;
- (\*\*) if for  $k \geq 1$  there exists a multiindex  $i(k) \in \mathfrak{J}_k$  such that  $b_{i(k)} = 0$ , then  $b_{i(k-1),j} \neq 0$  for  $1 \leq j \leq 2$  and  $j \neq i_k$ ,

gives rise to sequence  $\{s_{i(k)}(\mathbf{w}_{i(k)}^{(1)})\}_{i(k) \in \mathfrak{J}_k, k \in \mathbb{N}_0}$  herewith  $\mathbf{w}_0^{(1)} = \mathbf{w}^{(1)}$  and  $\{S_k(\mathbf{w}^{(k+1)})\}_{k \in \mathbb{N}_0}$  of two-dimensional linear fractional transformations

$$s_0(\mathbf{w}^{(1)}) = b_0 + w_1 + w_2, \quad v_{i(k)} = s_{i(k)}(\mathbf{w}_{i(k)}^{(1)}) = \frac{a_{i(k)}}{b_{i(k)} + w_{i(k),1} + w_{i(k),2}}, \quad i(k) \in \mathfrak{I}_k, k \geq 1,$$

and

$$S_0(\mathbf{w}^{(1)}) = s_0(\mathbf{w}^{(1)}), \quad S_k(\mathbf{w}^{(k+1)}) = S_{k-1}(\mathbf{v}^{(k)}), \quad k \geq 1,$$

and to a sequence  $\{f_k\}_{k \in \mathbb{N}_0}$ , given by

$$f_k = S_k(\mathbf{0}^{(k+1)}) \in \widehat{\mathbb{C}}, \quad k \geq 0,$$

where  $\mathbf{0}^{(k+1)} = (0, 0, \dots, 0)$  is a vector in  $\mathbb{C}^{2^{k+1}}$ .

**Definition 1.** *The ordered pair*

$$\langle \langle \{a_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}}, \{b_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}_0} \rangle, \{f_k\}_{k \in \mathbb{N}_0} \rangle$$

is the branched continued fraction denoted by symbols

$$b_0 + \sum_{i_1=1}^2 \frac{a_{i_1(1)}}{b_{i_1(1)} + \sum_{i_2=1}^2 \frac{a_{i_2(2)}}{b_{i_2(2)} + \dots + \sum_{i_k=1}^2 \frac{a_{i_k(k)}}{b_{i_k(k)} + \dots}}$$

The numbers  $b_0, a_{i(k)}$ , and  $b_{i(k)}$  are called elements of the branched continued fraction. The value

$$f_k = b_0 + \sum_{i_1=1}^2 \frac{a_{i_1(1)}}{b_{i_1(1)} + \sum_{i_2=1}^2 \frac{a_{i_2(2)}}{b_{i_2(2)} + \dots + \sum_{i_k=1}^2 \frac{a_{i_k(k)}}{b_{i_k(k)}}}$$

is called the  $k$ th approximant of the branched continued fraction.

Note that a new and more general so-called sets approach to the concept of a branched continued fraction was proposed by Antonova in [26].

Next, considering the branched continued fractions, we admit confluent case where there are no constraints (\*). Without reducing the generality, we will give the following definitions with approximant sequences  $\{f_k\}_{k \in \mathbb{N}}$ .

**Definition 2.** *A branched continued fraction*

$$\langle \langle \{a_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}}, \{b_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}_0} \rangle, \{f_k\}_{k \in \mathbb{N}} \rangle \tag{11}$$

converges if, at most, a finite number of its approximants don't make sense and if the limit of its sequence of approximants

$$\lim_{n \rightarrow \infty} f_n$$

exists and is finite.

**Definition 3.** *A branched continued fraction (11) converges absolutely if its sequence of approximants such that*

$$\sum_{n=1}^{\infty} |f_{n+1} - f_n| < +\infty.$$

**Definition 4.** *A branched continued fraction*

$$\langle \langle \{\widehat{a}_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}}, \{\widehat{b}_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}_0} \rangle, \{\widehat{f}_k\}_{k \in \mathbb{N}} \rangle$$

is a majorant of a branched continued fraction (11) if there exist a natural number  $n_0$  and a positive constant  $M$  such that for  $n \geq n_0$  and  $k \geq 1$  the following relation holds

$$|f_{n+k} - f_n| \leq M|\widehat{f}_{n+k} - \widehat{f}_n|.$$

Again, without reducing the generality, let us put  $b_0 = 0$ .

**Definition 5.** A convergence set  $\Omega$  is a set  $\Omega \neq \emptyset$  and  $\Omega \subseteq \mathbb{C} \times \mathbb{C}$  such that: if  $\langle a_{i(k)}, b_{i(k)} \rangle \in \Omega$  for all  $i(k) \in \mathcal{I}_k, k \geq 1$ , then a branched continued fraction

$$\langle \langle \{a_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}}, \{b_{i(k)}\}_{i(k) \in \mathcal{I}_k, k \in \mathbb{N}} \rangle, \{f_k\}_{k \in \mathbb{N}} \rangle \tag{12}$$

converges.

**Definition 6.** A uniform convergence set  $\Omega$  is a convergence set to which there corresponds a sequence  $\{\varepsilon_n\}_{n \in \mathbb{N}}$  of positive numbers depending only on  $\Omega$  and converging to 0 such that

$$|f_{n+k} - f_n| \leq \varepsilon_n, \quad n \geq 1, k \geq 1,$$

for every branched continued fraction (12) with all  $\langle a_{i(k)}, b_{i(k)} \rangle \in \Omega$ .

Reasoning similarly as in the proof of Theorem 2 [24], we will prove the following result:

**Theorem 2.** Suppose that  $m_{0,k}, k \geq 1$ , are constants such that

$$0 < m_{0,k} \leq 1, \quad k \geq 1. \tag{13}$$

Then, the following statements hold:

(A) The branched continued fraction

$$1 - z_{1,0} + \frac{m_{0,1}z_{0,1}}{1 + \frac{m_{0,2}(1 - m_{0,1})z_{0,2}}{1 - (1 - m_{0,2})z_{1,2} + \frac{m_{0,3}(1 - m_{0,2})z_{0,3}}{1 + \frac{m_{0,4}(1 - m_{0,3})z_{0,4}}{1 - (1 - m_{0,4})z_{1,4} + \frac{m_{0,5}(1 - m_{0,4})z_{0,5}}{1 + \dots}}}}}} \tag{14}$$

converges absolutely and uniformly for

$$|z_{1,2k}| \leq \frac{1}{2}, \quad |z_{0,2k+1}| \leq \frac{1}{2} \quad \text{and} \quad |z_{0,2k+2}| \leq 1, \quad k \geq 0; \tag{15}$$

(B) The values of the branched continued fraction (14) and of its approximants are in the closed disk

$$|w - 1| \leq 1. \tag{16}$$

**Proof.** Let us show that the majorant of branched continued fraction (14) is

$$\frac{1}{2} - \frac{\frac{m_{0,1}}{2}}{1 - \frac{m_{0,2}(1 - m_{0,1})}{1 - \frac{(1 - m_{0,2})}{2} - \frac{m_{0,3}(1 - m_{0,2})}{1 - \frac{m_{0,4}(1 - m_{0,3})}{1 - \frac{(1 - m_{0,4})}{2} - \frac{m_{0,5}(1 - m_{0,4})}{1 - \dots}}}}. \tag{17}$$

We set

$$Q_n^{(n)} = \widehat{Q}_n^{(n)} = 1, \quad n \geq 1, \tag{18}$$

and

$$Q_k^{(n)} = 1 - (1 - m_{0,k})\delta(k) + \frac{m_{0,k+1}(1 - m_{0,k})z_{0,k+1}}{1 + \dots - (1 - m_{0,n-2})\delta(n - 2) + \frac{m_{0,n-1}(1 - m_{0,n-2})z_{0,n-1}}{1 - (1 - m_{0,n-1})\delta(n - 1) + m_{0,n}(1 - m_{0,n-1})z_{0,n}}},$$

$$\widehat{Q}_k^{(n)} = 1 - (1 - m_{0,k})\widehat{\delta}(k) - \frac{m_{0,k+1}(1 - m_{0,k})\frac{1 + 2\widehat{\delta}(k + 1)}{2}}{1 - \dots - (1 - m_{0,n-2})\widehat{\delta}(n - 2) - \frac{m_{0,n-1}(1 - m_{0,n-2})\frac{1 + 2\widehat{\delta}(n - 1)}{2}}{1 - (1 - m_{0,n-1})\widehat{\delta}(n - 1) - m_{0,n}(1 - m_{0,n-1})\frac{1 + 2\widehat{\delta}(n)}{2}}},$$

where,  $n \geq 2, 1 \leq k \leq n - 1$ , and, for  $x \in \mathbb{N}$ ,

$$\delta(x) = \begin{cases} z_{1,x}, & \text{if } x \text{ is even,} \\ 0, & \text{if } x \text{ is odd,} \end{cases} \quad \widehat{\delta}(x) = \begin{cases} 1/2, & \text{if } x \text{ is even,} \\ 0, & \text{if } x \text{ is odd.} \end{cases} \tag{19}$$

Then

$$Q_k^{(n)} = 1 - (1 - m_{0,k})\delta(k) + \frac{m_{0,k+1}(1 - m_{0,k})z_{0,k+1}}{Q_{k+1}^{(n)}}, \tag{20}$$

$$\widehat{Q}_k^{(n)} = 1 - (1 - m_{0,k})\widehat{\delta}(k) - \frac{m_{0,k+1}(1 - m_{0,k})(1 + 2\widehat{\delta}(k + 1))}{2\widehat{Q}_{k+1}^{(n)}}, \tag{21}$$

where  $n \geq 2, 1 \leq k \leq n - 1$ .

Thus, for each  $n \geq 1$  we write the  $n$ th approximants of (14) and (17) as

$$f_n = 1 - z_{1,0} + \frac{m_{0,1}z_{0,1}}{Q_1^{(n)}}, \quad \widehat{f}_n = \frac{1}{2} - \frac{m_{0,1}}{2\widehat{Q}_1^{(n)}},$$

respectively.

Using relations (13), (15), and (18)–(21), for an arbitrary  $n \geq 1$  by induction on  $k, 1 \leq k \leq n$ , we show that

$$|Q_k^{(n)}| \geq \widehat{Q}_k^{(n)} \geq m_{0,k}. \quad (22)$$

For  $k = n$  inequalities (22) are obvious (see, (13) and (18)). By induction hypothesis that (22) hold for  $k = r + 1$ , where  $r + 1 \leq n$ , we prove (22) for  $k = r$ . Indeed, use of inequalities (15) and (19)–(21) lead to

$$\begin{aligned} |Q_r^{(n)}| &= \left| 1 - (1 - m_{0,r})\delta(r) + \frac{m_{0,r+1}(1 - m_{0,r})z_{0,r+1}}{Q_{r+1}^{(n)}} \right| \\ &\geq 1 - (1 - m_{0,r})|\delta(r)| - \frac{m_{0,r+1}(1 - m_{0,r})|z_{0,r+1}|}{|Q_{r+1}^{(n)}|} \\ &\geq 1 - (1 - m_{0,r})\widehat{\delta}(r) - \frac{m_{0,r+1}(1 - m_{0,r})(1 + 2\widehat{\delta}(r + 1))}{2\widehat{Q}_{r+1}^{(n)}} \\ &= \widehat{Q}_r^{(n)}. \end{aligned}$$

It is easy to see that

$$\widehat{\delta}(r) + \frac{1 + 2\widehat{\delta}(r + 1)}{2} = 1.$$

By virtue of estimates (22),  $\widehat{Q}_{r+1}^{(n)} \neq 0$ . Therefore, replacing  $m_{0,r+1}$  by  $\widehat{Q}_{r+1}^{(n)}$ , inequalities (22) are obtained for  $k = r$ .

From (13) and (22) it follows that  $Q_k^{(n)} \neq 0$  and  $\widehat{Q}_k^{(n)} > 0$  for all  $n \geq 1$  and  $1 \leq k \leq n$ . Applying the method suggested in ([25], p. 28) and the relations (18) and (20) we find the formula for the difference of two approximants of the branched continued fraction (14). For  $n \geq 1$  and  $k \geq 1$  on the first step we obtain

$$\begin{aligned} f_{n+k} - f_n &= 1 - z_{1,0} + \frac{m_{0,1}z_{0,1}}{Q_1^{(n+k)}} - \left( 1 - z_{1,0} + \frac{m_{0,1}z_{0,1}}{Q_1^{(n)}} \right) \\ &= -\frac{m_{0,1}z_{0,1}}{Q_1^{(n+k)}Q_1^{(n)}} (Q_1^{(n+k)} - Q_1^{(n)}). \end{aligned}$$

Let  $r$  be arbitrary integer number, moreover  $1 \leq r \leq n - 1$ ,  $n \geq 2$ . Then for  $n \geq 2$  and  $k \geq 1$  we have

$$\begin{aligned} Q_r^{(n+k)} - Q_r^{(n)} &= 1 - (1 - m_{0,r})\delta(r) + \frac{m_{0,r+1}(1 - m_{0,r})z_{0,r+1}}{Q_{r+1}^{(n+k)}} \\ &\quad - \left( 1 - (1 - m_{0,r})\delta(r) + \frac{m_{0,r+1}(1 - m_{0,r})z_{0,r+1}}{Q_{r+1}^{(n)}} \right) \\ &= -\frac{m_{0,r+1}(1 - m_{0,r})z_{0,r+1}}{Q_{r+1}^{(n+k)}Q_{r+1}^{(n)}} (Q_{r+1}^{(n+k)} - Q_{r+1}^{(n)}). \end{aligned} \quad (23)$$

Applying recurrence relation (23) and taking into account that

$$Q_n^{(n+k)} - Q_n^{(n)} = -(1 - m_{0,n})\delta(n) + \frac{m_{0,n+1}(1 - m_{0,n})z_{0,n+1}}{Q_{n+1}^{(n+k)}},$$

after  $n$ th step we obtain

$$f_{n+k} - f_n = (-1)^{n+1} \left( (1 - m_{0,n})\delta(n) - \frac{m_{0,n+1}(1 - m_{0,n})z_{0,n+1}}{Q_{n+1}^{(n+k)}} \right) \prod_{r=1}^n \frac{m_{0,r}(1 - m_{0,r-1})z_{0,r}}{Q_r^{(n+k)}Q_r^{(n)}},$$

where  $m_{0,0} = 0$ .

Using (19) and (22), we get the following

$$\begin{aligned} |f_{n+k} - f_n| &\leq \prod_{r=1}^n \frac{m_{0,r}(1 - m_{0,r-1})|z_{0,r}|}{|Q_r^{(n+k)}||Q_r^{(n)}|} \\ &\quad \times \left( (1 - m_{0,n})|\delta(n)| + \frac{m_{0,n+1}(1 - m_{0,n})|z_{0,n+1}|}{|Q_{n+1}^{(n+k)}|} \right) \\ &\leq \frac{1}{2^n} \prod_{r=1}^n \frac{m_{0,r}(1 - m_{0,r-1})(1 + 2\widehat{\delta}_{0,r})}{\widehat{Q}_r^{(n+k)}\widehat{Q}_r^{(n)}} \\ &\quad \times \left( (1 - m_{0,n})\widehat{\delta}(n) + \frac{m_{0,n+1}(1 - m_{0,n})(1 + 2\widehat{\delta}_{0,n+1})}{2\widehat{Q}_{n+1}^{(n+k)}} \right) \\ &= -(\widehat{f}_{n+k} - \widehat{f}_n), \end{aligned}$$

where  $n \geq 1, k \geq 1$ , and  $m_{0,0} = 0$ . Thus,

$$|f_{n+k} - f_n| \leq \widehat{f}_n - \widehat{f}_{n+k}, \quad n \geq 1, k \geq 1. \quad (24)$$

It follows that the sequence  $\{\widehat{f}_n\}_{n \in \mathbb{N}}$  is monotonically decreasing and due to inequalities (22) is bounded from below. Indeed, for  $n \geq 1$  we have

$$\begin{aligned} \widehat{f}_n &= \frac{1}{2} - \frac{m_{0,1}}{2\widehat{Q}_1^{(n)}} \\ &\geq 0. \end{aligned}$$

Therefore, there exists a limit

$$\widehat{f} = \lim_{n \rightarrow \infty} \widehat{f}_n.$$

Now, using the relation (24), we obtain for  $k \geq 1$

$$\begin{aligned} \sum_{n=1}^k |f_{n+1} - f_n| &\leq - \sum_{n=1}^k (\widehat{f}_{n+1} - \widehat{f}_n) \\ &= \frac{1 - m_{0,1}}{2} - \widehat{f}_{k+1}. \end{aligned}$$

If  $k \rightarrow \infty$  it follows that the branched continued fraction (14) converges absolutely and uniformly for  $z_{1,2k}$  and  $z_{0,k}$ ,  $k \geq 1$ , which satisfies the inequalities (15). This proves (A).

Finally, by (22) for any  $n \geq 1$  we obtain

$$\begin{aligned} |f_n - 1| &\leq |z_{1,0}| + \frac{m_{0,1}|z_{0,1}|}{|Q_1^{(n)}|} \\ &\leq \frac{1}{2} + \frac{1}{2} \\ &= 1, \end{aligned}$$

which proves (B).  $\square$

Note that Theorem 2 is an analogue to Theorem 11.1 in [27]. Moreover, it can be proved in another way, using a generalization of the Sleshinsky-Pringsheim criterion ([28], Proposition 1).

We will also need the convergence continuation theorem, which follows from Theorem 2.17 [25] (see also ([27], Theorem 24.2)).

**Theorem 3** (Convergence Continuation Theorem). *Suppose that  $\{f_n(\mathbf{z})\}_{n \in \mathbb{N}}$  is a sequence of functions holomorphic in the domain  $\Omega$ ,  $\Omega \subset \mathbb{C}^2$ , uniformly bounded on every compact subset of  $\Omega$ . Suppose that this sequence converges at each point of the set  $\Theta$ ,  $\Theta \subset \Omega$ , which is the real neighborhood of the point  $\mathbf{z}^0$  in  $\Omega$ . Then the sequence  $\{f_n(\mathbf{z})\}_{n \in \mathbb{N}}$  converges uniformly on every compact subset of the domain  $\Omega$  to a function holomorphic in this domain.*

Next, we recall the necessary concepts of the PC method in the theory of branched continued fractions, which will be used to establish the analytical continuation of the function (see, [29,30]).

Let

$$L(\mathbf{z}) = \sum_{p,q=0}^{+\infty} a_{p,q} z_1^p z_2^q,$$

where  $a_{p,q} \in \mathbb{C}$ ,  $p \geq 0$ ,  $q \geq 0$ ,  $\mathbf{z} \in \mathbb{C}^2$ , be a formal double power series at  $\mathbf{z} = \mathbf{0}$ . Let  $f(\mathbf{z})$  be function holomorphic in a neighbourhood of the origin  $\mathbf{z} = \mathbf{0}$ . Let the mapping  $\Lambda : f(\mathbf{z}) \rightarrow \Lambda(f)$  associate with  $f(\mathbf{z})$  its Taylor expansion in a neighbourhood of the origin.

**Definition 7.** *A sequence  $\{f_n(\mathbf{z})\}_{n \in \mathbb{N}}$  of functions holomorphic at the origin corresponds to a formal double power series  $L(\mathbf{z})$  at  $\mathbf{z} = \mathbf{0}$  if*

$$\lim_{n \rightarrow +\infty} \lambda(L - \Lambda(f_n)) = +\infty,$$

where  $\lambda$  is the function defined as follows:  $\lambda : \mathbb{L} \rightarrow \mathbb{Z}_{\geq 0} \cup \{+\infty\}$ ; if  $L(\mathbf{z}) \equiv 0$  then  $\lambda(L) = +\infty$ ; if  $L(\mathbf{z}) \not\equiv 0$  then  $\lambda(L) = n$ , where  $n$  is the smallest degree of homogeneous terms for which  $a_{p,q} \neq 0$ , that is  $n = p + q$ .

If  $\{f_n(\mathbf{z})\}_{n \in \mathbb{N}}$  corresponds at  $\mathbf{z} = \mathbf{0}$  to a formal double power series  $L(\mathbf{z})$ , then the order of correspondence of  $f_n(\mathbf{z})$  is defined to be

$$v_n = \lambda(L - \Lambda(f_n)).$$

By the definition of  $\lambda$ , the series  $L(\mathbf{z})$  and  $\Lambda(f_n)$  agree for all homogeneous terms up to and including degree  $(v_n - 1)$ .

**Definition 8.** *A branched continued fraction whose elements are polynomials in the space  $\mathbb{C}^2$  corresponds to a formal double power series  $L(\mathbf{z})$  at  $\mathbf{z} = \mathbf{0}$  if its sequence of approximants corresponds to  $L(\mathbf{z})$ .*

Finally, we present the well-known Weierstrass' theorem ([31], p. 23) and the principle of analytic continuation ([32], p. 39).

**Theorem 4** (Weierstrass' Theorem). *Suppose that  $\{f_n(\mathbf{z})\}_{n \in \mathbb{N}}$  is a sequence of holomorphic functions in a domain  $\Omega$ ,  $\Omega \subset \mathbb{C}^2$ , converges to a function  $f(\mathbf{z})$  uniformly on each compact subset in the domain  $\Omega$ . Then  $f(\mathbf{z})$  is a holomorphic in  $\Omega$ , and for any  $p \geq 0$ ,  $q \geq 0$ ,*

$$\frac{\partial^{p+q} f_n(\mathbf{z})}{\partial z_1^p \partial z_2^q} \rightarrow \frac{\partial^{p+q} f(\mathbf{z})}{\partial z_1^p \partial z_2^q} \quad \text{as } n \rightarrow +\infty$$

on each compact subset in the domain  $\Omega$ .

**Theorem 5** (Principle of Analytic Continuation). *Suppose that  $f_1(\mathbf{z})$  and  $f_2(\mathbf{z})$  are functions holomorphic in the domains  $\Omega_1$ ,  $\Omega_1 \subset \mathbb{C}^2$ , and  $\Omega_2$ ,  $\Omega_2 \subset \mathbb{C}^2$ , respectively, and suppose that  $\Omega_1 \cap \Omega_2$  is a domain. Next, suppose that in the real neighborhood of the point  $\mathbf{z}^0$ ,  $\mathbf{z}^0 \in \Omega_1 \cap \Omega_2$ , the functions  $f_1(\mathbf{z})$  and  $f_2(\mathbf{z})$  coincide. Then these functions are an analytic continuation of one another, i.e., there is a unique function  $f(\mathbf{z})$  that is holomorphic in  $\Omega_1 \cup \Omega_2$  and coincides with  $f_1(\mathbf{z})$  in  $\Omega_1$  and with  $f_2(\mathbf{z})$  in  $\Omega_2$ .*

### 3. Convergence and Analytical Continuation

**Proof of Theorem 1.** We prove (A). Let

$$Q_n^{(n)}(\mathbf{z}) = 1, \quad n \geq 1, \quad (25)$$

and

$$Q_{2k-1}^{(2n)}(\mathbf{z}) = 1 + \frac{h_{2k}z_2}{1 - z_1 + \frac{h_{2k+1}z_2}{1 + \dots - z_1 + \frac{h_{2n-1}z_2}{1 + h_{2n}z_2}}},$$

$$Q_{2k-2}^{(2n)}(\mathbf{z}) = 1 - z_1 + \frac{h_{2k-1}z_2}{1 + \frac{h_{2k}z_2}{1 + \dots - z_1 + \frac{h_{2n-1}z_2}{1 + h_{2n}z_2}}},$$

$$Q_{2k-1}^{(2n+1)}(\mathbf{z}) = 1 + \frac{h_{2k}z_2}{1 - z_1 + \frac{h_{2k+1}z_2}{1 + \dots + \frac{h_{2n}z_2}{1 - z_1 + h_{2n+1}z_2}}},$$

$$Q_{2k}^{(2n+1)}(\mathbf{z}) = 1 - z_1 + \frac{h_{2k+1}z_2}{1 + \frac{h_{2k+2}z_2}{1 + \dots + \frac{h_{2n}z_2}{1 - z_1 + h_{2n+1}z_2}}},$$

where  $n \geq 1, 1 \leq k \leq n$ . Then the following relations hold

$$Q_{2k-1}^{(2n)}(\mathbf{z}) = 1 + \frac{h_{2k}z_2}{Q_{2k}^{(2n)}(\mathbf{z})}, \quad Q_{2k-2}^{(2n)}(\mathbf{z}) = 1 - z_1 + \frac{h_{2k-1}z_2}{Q_{2k-1}^{(2n)}(\mathbf{z})}, \quad (26)$$

and

$$Q_{2k-1}^{(2n+1)}(\mathbf{z}) = 1 + \frac{h_{2k}z_2}{Q_{2k}^{(2n+1)}(\mathbf{z})}, \quad Q_{2k}^{(2n+1)}(\mathbf{z}) = 1 - z_1 + \frac{h_{2k+1}z_2}{Q_{2k+1}^{(2n+1)}(\mathbf{z})}, \quad (27)$$

where  $n \geq 1, 1 \leq k \leq n$ , and, thus, for each  $n \geq 1$  we write the  $n$ th approximants of (5) as

$$f_n(\mathbf{z}) = 1 - z_1 + \frac{h_1z_2}{Q_1^{(n)}(\mathbf{z})}. \quad (28)$$

Let  $n$  be an arbitrary natural number,  $\arg(z_2) = \alpha$ , and  $\mathbf{z}$  be an arbitrary fixed point from (7). By induction on  $k, 1 \leq k \leq n$ , we show the following

$$\operatorname{Re}(Q_{2k-1}^{(2n)}(\mathbf{z})e^{-i\alpha/2}) > \frac{\cos(\alpha/2)}{2} \geq c > 0 \quad (29)$$

and

$$\operatorname{Re}(Q_{2k-1}^{(2n+1)}(\mathbf{z})e^{-i\alpha/2}) > \frac{\cos(\alpha/2)}{2} \geq c > 0. \quad (30)$$

By virtue of an arbitrary fixed point  $\mathbf{z}, \mathbf{z} \in \Omega_{p,q}$  it follows that for its arbitrary neighborhood, there exists  $\delta, 0 < \delta \leq \pi/2$ , such that  $|\alpha/2| \leq \pi/2 - \delta$  and, thus,

$$\cos(\alpha/2) \geq \cos(\pi/2 - \delta) = \sin(\delta) = 2c > 0.$$

Let us prove the inequalities (29). From (26) we have

$$Q_{2k-1}^{(2n)}(\mathbf{z})e^{-i\alpha/2} = e^{-i\alpha/2} + \frac{h_{2k}z_2e^{-i\alpha}}{Q_{2k}^{(2n)}(\mathbf{z})e^{-i\alpha/2}}, \quad (31)$$

where  $1 \leq k \leq n$ , and

$$Q_{2k}^{(2n)}(\mathbf{z})e^{-i\alpha/2} = e^{-i\alpha/2} - \frac{z_1e^{-2i\alpha/2}}{e^{-i\alpha/2}} + \frac{h_{2k+1}z_2e^{-i\alpha}}{Q_{2k+1}^{(2n)}(\mathbf{z})e^{-i\alpha/2}}, \quad (32)$$

where  $1 \leq k \leq n-1$ . Using (3), (7), (25), (31), and Corollary 2 in [29], for  $k = n$  we obtain

$$\begin{aligned} \operatorname{Re}(Q_{2n-1}^{(2n)}(\mathbf{z})e^{-i\alpha/2}) &\geq \cos(\alpha/2) - \frac{(|h_{2n}| - \operatorname{Re}(h_{2n}))|z_2|}{2 \operatorname{Re}(Q_{2n}^{(2n)}(\mathbf{z})e^{-i\alpha/2})} \\ &> \cos(\alpha/2) - \frac{2pq}{2 \cos(\alpha/2)} \frac{1 + \cos(\alpha)}{4p} \\ &> \cos(\alpha/2) - \frac{\cos(\alpha/2)}{2} \\ &= \frac{\cos(\alpha/2)}{2}. \end{aligned}$$

Let the inequalities (29) hold for  $k = r+1$  such that  $r+1 \leq n$ . Then, by (3), (7), Corollary 2 in [29], and the induction hypothesis, from (31) and (32) for  $k = r$  we have

$$\begin{aligned} \operatorname{Re}(Q_{2r}^{(2n)}(\mathbf{z})e^{-i\alpha/2}) &\geq \cos(\alpha/2) - \operatorname{Re}(z_1e^{-i\alpha/2}) - \frac{(|h_{2r+1}| - \operatorname{Re}(h_{2r+1}))|z_2|}{2 \operatorname{Re}(Q_{2r+1}^{(2n)}(\mathbf{z})e^{-i\alpha/2})} \\ &> \cos(\alpha/2) - (1-q)\cos(\alpha/2) - \frac{2pq}{2 \cos(\alpha/2)} \frac{1 + \cos(\alpha)}{4p} \\ &= \cos(\alpha/2) - (1-q)\cos(\alpha/2) - \frac{q \cos(\alpha/2)}{2} \\ &= \frac{q \cos(\alpha/2)}{2} \end{aligned}$$

and

$$\begin{aligned} \operatorname{Re}(Q_{2r-1}^{(2n)}(\mathbf{z})e^{-i\alpha/2}) &\geq \cos(\alpha/2) - \frac{(|h_{2r}| - \operatorname{Re}(h_{2r}))|z_2|}{2 \operatorname{Re}(Q_{2r}^{(2n)}(\mathbf{z})e^{-i\alpha/2})} \\ &> \cos(\alpha/2) - \frac{2pq}{2q \cos(\alpha/2)} \frac{1 + \cos(\alpha)}{4p} \\ &= \cos(\alpha/2) - \frac{\cos(\alpha/2)}{2} \\ &= \frac{\cos(\alpha/2)}{2}, \end{aligned}$$

respectively.

Similarly, we obtain the inequalities (30).

Thus,  $Q_1^{(n)}(\mathbf{z}) \neq 0$  for  $n \geq 1$  and for all  $\mathbf{z} \in \Omega_{p,q}$ , i.e., that each approximant (28) is a holomorphic function in the domain (7).

Let  $\Gamma$  be an arbitrary compact subset of (7). Then there exists an open bi-disk

$$\Theta_r = \{\mathbf{z} \in \mathbb{C}^2 : |z_1| < r, |z_2| < r\}$$

of radius  $r$ ,  $r > 0$ , such that  $\Gamma \subset \Theta_r$ . Moreover, for any  $n \geq 1$  and  $\mathbf{z} \in \Omega_{p,q} \cap \Theta_r$  from (25), (29), and (30) we have

$$\begin{aligned} |f_n(\mathbf{z})| &\leq 1 + r + \frac{|h_1|r}{\operatorname{Re}(Q_1^{(n)}(\mathbf{z})e^{-i\alpha/2})} \\ &< 1 + r + \frac{2hr}{\cos(\alpha/2)} \\ &= C(\Omega_{p,q} \cap \Theta_r), \end{aligned}$$

where  $h$  is defined in (9), i.e., the sequence  $\{f_n(\mathbf{z})\}_{n \in \mathbb{N}}$  is uniformly bounded on every compact subset of  $\Omega_{p,q}$ .

It is obvious that for every  $l$  such that

$$0 < l < \min \left\{ \frac{1-d}{2}, \frac{d(1-d)}{2h}, 1-q, \frac{1}{4p} \right\}, \quad (33)$$

where  $p$  and  $q$  are defined in (4),  $h$  and  $d$  are defined in (9), the domain

$$Y_l = \{\mathbf{z} \in \mathbb{R}^2 : 0 < z_1 < l, 0 < z_2 < l\}$$

contained in (7), in particular  $Y_{l/2} \subset \Omega_{p,q}$ . Using (9) and (33), for any  $\mathbf{z} \in Y_l$ ,  $Y_l \subset \Omega_{p,q}$ , we obtain that

$$0 < z_1 < \frac{1-d}{2}$$

and for any  $k \geq 1$

$$\begin{aligned} |h_k z_2| &< \frac{hd(1-d)}{2h} \\ &= \frac{d(1-d)}{2}, \end{aligned}$$

i.e., the elements of (5) satisfy the Theorem 1, with  $m_{0,k} = d$ ,  $k \geq 1$ . It means that branched continued fraction (5) converges for all  $\mathbf{z} \in Y_l$ ,  $Y_l \subset \Omega_{p,q}$ . Therefore, according to the Theorem 3, the convergence of (5) is uniform on compact subsets of the domain (7).

From (A) of Theorem 1, with  $m_{0,k} = d$ ,  $k \geq 1$ ,  $0 < d < 1$ , it follows that the branched continued fraction (5) converges for all  $\mathbf{z} \in \Omega^{d,h}$ , where  $\Omega^{d,h}$  is defined by (8), and from (B) of the same theorem it follows that all approximants of (5) lie in the closed disk (16) if  $\mathbf{z} \in \Omega^{d,h}$ . Hence, by Theorem 3, the branched continued fraction (5) converges uniformly on compact subsets of the domain (8). This is a complete proof of (A).

Now, we prove (B). Let

$$\begin{aligned} G_{2n}^{(2n)}(\mathbf{z}) &= \frac{F_2(\alpha + n, \beta, \beta' + n; \beta, \gamma' + 2n; z_1, -z_2)}{F_2(\alpha + n + 1, \beta, \beta' + n; \beta, \gamma' + 2n + 1; z_1, -z_2)}, \\ G_{2n+1}^{(2n+1)}(\mathbf{z}) &= \frac{F_2(\alpha + n + 1, \beta, \beta' + n; \beta, \gamma' + 2n + 1; z_1, -z_2)}{F_2(\alpha + n + 1, \beta, \beta' + n + 1; \beta, \gamma' + 2n + 2; z_1, -z_2)}. \end{aligned}$$

and

$$\begin{aligned}
 G_{2k-1}^{(2n)}(\mathbf{z}) &= 1 + \frac{h_{2k}z_2}{1 - z_1 + \frac{h_{2k+1}z_2}{1 + \dots - z_1 + \frac{h_{2n-1}z_2}{G_{2n}^{(2n)}(\mathbf{z})}}}, \\
 G_{2k-2}^{(2n)}(\mathbf{z}) &= 1 - z_1 + \frac{h_{2k-1}z_2}{1 + \frac{h_{2k}z_2}{1 + \dots - z_1 + \frac{h_{2n-1}z_2}{G_{2n}^{(2n)}(\mathbf{z})}}}, \\
 G_{2k-1}^{(2n+1)}(\mathbf{z}) &= 1 + \frac{h_{2k}z_2}{1 - z_1 + \frac{h_{2k+1}z_2}{1 + \dots + \frac{h_{2n}z_2}{G_{2n+1}^{(2n+1)}(\mathbf{z})}}}, \\
 G_{2k}^{(2n+1)}(\mathbf{z}) &= 1 - z_1 + \frac{h_{2k+1}z_2}{1 + \frac{h_{2k+2}z_2}{1 + \dots + \frac{h_{2n}z_2}{G_{2n+1}^{(2n+1)}(\mathbf{z})}}},
 \end{aligned}$$

where  $n \geq 1$  and  $1 \leq k \leq n$ . Then

$$G_{2k-1}^{(2n)}(\mathbf{z}) = 1 + \frac{h_{2k}z_2}{G_{2k}^{(2n)}(\mathbf{z})}, \quad G_{2k-2}^{(2n)}(\mathbf{z}) = 1 - z_1 + \frac{h_{2k-1}z_2}{G_{2k-1}^{(2n)}(\mathbf{z})}, \tag{34}$$

and

$$G_{2k-1}^{(2n+1)}(\mathbf{z}) = 1 + \frac{h_{2k}z_2}{G_{2k}^{(2n+1)}(\mathbf{z})}, \quad G_{2k}^{(2n+1)}(\mathbf{z}) = 1 - z_1 + \frac{h_{2k+1}z_2}{G_{2k+1}^{(2n+1)}(\mathbf{z})}, \tag{35}$$

where  $n \geq 1, 1 \leq k \leq n$ .

It follows (see also [22]) that for each  $n \geq 1$

$$\begin{aligned}
 \frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} &= 1 - z_1 + \frac{h_1z_2}{1 + \frac{h_2z_2}{1 + \dots - z_1 + \frac{h_{2n-1}z_2}{1 + \frac{h_{2n}z_2}{G_{2n}^{(2n)}(\mathbf{z})}}}} \\
 &= 1 - z_1 + \frac{h_1z_2}{G_1^{(2n)}(\mathbf{z})}
 \end{aligned}$$

and

$$\begin{aligned}
 \frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} &= 1 - z_1 + \frac{h_1z_2}{1 + \frac{h_2z_2}{1 + \dots + \frac{h_{2n}z_2}{1 - z_1 + \frac{h_{2n+1}z_2}{G_{2n+1}^{(2n+1)}(\mathbf{z})}}}} \\
 &= 1 - z_1 + \frac{h_1z_2}{G_1^{(2n+1)}(\mathbf{z})}.
 \end{aligned}$$

Since  $G_k^{(n)}(\mathbf{0}) = 1$  and  $Q_k^{(n)}(\mathbf{0}) = 1$  for any  $1 \leq k \leq n, n \geq 1$ , then the  $1/G_k^{(n)}(\mathbf{z})$  and  $1/Q_k^{(n)}(\mathbf{z})$  have formal Taylor expansions in a neighborhood of the origin. It is clear that  $G_k^{(n)}(\mathbf{z}) \neq 0$  and  $Q_k^{(n)}(\mathbf{z}) \neq 0$  for all  $1 \leq k \leq n$  and  $n \geq 1$ . Applying the method suggested in ([25], p. 28) and (25)–(27), (34), and (35), for each  $n \geq 1$  we have

$$\frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} - f_{2n-1}(\mathbf{z}) = -\frac{h_{2n} z_2^{2n}}{G_{2n}^{(2n)}(\mathbf{z})} \prod_{r=1}^{2n-1} \frac{h_r}{G_r^{(2n)}(\mathbf{z}) Q_r^{(2n-1)}(\mathbf{z})}$$

and

$$\frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} - f_{2n}(\mathbf{z}) = -z_2^{2n} \left( z_1 - \frac{h_{2n+1} z_2}{G_{2n+1}^{(2n+1)}(\mathbf{z})} \right) \prod_{r=1}^{2n} \frac{h_r}{G_r^{(2n+1)}(\mathbf{z}) Q_r^{(2n)}(\mathbf{z})}.$$

Hence, in a neighborhood of origin for any  $n \geq 1$ , we have

$$\Lambda \left( \frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} \right) - \Lambda(f_n) = \sum_{\substack{p+q \geq n+1 \\ p \geq 0, q \geq 0}} a_{p,q}^{(n)} z_1^p z_2^q,$$

where  $a_{p,q}^{(n)}, p \geq 0, q \geq 0, p + q \geq n + 1$ , are some coefficients. It follows that

$$\lambda \left( \Lambda \left( \frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} \right) - \Lambda(f_n) \right) = n + 1$$

tends monotonically to  $+\infty$  as  $n \rightarrow +\infty$ .

Thus, the branched continued fraction (5) corresponds at  $\mathbf{z} = \mathbf{0}$  to a formal double power series

$$\Lambda \left( \frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} \right).$$

Let  $\Delta$  be a neighborhood of the origin contained (6), and in which

$$\Lambda \left( \frac{F_2(\alpha, \beta, \beta'; \beta, \gamma'; z_1, -z_2)}{F_2(\alpha + 1, \beta, \beta'; \beta, \gamma' + 1; z_1, -z_2)} \right) = \sum_{p,q=0}^{+\infty} a_{p,q} z_1^p z_2^q. \tag{36}$$

From (A), it follows that the sequence of approximants of (5) converges uniformly on each compact subset of  $\Delta$  to function  $f(\mathbf{z})$  holomorphic in the domain  $\Delta$ . Then, by Theorem 3, for arbitrary  $p + q, p \geq 0, q \geq 0$ , we have

$$\frac{\partial^{p+q} f_n(\mathbf{z})}{\partial z_1^p \partial z_2^q} \rightarrow \frac{\partial^{p+q} f(\mathbf{z})}{\partial z_1^p \partial z_2^q} \text{ as } n \rightarrow +\infty$$

on each compact subset of  $\Delta$ . And now, due to the above proven, the expansion of each approximant of (5) into formal double power series and series (36) agree for all homogeneous terms up to and including degree  $n$ . Then, for any  $p + q, p \geq 0, q \geq 0$ , we have

$$\lim_{n \rightarrow +\infty} \left( \frac{\partial^{p+q} f_n(\mathbf{0})}{\partial z_1^p \partial z_2^q} \right) = \frac{\partial^{p+q} f}{\partial z_1^p \partial z_2^q}(\mathbf{0}) = p!q! a_{p,q}.$$

Hence,

$$f(\mathbf{z}) = \sum_{p,q=0}^{+\infty} \frac{1}{p!q!} \left( \frac{\partial^{p+q} f}{\partial z_1^p \partial z_2^q}(\mathbf{0}) \right) z_1^p z_2^q = \sum_{p,q=0}^{+\infty} a_{p,q} z_1^p z_2^q$$

for all  $\mathbf{z} \in \Delta$ .

Finally, (B) follows from Theorem 4.  $\square$

Note that in the same way the domains of the analytical continuation of the ratios of Horn’s hypergeometric functions  $H_4$ ,  $H_6$ , and  $H_7$  were obtained in the works [33–35], respectively. Another approach using the PF Method (see, [29]) is applied in [36].

**Corollary 1.** Suppose that  $\beta'$  and  $\gamma'$  are complex constants such that satisfy inequality (3), where

$$h_1 = \frac{\beta'}{\gamma'}, \quad h_{2k} = \frac{k(\gamma' - \beta' + k - 1)}{(\gamma' + 2k - 2)(\gamma' + 2k - 1)}, \quad \text{and} \quad h_{2k+1} = \frac{(\beta' + k)(\gamma' + k - 1)}{(\gamma' + 2k - 1)(\gamma' + 2k)}, \quad k \geq 1, \quad (37)$$

herewith  $\gamma' \notin \{0, -1, -2, \dots\}$ , and where  $p > 0$  and  $0 < q < 1$ . Then the branched continued fraction


$$\frac{1}{1 - z_1 + \frac{\frac{\beta'}{\gamma'} z_2}{1 + \frac{(\gamma' - \beta') z_2}{(\beta' + 1)\gamma'}}} \quad (38)$$

$$1 - z_1 + \frac{\frac{(\gamma' - \beta') z_2}{(\beta' + 1)\gamma'}}{1 - z_1 + \frac{(\gamma' + 1)(\gamma' + 2) z_2}{2(\gamma' - \beta' + 1)}}$$

$$1 + \frac{\frac{2(\gamma' - \beta' + 1) z_2}{(\gamma' + 2)(\gamma' + 3)}}{1 + \dots}$$

converges uniformly on every compact subset of the domain (6) to the function  $f(\mathbf{z})$  holomorphic in this domain, in addition, the function  $f(\mathbf{z})$  is an analytic continuation of the function  $F_2(1, \beta, \beta'; \beta, \gamma'; z_1, -z_2)$  in the domain  $\Omega_{p,q}^{h,l}$ .

Graphical illustrations of domains for variables  $z_1$  and  $z_2$  in (6) are shown in Figure 1a–c.



**Figure 1.** Domains for variables  $z_1$  and  $z_2$  in (6).

By using Theorem 1, we obtain the following result:

**Theorem 6.** Suppose that  $\alpha$ ,  $\beta'$ , and  $\gamma'$  are real constants such that

$$0 < h_k \leq h, \quad k \geq 1, \quad (39)$$

where  $h_k, k \geq 1$ , are given in (2) herewith  $\gamma' \notin \{0, -1, -2, \dots\}$ ,  $h$  is a positive number. Then the branched continued fraction (5) converges uniformly on every compact subset of the domain

$$\Phi_h = \left\{ \mathbf{z} \in \mathbb{C}^2 : z_1 \notin [1, +\infty), z_2 \notin \left( -\infty, -\frac{1}{8h} \right] \right\} \quad (40)$$

to the function  $f(\mathbf{z})$  holomorphic in this domain, in addition, the function  $f(\mathbf{z})$  is an analytic continuation of the function (10) in the domain  $\Phi_h$ .

**Proof.** If  $h_k > 0$  for all  $k \geq 1$ , then it is clear that inequality (3) is true for all  $p > 0$ . Let  $\Gamma$  be an arbitrary compact subset of the domain (40). Since  $0 < q < 1$  and  $0 < d < 1$ , the inclusions  $\Gamma \subseteq \Omega_{p,q}^{d,h} \subseteq \Phi_h$  hold for  $d = 1/2$  and certain fairly small  $p$  and  $q$  for which the set  $\Omega_{p,q}^{d,h}$  is the domain (6). Thus, this theorem is a direct corollary of Theorem 1.  $\square$

**Corollary 2.** Suppose that  $\alpha$ ,  $\beta'$ , and  $\gamma'$  are real constants such that satisfy inequality (39), where  $h_k, k \geq 1$ , are given in (37) herewith  $\gamma' \notin \{0, -1, -2, \dots\}$ ,  $h$  is a positive number. Then the branched continued fraction (38) converges uniformly on every compact subset of the domain (40) to the function  $f(\mathbf{z})$  holomorphic in this domain, in addition, the function  $f(\mathbf{z})$  is an analytic continuation of the function  $F_2(1, \beta, \beta'; \beta, \gamma'; \mathbf{z})$  in the domain  $\Phi_h$ .

Note that (40) is the Cartesian product of the plane cut along the real axis from 1 to  $+\infty$  and the plane cut along the real axis from  $-1/(8h)$  to  $-\infty$ , where  $h$  is a positive number satisfying (39).

By using Theorem 5, we directly obtain the following result from Theorem 2.2 in [23] and Theorem 6:

**Theorem 7.** Suppose that  $\alpha$ ,  $\beta'$ , and  $\gamma'$  are real constants such that satisfy inequality (39), where  $h_k, k \geq 1$ , are given in (2) herewith  $\gamma' \notin \{0, -1, -2, \dots\}$ ,  $h$  is a positive number. Then the branched continued fraction (1) converges uniformly on every compact subset of the domain

$$\Xi_h = \left\{ \mathbf{z} \in \mathbb{C}^2 : z_1 \notin [1, +\infty), z_2 \notin \left[ \frac{1}{4h}, +\infty \right) \right\} \quad (41)$$

to the function  $f(\mathbf{z})$  holomorphic in this domain, in addition, the function  $f(\mathbf{z})$  is an analytic continuation of the function on the left side of (1) in the domain  $\Xi_h$ .

**Corollary 3.** Suppose that  $\alpha$ ,  $\beta'$ , and  $\gamma'$  are real constants such that satisfy inequality (39), where  $h_k, k \geq 1$ , are given in (37) herewith  $\gamma' \notin \{0, -1, -2, \dots\}$ ,  $h$  is a positive number. Then the branched continued fraction

$$1 - z_1 - \frac{1}{1 - \frac{\frac{\beta'}{\gamma'} z_2}{(\gamma' - \beta') z_2}} = \frac{1}{1 - \frac{(\beta' + 1)\gamma'}{(\gamma' + 1)(\gamma' + 2)} z_2} = \frac{1}{1 - \frac{2(\gamma' - \beta' + 1)}{(\gamma' + 2)(\gamma' + 3)} z_2} = \dots$$

converges uniformly on every compact subset of the domain (41) to the function  $f(\mathbf{z})$  holomorphic in the this domain, in addition, the function  $f(\mathbf{z})$  is an analytic continuation of the function  $F_2(1, \beta, \beta'; \beta, \gamma'; \mathbf{z})$  in the domain  $\Xi_h$ .

#### 4. Examples

As an example, by Corollary 3 we get

$$\begin{aligned} \ln\left(1 + \frac{z_2}{1 + z_1}\right) &= z_2 F_2(1, \beta, 1; \beta, 2; -z_1, -z_2) \\ &= \frac{z_2}{1 + z_1 + \frac{\frac{1}{2}z_2}{1 + \frac{\frac{1}{6}z_2}{1 + z_1 + \frac{\frac{1}{3}z_2}{1 + \frac{\frac{1}{5}z_2}{1 + \dots}}}}}, \end{aligned} \quad (42)$$

where the branched continued fraction converges and represents a single-valued branch of the analytic function on the left side of (42) in the domain

$$\Xi_{1/2} = \left\{ \mathbf{z} \in \mathbb{C}^2 : z_1 \notin (-\infty, -1], z_2 \notin \left(-\infty, -\frac{1}{2}\right] \right\}.$$

One more example, by Corollary 3 we obtain

$$\begin{aligned} \arctan \sqrt{\frac{z_2}{1 + z_1}} &= \sqrt{z_2(1 + z_1)} F_2\left(1, \beta, \frac{1}{2}; \beta, \frac{3}{2}; -z_1, -z_2\right) \\ &= \frac{\sqrt{z_2(1 + z_1)}}{1 + z_1 + \frac{\frac{1}{3}z_2}{1 + \frac{\frac{4}{15}z_2}{1 + \frac{\frac{9}{35}z_2}{1 + \frac{\frac{16}{63}z_2}{1 + \dots}}}}}, \end{aligned} \quad (43)$$

where the branched continued fraction converges and represents a single-valued branch of the analytic function on the left side of (43) in the domain

$$\Xi_{1/3} = \left\{ \mathbf{z} \in \mathbb{C}^2 : z_1 \notin (-\infty, -1], z_2 \notin \left(-\infty, -\frac{3}{4}\right] \right\}.$$

#### 5. Conclusions

In this paper, we discussed the representation and extension of the analytic functions due branched continued fractions as a special family of functions. Our results are new symmetric domains of analytical extension of the Appell's hypergeometric function  $F_2$  with certain conditions on its real and complex parameters. In particular, we obtained the domain of analytical continuation, which is the Cartesian product of the plane cut along the real axis from 1 to  $+\infty$  and the plane cut along the real axis from  $-1/(4h)$  to  $-\infty$ , where and  $h$  is a positive number satisfying (39). However, the problem of establishing

the domains of the analytical extension of the Appell's hypergeometric functions  $F_2$  with arbitrary parameters remains open.

The results of the study of branched continued fraction expansions of the Appell's hypergeometric functions  $F_1$ ,  $F_3$  and  $F_4$  can be found in [37–40].

Further studies of branched continued fraction expansions consist in the use of new parabolic [41–43] and angular [44,45] domains of convergence of branched continued fractions. Other directions of research are truncation errors analysis [46–50] and computational stability [51–53]. Finally, taking into account the efficiency of approximation of functions by branched continued fractions [23,34,36] and the breadth of application of hypergeometric function, the applied direction of research is natural and intriguing.

**Funding:** This research received no external funding.

**Data Availability Statement:** Data are contained within the article.

**Acknowledgments:** The authors were partially supported by the Ministry of Education and Science of Ukraine, project registration number 0123U101791.

**Conflicts of Interest:** The author declare no conflicts of interest.

## References

- Appell, P. Sur les séries hyper géométriques de deux variables et sur des équations différentielles lineaires aux dérivées partielles. *CR Acad. Sci. Paris* **1880**, *90*, 296–298. 731–734.
- Appell, P. Sur les fonctions hyper géométriques de deux variables. *J. Math. Pures Appl.* **1982**, *8*, 173–216.
- Mazars, M. Canonical partition functions of freely jointed chains. *J. Phys. A Math. Gen.* **1998**, *31*, 1949–1964. [[CrossRef](#)]
- Hutchinson, T.P. Compound gamma bivariate distributions. *Metrika* **1981**, *28*, 263–271. [[CrossRef](#)]
- Bentalha, Z.E.A. Representation of the Coulomb matrix elements by means of Appell hypergeometric function  $F_2$ . *Math. Phys. Anal. Geom.* **2018**, *21*, 10. [[CrossRef](#)]
- Shpot, M.A. A massive Feynman integral and some reduction relations for Appell functions. *J. Math. Phys.* **2007**, *48*, 123512. [[CrossRef](#)]
- Kol, B.; Shir, R. The propagator seagull: General evaluation of a two loop diagram. *J. High Energy Phys.* **2019**, *2019*, 83. [[CrossRef](#)]
- Tarasov, V.F. Multipole matrix elements for DH-systems and their asymptotics. *Int. J. Mod. Phys. B* **1995**, *9*, 2699–2718. [[CrossRef](#)]
- Brychkov, Y.A.; Saad, N. On some formulas for the Appell function  $F_2(a, b, b'; c, c'; w; z)$ . *Integral Transforms Spec. Funct.* **2013**, *25*, 111–123. [[CrossRef](#)]
- Wang, X. Recursion formulas for Appell functions. *Integral Transforms Spec. Funct.* **2012**, *23*, 421–433. [[CrossRef](#)]
- Opps, S.B.; Saad, N.; Srivastava, H.M. Some reduction and transformation formulas for the Appell hypergeometric function  $F_2$ . *J. Math. Anal. Appl.* **2005**, *302*, 180–195. [[CrossRef](#)]
- Hahne, G.E. Analytic continuation of Appell's hypergeometric series  $F_2$  to the vicinity of the singular point  $x = 1, y = 1$ . *J. Math. Phys.* **1969**, *10*, 524–531. [[CrossRef](#)]
- Jaeger, J.C. A continuation formula for Appell's function  $F_2$ . *J. Lond. Math. Soc.* **1938**, *13*, 254. [[CrossRef](#)]
- Sud, K.; Wright, L.E. A new analytic continuation of Appell's hypergeometric series  $F_2$ . *J. Math. Phys.* **1976**, *17*, 1719–1721. [[CrossRef](#)]
- Mimachi, K. Integral representations of Appell's  $F_2, F_3$ , Horn's  $H_2$  and Olsson's  $F_p$  functions. *Kyushu J. Math.* **2020**, *74*, 1–13. [[CrossRef](#)]
- Slater, L.J. The integration of hypergeometric functions. *Math. Proc. Camb. Philos. Soc.* **1955**, *51*, 288–296. [[CrossRef](#)]
- Chelo, F.; López, J.L. The Appell's function  $F_2$  for large values of its variables. *Q. Appl. Math.* **2010**, *68*, 701–712. [[CrossRef](#)]
- Hang, P.-C.; Luo, M.-J. Asymptotics of Saran's hypergeometric function  $F_K$ . *J. Math. Anal. Appl.* **2025**, *541*, 128707. [[CrossRef](#)]
- Minton, B.M. A note on the asymptotic expansion of the second Appell function. *Math. Proc. Camb. Philos. Soc.* **1968**, *64*, 1055. [[CrossRef](#)]
- Luo, M.-J.; Raina, R.K. On certain results related to the hypergeometric function  $F_K$ . *J. Math. Anal. Appl.* **2021**, *504*, 125439. [[CrossRef](#)]
- Ananthanarayan, B.; Bera, S.; Friot, S.; Marichev, O.; Pathak, T. On the evaluation of the Appell  $F_2$  double hypergeometric function. *Comput. Phys. Commun.* **2023**, *284*, 108589. [[CrossRef](#)]
- Bodnar, D.I. Expansion of a ratio of hypergeometric functions of two variables in branching continued fractions. *J. Math. Sci.* **1993**, *64*, 1155–1158. [[CrossRef](#)]
- Antonova, T.; Cesarano, C.; Dmytryshyn, R.; Sharyn, S. An approximation to Appell's hypergeometric function  $F_2$  by branched continued fraction. *Dolomites Res. Notes Approx.* **2024**, *17*, 22–31. [[CrossRef](#)]
- Dmytryshyn, R.I. Convergence of multidimensional A- and J-fractions with independent variables. *Comput. Methods Funct. Theory* **2022**, *22*, 229–242. [[CrossRef](#)]
- Bodnar, D.I. *Branched Continued Fractions*; Naukova Dumka: Kyiv, Ukraine, 1986. (In Russian)

26. Antonova, T. On structure of branched continued fractions. *Carpathian Math. Publ.* **2024**, *16*, 391–400. [[CrossRef](#)]
27. Wall, H.S. *Analytic Theory of Continued Fractions*; D. Van Nostrand Co.: New York, NY, USA, 1948.
28. Antonova, T.M. On convergence of branched continued fraction expansions of Horn's hypergeometric function  $H_3$  ratios. *Carpathian Math. Publ.* **2021**, *13*, 642–650. [[CrossRef](#)]
29. Antonova, T.; Dmytryshyn, R.; Goran, V. On the analytic continuation of Lauricella-Saran hypergeometric function  $F_K(a_1, a_2, b_1, b_2; a_1, b_2, c_3; z)$ . *Mathematics* **2023**, *11*, 4487. [[CrossRef](#)]
30. Cuyt, A.A.M.; Petersen, V.; Verdonk, B.; Waadeland, H.; Jones, W.B. *Handbook of Continued Fractions for Special Functions*; Springer: Dordrecht, The Netherlands, 2008.
31. Shabat, B.V. *Introduce to Complex Analysis. Part II. Functions of Several Variables*; American Mathematical Society: Providence, RI, USA, 1992.
32. Vladimirov, V.S. *Methods of the Theory of Functions of Many Complex Variables*; The MIT Press: Cambridge, MA, USA, 1966.
33. Dmytryshyn, R.; Lutsiv, I.-A.; Dmytryshyn, M. On the analytic extension of the Horn's hypergeometric function  $H_4$ . *Carpathian Math. Publ.* **2024**, *16*, 32–39. [[CrossRef](#)]
34. Antonova, T.; Dmytryshyn, R.; Sharyn, S. Branched continued fraction representations of ratios of Horn's confluent function  $H_6$ . *Constr. Math. Anal.* **2023**, *6*, 22–37. [[CrossRef](#)]
35. Hladun, V.; Rusyn, R.; Dmytryshyn, M. On the analytic extension of three ratios of Horn's confluent hypergeometric function  $H_7$ . *Res. Math.* **2024**, *32*, 60–70. [[CrossRef](#)]
36. Antonova, T.; Dmytryshyn, R.; Sharyn, S. Generalized hypergeometric function  ${}_3F_2$  ratios and branched continued fraction expansions. *Axioms* **2021**, *10*, 310. [[CrossRef](#)]
37. Bodnar, D.I.; Manzii, O.S. Expansion of the ratio of Appel hypergeometric functions  $F_3$  into a branching continued fraction and its limit behavior. *J. Math. Sci.* **2001**, *107*, 3550–3554. .:1011977720316. [[CrossRef](#)]
38. Bodnar, D.I. Multidimensional C-fractions. *J. Math. Sci.* **1998**, *90*, 2352–2359. [[CrossRef](#)]
39. Hladun, V.R.; Hoyenko, N.P.; Manzij, O.S.; Ventyk, L. On convergence of function  $F_4(1, 2; 2, 2; z_1, z_2)$  expansion into a branched continued fraction. *Math. Model. Comput.* **2022**, *9*, 767–778. [[CrossRef](#)]
40. Manzii, O.S. On the approximation of an Appell hypergeometric function by a branched continued fraction. *J. Math. Sci.* **1998**, *90*, 2376–2380. [[CrossRef](#)]
41. Dmytryshyn, R.; Antonova, T.; Dmytryshyn, M. On the analytic extension of the Horn's confluent function  $H_6$  on domain in the space  $\mathbb{C}^2$ . *Constr. Math. Anal.* **2024**, *accept.*
42. Bodnar, D.I.; Bilanyk, I.B. Parabolic convergence regions of branched continued fractions of the special form. *Carpathian Math. Publ.* **2021**, *13*, 619–630. [[CrossRef](#)]
43. Bodnar, D.I.; Bilanyk, I.B. Two-dimensional generalization of the Thron-Jones theorem on the parabolic domains of convergence of continued fractions. *Ukr. Math. J.* **2023**, *74*, 1317–1333. [[CrossRef](#)]
44. Bodnar, D.I.; Bilanyk, I.B. Estimation of the rates of pointwise and uniform convergence of branched continued fractions with inequivalent variables. *J. Math. Sci.* **2022**, *265*, 423–437. [[CrossRef](#)]
45. Bodnar, D.I.; Bilanyk, I.B. On the convergence of branched continued fractions of a special form in angular domains. *J. Math. Sci.* **2020**, *246*, 188–200. [[CrossRef](#)]
46. Antonova, T.M.; Dmytryshyn, M.V.; Vozna, S.M. Some properties of approximants for branched continued fractions of the special form with positive and alternating-sign partial numerators. *Carpathian Math. Publ.* **2018**, *10*, 3–13. [[CrossRef](#)]
47. Antonova, T.M.; Sus', O.M. Sufficient conditions for the equivalent convergence of sequences of different approximants for two-dimensional continued fractions. *J. Math. Sci.* **2018**, *228*, 1–10. [[CrossRef](#)]
48. Antonova, T.M.; Sus', O.M.; Vozna, S.M. Convergence and estimation of the truncation error for the corresponding two-dimensional continued fractions. *Ukr. Math. J.* **2022**, *74*, 501–518. [[CrossRef](#)]
49. Bodnar, D.I.; Bodnar, O.S.; Bilanyk, I.B. A truncation error bound for branched continued fractions of the special form on subsets of angular domains. *Carpathian Math. Publ.* **2023**, *15*, 437–448. [[CrossRef](#)]
50. Bodnar, O.S.; Dmytryshyn, R.I.; Sharyn, S.V. On the convergence of multidimensional S-fractions with independent variables. *Carpathian Math. Publ.* **2020**, *12*, 353–359. [[CrossRef](#)]
51. Dmytryshyn, R.; Cesarano, C.; Lutsiv, I.-A.; Dmytryshyn, M. Numerical stability of the branched continued fraction expansion of Horn's hypergeometric function  $H_4$ . *Mat. Stud.* **2024**, *61*, 51–60. [[CrossRef](#)]
52. Hladun, V.R.; Bodnar, D.I.; Rusyn, R.S. Convergence sets and relative stability to perturbations of a branched continued fraction with positive elements. *Carpathian Math. Publ.* **2024**, *16*, 16–31. [[CrossRef](#)]
53. Hladun, V.R. Some sets of relative stability under perturbations of branched continued fractions with complex elements and a variable number of branches. *J. Math. Sci.* **2016**, *215*, 11–25. [[CrossRef](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.