



*symmetry*



Review

---

# Time–Energy Uncertainty Relation for Neutrino Oscillations: Historical Development, Applications, and Future Prospects

---

Giuseppe Gaetano Luciano and Luca Smaldone

Special Issue

Symmetry in Neutrino Physics and Astrophysics

Edited by



Prof. Dr. Célio Adrega De Moura Junior and Dr. Fernando Rossi-Torres



<https://doi.org/10.3390/sym15112032>

Review

# Time–Energy Uncertainty Relation for Neutrino Oscillations: Historical Development, Applications, and Future Prospects

Giuseppe Gaetano Luciano <sup>1,\*</sup>  and Luca Smaldone <sup>2,3</sup> 

- <sup>1</sup> Department of Chemistry, Physics and Environmental and Soil Sciences, Escola Politècnica Superior, Universitat de Lleida, Av. Jaume II, 69, 25001 Lleida, Spain
- <sup>2</sup> Dipartimento di Fisica, Università degli Studi di Salerno, Via Giovanni Paolo II, 132, 84084 Fisciano, SA, Italy; lsmaldone@unisa.it
- <sup>3</sup> INFN Gruppo Collegato di Salerno-Sezione di Napoli, c/o Dipartimento di Fisica, Università degli Studi di Salerno, Via Giovanni Paolo II, 132, 84084 Fisciano, SA, Italy
- \* Correspondence: giuseppegatano.luciano@udl.cat

**Abstract:** The time–energy uncertainty relation (TEUR) plays a fundamental role in quantum mechanics, as it allows the grasping of peculiar aspects of a variety of phenomena based on very general principles and symmetries of the theory. Using the Mandelstam–Tamm method, TEUR has recently been derived for neutrino oscillations by connecting the uncertainty in neutrino energy with the characteristic timescale of oscillations. Interestingly, the suggested interpretation of neutrinos as unstable-like particles has proved to naturally emerge in this context. Further aspects were later discussed in semiclassical gravity theory, by computing corrections to the neutrino energy uncertainty in a generic stationary curved spacetime, and in quantum field theory, where the clock observable turns out to be identified with the non-conserved flavor charge operator. In the present work, we give an overview on the above achievements. In particular, we analyze the implications of TEUR and explore the impact of gravitational and non-relativistic effects on the standard condition for neutrino oscillations.

**Keywords:** time–energy uncertainty relation; quantum theory; general relativity; neutrino; flavor oscillations; unstable particles



check for updates

**Citation:** Luciano, G.G.; Smaldone, L. Time–Energy Uncertainty Relation for Neutrino Oscillations: Historical Development, Applications, and Future Prospects. *Symmetry* **2023**, *15*, 2032. <https://doi.org/10.3390/sym15112032>

Academic Editors: Antonia Di Crescenzo, Célio Adrega De Moura Junior and Sergei D. Odintsov

Received: 19 October 2023  
Revised: 3 November 2023  
Accepted: 7 November 2023  
Published: 8 November 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Uncertainty relations constitute a cornerstone of quantum theory. For example, the Heisenberg principle sets a fundamental lower limit on how precisely we can perform simultaneous measurements of position and momentum of any quantum system. A direct consequence is that an exactly specified (classical-like) phase-space trajectory cannot be defined for these systems. Another such pair of incompatible variables are energy and time. An informal meaning of the time–energy uncertainty relation (TEUR) is that quantum states that only exist for a relatively short time cannot have a sharply determined energy.

Despite formal similarities, the analogy between TEUR and the Heisenberg principle cannot be pursued all the way because of the lack of a time operator in quantum mechanics (QM). In fact, the proper formalization and general validity of TEUR have been controversial issues since the advent of quantum theory. For a comprehensive review on the subject, one can refer to [1], while a recent discussion on the non-uniqueness of TEUR formulation appears in [2]. In what follows, we shall basically resort to the most general Mandelstam–Tamm version of TEUR, which is based on the only assumption that the evolution of Heisenberg operators in quantum theory is ruled by the Hamiltonian  $H$  [3]. Using natural units  $\hbar = c = G = 1$ , TEUR in this framework takes the form

$$\Delta E \Delta t \geq \frac{1}{2}, \quad (1)$$

where  $\Delta E \equiv \sigma_H$  is interpreted as the standard deviation of the energy on a generic quantum state  $|\psi\rangle$ , and

$$\Delta t \equiv \sigma_O / \left| \frac{d\langle O(t) \rangle}{dt} \right|. \quad (2)$$

Here,  $O(t)$  denotes the “clock observable”, whose dynamics quantifies temporal changes of the system, and  $\Delta t$  is the characteristic timescale over which the mean value  $\langle O(t) \rangle \equiv \langle \psi | O(t) | \psi \rangle = \langle \psi(t) | O | \psi(t) \rangle$  varies by a standard deviation.

One of the most common applications of TEUR concerns the understanding of the decay of excited states of atoms. In this scenario, the minimum time it takes for an atom to decay to its ground state and radiate is related to the energy uncertainty of the excited state [4]. Based on the Mandelstam–Tamm method, in [5], TEUR was derived for neutrino flavor oscillations in the QM picture.

Neutrino oscillation is a phenomenon characterized by a finite timescale, the oscillation time [6], and thus, particularly suited for a description in terms of Equation (1). By identifying  $\Delta t$  with the time during which a significant change in flavor content happens for a given neutrino state, the authors of [5] came to the conclusion that a non-vanishing neutrino energy uncertainty  $\Delta E$  is actually needed for oscillations to occur.

Interesting advances in the study of TEUR for neutrino oscillations were later achieved from both the phenomenological and theoretical points of view. In the former context, the applicability of TEUR to the Mössbauer neutrino experiment was analyzed with conflicting results [7,8]. On the theoretical side, a generalized TEUR was obtained in [9] for a generic stationary curved spacetime, while in [10], Equation (1) was recast in terms of a flavor–energy uncertainty relation by using the quantum field theoretical (QFT) formalism of [11,12]. Remarkably, the QFT approach of [10] brings out the interpretation of TEUR for flavor neutrinos as fundamental bounds on energy variances, in close analogy with the case of unstable particles. All of the above considerations make it clear that a careful study of TEUR might reveal valuable information about the true nature of neutrinos and the related mechanism of flavor oscillations, which are still subject of active investigation.

Based on the outlined premises, here we review recent progress in the description of TEUR for neutrino oscillations. After a brief historical overview, we summarize basic concepts and results in the literature, with focus on the future prospects and research lines that can be inspired by the present analysis. The rest of the paper is structured as follows. In Section 2, we go through the derivation of the original QM TEUR for both neutrinos and unstable particles. Section 3 is devoted to exploring gravitational effects on TEUR, while the extension to QFT is investigated in Section 4. Conclusions and discussion are presented in Section 5.

## 2. TEUR for Oscillating and Unstable Particles

Let us start by presenting TEUR for neutrino oscillations as first derived by Bilenky [5]. The result is compared to the corresponding formula for unstable particles [13,14], emphasizing conceptual similarities and differences.

### 2.1. TEUR for Neutrino Oscillations (*à-la* Bilenky)

Following [5], we set  $O = P_l = |v_l\rangle\langle v_l|$  in the Mandelstam–Tamm inequality (1), where  $|v_l\rangle$  ( $l = e, \mu, \tau$ ) is the flavor neutrino state ( $\langle v_l | v_{l'} \rangle = \delta_{ll'}$ ) and  $P_l$  is the projection operator on this state. It is straightforward to check that the average value of  $P_l$  on a generic state  $|\psi(t)\rangle$  is nothing but the probability to find the flavor neutrino  $v_l$  in  $|\psi(t)\rangle$ , i.e.,  $\langle P_l(t) \rangle = |\langle v_l | \psi(t) \rangle|^2$ . Hence, assuming  $|\psi(0)\rangle = |v_l\rangle$ , one has that  $\langle P_l(t) \rangle$  coincides with the survival probability  $P_{v_l \rightarrow v_l}(t)$  of the neutrino  $v_l$  at time  $t$ . Clearly, the following relations hold true:

$$P_{v_l \rightarrow v_l}(0) = 1, \quad P_{v_l \rightarrow v_l}(t) \leq 1 \text{ for any } t > 0. \quad (3)$$

Projection operators are idempotent, which means they satisfy  $P^2 = P$ . From this condition, it follows that the standard deviation  $\Delta P_l$  obeys

$$\Delta P_l(t) = \sqrt{P_{\nu_l \rightarrow \nu_l}(t) - P_{\nu_l \rightarrow \nu_l}^2(t)}. \quad (4)$$

By substitution into Equation (1), we derive

$$\Delta E \geq \frac{1}{2} \frac{\left| \frac{d}{dt} P_{\nu_l \rightarrow \nu_l}(t) \right|}{\sqrt{P_{\nu_l \rightarrow \nu_l}(t) - P_{\nu_l \rightarrow \nu_l}^2(t)}}. \quad (5)$$

Let us now consider the survival probability  $P_{\nu_l \rightarrow \nu_l}(t)$  in the time interval  $0 \leq t \leq t_{1min}$ , where  $t_{1min}$  is the time when  $P_{\nu_l \rightarrow \nu_l}(t)$  reaches the first minimum. In this range,  $P_{\nu_l \rightarrow \nu_l}(t)$  is a monotonically decreasing function [5,7,8,15–17]. By integration of Equation (5) from 0 to  $t$ , we finally get

$$\Delta E t \geq \frac{1}{2} \left[ \frac{\pi}{2} - \sin^{-1}(2P_{\nu_l \rightarrow \nu_l}(t) - 1) \right]. \quad (6)$$

It is useful to apply the above inequality to some specific cases of experimental interest. For instance, atmospheric neutrinos are produced by cosmic rays interacting in the upper atmosphere of the Earth. In turn, the resulting pions and kaons decay into muon neutrinos and muons, many of which give rise to electrons and a muon neutrino–electron neutrino pair. A crude estimation indicates that muon neutrinos are roughly two times more abundant than electron neutrinos [18,19]. In the atmospheric Long Baseline region, the survival probability  $P_{\nu_\mu \rightarrow \nu_\mu}(t)$  is driven by the neutrino-mass-squared difference  $\Delta m_{23}^2$  (for more details on the theoretical treatment of neutrino mixing, see Sections 3 and 4). By taking into account that  $P_{\nu_\mu \rightarrow \nu_\mu}(t_{1min}^{(23)}) \simeq 0$ , where  $t_{1min}^{(23)} = 2\pi E / \Delta m_{23}^2$ , we obtain the following TEUR for ultrarelativistic neutrinos ( $L \simeq t$ ) [5]

$$\Delta E t_{osc}^{(23)} \geq \pi, \quad (7)$$


where  $t_{osc}^{(23)} = 2t_{1min}^{(23)}$  denotes the period of oscillations in the atmospheric Long Baseline region. As explained in [5], the above relation provides a necessary condition for atmospheric neutrino oscillations to be observed.

As a further example, we consider the  $\Delta m_{23}^2$ -driven survival probability of  $\bar{\nu}_e$ . According to the CHOOZ experiment [20], this probability is close to the maximal value of unity. Using the condition  $P_{\bar{\nu}_e \rightarrow \bar{\nu}_e}(t_{1min}^{(23)}) = 1 - \sin^2 2\theta_{13}$  along with the constraint  $\sin^2 2\theta_{13} \lesssim 2 \times 10^{-1}$ , where  $\theta_{13}$  is the mixing angle, Equation (6) becomes (up to terms of order  $\sin^2 2\theta_{13}$ )

$$\Delta E t_{osc}^{(23)} \geq 2 \sin 2\theta_{13}, \quad (8)$$

which is less stringent than the bound (7).

An interesting experimental application of TEUR in the high-energy regime concerns Mössbauer neutrino effects [7,8,15,16,21–24]. Mössbauer neutrinos could be produced and detected in recoilless Mössbauer transitions



with energy  $E_\nu \simeq 18.6$  keV and relative uncertainty  $\Delta E_\nu / E_\nu \simeq 4.5 \times 10^{-16}$ . Flavor oscillations of Mössbauer neutrinos were studied in [22] by using the macroscopic Feynman-diagram approach [22,25–28]. This result was, however, challenged by Bilenky et al. [7], who estimated that the minimum uncertainty needed for Mössbauer oscillations to occur according to TEUR should be  $\Delta E_\nu / E_\nu \gtrsim 10^{-12}$  for atmospheric neutrinos,  $\Delta E_\nu / E_\nu \gtrsim 10^{-13}$  for reactor neutrinos, and  $\Delta E_\nu / E_\nu \gtrsim 10^{-14}$  for solar neutrinos. This controversy was later debated in [8,15,16,22]. In particular, a connection between this issue and the possible non-stationary nature of oscillations was investigated in [15,16]. The conclusion was that,

if time plays an active role in neutrino oscillations, one cannot escape from the implications of TEUR. Conversely, if oscillations are a stationary phenomenon, the viewpoint of Akhmedov [22] would be the correct one. In the absence of a fundamental and widely shared theory of neutrino oscillations, valuable hints could be provided by the possible future detection of Mössbauer neutrinos.

### 2.2. TEUR and Unstable Particles

We here digress briefly to discuss TEUR for unstable particles. Let us stress that such systems provide one of the most interesting examples in which TEUR is typically applied, see, e.g., [29]. Toward this end, we stick to the original treatment of [13] and the recent review in [14].

The analysis of [13], which inspired the derivation of TEUR for neutrino oscillations, is built upon the same considerations leading to Equation (5), with  $P_{\nu_l \rightarrow \nu_l}$  being now replaced by the so-called quantum non-decay probability  $P_t$ . For convenience, let us recast the inequality (5) as

$$\left| \frac{d}{dt} P_t \right| \leq 2\Delta E \sqrt{P_t - P_t^2}, \tag{10}$$

which allows identification of the particle half-life  $T_h$ , for which  $P_t = 1/2$  and the right-hand side attains its maximum value. Then, Equation (10) gives the weaker inequality

$$\left| \frac{d}{dt} P_t \right| \leq \Delta E, \tag{11}$$

which sets a time-limit to the instability of decaying quantum systems (see below for more quantitative discussion).

Since for decaying particles  $P_{t=0} = 1$  and  $P_{t \rightarrow \infty} \rightarrow 0$ , we can infer the following features that are likely to be of interest in the asymptotic regimes of decay

$$\left| \frac{d}{dt} P_t \right| = 0, \quad \text{for } t = 0, \tag{12}$$

$$\frac{d}{dt} \cos^{-1} P_t \leq 2\Delta E, \quad \text{for } t \rightarrow 0, \tag{13}$$

$$\frac{d}{dt} P_t^{1/2} \leq \Delta E, \quad \text{for } t \rightarrow \infty. \tag{14}$$

These can be unified as

$$\frac{d}{dt} \cos^{-1} P_t^{1/2} \leq \Delta E, \quad \text{for } 0 \leq t \leq \infty. \tag{15}$$

Integration of the the latter inequality gives

$$\Delta E t \geq \cos^{-1} P_t^{1/2}, \tag{16}$$

which is, in fact, the analog of Equation (6) for unstable particles. At this stage, it becomes clear that the parallelism between neutrinos and unstable particles has a merely formal meaning—indeed, although a TEUR can be associated with both systems, the phenomenon of oscillations is not strictly equivalent to a decay, since the oscillation probability does not vanish at asymptotically long time. Notice also that Equation (16) fixes the minimum time-limit to  $\pi/(2\Delta E)$ .

Now, for  $t$  equal to the half-life  $T_h$  defined earlier, Equation (16) leads straightforwardly to

$$\Delta E T_h \geq \frac{\pi}{4} \quad \text{for } 0 < T_h \leq \frac{\pi}{2\Delta E}. \tag{17}$$

Conversely, if  $T_h > \pi/(2\Delta E)$ , the non-negativity of  $P_t$  allows us to write

$$P_t \geq 0 > 1 - \frac{2\Delta E t}{\pi}, \quad \text{for } t > \frac{\pi}{2\Delta E}, \quad (18)$$

which again implies the inequality (17). Combining these two results, we finally obtain the Mandelstam–Tamm version of TEUR for a decaying quantum system

$$\Delta E T_h \geq \frac{\pi}{4}. \quad (19)$$

The above derivation has been recently revised in [14], leading to a more stringent bound. Based on the approach of [30], the basic idea is to regard a system of unstable particles as an open quantum system. This entails that a comprehensive analysis cannot be limited to the system alone, but must necessarily involve its surrounding “environment” too. In a field theoretical language, such a prescription amounts to saying that it is not sufficient to describe the system as excitations of some fields on a single vacuum state, but one should take into account extra (artificial) degrees of freedom. A typical strategy consists of doubling each physical degree of freedom, resulting into an enlargement of the Hilbert space. Although this construction was originally proposed to develop a QFT at finite temperature—Thermo Field Dynamics [31–33]—various applications have been later considered in QFT on curved spacetime [34] and the quantum brain model [35], among others.

For practical purposes, let us consider a set of canonical fermionic operators  $a_{\mathbf{k}}, a_{\mathbf{k}}^\dagger$ , which annihilate and create a decaying particle of momentum  $\mathbf{k}$  at  $t = 0$ , respectively. The associated vacuum is denoted by  $|0\rangle$ , i.e.,  $a_{\mathbf{k}}|0\rangle = 0$ . Following the standard notation [31,32], we indicate the corresponding set of *doubled* operators by  $\tilde{a}_{\mathbf{k}}, \tilde{a}_{\mathbf{k}}^\dagger$  and the related vacuum by  $|\tilde{0}\rangle$ , so that  $\tilde{a}_{\mathbf{k}}|\tilde{0}\rangle = 0$ . Notice that such new operators obey the same (anti-) commutation relations as  $a_{\mathbf{k}}, a_{\mathbf{k}}^\dagger$ , i.e.,  $\{a_{\mathbf{k}}, a_{\mathbf{p}}^\dagger\} = \{\tilde{a}_{\mathbf{k}}, \tilde{a}_{\mathbf{p}}^\dagger\} = \delta(\mathbf{k} - \mathbf{p})$ .

Using the above tools, it is natural to define an enlarged Fock space having the tensor-product structure between the Fock spaces associated with  $a_{\mathbf{k}}$  and  $\tilde{a}_{\mathbf{k}}$ , respectively. In this setting, the underlying vacuum state  $|0\rangle\rangle = |0\rangle \otimes |\tilde{0}\rangle$  is defined in such a way that

$$a_{\mathbf{k}}|0\rangle\rangle = \tilde{a}_{\mathbf{k}}|0\rangle\rangle = 0. \quad (20)$$

The core idea of [14] is to construct a state  $|0(\varphi)\rangle$ , such that the expectation value of the number operator  $N_{\mathbf{k}} = a_{\mathbf{k}}^\dagger a_{\mathbf{k}}$  satisfies

$$\mathcal{N}_{\mathbf{k}}(t) \equiv \langle 0(\varphi) | N_{\mathbf{k}} | 0(\varphi) \rangle = \exp(-\Gamma_{\mathbf{k}} t), \quad (21)$$

where  $\Gamma_{\mathbf{k}}$  is the inverse lifetime of the unstable particle (the physical meaning of the parameter  $\varphi$  will become clear later). Inspired by Thermo Field Dynamics [31], one can introduce the new set of ladder operators via the Bogoliubov transformations

$$a_{\mathbf{k}}(\varphi) = \cos \varphi_{\mathbf{k}} a_{\mathbf{k}} - \sin \varphi_{\mathbf{k}} \tilde{a}_{\mathbf{k}}^\dagger, \quad (22)$$

$$\tilde{a}_{\mathbf{k}}(\varphi) = \cos \varphi_{\mathbf{k}} \tilde{a}_{\mathbf{k}} + \sin \varphi_{\mathbf{k}} a_{\mathbf{k}}^\dagger. \quad (23)$$

The related vacuum is defined by the usual condition  $a_{\mathbf{k}}(\varphi) |0(\varphi)\rangle = \tilde{a}_{\mathbf{k}}(\varphi) |0(\varphi)\rangle = 0$ . By explicit computation, it is possible to show that [31]

$$|0(\varphi)\rangle = \prod_{\mathbf{k}} \left( \cos \varphi_{\mathbf{k}} + \sin \varphi_{\mathbf{k}} a_{\mathbf{k}}^\dagger \tilde{a}_{\mathbf{k}}^\dagger \right) |0\rangle\rangle. \quad (24)$$

Combining Equations (21) and (24), we infer

$$\mathcal{N}_{\mathbf{k}}(t) = \sin^2 \varphi_{\mathbf{k}}(t) = \exp(-\Gamma_{\mathbf{k}} t), \quad (25)$$

which displays the meaning of  $\varphi_{\mathbf{k}} \equiv \varphi_{\mathbf{k}}(t)$  in terms of the inverse lifetime  $\Gamma_{\mathbf{k}}$ .

To have a more physical intuition on  $|0(\varphi)\rangle$ , let us consider the asymptotic regions  $t \rightarrow 0$  and  $t \rightarrow \infty$ . In the first case, we have

$$|0(\varphi)\rangle|_{t=0} = \prod_{\mathbf{k}} |0(\varphi_{\mathbf{k}})\rangle, \quad (26)$$

where

$$|0(\varphi_{\mathbf{k}})\rangle = a_{\mathbf{k}}^{\dagger} \bar{a}_{\mathbf{k}}^{\dagger} |0\rangle, \quad (27)$$

is the state describing an unstable particle of momentum  $\mathbf{k}$ . Conversely, the limit

$$|0(\varphi)\rangle|_{t=\infty} = |0\rangle, \quad (28)$$

gives the zero-particle state.

In order to extract TEUR, we now switch to the Heisenberg representation. In this picture, the expectation value (25) is kept unchanged, while the vacuum  $|0(\varphi)\rangle$  must be set as in Equation (26). In turn, the number operator  $N_{\mathbf{k}}$  defined above in Equation (21) gains a non-trivial time-dependence. With this setting, we can choose  $O(t) = N_{\mathbf{k}}(t)$  in Equations (1) and (2) to obtain

$$\sigma_N^2 = \langle 0(\varphi) | N_{\mathbf{k}}^2(t) | 0(\varphi) \rangle - \langle 0(\varphi) | N_{\mathbf{k}}(t) | 0(\varphi) \rangle^2 = \mathcal{N}_{\mathbf{k}}(t)(1 - \mathcal{N}_{\mathbf{k}}(t)), \quad (29)$$

which yields

$$\left| \frac{d\mathcal{N}_{\mathbf{k}}(t)}{dt} \right| \leq 2\Delta E \sqrt{\mathcal{N}_{\mathbf{k}}(t)(1 - \mathcal{N}_{\mathbf{k}}(t))}. \quad (30)$$

Notice that this inequality is the same as Equation (10), with the non-decay probability  $P_t$  being now formally replaced by the vacuum density  $\mathcal{N}_{\mathbf{k}}(t)$ . For the characteristic time  $T_h$  that maximizes the right side, we have  $\mathcal{N}_{\mathbf{k}}(T_h) = 1/2$  and

$$\Delta E \geq \left| \frac{d\mathcal{N}_{\mathbf{k}}(t)}{dt} \right|, \quad (31)$$

to be compared with Equation (11).

The relation (31) can be further manipulated by resorting to the triangular inequality and integrating both sides from 0 to  $T$  to obtain

$$\Delta E T \geq \int_0^T dt \left| \frac{d\mathcal{N}_{\mathbf{k}}(t)}{dt} \right| \geq \left| \int_0^T dt \frac{d\mathcal{N}_{\mathbf{k}}(t)}{dt} \right|, \quad (32)$$

which yields

$$\Delta E T \geq 1 - \mathcal{N}_{\mathbf{k}}(T). \quad (33)$$

At  $T = T_h$ , we then get the Heisenberg-like lower bound

$$\Delta E T_h \geq \frac{1}{2}, \quad (34)$$

that is stronger than the one in Equation (19).

As a final remark, it is worth noting from Equation (25) that  $T_h = \log 2 / \Gamma_{\mathbf{k}} = \tau_{\mathbf{k}} \log 2$ , where  $\tau_{\mathbf{k}} = 1 / \Gamma_{\mathbf{k}}$  is the particle lifetime. Therefore, TEUR (34) can be rearranged as

$$\Delta E \geq \frac{1}{\tau_{\mathbf{k}} \log 4}. \quad (35)$$

This inequality sets an intrinsic lower bound on the energy distribution width of unstable quantum particles. In the same way, by continuing the analogy between neutrinos and unstable particles, we can interpret Equation (5) as a fundamental limit on the width of the neutrino energy distribution.



### 3. Gravitational Effects on TEUR

In [9], TEUR for neutrino oscillations has been derived for a generic stationary curved spacetime to study how gravity affects the characteristic oscillation length. To review such a formalism, we first take a step back and consider the Bilenky-like (Minkowski) form (6) of TEUR, here rewritten equivalently as

$$2\Delta E \sqrt{P_{\nu_l \rightarrow \nu_l}(x) - P_{\nu_l \rightarrow \nu_l}^2(x)} \geq \left| \frac{d}{dt} P_{\nu_l \rightarrow \nu_l}(x) \right|, \tag{36}$$

where, for more generality, we have assumed that the neutrino survival probability  $P_{\nu_l \rightarrow \nu_l}$  depends on  $t$  through the spacetime coordinate  $x \equiv x(t)$ . For simplicity, from now on we restrict to a toy model involving two flavors only, but the same considerations can be extended to the more realistic case of three neutrino generations.

As observed in Section 2.2, the left side of Equation (36) reaches its maximum when  $P_{\nu_l \rightarrow \nu_l} = 1/2$ . Integrating both sides between some initial time  $t_0$  (such that  $P_{\nu_l \rightarrow \nu_l}(x(t_0)) = 1$ ) and  $t$ , and employing once more the triangular inequality, we are led to [9]

$$\Delta E T \geq P_{\nu_l \rightarrow \nu_l}(x(t)), \tag{37}$$

where  $T \equiv \int_{t_0}^t dt' = t - t_0$  is the neutrino flight-time and  $P_{\nu_l \rightarrow \nu_l}(x(t)) = 1 - P_{\nu_l \rightarrow \nu_l}(x(t))$  is the oscillation probability.

Gravitational corrections to flavor oscillations typically enter the neutrino phase [36–53]. To find out the form of these extra terms, we write explicitly the generic flavor eigenstate  $| \nu_l \rangle$  as a mixture of the mass eigenstates  $| \nu_k \rangle$  ( $k = 1, 2$ ) according to [6]

$$| \nu_l(x) \rangle = \sum_{k=1,2} U_{lk}(\theta) | \nu_k(x) \rangle, \quad l = e, \mu, \tag{38}$$

where  $U_{lk}(\theta)$  is the generic element of the mixing matrix [6].

Since the states  $| \nu_k(x) \rangle$  are Hamiltonian eigenstates, in the standard quantum theory using a flat background, their evolution is governed by the usual relation

$$| \nu_k(x) \rangle = \exp[-i\varphi_k(x)] | \nu_k \rangle, \quad (k = 1, 2), \tag{39}$$

where

$$\varphi_k(x) \equiv E_k t - \mathbf{p}_k \cdot \mathbf{x} \tag{40}$$

denotes the QM phase of the  $k^{th}$  eigenstate having mass  $m_k$ , energy  $E_k$ , and three-momentum  $\mathbf{p}_k$ , respectively, (not to be confused with the parameter  $\varphi_k$  appearing in the Bogoliubov transformations (22) and (23)). These three quantities are connected via the flat mass-shell relation  $E_k^2 = m_k^2 + |\mathbf{p}_k|^2$ .

For relativistic neutrinos traveling along the  $x$ -axis from the source point  $x_0$  to the detector point  $x$  (with  $x > x_0$ ), the phase (40) can be simplified to

$$\varphi_k \simeq \frac{m_k^2}{2E} L_p, \tag{41}$$

where we have assumed the mass eigenstates to have roughly the same energy  $E$ . Furthermore, we have denoted the proper distance traveled by neutrinos by  $L_p = x - x_0$ , which can be approximated to the flight-time  $T$  in the relativistic limit.

Now, substitution of Equation (41) into (38) and (39) gives

$$| \nu_l(x) \rangle = \sum_{k=1,2} U_{lk}(\theta) \exp\left(-i \frac{m_k^2}{2E} L_p\right) | \nu_k \rangle, \quad l = e, \mu. \tag{42}$$



In turn, the probability that a neutrino  $|ν_l\rangle$  undergoes a flavor transition to  $|ν_{l'}\rangle$  after propagating over the distance  $L_p$  is [9]

$$\begin{aligned}
 P_{ν_l \rightarrow ν_{l'}}(x) &= |\langle ν_{l'}(x) | ν_l(x_0) \rangle|^2 \\
 &= \sin^2(2\theta) \sin^2\left(\frac{\varphi_{12}}{2}\right) \\
 &= \sin^2(2\theta) \sin^2\left(\frac{\pi L_p}{L^{osc}}\right), \quad l \neq l',
 \end{aligned}
 \tag{43}$$

where in the second step  $\varphi_{12} \equiv \varphi_1 - \varphi_2$  denotes the phase-shift acquired by the mass eigenstates during the propagation, while in the last step we have introduced the characteristic oscillation length  $L^{osc}$

$$L^{osc} \equiv \frac{4\pi E}{\Delta m^2}, \quad \Delta m^2 = |m_1^2 - m_2^2|.
 \tag{44}$$

Accordingly, TEUR in Equation (37) takes the form

$$\Delta E T \geq \sin^2(2\theta) \sin^2\left(\frac{\pi L_p}{L^{osc}}\right).
 \tag{45}$$

Since this inequality holds for any spacetime point, we can set  $L_p = L^{osc}/2$  so as to maximize the right side and obtain the neutrino oscillation condition

$$\Delta E \geq \frac{2 \sin^2(2\theta)}{L^{osc}}.
 \tag{46}$$

Two comments are in order here: First, we notice that for  $\theta = 0$  and/or  $L^{osc} \rightarrow \infty$ , the lower bound on the neutrino energy uncertainty vanishes, as expected in the absence of mixing and/or oscillations. Moreover, in [54],  $\Delta E$  is interpreted as the minimum energy transferred to neutrinos in scattering processes, which is necessary to reveal them once the oscillation occurred.

### 3.1. TEUR in Curved Spacetime

Let us extend the above formalism to stationary curved spacetime. For this kind of metrics, we remark that TEUR can be consistently defined, since there always exists a global time-like Killing vector field  $K^\mu$ , such that  $K \equiv \int K_\mu T^{\mu\nu} d\Sigma_\nu$  does not depend on the space-like hypersurface  $\Sigma$ , where  $T^{\mu\nu}$  is the (conserved) stress-energy tensor [55]. We are then allowed to introduce a coordinate  $t$ , such that the metric is independent of it and with respect to which  $K^\mu = (1, 0, 0, 0)$ . One has  $K \equiv \int \sqrt{-g} : T^0_0 : d^3x = H$ , where  $: T^0_0 :$  denotes the normal-ordered operator,  $g$  is the determinant of the metric tensor  $g^{\mu\nu}$ , and  $H$  is the Hamiltonian of the considered system. More generally, the role of the above Killing vector can be taken by the Kodama vector for dynamical metrics [56].

To generalize TEUR for flavor oscillations to curved spacetime, it proves convenient to rephrase the evolution (42) of the neutrino state in a manifestly covariant way as [37]

$$|ν_l(\lambda)\rangle = \sum_{k=1,2} U_{lk}(\theta) \exp(-i\Phi) |ν_k\rangle, \quad l = e, \mu,
 \tag{47}$$

where  $\lambda$  is the neutrino world-line parameter and  $\Phi$  generalizes the definition (40) of neutrino phase to

$$\Phi = \int_{\lambda_0}^\lambda P_\mu \frac{dx^\mu_{null}}{d\lambda'} d\lambda'.
 \tag{48}$$

Notice that  $P_\mu$  and  $dx^\mu_{null}/d\lambda$  in the above integral are the generator of spacetime translations of neutrino mass states and the null tangent vector to the neutrino world line, respectively. For  $g_{\mu\nu} = \eta_{\mu\nu}$ , one can check that Equation (48) reproduces the evolution (42) in flat spacetime, as expected [9].

For a generic stationary curved spacetime, the operator  $P_\mu$  is defined by the generalized mass-shell relation [37]

$$\left(P^\mu + \frac{A^\mu \gamma^5}{2}\right) \left(P_\mu + \frac{A_\mu \gamma^5}{2}\right) = M^2, \tag{49}$$

where  $M = \text{diag}[m_1, m_2]$  is the diagonal neutrino mass-matrix and  $\gamma_\mu$  are the Dirac matrices obeying the Clifford algebra  $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$  ( $\gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3$ ). Furthermore, the potential-like term

$$A^\mu = \frac{\sqrt{g}}{4} e_a^\mu \epsilon^{\hat{a}\hat{b}\hat{c}\hat{d}} (\partial_\sigma e_{\hat{b}\nu} - \partial_\nu e_{\hat{b}\sigma}) e_c^\nu e_d^\sigma \tag{50}$$

enters the spin-connection in the Dirac equation on curved spacetime [9,37]. Here,  $e_a^\mu$  are the tetrads connecting general curvilinear and locally inertial coordinates, denoted by Greek and (hatted) Latin indices, respectively. The symbol  $\epsilon^{\hat{a}\hat{b}\hat{c}\hat{d}}$  stands for the totally antisymmetric Levi-Civita tensor of component  $\epsilon^{\hat{0}\hat{1}\hat{2}\hat{3}} = +1$ , while  $\partial$  indicates ordinary derivative.

As shown in [37], the integral in (48) can be simplified for relativistic neutrinos moving along a null trajectory by neglecting the  $\mathcal{O}(A^2)$  and  $\mathcal{O}(AM^2)$  terms. Under these assumptions, we obtain [9]

$$\Phi = \int_{\lambda_0}^{\lambda} \left( \frac{M^2}{2} - \frac{dx_{null}^\mu}{d\lambda'} A_\mu \gamma^5 \right) d\lambda', \tag{51}$$

which can be further manipulated by rewriting the line element  $d\lambda$  in terms of the differential proper distance  $d\ell$  at constant time for null trajectories as

$$d\lambda = d\ell \left( -g_{ij} \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda} \right)^{-\frac{1}{2}} = d\ell \left[ g_{00} \left( \frac{dt}{d\lambda} \right)^2 + 2g_{0i} \frac{dt}{d\lambda} \frac{dx^i}{d\lambda} \right]^{-\frac{1}{2}}. \tag{52}$$

Equation (51) provides the crucial ingredient toward reformulating TEUR in curved spacetime. To this end, we recast Equation (37) in terms of  $\lambda$  to obtain

$$\Delta E T(\lambda) \geq \mathcal{P}_{\nu_l \rightarrow \nu_{l'}}(x(\lambda)), \tag{53}$$

where the neutrino flight-time is given by  $T(\lambda) \equiv t(\lambda) - t(\lambda_0)$  for a fixed  $\lambda_0$ . From Equation (43), it follows that

$$\Delta E \geq \frac{\sin^2(2\theta)}{T(\lambda)} \sin^2 \left[ \frac{\varphi_{12}(x(\lambda))}{2} \right], \quad \varphi_{12} \equiv \varphi_1 - \varphi_2, \tag{54}$$

where the neutrino phase  $\varphi_k$  ( $k = 1, 2$ ) is now generally defined as the eigenvalue of  $\Phi$  in Equation (51) with respect to the  $k^{\text{th}}$  mass eigenstate, i.e.,  $\Phi|v_k\rangle = \varphi_k|v_k\rangle$ .

Before moving onto the evaluation of the above inequality for some specific metrics, it is worth observing that  $\Delta E$  appearing in Equation (54) represents the uncertainty on the neutrino energy as measured by an asymptotic observer (see [9] for more details). However, the truly measurable quantity in neutrino oscillation experiments is the local energy uncertainty  $\Delta E_\ell$ , which represents the uncertainty for a local observer temporarily at rest in the curved background (and, thus, with respect to the oscillation experiment). The connection between these two quantities is established via the vierbein fields

$$E_\ell(x) \equiv P_\delta(x) = e_\delta^\nu(x) P_\nu. \tag{55}$$

Clearly, this transformation must also be implemented into the expression of the phase-shift  $\varphi_{12}$ , along with the usage of the proper distance traveled by neutrinos. This is an essential step to make a consistent comparison with the TEUR (46) in the limit when  $g_{\mu\nu} \rightarrow \eta_{\mu\nu}$ .

### 3.1.1. Schwarzschild Spacetime

The Schwarzschild metric in isotropic coordinates  $(t, x, y, z)$  and (linearized) weak-field approximation is

$$ds^2 = (1 + 2\phi)dt^2 - (1 - 2\phi)(dx^2 + dy^2 + dz^2), \tag{56}$$

where  $\phi(r) = -M/r \equiv -M/\sqrt{x^2 + y^2 + z^2}$  is the gravitational potential and  $M$  the spherically symmetric source mass. The only non-trivial components of the tetrads are  $e_0^0 = 1 - \phi$  and  $e_j^i = (1 + \phi)\delta_j^i$ , where  $\delta_j^i$  is the Kronecker delta.

Because of the particular symmetry of the metric, the potential  $A_\mu$  in Equation (50) vanishes. From Equations (51) and (52), the resulting neutrino phase-shift  $\varphi_{12}$  takes the simple form [9]

$$\varphi_{12} = \frac{\Delta m^2}{2} \int_{\ell_0}^{\ell} \frac{1 + \phi}{E} d\ell', \tag{57}$$

with  $E \equiv P_0 = g_{00} dt/d\lambda$ .

If we set the frame axes so that the neutrino propagates radially (e.g., along the  $x$ -direction), we have  $d\ell = (1 - \phi) dx$ . Then, by taking into account that  $E$  is constant along a null trajectory, the integral (57) can be solved straightforwardly to give the following form of TEUR

$$\Delta E \geq \frac{\sin^2(2\theta)}{T} \sin^2 \left[ \frac{\Delta m^2(x - x_0)}{4E} \right]. \tag{58}$$

In terms of the proper distance

$$L_p \equiv \int_{x_0}^x \sqrt{-g_{11}} dx' = x - x_0 + M \ln \left( \frac{x}{x_0} \right), \tag{59}$$

and the local energy (55), we obtain

$$\frac{\Delta E_\ell}{1 - \phi} \geq \frac{\sin^2(2\theta)}{T} \sin^2 \left( \frac{\pi L_p}{L_\phi^{osc}} \right), \tag{60}$$

where we have introduced the Schwarzschild-corrected proper oscillation length (at the leading order in  $\phi$ ) [39]

$$L_\phi^{osc} \equiv \frac{4\pi E_\ell}{\Delta m^2} \left[ 1 + \phi + \frac{M}{x - x_0} \ln \left( \frac{x}{x_0} \right) \right]. \tag{61}$$

Here, the gravitational potential is to be regarded as  $\phi(x) = -M/x$ .

As emphasized in the text below Equation (45), the right side of the inequality (61) is maximized for  $L_p = L_\phi^{osc}/2$ , which in turn gives the following relation between the time interval  $T$  and the oscillation length  $L_\phi^{osc}$  [9]

$$T = \frac{L_\phi^{osc}}{2} \left[ 1 + \frac{M}{x - x_0} \ln \left( \frac{x}{x_0} \right) \right]. \tag{62}$$

By plugging into Equation (60), we can finally cast the TEUR in Schwarzschild spacetime as

$$\Delta E_\ell \geq \frac{2 \sin^2(2\theta)}{L_{eff}^{osc}(M)}, \tag{63}$$

where

$$L_{eff}^{osc}(M) \equiv L_\phi^{osc} \left[ 1 + \phi + \frac{M}{x - x_0} \ln \left( \frac{x}{x_0} \right) \right] \tag{64}$$

is the *effective oscillation length*.

The last form of TEUR (63) allows for a direct comparison with the corresponding flat expression (46). Remarkably, we can see that gravitational corrections increase  $L_{eff}^{osc}$  with respect to the standard oscillation length (44), resulting into a more stringent bound on  $\Delta E_\ell$ . For consistency, one can check that  $L_{eff}^{osc}(M) \rightarrow L^{osc}$  for  $M \rightarrow 0$  and the inequality (46) is recovered. Conversely, in the limit  $L_{eff}^{osc}(M) \rightarrow \infty$ , the right side of Equation (63) vanishes, in line with the discussion below Equation (46).

### 3.1.2. Lense–Thirring Spacetime

We can now apply the same line of reasoning to extract TEUR in Lense–Thirring background, which can be interpreted as the exterior gravitational field of a rotating spherical source. Notice also that this metric is usually adopted to describe gravitomagnetic frame-dragging effects, see for instance [57].

In the weak-field approximation, the line element on the equatorial plane reads

$$ds^2 = (1 + 2\phi)dt^2 - \frac{\phi\Omega}{r^2}(x dy - y dx) dt - (1 - 2\phi)(dx^2 + dy^2 + dz^2), \quad (65)$$

where  $\Omega \equiv 4R^2\omega/5$ ,  $\omega$  is the angular velocity of the source of radius  $R$  and  $\phi$  the gravitational potential defined below Equation (56). Without loss of generality, we assume the rotation takes place around the  $z$  axis.

Following the same calculations as in Section 3.1.1, we now arrive at [9]

$$\Delta E_\ell \geq \frac{2 \sin^2(2\theta)}{L_{eff}^{osc}(M, \Omega)}, \quad (66)$$

where the effective oscillation length exhibits a non-trivial dependence on both the mass and angular velocity of the source as

$$L_{eff}^{osc}(M, \Omega) \equiv L^{osc} \left[ 1 + \frac{M}{x - x_0} \ln \left( \frac{r + x}{r_0 + x_0} \right) + \frac{M\Omega}{2b(x - x_0)} \left( \frac{x}{r} - \frac{x_0}{r_0} \right) + \phi \left( 1 + \frac{\Omega b}{r^2} \right) \right], \quad (67)$$

with  $b > R$  being the impact parameter. As discussed for Schwarzschild geometry, the effective oscillation length is increased with respect to the flat case. The latter is however restored for  $M, \Omega \rightarrow 0$ , as it should be.

### 3.1.3. Rindler Spacetime

As a final example, we discuss TEUR for an observer undergoing constant proper acceleration  $a > 0$  (a Rindler observer). It is interesting to note that, although the corresponding Rindler metric is flat, it well-describes the static gravitational field that would be appropriate for a spacetime with fictitious “infinite flat earth” [58]. Assuming the acceleration along the  $x$ -axis, the (linearized) line element in Rindler–Fermi coordinates is [59]

$$ds^2 = (1 + ax)dt^2 - (dx^2 + dy^2 + dz^2). \quad (68)$$

TEUR for a Rindler observer can be cast in the same form as Equation (63) (or Equation (66)), with the acceleration-dependent oscillation length being now defined by [9]

$$L_{eff}^{osc}(a) \equiv L^{osc} \left[ 1 + \frac{a}{4}(x - x_0) \right]. \quad (69)$$

Once again, the effective oscillation length is increased with respect to the flat case, giving rise to a more stringent bound on the energy uncertainty  $\Delta E_\ell$ . As expected, the standard formula in Minkowski spacetime is reproduced for  $a \rightarrow 0$ . Conversely, the limit of  $L_{eff}^{osc}(a) \rightarrow \infty$  for fixed  $a$  cannot be considered in the present case, as it invalidates the linearized approximation under which Equation (69) is derived [59].

#### 4. TEUR in Quantum Field Theory

We start this section by reviewing the TEUR in the Pontecorvo formalism for neutrino mixing and oscillations [60–64] as developed in [10]. This analysis gives similar results to the ones shown in Sections 2 and 3. Although the Pontecorvo approach is experimentally successful, it has however several conceptual drawbacks and is widely recognized to be only applicable to relativistic neutrinos [11,65–69]. These issues motivate us to introduce the main aspects of the so-called *flavor Fock space* approach to neutrino mixing in QFT [11,70–77], which extends the Pontecorvo formalism to the non-relativistic domain. We then show how TEUR can be studied in this generalized framework [10]. For consistency with the original notation [11], in what follows, we label neutrino fields with definite masses by Latin indices, while Greek indices are reserved for definite flavor fields.

##### 4.1. Time–Energy Uncertainty Relations for Neutrino Oscillations: Pontecorvo Flavor States

In Section 2.2, we re-formulated the TEUR for unstable particles in a second quantization language. The same procedure can be repeated for the Minkowski form of TEUR for neutrino oscillations.

The fields  $v_j(x)$ ,  $j = 1, 2$  describe Dirac neutrinos with definite masses  $m_j$ :

$$v_j(x) = \sum_r \int \frac{d^3k}{(2\pi)^{\frac{3}{2}}} \left[ u_{\mathbf{k},j}^r(t) \alpha_{\mathbf{k},j}^r + v_{-\mathbf{k},j}^r(t) \beta_{-\mathbf{k},j}^{r\dagger} \right] e^{i\mathbf{k}\cdot\mathbf{x}}, \quad (70)$$

with  $x \equiv (\mathbf{x}, t)$ ,  $u_{\mathbf{k},j}^r(t) = e^{-i\omega_{\mathbf{k},j}t} u_{\mathbf{k},j}^r$ ,  $v_{\mathbf{k},j}^r(t) = e^{i\omega_{\mathbf{k},j}t} v_{\mathbf{k},j}^r$ , and  $\omega_{\mathbf{k},j} = \sqrt{|\mathbf{k}|^2 + m_j^2}$ . The ladder operators  $\alpha_{\mathbf{k},j}^r$  and  $\beta_{\mathbf{k},j}^r$  annihilate the massive neutrino vacuum  $|0\rangle_{1,2}$ , i.e.,

$$\alpha_{\mathbf{k},j}^r |0\rangle_{1,2} = 0 = \beta_{\mathbf{k},j}^r |0\rangle_{1,2}. \quad (71)$$

The anticommutation relations are

$$\{v_i^\alpha(x), v_j^{\beta\dagger}(y)\}_{t_x=t_y} = \delta^3(\mathbf{x} - \mathbf{y}) \delta_{\alpha\beta} \delta_{ij}, \quad (72)$$

$$\{\alpha_{\mathbf{k},i}^r, \alpha_{\mathbf{q},j}^{s\dagger}\} = \delta_{\mathbf{kq}} \delta_{rs} \delta_{ij}, \quad \{\beta_{\mathbf{k},i}^r, \beta_{\mathbf{q},j}^{s\dagger}\} = \delta_{\mathbf{kq}} \delta_{rs} \delta_{ij}, \quad (73)$$

while the orthonormality and completeness relations are fixed as

$$u_{\mathbf{k},i}^{r\dagger} u_{\mathbf{k},i}^s = v_{\mathbf{k},i}^{r\dagger} v_{\mathbf{k},i}^s = \delta_{rs}, \quad u_{\mathbf{k},i}^{r\dagger} v_{-\mathbf{k},i}^s = 0, \quad \sum_r (u_{\mathbf{k},i}^{r\alpha*} u_{\mathbf{k},i}^{r\beta} + v_{-\mathbf{k},i}^{r\alpha*} v_{-\mathbf{k},i}^{r\beta}) = \delta_{\alpha\beta}. \quad (74)$$

Similarly to the case of unstable particles, we define the number operator for flavor (Pontecorvo) neutrinos as

$$N_{P,\sigma}(t) = \sum_{\mathbf{k},r} \alpha_{P,\mathbf{k},\sigma}^{r\dagger}(t) \alpha_{P,\mathbf{k},\sigma}^r(t), \quad (75)$$

where the flavor ladder operators  $\alpha_{P,\mathbf{k},\sigma}^r$  are just linear combinations of the mass operators

$$\alpha_{P,\mathbf{k},\sigma}^r = \sum_{j=1}^2 U_{\sigma j} \alpha_{\mathbf{k},j}^r. \quad (76)$$

Then, Pontecorvo flavor states can be built as

$$|v_{\mathbf{k},\sigma}^r\rangle_P = \alpha_{P,\mathbf{k},\sigma}^{r\dagger} |0\rangle_{1,2}, \quad (77)$$

which coincide with Equation (42) of the QM treatment when one assumes equal momenta.

The flavor survival probability can be thus computed as the expectation value of the number operator on a reference flavor state

$$\mathcal{P}_{\sigma \rightarrow \sigma}(t) = \langle N_{P,\sigma}(t) \rangle_{\sigma} = 1 - \sin^2(2\theta) \sin^2\left(\frac{\Omega_{\mathbf{k}}^- t}{2}\right), \tag{78}$$

where, in this subsection,  $\langle \dots \rangle_{\sigma} = {}_P \langle v_{\mathbf{k},\sigma}^r | \dots | v_{\mathbf{k},\sigma}^r \rangle_P$ . Moreover, we have introduced the shorthand notation  $\Omega_{\mathbf{k}}^{\pm} \equiv \omega_{\mathbf{k},1} \pm \omega_{\mathbf{k},2}$ .

It is clear that the number of flavor neutrinos is a good candidate for being clock-observable, i.e.,  $O(t) = N_{P,\sigma}(t)$ . Noting that

$$\sigma_N^2 = \langle N_{P,\sigma}^2(t) \rangle_{\sigma} - \langle N_{P,\sigma}(t) \rangle_{\sigma}^2 = \mathcal{P}_{\sigma \rightarrow \sigma}(t)(1 - \mathcal{P}_{\sigma \rightarrow \sigma}(t)),$$

we get

$$\left| \frac{d\mathcal{P}_{\sigma \rightarrow \sigma}(t)}{dt} \right| \leq 2\Delta E \sqrt{\mathcal{P}_{\sigma \rightarrow \sigma}(t)(1 - \mathcal{P}_{\sigma \rightarrow \sigma}(t))}, \tag{79}$$

which is in fact equivalent to Equation (36). An important remark is that  $\sigma_N$  quantifies the dynamic flavor entanglement of the neutrino state (77) (see [78,79]) and it naturally appears in the study of Leggett–Garg inequalities in the context of neutrino oscillations [80,81].

The inequality (79) can be made simpler by following similar steps to those below Equation (32). In so doing, we arrive to

$$\Delta E T \geq \mathcal{P}_{\sigma \rightarrow \rho}(T), \quad \sigma \neq \rho, \tag{80}$$

where  $\mathcal{P}_{\sigma \rightarrow \rho}(t) = 1 - \mathcal{P}_{\sigma \rightarrow \sigma}(t)$  is the flavor oscillation probability at time  $t$ . For  $T = T_h$  such that  $\mathcal{P}_{\sigma \rightarrow \rho}(T_h) = \frac{1}{2}$ , we finally get the analog of Equation (34) for neutrino oscillations.

#### 4.2. Flavor Fock Space Approach to Neutrino Mixing and Oscillations

Consider the weak decay of a  $W^+$  boson,  $W^+ \rightarrow e^+ + \nu_e$ . This process can be described by the Lagrangian  $\mathcal{L} = \mathcal{L}_\nu + \mathcal{L}_l + \mathcal{L}_{int}$  with

$$\mathcal{L}_\nu = \bar{\nu}(i\gamma_\mu \partial^\mu - M_\nu)\nu, \tag{81}$$

$$\mathcal{L}_l = \bar{l}(i\gamma_\mu \partial^\mu - M_l)l, \tag{82}$$

$$\mathcal{L}_{int} = \frac{g}{2\sqrt{2}} \left[ W_\mu^+ \bar{\nu} \gamma^\mu (1 - \gamma^5) l + h.c. \right], \tag{83}$$

where  $\nu = (\nu_e, \nu_\mu)^T$ ,  $l = (e, \mu)^T$ , and

$$M_\nu = \begin{pmatrix} m_e & m_{e\mu} \\ m_{e\mu} & m_\mu \end{pmatrix}, \quad M_l = \begin{pmatrix} \tilde{m}_e & 0 \\ 0 & \tilde{m}_\mu \end{pmatrix}. \tag{84}$$

The off-diagonal components of  $M_\nu$  give rise to the neutrino mixing terms in  $\mathcal{L}_\nu$ . These terms can be diagonalized by means of the field mixing transformation [60–64]

$$\nu_\sigma(x) = \sum_j U_{\sigma j} \nu_j(x), \quad \sigma = e, \mu; \quad j = 1, 2, \tag{85}$$

where  $U$  is the mixing matrix

$$U = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}, \tag{86}$$

with  $\tan 2\theta = 2m_{e\mu}/(m_\mu - m_e)$ , so that

$$\begin{aligned} \mathcal{L} &= \sum_j \bar{\nu}_j (i\gamma_\mu \partial^\mu - m_j) \nu_j + \sum_\sigma \bar{l} (i\gamma_\mu \partial^\mu - \tilde{m}_\sigma) l \\ &+ \frac{g}{2\sqrt{2}} \sum_{\sigma,j} \left[ W_\mu^+(x) \bar{\nu}_j U_{j\sigma}^* \gamma^\mu (1 - \gamma^5) l_\sigma + h.c. \right]. \end{aligned} \tag{87}$$

This fact led to the development of the idea that neutrino flavor states can be defined as linear combinations of neutrino mass states, as discussed in the previous sections. However, the field mixing transformation (85) does not imply the definition (77). Then, the question arises as to how to consistently define the flavor states  $|\nu_\sigma\rangle$  in QFT.

In order to reveal this, we start from the observation that the Lagrangian  $\mathcal{L}$  is invariant under the action of the global  $U(1)$  transformations  $\nu \rightarrow e^{i\alpha} \nu$  and  $l \rightarrow e^{i\alpha} l$ , leading to the conservation of the total flavor charge  $Q_l^{tot}$  corresponding to the lepton-number conservation [64]. This can be written in terms of the flavor charges for neutrinos and charged leptons

$$Q_l^{tot} = \sum_{\sigma=e,\mu} Q_\sigma^{tot}(t), \quad Q_\sigma^{tot}(t) = Q_{\nu_\sigma}(t) + Q_\sigma, \tag{88}$$

with

$$\begin{aligned} Q_e &= \int d^3\mathbf{x} e^\dagger(x) e(x), & Q_{\nu_e}(t) &= \int d^3\mathbf{x} \nu_e^\dagger(x) \nu_e(x), \\ Q_\mu &= \int d^3\mathbf{x} \mu^\dagger(x) \mu(x), & Q_{\nu_\mu}(t) &= \int d^3\mathbf{x} \nu_\mu^\dagger(x) \nu_\mu(x). \end{aligned} \tag{89}$$

Noticing that  $[\mathcal{L}_{int}(\mathbf{x}, t), Q_\sigma^{tot}(t)] = 0$ , we see that neutrinos are produced and detected with a definite flavor [6,66]. However,  $[(\mathcal{L}_\nu + \mathcal{L}_l)(\mathbf{x}, t), Q_\sigma^{tot}(t)] \neq 0$ , leading to the flavor oscillation phenomenon.

The second key observation is that the field mixing transformation (85) can be exactly rewritten as [11,70]

$$\nu_e^\alpha(x) = G_\theta^{-1}(t) \nu_1^\alpha(x) G_\theta(t), \tag{90}$$

$$\nu_\mu^\alpha(x) = G_\theta^{-1}(t) \nu_2^\alpha(x) G_\theta(t), \tag{91}$$

where the *mixing generator* reads

$$G_\theta(t) = \exp[\theta(S_+(t) - S_-(t))], \tag{92}$$

$$S_+(t) \equiv \int d^3\mathbf{x} \nu_1^\dagger(x) \nu_2(x), \quad S_-(t) \equiv \int d^3\mathbf{x} \nu_2^\dagger(x) \nu_1(x). \tag{93}$$

In fact, from the above equations we get, e.g., for  $\nu_e$

$$\frac{d^2}{d\theta^2} \nu_e^\alpha = -\nu_e^\alpha, \tag{94}$$

with the initial conditions

$$\nu_e^\alpha|_{\theta=0} = \nu_1^\alpha, \quad \left. \frac{d}{d\theta} \nu_e^\alpha \right|_{\theta=0} = \nu_2^\alpha. \tag{95}$$

The key point is that the vacuum  $|0\rangle_{1,2}$  is not invariant under the action of the mixing generator  $G_\theta(t)$ . In fact, one has

$$|0(t)\rangle_{e,\mu} \equiv G_\theta^{-1}(t) |0\rangle_{1,2} = e^{-\theta(S_+(t) - S_-(t))} |0\rangle_{1,2}. \tag{96}$$



The state (96) is known as a *flavor vacuum* because it is annihilated by the flavor ladder operators  $\alpha_\sigma(t)$  and  $\beta_\sigma(t)$  defined by

$$\alpha_e(t)|0(t)\rangle_{e,\mu} \equiv G_\theta^{-1}(t)\alpha_1 G_\theta(t) G_\theta^{-1}(t)|0\rangle_{1,2} = 0, \quad (97)$$

and similarly for  $\beta_\sigma(t)$ . Their explicit form is

$$\alpha_{\mathbf{k},e}^r(t) = \cos\theta \alpha_{\mathbf{k},1}^r + \sin\theta \left( U_{\mathbf{k}}^*(t) \alpha_{\mathbf{k},2}^r + \epsilon^r V_{\mathbf{k}}(t) \beta_{-\mathbf{k},2}^{r\dagger} \right), \quad (98)$$

$$\alpha_{\mathbf{k},\mu}^r(t) = \cos\theta \alpha_{\mathbf{k},2}^r - \sin\theta \left( U_{\mathbf{k}}(t) \alpha_{\mathbf{k},1}^r - \epsilon^r V_{\mathbf{k}}(t) \beta_{-\mathbf{k},1}^{r\dagger} \right), \quad (99)$$

$$\beta_{-\mathbf{k},e}^r(t) = \cos\theta \beta_{-\mathbf{k},1}^r + \sin\theta \left( U_{\mathbf{k}}^*(t) \beta_{-\mathbf{k},2}^r - \epsilon^r V_{\mathbf{k}}(t) \alpha_{\mathbf{k},2}^{r\dagger} \right), \quad (100)$$

$$\beta_{-\mathbf{k},\mu}^r(t) = \cos\theta \beta_{-\mathbf{k},2}^r - \sin\theta \left( U_{\mathbf{k}}(t) \beta_{-\mathbf{k},1}^r + \epsilon^r V_{\mathbf{k}}(t) \alpha_{\mathbf{k},1}^{r\dagger} \right). \quad (101)$$

In these equations, we have defined  $\epsilon^r \equiv (-1)^r$ , while  $U_{\mathbf{k}}$  and  $V_{\mathbf{k}}$  are the *Bogoliubov coefficients*

$$U_{\mathbf{k}}(t) \equiv u_{\mathbf{k},2}^{r\dagger} u_{\mathbf{k},1}^r e^{i(\omega_{\mathbf{k},2}-\omega_{\mathbf{k},1})t} = |U_{\mathbf{k}}| e^{i(\omega_{\mathbf{k},2}-\omega_{\mathbf{k},1})t}, \quad (102)$$

$$V_{\mathbf{k}}(t) \equiv \epsilon^r u_{\mathbf{k},1}^{r\dagger} v_{-\mathbf{k},2}^r e^{i(\omega_{\mathbf{k},2}+\omega_{\mathbf{k},1})t} = |V_{\mathbf{k}}| e^{i(\omega_{\mathbf{k},2}+\omega_{\mathbf{k},1})t}, \quad (103)$$

where the time-independent part of the coefficients is given by

$$\begin{aligned} |U_{\mathbf{k}}| &\equiv u_{\mathbf{k},2}^{r\dagger} u_{\mathbf{k},1}^r = v_{-\mathbf{k},1}^{r\dagger} v_{-\mathbf{k},2}^r \\ &= \left( \frac{\omega_{\mathbf{k},1} + m_1}{2\omega_{\mathbf{k},1}} \right)^{\frac{1}{2}} \left( \frac{\omega_{\mathbf{k},2} + m_2}{2\omega_{\mathbf{k},2}} \right)^{\frac{1}{2}} \left( 1 + \frac{\mathbf{k}^2}{(\omega_{\mathbf{k},1} + m_1)(\omega_{\mathbf{k},2} + m_2)} \right), \end{aligned} \quad (104)$$

$$\begin{aligned} |V_{\mathbf{k}}| &= \epsilon^r u_{\mathbf{k},1}^{r\dagger} v_{-\mathbf{k},2}^r = -\epsilon^r u_{\mathbf{k},2}^{r\dagger} v_{-\mathbf{k},1}^r \\ &= \frac{|\mathbf{k}|}{\sqrt{4\omega_{\mathbf{k},1}\omega_{\mathbf{k},1}}} \left( \sqrt{\frac{\omega_{\mathbf{k},2} + m_2}{\omega_{\mathbf{k},1} + m_1}} - \sqrt{\frac{\omega_{\mathbf{k},1} + m_1}{\omega_{\mathbf{k},2} + m_2}} \right). \end{aligned} \quad (105)$$

Notice that  $|U_{\mathbf{k}}|^2 + |V_{\mathbf{k}}|^2 = 1$ . It is straightforward to check that in the relativistic limit,  $\omega_{\mathbf{k},j} \approx |\mathbf{k}|$ ,  $|U_{\mathbf{k}}| \rightarrow 1$  and  $|V_{\mathbf{k}}| \rightarrow 0$ . Furthermore,  $|V_{\mathbf{k}}| = 0$  when  $m_1 = m_2$  and/or  $\theta = 0$ , i.e., when no mixing occurs.  $|V_{\mathbf{k}}|^2$  reaches a maximum at  $|\mathbf{k}| = \sqrt{m_1 m_2}$  with  $|V_{\mathbf{k}}|_{max}^2 \rightarrow 1/2$  for  $\frac{(m_2-m_1)^2}{m_1 m_2} \rightarrow \infty$ , and  $|V_{\mathbf{k}}|^2 \simeq \frac{(m_2-m_1)^2}{4|\mathbf{k}|^2}$  for  $|\mathbf{k}| \gg \sqrt{m_1 m_2}$  at the first non-vanishing order.

The flavor fields can be thus expanded as

$$v_e(x) = \sum_{\mathbf{k},r} \frac{e^{i\mathbf{k}\cdot\mathbf{x}}}{\sqrt{V}} \left[ u_{\mathbf{k},1}^r(t) \alpha_{\mathbf{k},e}^r(t) + v_{-\mathbf{k},1}^r(t) \beta_{-\mathbf{k},e}^{r\dagger}(t) \right], \quad (106)$$

$$v_\mu(x) = \sum_{\mathbf{k},r} \frac{e^{i\mathbf{k}\cdot\mathbf{x}}}{\sqrt{V}} \left[ u_{\mathbf{k},2}^r(t) \alpha_{\mathbf{k},\mu}^r(t) + v_{-\mathbf{k},2}^r(t) \beta_{-\mathbf{k},\mu}^{r\dagger}(t) \right], \quad (107)$$

where  $\sum_{\mathbf{k},r}$  is a simplified notation for the superposition over all field modes of momentum  $\mathbf{k}$  and polarization  $r$ .

A *flavor Hilbert space* (at some reference time, say,  $t = 0$ ) is defined as

$$\mathcal{H}_{e,\mu} = \left\{ \alpha_{e,\mu}^\dagger, \beta_{e,\mu}^\dagger, |0\rangle_{e,\mu} \right\}, \quad (108)$$

with  $|0\rangle_{e,\mu} \equiv |0(t = 0)\rangle_{e,\mu}$ . This is a different Hilbert space with respect to the one of mass-neutrino states. In fact, one can verify that

$$\lim_{V \rightarrow \infty} {}_{1,2} \langle 0|0(t)\rangle_{e,\mu} = \lim_{V \rightarrow \infty} e^{V \int \frac{d^3\mathbf{k}}{(2\pi)^3} \ln(1 - \sin^2 \theta |V_{\mathbf{k}}|^2)} = 0, \tag{109}$$

i.e., flavor and massive fields belong to unitarily inequivalent representations of the anti-commutation relations.

The previous discussions suggest that flavor states  $|\nu_{\mathbf{k},\sigma}^r\rangle$  can be built as one particle states of the flavor Fock space

$$|\nu_{\mathbf{k},\sigma}^r\rangle = \alpha_{\mathbf{k},\sigma}^{r+} |0\rangle_{e,\mu}, \tag{110}$$

and similarly for the antineutrino ( $|\beta_{\mathbf{k},\sigma}^r\rangle \equiv \beta_{\mathbf{k},\sigma}^{r+} |0\rangle_{e,\mu}$ ). One can prove that these states are exact eigenstates of the charge operators at the reference (production/detection) time, i.e.,

$$Q_{\nu_\sigma}(0) |\nu_{\mathbf{k},\sigma}^r\rangle = |\nu_{\mathbf{k},\sigma}^r\rangle. \tag{111}$$

In this approach, the flavor oscillation probability is computed by taking the expectation value of the time-dependent flavor charges with respect to a reference time flavor state [82]

$$Q_{\sigma \rightarrow \rho}(t) = \langle Q_{\nu_\rho}(t) \rangle_\sigma, \tag{112}$$

where  $\langle \dots \rangle_\sigma \equiv \langle \nu_{\mathbf{k},\sigma}^r | \dots | \nu_{\mathbf{k},\sigma}^r \rangle$ , which gives

$$Q_{\sigma \rightarrow \rho}(t) = \sin^2(2\theta) \left[ |U_{\mathbf{k}}|^2 \sin^2\left(\frac{\Omega_{\mathbf{k}}^- t}{2}\right) + |V_{\mathbf{k}}|^2 \sin^2\left(\frac{\Omega_{\mathbf{k}}^+ t}{2}\right) \right], \quad \sigma \neq \rho, \tag{113}$$

$$Q_{\sigma \rightarrow \sigma}(t) = 1 - Q_{\sigma \rightarrow \rho}(t), \quad \sigma \neq \rho. \tag{114}$$

Notice the presence of the term proportional to  $|V_{\mathbf{k}}|^2$  in the oscillation probability Equation (113), which introduces fast oscillations that are not present in the usual QM formula (see Equation (78)). As already mentioned,  $|V_{\mathbf{k}}|^2 \rightarrow 0$  in the relativistic limit  $|\mathbf{k}| \gg m_j$ ,  $j = 1, 2$ , and the oscillation formula reduces to the standard result, as it should be. It has been proven in the simple case of scalar field mixing that the above oscillation formula is the time component of a Lorentz-covariant formula, though the flavor vacuum breaks the Lorentz invariance [83]. Furthermore, connections between implications of the QFT treatment of mixing and extended (Tsallis-like) statistics have been explored in [84].

#### 4.3. Time–Energy Uncertainty Relation for Neutrino Oscillations in QFT

It is clear that in the QFT treatment, lepton charges are natural candidates for clock observables. In fact, starting from

$$[Q_{\nu_\sigma}(t), H] = i \frac{dQ_{\nu_\sigma}(t)}{dt} \neq 0, \tag{115}$$

we find the *flavor–energy* uncertainty relation

$$\sigma_H \sigma_Q \geq \frac{1}{2} \left| \frac{dQ_{\sigma \rightarrow \sigma}(t)}{dt} \right|. \tag{116}$$

Proceeding as in the Pontecorvo case, one finds

$$\left| \frac{dQ_{\sigma \rightarrow \sigma}(t)}{dt} \right| \leq \Delta E. \tag{117}$$

From Equation (116), we get the simple form for TEUR

$$\Delta E T \geq \mathcal{Q}_{\sigma \rightarrow \rho}(T), \quad \sigma \neq \rho. \tag{118}$$

When  $m_i/|\mathbf{k}| \rightarrow 0$ , i.e., in the relativistic case, we get

$$|U_{\mathbf{k}}|^2 \approx 1 - \varepsilon(\mathbf{k}), \quad |V_{\mathbf{k}}|^2 \approx \varepsilon(\mathbf{k}), \tag{119}$$

with  $\varepsilon(\mathbf{k}) \equiv (m_1 - m_2)^2/4|\mathbf{k}|^2$ . In the same limit

$$\Omega_{\mathbf{k}}^- \approx \frac{\delta m^2}{4|\mathbf{k}|} = \frac{\pi}{L^{osc}}, \quad \Omega_{\mathbf{k}}^+ \approx |\mathbf{k}|. \tag{120}$$

Evaluating the inequality (118) at the leading order and for  $T \approx L = L^{osc}/2$ , one finds

$$\Delta E \geq \frac{2 \sin^2(2\theta)}{L^{osc}}. \tag{121}$$

Then, in such a limit, we recover the result (46). As we said above, such a neutrino TEUR is usually regarded as a condition for flavor oscillations to occur [5,85]. In other words, if one managed to measure with a great accuracy the neutrino energy/masses, it could be inferred which massive neutrino was produced in the weak interaction. Then, the oscillation would not occur, as remarked by Bilenky et al. [7] (see also the discussion on Mössbauer neutrinos in Section 2.1). This reasoning is based on the idea that flavor neutrinos are just a superposition of the “physical” massive neutrinos. However, the QFT approach bring us to a different interpretation of the inequality (121). In fact, note that Equation (109) implies

$$\lim_{V \rightarrow \infty} \langle v_{\mathbf{k},i}^r | v_{\mathbf{k},\sigma}^r \rangle = 0, \quad i = 1, 2, \tag{122}$$

i.e., neutrino flavor eigenstates, which are produced in charged current weak decays, *cannot* be generally written as a linear superposition of single-particle massive neutrino states. However, for Pontecorvo states (77), which are a good approximation of the exact flavor states in the relativistic regime, it holds

$$\lim_{V \rightarrow \infty} \langle v_{\mathbf{k},1}^r | v_{\mathbf{k},e}^r \rangle_P = \cos \theta. \tag{123}$$

This apparent contradiction is resolved by observing that

$$\lim_{m_i/|\mathbf{k}| \rightarrow 0} \lim_{V \rightarrow \infty} \neq \lim_{V \rightarrow \infty} \lim_{m_i/|\mathbf{k}| \rightarrow 0}, \tag{124}$$

i.e., the relativistic  $m_i/|\mathbf{k}| \rightarrow 0$  limit cannot be exchanged with the “thermodynamical” limit. The relativistic approximation has to be considered just as a single-particle approximation, which does not take into account the intrinsic multi-particle nature of QFT. The relation (122) should be thus understood as

$$\langle v_{\mathbf{k},i}^r | v_{\mathbf{k},\sigma}^r \rangle = {}_{1,2} \langle 0_{\mathbf{k}} | a_{\mathbf{k},1}^r a_{\mathbf{k},e}^{r\dagger} | 0_{\mathbf{k}} \rangle_{e,\mu} \prod_{\mathbf{p} \neq \mathbf{k}} {}_{1,2} \langle 0_{\mathbf{p}} | 0_{\mathbf{p}} \rangle_{e,\mu}, \tag{125}$$

where we have used the fact that the Hilbert spaces for both massive and flavor fields have a tensor product structure [86]. The first factor on the r.h.s. corresponds to (123), and it is finite. However, as said above, this corresponds to a selection of a single particle sub-space from the complete Hilbert space. In other words, beyond the QM single particle view, the Pontecorvo definition of neutrino state does not work anymore. Then, the inequality (121) should be regarded as a fundamental bound on the energy precision that can be reached experimentally: as in the case of unstable particles, flavor neutrinos have an intrinsic energy spread that is related to their “lifetime”, i.e.,  $L^{osc}$ . From this perspective, observations of

Mössbauer neutrinos could not occur, since the required energy uncertainty would be smaller than the minimum value allowed by TEUR (see the discussion in Section 2.1).

Let us now consider the exact oscillation formula (113) at the first order in  $\varepsilon(\mathbf{k})$ :

$$\mathcal{Q}_{\sigma \rightarrow \rho}(t) \approx \sin^2(2\theta) \left[ \sin^2\left(\frac{\pi t}{L^{osc}}\right) (1 - \varepsilon(\mathbf{k})) + \varepsilon(\mathbf{k}) \sin^2(|\mathbf{k}|t) \right], \quad \sigma \neq \rho. \quad (126)$$

Evaluating the inequality (118) at  $T = L^{osc}/2$  we get

$$\Delta E \geq \frac{2 \sin^2 2\theta}{L^{osc}} \left[ 1 - \varepsilon(\mathbf{k}) \cos^2\left(\frac{|\mathbf{k}|L^{osc}}{2}\right) \right]. \quad (127)$$

We thus find that the energy bound is lowered with respect to (121). If we set  $m_1 = 0.0497 \text{ eV}$ ,  $m_2 = 0.0504 \text{ eV}$  for neutrino masses in the inverted hierarchy [87] and  $|\mathbf{k}| = 1 \text{ MeV}$ , we obtain  $\varepsilon(\mathbf{k}) = 2 \times 10^{-19}$ , which reveals that such a correction is negligible in the relativistic regime and Equation (127) can be naively approximated by the quantum mechanical version of TEUR.

The situation changes if one looks at the non-relativistic regime, e.g.,  $|\mathbf{k}| = \sqrt{m_1 m_2}$ . In such a case,

$$|U_{\mathbf{k}}|^2 = \frac{1}{2} + \frac{\xi}{2} = 1 - |V_{\mathbf{k}}|^2, \quad (128)$$

$$\xi = \frac{2\sqrt{m_1 m_2}}{m_1 + m_2}, \quad (129)$$

and we can rewrite the TEUR (118) in the form

$$\Delta E T \geq \frac{\sin^2 2\theta}{2} [1 - \cos(\tilde{\omega}_1 T) \cos(\tilde{\omega}_2 T) - \xi \sin(\tilde{\omega}_1 T) \sin(\tilde{\omega}_2 T)], \quad (130)$$

where  $\tilde{\omega}_j = \sqrt{m_j(m_1 + m_2)}$ . In order to compare it with the previous cases, we evaluate such an expression at  $T = \tilde{L}^{osc}/4$ , with  $\tilde{L}^{osc} = 4\pi\sqrt{m_1 m_2}/\delta m^2$ , obtaining

$$\Delta E \geq \frac{2 \sin^2 2\theta}{\tilde{L}^{osc}} (1 - \chi). \quad (131)$$

Here, we have introduced the shorthand notation

$$\chi = \xi \sin(\tilde{\omega}_1 \tilde{L}^{osc}/4) \sin(\tilde{\omega}_2 \tilde{L}^{osc}/4) + \cos(\tilde{\omega}_1 \tilde{L}^{osc}/4) \cos(\tilde{\omega}_2 \tilde{L}^{osc}/4). \quad (132)$$

Using the same values for neutrino masses as above, we estimate  $\chi = 0.1$ , which implies that the original bound on the neutrino energy is now decreased by 10%.

## 5. Conclusions and Discussion

We have discussed recent advances in the study of TEUR for neutrino flavor oscillations. In particular, we have investigated how the the original inequality Equation (6) derived by Bilenky changes in the presence of gravity and in a field-theoretical picture, respectively. In the first case and for some specific background metrics, it has been shown that gravitational corrections can be rearranged so as to leave the form of the generalized TEUR unchanged, provided one defines an effective (gravity-dependent) oscillation length (see Section 3). Conversely, the QFT formulation of TEUR relies upon the identification of the (non-conserved) flavor charge operator with the clock observable. In the latter context, we have argued that the interpretation of neutrinos as “unstable” particles does naturally emerge, the lifetime being related to the characteristic oscillation length (see the discussion below Equation (125)). In passing, we mention that the analogy of oscillating neutrinos with unstable particles has been pointed out also in a recent study [88], in which the oscillation

formula (113) has been independently obtained by considering mixing as an interaction and employing the usual perturbation expansion in the Dirac picture. In that work, the role of TEUR is fundamental because it requires the use of finite-time QFT in order to study flavor oscillations, which would be otherwise spoiled in the  $S$ -matrix formalism ( $t \rightarrow \infty$ ).

Further aspects are yet to be explored. In a recent analysis [44], neutrino oscillations have been addressed in extended theories of gravity as a testing ground for the violation of the strong equivalence principle (SEP). Potential violation effects manifest themselves in the form of a generalized oscillation length depending on the SEP parameter. By plugging into the gravity-modified TEUR, one could exploit this result to delve deeper into the interplay between neutrino oscillations and SEP, and possibly constrain SEP corrections. Moreover, a link between cosmic-scale phenomena and the QFT properties of field mixing has been analyzed in [89] in connection with the suggestive interpretation of the flavor vacuum condensate as a dark energy candidate (see also [90–92] for other similar dark energy models). It would be interesting to extend our study to this context in the effort to improve the current bounds on neutrino masses through dark energy constraints. In addition, growing interest is being aroused by the study of quantum correlations and quantum coherence in neutrino oscillations [78,93–97]. Specifically, in [95], the complete complementarity relations have been applied to neutrino flavor oscillations to fully characterize the quantumness of such a phenomenon. In line with previous studies, the result has been found that quantum correlations still survive after the complete spatial separation of the wave packets composing a flavor state, revealing that the quantum nature of mixed neutrinos goes beyond the pure flavor oscillations. It is our intent to extend the TEUR paradigm to the quantum-information theoretic analysis of oscillations to explore the time non-classicality of this phenomenon in a relativistic domain. Preliminary results in this direction appear in [81]. These and some other research lines are under active investigation and will be presented elsewhere.

**Author Contributions:** Conceptualization, investigation, methodology, validation, supervision, writing and original draft preparation, G.G.L. and L.S. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Data Availability Statement:** All data used in the present analysis have already been freely released and published by the corresponding research teams. In our text, we properly give all necessary references to these works and hence, no further data deposit is needed.

**Acknowledgments:** G.G.L. acknowledges the Spanish “Ministerio de Universidades” for the award of a Maria Zambrano fellowship and funding received from the European Union—NextGenerationEU. He also acknowledges participation in the COST Association Action CA18108 “Quantum Gravity Phenomenology in the Multimessenger Approach” and LISA Cosmology Working group.

**Conflicts of Interest:** The authors declare no conflict of interest.

## Abbreviations

The following abbreviations are used in this manuscript:

|      |                                  |
|------|----------------------------------|
| TEUR | Time–Energy Uncertainty Relation |
| QFT  | Quantum Field Theory             |
| QM   | Quantum Mechanics                |

## References

1. Bauer, M.; Mello, P.A. The Time-Energy Uncertainty Relation. *Ann. Phys.* **1978**, *111*, 38–60. [[CrossRef](#)]
2. Busch, P. *The Time-Energy Uncertainty Relation*; Springer: Berlin/Heidelberg, Germany, 2001. [[CrossRef](#)]
3. Mandelstam, L.; Tamm, I. The Uncertainty Relation Between Energy and Time in Non-relativistic Quantum Mechanics. *J. Phys. Mosc.* **1945**, *9*, 249. [[CrossRef](#)]
4. Wigner, E.P. On the Time-Energy Uncertainty Relation. In *Special Relativity and Quantum Theory: A Collection of Papers on the Poincaré Group*; Noz, M.E., Kim, Y.S., Eds.; Springer: Dordrecht, The Netherlands, 1988; pp. 199–209. [[CrossRef](#)]

5. Bilenky, S.M. Neutrino oscillations and time-energy uncertainty relation. *arXiv* **2005**, arXiv:hep-ph/0605228.
6. Giunti, C.; Kim, C. *Fundamentals of Neutrino Physics and Astrophysics*; OUP Oxford: Oxford, UK, 2007.
7. Bilenky, S.M.; von Feilitzsch, F.; Potzel, W. Time-energy uncertainty relations for neutrino oscillations and the Mössbauer neutrino experiment. *J. Phys. Nucl. Part. Phys.* **2008**, *35*, 095003. [[CrossRef](#)]
8. Akhmedov, E.K.; Kopp, J.; Lindner, M. On application of the time-energy uncertainty relation to Mossbauer neutrino experiments. *J. Phys. G* **2009**, *36*, 078001. .
9. Blasone, M.; Lambiase, G.; Luciano, G.G.; Petrucciello, L.; Smaldone, L. Time-energy uncertainty relation for neutrino oscillations in curved spacetime. *Class. Quant. Grav.* **2020**, *37*, 155004. [[CrossRef](#)]
10. Blasone, M.; Jizba, P.; Smaldone, L. Flavor-energy uncertainty relations for neutrino oscillations in quantum field theory. *Phys. Rev. D* **2019**, *99*, 016014. [[CrossRef](#)]
11. Blasone, M.; Vitiello, G. Quantum field theory of fermion mixing. *Ann. Phys.* **1995**, *244*, 283–311; Erratum in *Ann. Phys.* **1996**, *249*, 363–364. [[CrossRef](#)]
12. Blasone, M.; Capolupo, A.; Romei, O.; Vitiello, G. Quantum field theory of boson mixing. *Phys. Rev. D* **2001**, *63*, 125015, [[hep-ph/0102048](#)]. [[CrossRef](#)]
13. Bhattacharyya, K. Quantum decay and the Mandelstam-Tamm-energy inequality. *J. Phys. A Math. Gen.* **1983**, *16*, 2993. [[CrossRef](#)]
14. Blasone, M.; Jizba, P.; Smaldone, L. Flavor neutrinos as unstable particles. *J. Phys. Conf. Ser.* **2020**, *1612*, 012004. [[CrossRef](#)]
15. Bilenky, S.M.; von Feilitzsch, F.; Potzel, W. Different Schemes of Neutrino Oscillations in Mossbauer Neutrino Experiment. *arXiv* **2008**, arXiv:0804.3409.
16. Bilenky, S.M.; von Feilitzsch, F.; Potzel, W. Reply to the Comment on ‘On application of the time-energy uncertainty relation to Moessbauer neutrino experiments’ by E Kh Akhmedov, J Kopp and M Lindner. *J. Phys. G* **2009**, *36*, 078002. [[CrossRef](#)]
17. Bilenky, S.M.; von Feilitzsch, F.; Potzel, W. Neutrino Oscillations and Uncertainty Relations. *J. Phys. G* **2011**, *38*, 115002. [[CrossRef](#)]
18. Aliu, E.; Andringa, S.; Aoki, S.; Argyriades, J.; Asakura, K.; Ashie, R.; Berns, H.; Bhang, H.; Blondel, A.; Borghi, S.; et al. Evidence for muon neutrino oscillation in an accelerator-based experiment. *Phys. Rev. Lett.* **2005**, *94*, 081802. [[CrossRef](#)] [[PubMed](#)]
19. Ashie, Y.; Hosaka, J.; Ishihara, K.; Itow, Y.; Kameda, J.; Koshio, Y.; Minamino, A.; Mitsuda, C.; Miura, M.; Moriyama, S.; et al. Evidence for an oscillatory signature in atmospheric neutrino oscillation. *Phys. Rev. Lett.* **2004**, *93*, 101801. [[CrossRef](#)] [[PubMed](#)]
20. Apollonio, M.; Baldini, A.; Bemporad, C.; Caffau, E.; Cei, F.; Declais, Y.; De Kerret, H.; Dieterle, B.; Etenko, A.; George, J.; et al. Limits on neutrino oscillations from the CHOOZ experiment. *Phys. Lett. B* **1999**, *466*, 415–430. [[CrossRef](#)]
21. Kells, W.P.; Schiffer, J.P. Possibility of observing recoilless resonant neutrino absorption. *Phys. Rev. C* **1983**, *28*, 2162–2164. [[CrossRef](#)]
22. Akhmedov, E.K.; Kopp, J.; Lindner, M. Oscillations of Mossbauer neutrinos. *J. High Energy Phys.* **2008**, *2008*, 005. [[CrossRef](#)]
23. Machado, P.A.N.; Nunokawa, H.; Pereira dos Santos, F.A.; Zukanovich Funchal, R. Testing Nonstandard Neutrino Properties with a Mössbauer Oscillation Experiment. *J. High Energy Phys.* **2011**, *11*, 136. [[CrossRef](#)]
24. Raghavan, R.S.; Minic, D.; Takeuchi, T.; Tze, C.H. Using Neutrinos to test the Time-Energy Uncertainty Relation in an Extreme Regime. *arXiv* **2012**, arXiv:1210.5639.
25. Giunti, C.; Kim, C.W.; Lee, J.A.; Lee, U.W. On the treatment of neutrino oscillations without resort to weak eigenstates. *Phys. Rev. D* **1993**, *48*, 4310–4317. [[CrossRef](#)] [[PubMed](#)]
26. Beuthe, M. Oscillations of neutrinos and mesons in quantum field theory. *Phys. Rept.* **2003**, *375*, 105–218. [[CrossRef](#)]
27. Naumov, D.V.; Naumov, V.A. Relativistic wave packets in a field theoretical approach to neutrino oscillations. *Russ. Phys. J.* **2010**, *53*, 549–574. [[CrossRef](#)]
28. Naumov, D.V.; Naumov, V.A. A Diagrammatic treatment of neutrino oscillations. *J. Phys. G* **2010**, *37*, 105014. [[CrossRef](#)]
29. Griffiths, D. *Introduction to Quantum Mechanics*; Cambridge University Press: Cambridge, UK, 2017.
30. De Filippo, S.; Vitiello, G. Vacuum Structure for Unstable Particles. *Lett. Nuovo Cim.* **1977**, *19*, 92. [[CrossRef](#)]
31. Umezawa, H.; Matsumoto, H.; Tachiki, M. *Thermo Field Dynamics and Condensed States*; North-Holland Publishing Company: Amsterdam, The Netherlands, 1982.
32. Umezawa, H. *Advanced Field Theory: Micro, Macro, and Thermal Physics*; American Inst. of Physics: Woodbury, NY, USA, 1993.
33. Blasone, M.; Jizba, P.; Luciano, G.G. Unified formalism for Thermal Quantum Field Theories: A geometric viewpoint. *Ann. Phys.* **2018**, *397*, 213–233. [[CrossRef](#)]
34. Israel, W. Thermo field dynamics of black holes. *Phys. Lett. A* **1976**, *57*, 107–110. [[CrossRef](#)]
35. Vitiello, G. Dissipation and memory capacity in the quantum brain model. *Int. J. Mod. Phys. B* **1995**, *9*, 973. [[CrossRef](#)]
36. Ahluwalia, D.V.; Burgard, C. Gravitationally induced quantum mechanical phases and neutrino oscillations in astrophysical environments. *Gen. Rel. Grav.* **1996**, *28*, 1161–1170. [[CrossRef](#)]
37. Cardall, C.Y.; Fuller, G.M. Neutrino oscillations in curved space-time: An Heuristic treatment. *Phys. Rev. D* **1997**, *55*, 7960–7966. [[CrossRef](#)]
38. Blasone, M.; Lambiase, G.; Luciano, G.G.; Petrucciello, L. Neutrino oscillations in accelerated frames. *Europhys. Lett.* **2018**, *124*, 51001. [[CrossRef](#)]
39. Fornengo, N.; Giunti, C.; Kim, C.; Song, J. Gravitational effects on the neutrino oscillation. *Phys. Rev. D* **1997**, *56*, 1895–1902. [[CrossRef](#)]
40. Capozziello, S.; Lambiase, G. Neutrino oscillations in Brans-Dicke theory of gravity. *Mod. Phys. Lett. A* **1999**, *14*, 2193. [[CrossRef](#)]
41. Capozziello, S.; De Laurentis, M. Extended Theories of Gravity. *Phys. Rept.* **2011**, *509*, 167–321. [[CrossRef](#)]



42. Piriz, D.; Roy, M.; Wudka, J. Neutrino oscillations in strong gravitational fields. *Phys. Rev. D* **1996**, *54*, 1587–1599. [[CrossRef](#)] [[PubMed](#)]
43. Lambiase, G. Pulsar kicks induced by spin flavor oscillations of neutrinos in gravitational fields. *Mon. Not. Roy. Astron. Soc.* **2005**, *362*, 867–871. [[CrossRef](#)]
44. Buoninfante, L.; Luciano, G.G.; Petruzzello, L.; Smaldone, L. Neutrino oscillations in extended theories of gravity. *Phys. Rev. D* **2020**, *101*, 024016. [[CrossRef](#)]
45. Dvornikov, M. Neutrino interaction with background matter in a non-inertial frame. *Mod. Phys. Lett. A* **2015**, *30*, 1530017. [[CrossRef](#)]
46. Bhattacharya, T.; Habib, S.; Mottola, E. Gravitationally induced neutrino oscillation phases in static space-times. *Phys. Rev. D* **1999**, *59*, 067301. [[CrossRef](#)]
47. Luciano, G.G.; Petruzzello, L. Testing gravity with neutrinos: From classical to quantum regime. *Int. J. Mod. Phys. D* **2020**, *29*, 2043002. [[CrossRef](#)]
48. Lambiase, G.; Papini, G.; Punzi, R.; Scarpetta, G. Neutrino optics and oscillations in gravitational fields. *Phys. Rev. D* **2005**, *71*, 073011. [[CrossRef](#)]
49. Visinelli, L. Neutrino flavor oscillations in a curved space-time. *Gen. Rel. Grav.* **2015**, *47*, 62. [[CrossRef](#)]
50. Lambiase, G.; Mastrototaro, L.; Visinelli, L. Gravitational waves and neutrino oscillations in Chern-Simons axion gravity. *J. Cosmol. Astropart. Phys.* **2023**, *1*, 11. [[CrossRef](#)]
51. Luciano, G.G.; Blasone, M. Gravitational Effects on Neutrino Decoherence in the Lense–Thirring Metric. *Universe* **2021**, *7*, 417. [[CrossRef](#)]
52. Capolupo, A.; Lambiase, G.; Quaranta, A. Neutrinos in curved spacetime: Particle mixing and flavor oscillations. *Phys. Rev. D* **2020**, *101*, 095022. [[CrossRef](#)]
53. Capolupo, A.; Quaranta, A.; Setaro, P.A. Boson mixing and flavor oscillations in curved spacetime. *Phys. Rev. D* **2022**, *106*, 043013. [[CrossRef](#)]
54. Ahluwalia, D.V. Notes on the kinematic structure of the three flavor neutrino oscillation framework. *Int. J. Mod. Phys. A* **1997**, *12*, 5081–5102. [[CrossRef](#)]
55. DeWitt, B.S. Quantum Field Theory in Curved Space-Time. *Phys. Rept.* **1975**, *19*, 295–357. [[CrossRef](#)]
56. Kodama, H. Conserved Energy Flux for the Spherically Symmetric System and the Back Reaction Problem in the Black Hole Evaporation. *Prog. Theor. Phys.* **1980**, *63*, 1217. [[CrossRef](#)]
57. Ciufolini, I.; Wheeler, J. *Gravitation and Inertia*; Princeton Series in Physics; Princeton University Press: Princeton, NJ, USA, 1995.
58. Sciamia, D.W.; Candelas, P.; Deutsch, D. Quantum field theory, horizons and thermodynamics. *Adv. Phys.* **1981**, *30*, 327–366. [[CrossRef](#)]
59. Misner, C.W.; Thorne, K.S.; Wheeler, J.A. *Gravitation*; W. H. Freeman: San Francisco, CA, USA, 1973.
60. Gribov, V.N.; Pontecorvo, B. Neutrino astronomy and lepton charge. *Phys. Lett. B* **1969**, *28*, 493. [[CrossRef](#)]
61. Bilenky, S.M.; Pontecorvo, B. Quark-Lepton Analogy and Neutrino Oscillations. *Phys. Lett. B* **1976**, *61*, 248. [[CrossRef](#)]
62. Bilenky, S.M.; Pontecorvo, B. Again on Neutrino Oscillations. *Lett. Nuovo Cim.* **1976**, *17*, 569. [[CrossRef](#)]
63. Bilenky, S.M.; Pontecorvo, B. Lepton Mixing and the Solar Neutrino Puzzle. *Comments Nucl. Part. Phys.* **1977**, *7*, 149–152.
64. Bilenky, S.M.; Petcov, S.T. Massive Neutrinos and Neutrino Oscillations. *Rev. Mod. Phys.* **1987**, *59*, 671; Erratum in *Rev. Mod. Phys.* **1989**, *61*, 169; Erratum in *Rev. Mod. Phys.* **1988**, *60*, 575. [[CrossRef](#)]
65. Mannheim, P.D. Derivation of the formalism for neutrino matter oscillations from the neutrino relativistic field equations. *Phys. Rev. D* **1988**, *37*, 1935–1941. [[CrossRef](#)]
66. Giunti, C.; Kim, C.W.; Lee, U.W. Remarks on the weak states of neutrinos. *Phys. Rev. D* **1992**, *45*, 2414–2420. [[CrossRef](#)]
67. Ho, C.M. On Neutrino Flavor States. *J. High Energy Phys.* **2012**, *12*, 022. [[CrossRef](#)]
68. Lobanov, A.E. Particle quantum states with indefinite mass and neutrino oscillations. *Ann. Phys.* **2019**, *403*, 82–105. [[CrossRef](#)]
69. Fantini, G.; Gallo Rosso, A.; Vissani, F.; Zema, V. Introduction to the Formalism of Neutrino Oscillations. *Adv. Ser. Direct. High Energy Phys.* **2018**, *28*, 37–119. [[CrossRef](#)]
70. Alfinito, E.; Blasone, M.; Iorio, A.; Vitiello, G. Squeezed neutrino oscillations in quantum field theory. *Phys. Lett. B* **1995**, *362*, 91–96. [[CrossRef](#)]
71. Fujii, K.; Habe, C.; Yabuki, T. Note on the field theory of neutrino mixing. *Phys. Rev. D* **1999**, *59*, 113003. [[CrossRef](#)]
72. Hannabuss, K.C.; Latimer, D.C. The quantum field theory of fermion mixing. *J. Phys. A* **2000**, *33*, 1369–1373. [[CrossRef](#)]
73. Fujii, K.; Habe, C.; Yabuki, T. Remarks on flavor-neutrino propagators and oscillation formulas. *Phys. Rev. D* **2001**, *64*, 013011. [[CrossRef](#)]
74. Ji, C.R.; Mishchenko, Y. General theory of quantum field mixing. *Phys. Rev. D* **2002**, *65*, 096015. [[CrossRef](#)]
75. Hannabuss, K.C.; Latimer, D.C. Fermion mixing in quasifree states. *J. Phys. A* **2003**, *36*, L69. [[CrossRef](#)]
76. Lee, C.Y. Interactions and oscillations of coherent flavor eigenstates in beta decay. *Mod. Phys. Lett. A* **2020**, *35*, 2030015. [[CrossRef](#)]
77. Luciano, G.G. On the flavor/mass dichotomy for mixed neutrinos: A phenomenologically motivated analysis based on lepton charge conservation in neutron decay. *Eur. Phys. J. Plus* **2023**, *138*, 83. [[CrossRef](#)]
78. Blasone, M.; Dell’Anno, F.; De Siena, S.; Illuminati, F. Entanglement in neutrino oscillations. *Europhys. Lett.* **2009**, *85*, 50002. [[CrossRef](#)]



79. Alok, A.K.; Banerjee, S.; Sankar, S.U. Quantum correlations in terms of neutrino oscillation probabilities. *Nucl. Phys. B* **2016**, *909*, 65–72. [[CrossRef](#)]
80. Naikoo, J.; Kumari, S.; Banerjee, S.; Pan, A. Probing inequivalent forms of Leggett–Garg inequality in subatomic systems. *J. Phys. G* **2020**, *47*, 095004. [[CrossRef](#)]
81. Blasone, M.; Illuminati, F.; Petruzzello, L.; Simonov, K.; Smaldone, L. No-signaling-in-time as a condition for macrorealism: The case of neutrino oscillations. *Eur. Phys. J. C* **2023**, *83*, 688. [[CrossRef](#)]
82. Blasone, M.; Henning, P.A.; Vitiello, G. The Exact formula for neutrino oscillations. *Phys. Lett. B* **1999**, *451*, 140–145. [[CrossRef](#)]
83. Blasone, M.; Jizba, P.; Mavromatos, N.E.; Smaldone, L. Some nontrivial aspects of Poincaré and CPT invariance of flavor vacuum. *Phys. Rev. D* **2020**, *102*, 025021. [[CrossRef](#)]
84. Luciano, G.G.; Blasone, M. Nonextensive Tsallis statistics in Unruh effect for Dirac neutrinos. *Eur. Phys. J. C* **2021**, *81*, 995. [[CrossRef](#)]
85. Kayser, B. On the quantum mechanics of neutrino oscillation. *Phys. Rev. D* **1981**, *24*, 110–116. [[CrossRef](#)]
86. Berezin, J.; Berezin, F.; Firm, E.S.T.; Mugibayashi, N.; Jeffrey, A. *The Method of Second Quantization*; Pure and Applied Physics: A Series of Monographs and Textbooks. 24; Academic Press: Cambridge, MA, USA, 1966.
87. Tanabashi, M.; Grp, P.D.; Hagiwara, K.; Hikasa, K.; Nakamura, K.; Sumino, Y.; Takahashi, F.; Tanaka, J.; Agashe, K.; Aielli, G.; et al. Review of Particle Physics. *Phys. Rev. D* **2018**, *98*, 030001. [[CrossRef](#)]
88. Blasone, M.; Giacosa, F.; Smaldone, L.; Torrieri, G. Neutrino oscillations in the interaction picture. *Eur. Phys. J. C* **2023**, *83*, 736. [[CrossRef](#)]
89. Capolupo, A.; Quaranta, A. Boson mixing and flavor vacuum in the expanding Universe: A possible candidate for the dark energy. *Phys. Lett. B* **2023**, *840*, 137889. [[CrossRef](#)]
90. Addazi, A.; Capozziello, S.; Gan, Q.; Marcianò, A. Dark energy and neutrino superfluids. *Phys. Dark Univ.* **2022**, *37*, 101102. [[CrossRef](#)]
91. Carrillo González, M.; Liang, Q.; Sakstein, J.; Trodden, M. Neutrino-Assisted Early Dark Energy: Theory and Cosmology. *J. Cosmol. Astropart. Phys.* **2021**, *04*, 063. [[CrossRef](#)]
92. Amiri, H.R.; Salehi, A.; Noroozi, A.H. Constraining neutrino mass in dark energy dark matter interaction and comparison with 2018 Planck results. *Eur. Phys. J. C* **2021**, *81*, 479. [[CrossRef](#)]
93. Blasone, M.; Dell’Anno, F.; De Siena, S.; Illuminati, F. Neutrino flavor entanglement. *Nucl. Phys. B-Proc. Suppl.* **2013**, *237–238*, 320–322. [[CrossRef](#)]
94. Banerjee, S.; Alok, A.K.; Srikanth, R.; Hiesmayr, B.C. A quantum information theoretic analysis of three flavor neutrino oscillations. *Eur. Phys. J. C* **2015**, *75*, 487. [[CrossRef](#)]
95. Bittencourt, V.A.S.V.; Blasone, M.; De Siena, S.; Matrella, C. Complete complementarity relations for quantum correlations in neutrino oscillations. *Eur. Phys. J. C* **2022**, *82*, 566. [[CrossRef](#)]
96. Li, Y.W.; Li, L.J.; Song, X.K.; Wang, D.; Ye, L. Geuine tripartite entanglement in three-flavor neutrino oscillations. *Eur. Phys. J. C* **2022**, *82*, 799. [[CrossRef](#)]
97. Wang, G.J.; Li, Y.W.; Li, L.J.; Song, X.K.; Wang, D. Monogamy properties of quantum correlations in neutrino oscillations. *Eur. Phys. J. C* **2023**, *83*, 801. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.