



From geometry to cosmology: a pedagogical review of inflation in curvature, torsion, and extended gravity theories

Davood Momeni^{2,1,a} 

¹ Department of Physics and Pre-Engineering, Northeast Community College, Norfolk, NE 68701, USA

² Centre for Space Research, North-West University, Potchefstroom 2520, South Africa

Received: 29 May 2025 / Accepted: 30 August 2025
© The Author(s) 2025

Abstract We present a comprehensive and pedagogical review of inflationary cosmology in the context of a wide spectrum of modified theories of gravity. This includes curvature-based frameworks such as $f(R)$ gravity, Gauss–Bonnet gravity $f(G)$, and mixed curvature invariants like $f(R, G)$; torsion-based models including teleparallel gravity $f(T)$ and Einstein–Cartan theory; and non-metricity-inspired theories such as symmetric teleparallel gravity $f(Q)$. Scalar–tensor extensions are systematically explored where scalar fields couple not only to the Ricci scalar R , but also to torsion T , non-metricity Q , the Gauss–Bonnet invariant G , and the Ricci tensor $R_{\mu\nu}$. For each class of models, we derive and analyze inflationary dynamics under slow-roll approximations, highlighting the effects of geometry-induced modifications to the inflaton field equations. We examine background evolution, reheating mechanisms, and perturbation spectra, with explicit focus on observables including the spectral index n_s , the tensor-to-scalar ratio r , running α_s , and non-Gaussianity f_{NL} . Numerical predictions are compared with Planck 2018, BICEP/Keck, and upcoming CMB-S4 forecasts. The review includes auxiliary formulations such as conformal transformations, dynamical system reconstructions, and effective field theory embeddings. We explore both analytical solutions and numerical phase-space portraits, employing Bayesian model selection, machine learning classifiers, and Hubble-flow reconstruction techniques to compare the viability of models across observational parameter space. Special attention is paid to exotic inflationary scenarios motivated by string theory (e.g., brane inflation, axion monodromy), extra-dimensional constructions, mimetic gravity, non-local infinite-derivative models, and theories with varying fundamental constants à la Bekenstein. We also assess the formation of primordial black holes and stochastic gravitational waves as potential discriminants

among models. This work aims to serve as both a detailed theoretical resource and a phenomenological guide, offering an integrated, high-precision roadmap for navigating the inflationary landscape beyond general relativity. By connecting modified gravity frameworks with cosmological data and forward-looking missions (e.g., LiteBIRD, LISA), we seek to elucidate the next generation of viable inflationary paradigms.

Contents

1	Introduction
2	Observational constraints on inflation
3	$f(R)$ gravity and Starobinsky inflation
4	Extended $f(R)$ models
5	$f(G)$ Gauss–Bonnet inflation
6	$f(R, G)$ mixed models
7	$f(T)$ torsion gravity
8	Scalar field coupled to Ricci scalar
9	Scalar field coupled to Ricci tensor
10	Scalar field coupled to torsion tensor
11	Scalar field coupled to Gauss–Bonnet term
12	Einstein–Cartan models of inflation
13	Mimetic gravity and inflation
14	Non-local gravity and infinite derivative models
15	Palatini formalism in inflation
16	Metric-affine and $f(Q)$ gravity
17	Dynamical system approaches to modified gravity inflation
18	Reheating and preheating in modified gravity
19	Primordial black holes and gravitational waves in modified gravity
20	Inflation in braneworld and higher-dimensional theories

^a e-mail: dmomeni@northeast.edu (corresponding author)

21	Bayesian model selection and machine learning approaches to inflation
22	Summary of viable inflationary models in modified gravity
23	Tachyon inflation in modified gravity
24	Inflation in string-inspired gravity
25	Inflation in conformal and Weyl-invariant gravity
26	Phenomenological models of inflation
27	Inflation in MOdified Gravity with anisotropic tensor (MOG)
28	Inflation in cosmological relativity: Carmeli's theory
29	Inflationary insights from Jacob Bekenstein's theoretical frameworks
30	Inflation and extra dimensions
31	Comparative assessment and discriminators among models
32	Open problems and future directions in inflationary modified gravity
33	Open questions and future directions
34	Discussion and outlook
	Appendix
	Appendix A: Reheating and preheating in modified gravity
	Appendix B: Conformal transformations in modified gravity
	Appendix C: Perturbation equations in modified gravity
	Appendix D: Inflationary observables
	Appendix E: Dynamical system formulation
	Appendix F: Auxiliary function reconstructions
	References

1 Introduction

The theory of cosmic inflation has profoundly transformed our understanding of the early universe. Introduced in the early 1980s to address foundational issues in Big Bang cosmology, inflation posits a brief epoch of accelerated expansion that occurred just after the Planck time. This rapid expansion solves several problems inherent in the standard hot Big Bang model – most notably the horizon, flatness, and monopole problems – while simultaneously providing a mechanism for generating primordial density perturbations that seeded large-scale structure. The earliest proposals by Guth [1], Linde [2], and Albrecht and Steinhardt [3] marked the birth of a new paradigm that has since matured into a cornerstone of modern cosmology. The initial clues pointing toward an inflationary phase in the early universe were provided by precise measurements of temperature anisotropies in the cosmic microwave background (CMB). Landmark observations by COBE [4], WMAP [5], and Planck [6] solidified the inflationary paradigm, offering constraints on the amplitude and scale-dependence of primordial perturbations. The Planck 2018 analysis [7], along with more recent data from BICEP/Keck [8], continues to narrow the parameter

space for inflationary models, particularly favoring those predicting a low tensor-to-scalar ratio r and a nearly scale-invariant spectral index n_s .

The growing precision of observational cosmology, including upcoming surveys such as CMB-S4 [9], demands theoretical frameworks that go beyond canonical scalar field inflation. This has motivated a surge in interest in modified gravity theories, particularly those capable of generating inflation through geometric mechanisms. Starobinsky's seminal $f(R)$ model [10] stands out as an early example of inflation driven purely by higher-order curvature terms. Subsequent developments in Gauss–Bonnet gravity [11], unified frameworks [12], and teleparallel gravity [13] have further diversified the inflationary model space, linking gravitational modifications to observable predictions. These foundational studies form the backbone of this review, which aims to bridge theoretical innovation with empirical viability. Recent advances in observational cosmology have not only sharpened the statistical precision of inflationary parameters but also highlighted the need for theoretical models with minimal fine-tuning. The seven-year WMAP data release provided strong evidence for a flat universe with adiabatic, Gaussian initial conditions, laying groundwork for single-field slow-roll inflation scenarios with a red-tilted spectrum [5]. The Planck 2018 results have further refined these estimates, placing stringent limits on inflationary parameters such as the scalar spectral index and the tensor-to-scalar ratio [6,7]. These constraints disfavor many popular inflationary models and have elevated the status of geometrically driven frameworks – particularly those predicting low r values – in the search for realistic early-universe dynamics.

In tandem with data, theoretical innovation has been shaped by the need to embed inflation within a broader gravitational framework. Starobinsky's original $f(R)$ model [10] set the stage for understanding inflation as a manifestation of higher-order curvature corrections. This idea was expanded upon by Nojiri and Odintsov in their development of Gauss–Bonnet and unified modified gravity models [11,12], which further accommodate dark energy and late-time acceleration. The incorporation of torsion into cosmology via $f(T)$ gravity, as extensively reviewed by Cai et al. [13], has opened a new window for formulating inflation without scalar fields or with alternative reheating mechanisms. Meanwhile, forthcoming missions such as CMB-S4 [9] promise to probe gravitational waves and primordial non-Gaussianities with unprecedented sensitivity, enabling the discrimination among competing models and providing key tests for extended theories of gravity. In its simplest form, inflation is driven by a scalar field slowly rolling down a nearly flat potential. This setup naturally generates an almost scale-invariant power spectrum of adiabatic, Gaussian perturbations, a prediction remarkably consistent with observations. The Cosmic Microwave Background (CMB)

anisotropies observed by COBE, WMAP, and more recently, the Planck satellite [6], have provided overwhelming support for these predictions. Yet, despite its success, the simplest models of scalar-field inflation embedded in Einstein gravity face several theoretical challenges, including fine-tuning of the initial conditions, the choice of the inflaton potential, the ultraviolet (UV) sensitivity of scalar fields, and the lack of a natural embedding in high-energy theories.

These challenges have motivated an extensive exploration of inflation within modified gravity frameworks. Such models either generalize the Einstein–Hilbert action or reformulate gravity in a geometrically novel way, for example by employing torsion or non-metricity instead of curvature as the underlying source of gravitational dynamics. Among the earliest and most successful of these is the $f(R)$ theory of gravity, where the Ricci scalar R in the action is replaced by an arbitrary function $f(R)$. The seminal model proposed by Starobinsky [10], featuring an $R + \alpha R^2$ action, not only drives inflation without requiring an inflaton but is also one of the most favored models under current CMB constraints.

Building upon $f(R)$ gravity, further extensions have incorporated additional curvature invariants such as the Gauss–Bonnet term G , leading to $f(G)$ and $f(R, G)$ models. These higher-order theories are often motivated by low-energy effective actions in string theory and offer richer phenomenological dynamics, such as bouncing cosmologies, graceful exit from inflation, and distinctive gravitational wave signatures [11, 12]. Similarly, $f(T)$ gravity, based on the teleparallel formulation where torsion replaces curvature, has emerged as a viable platform for inflationary models with alternative geometric foundations [13]. In $f(R, G)$ models, bounces with future singularities have been classified in detail [14]. A smooth bounce in $f(R)$ gravity accommodating late-time acceleration has been constructed in [15].

Other classes of modified gravity relevant to inflation include Palatini formalism, mimetic gravity, non-local and infinite derivative theories, scalar–tensor theories with non-minimal couplings to Ricci curvature or torsion, and metric-affine models involving non-metricity and connection degrees of freedom. These extensions not only offer diverse dynamical mechanisms to realize inflation but often embed it in a broader cosmological evolution including dark energy, reheating, and late-time acceleration.

Motivation and scope of this work

This review is motivated by the growing need to unify the rapidly diversifying landscape of inflationary models inspired by modifications to general relativity. While several earlier reviews have focused on specific classes – such as $f(R)$ models or Gauss–Bonnet inflation – our aim is to offer a broader, integrative synthesis that places these developments in a comparative and observationally driven context.

We focus on the geometric origins of inflationary mechanisms, emphasizing how curvature, torsion, and non-metricity individually or collectively drive accelerated expansion in the early universe. The scope of this review extends from classical modified gravity theories to more speculative frameworks inspired by string theory, extra dimensions, and quantum gravity corrections.

The overarching goal is to organize the extensive array of inflationary models into a coherent structure that facilitates comparison, evaluation, and future development. Each model class is introduced with historical and theoretical context, followed by analysis of its dynamical behavior and observational viability. Specifically, the categories reviewed include:

- $f(R)$ gravity and Starobinsky-type inflation,
- Gauss–Bonnet and mixed curvature models: $f(G)$ and $f(R, G)$,
- Torsion-based models including $f(T)$ gravity and scalar–torsion interactions,
- Theories with scalar fields non-minimally coupled to curvature invariants,
- Frameworks beyond curvature: mimetic gravity, non-local models, and infinite derivative gravity,
- Non-Riemannian models such as metric-affine gravity and $f(Q)$ theories,
- Palatini and bimetric extensions with independent connection dynamics.

By systematically covering both the theoretical underpinnings and the phenomenological consequences of these models, we aim to provide researchers with a reliable reference for navigating the current and future landscape of inflation in modified gravity.

In each case, we review the essential theoretical setup, provide summary tables of viable parameter ranges, and discuss compatibility with current CMB observations – including constraints on the scalar spectral index n_s and tensor-to-scalar ratio r from Planck [7], BICEP/Keck [8], and forecasts from future missions such as LiteBIRD and CMB-S4 [9].

Our approach

A distinctive aspect of this review is the consistent application of the vielbein (or tetrad) formalism, which enables a geometrically unified treatment of curvature, torsion, and non-metricity. This approach proves particularly advantageous in teleparallel and metric-affine theories, where the geometric degrees of freedom are not limited to curvature alone but also encompass torsional and affine structures.

To capture the dynamical behavior of inflationary scenarios, we utilize dynamical systems techniques to study the phase-space structure and identify inflationary attractors and their stability properties. This allows us to classify models

based on their qualitative evolution and to pinpoint the conditions under which inflation arises naturally within each theory.

In parallel, we investigate post-inflationary dynamics, focusing on reheating and preheating processes and the potential formation of primordial black holes (PBHs) and gravitational waves. These features serve as observational probes into the viability and distinctiveness of each inflationary framework.

A comprehensive comparison is provided in the form of summary tables that assess each model's consistency with observational constraints, including predictions for spectral indices, tensor-to-scalar ratios, non-Gaussianity, and gravitational wave spectra. Bayesian model selection and emerging machine learning tools are explored as powerful frameworks for navigating high-dimensional parameter spaces and accelerating model discrimination.

This integrated methodology is designed to provide both a rigorous theoretical foundation and a practical, data-driven map for future exploration of inflation within modified gravitational theories.

Organization of the paper

This review is structured to guide the reader from observational foundations through theoretical diversity to phenomenological applications. We begin with a discussion of cosmological observations that shape the inflationary landscape, focusing on constraints from the cosmic microwave background and future surveys.

Subsequent sections explore major classes of inflationary models in modified gravity, including those derived from curvature (e.g., $f(R)$ and Gauss–Bonnet terms), torsion-based formulations, and hybrid scenarios that mix geometric ingredients. Special attention is given to scalar fields coupled to various geometric invariants, and to extensions like mimetic and non-local gravity, Einstein–Cartan models, and metric-affine frameworks.

We then turn to the analysis of inflationary dynamics using phase space methods, followed by discussions on post-inflationary physics such as reheating, primordial black hole production, and gravitational waves. The application of Bayesian statistics and machine learning to model evaluation and selection is also introduced.

The review culminates in a synthesis of results, with comparative tables summarizing the viability and distinct predictions of each model class. A concluding discussion highlights open questions and outlines directions for future exploration in inflationary cosmology.

2 Observational constraints on inflation

Inflation is a cornerstone of modern cosmology and, unlike many high-energy theories, it is directly testable through cosmological observables. The inflationary phase leaves clear imprints on the Cosmic Microwave Background (CMB), Large Scale Structure (LSS), and the primordial gravitational wave background, providing a powerful window into the physics of the early universe. High-precision measurements from Planck, BICEP/Keck, and upcoming missions such as LiteBIRD and CMB-S4 offer unprecedented sensitivity to inflationary parameters, including the scalar spectral index n_s , the tensor-to-scalar ratio r , and non-Gaussianity parameters such as f_{NL} .

These observational probes not only allow for discrimination between different inflationary models but also serve as tests of fundamental physics – probing the ultraviolet completion of gravity, the validity of the slow-roll paradigm, and the possible existence of new degrees of freedom. In this context, modified gravity theories such as $f(R)$, scalar–tensor, Gauss–Bonnet, $f(Q)$, and teleparallel formulations have emerged as viable frameworks to realize inflation. These theories often predict novel signatures, including scale-dependent gravitational wave spectra, modified sound speeds, and exotic reheating dynamics.

As cosmological data continue to improve in accuracy and breadth, the interface between inflation and modified gravity provides a unique testing ground for ideas at the intersection of cosmology, quantum gravity, and high-energy theory. In this review, we explore this rich landscape, presenting a comparative and pedagogical analysis of inflationary scenarios derived from extended gravitational actions, with particular attention to their observational viability and theoretical consistency.

CMB temperature and polarization

The primary inflationary observables extracted from the Cosmic Microwave Background (CMB) provide crucial insights into the dynamics of the early universe and the nature of the inflaton field. These include:

- **The scalar spectral index n_s :** This quantifies the deviation from exact scale invariance in the primordial power spectrum of curvature perturbations. A value of $n_s < 1$ indicates a red tilt, meaning that fluctuations on larger scales have slightly higher amplitude than those on smaller scales. This is a hallmark prediction of slow-roll inflation.
- **The running of the spectral index $\alpha_s = \frac{dn_s}{d \ln k}$:** This measures the scale dependence of the spectral tilt itself. Although most simple models predict a very small negative running, a significant detection could point to non-

Table 1 Planck 2018 + BICEP/Keck 2018 constraints (68% CL unless stated otherwise)

Parameter	Value
Scalar spectral index n_s	0.9649 ± 0.0042
Running $\alpha_s \equiv \frac{dn_s}{d \ln k}$	-0.0045 ± 0.0067
Tensor-to-scalar ratio $r_{0.002}$	< 0.056 (95% CL)
Amplitude $\log(10^{10} A_s)$	3.044 ± 0.014

trivial inflaton dynamics or higher-order corrections in the inflationary potential.

- **The tensor-to-scalar ratio r :** This parameter represents the amplitude of primordial gravitational waves relative to scalar perturbations. A detection of $r > 0$ would provide direct evidence of inflationary tensor modes and strongly constrain the energy scale of inflation.

These observables are reconstructed from a combination of temperature anisotropies (TT), E-mode polarization (EE), their cross-correlations (TE), and the B-mode polarization spectrum. The B-modes, in particular, are of paramount importance: while E-modes arise from scalar perturbations, B-modes can only be sourced (at leading order) by tensor perturbations, making them a smoking-gun signature of primordial gravitational waves.

Ongoing and upcoming experiments such as Planck, BICEP/Keck, LiteBIRD, and CMB-S4 are pushing the sensitivity of these measurements to unprecedented levels. They allow us to probe inflationary physics at energy scales near the grand unified theory (GUT) scale and to test whether inflation arises from standard single-field slow-roll models, from multi-field setups, or from more exotic mechanisms including those based on modified gravity.

Constraints from Planck and BICEP/Keck

Planck 2018 data [6, 7], combined with BICEP/Keck 2018 [8], yield the following best-fit values (Table 1):

These constraints disfavor many large-field inflation models with steep potentials (e.g., $V(\phi) \propto \phi^2$), while favoring flatter or plateau-like models that naturally predict low r and acceptable values of n_s . In particular:

- **$f(R)$ (Starobinsky) models** predict $n_s \sim 0.965$, $r \sim 0.003$: *excellent agreement*.
- **$f(T)$ models** can reconstruct viable potentials that fit within current bounds but depend sensitively on the chosen form of $f(T)$.
- **$f(R, G)$ and scalar–Gauss–Bonnet models** may require tuning to avoid ghost modes and satisfy $c_T = c$ from GW170817.

Table 2 Forecasted constraints from LiteBIRD and CMB-S4 (design goals)

Parameter	LiteBIRD	CMB-S4
$\sigma(n_s)$	~ 0.002	< 0.0015
$\sigma(r)$	~ 0.001	~ 0.001
$\sigma(\alpha_s)$	~ 0.003	~ 0.002
Non-Gaussianity f_{NL}	–	~ 1 (local)

Model forecasting: LiteBIRD and CMB-S4

Future missions like **LiteBIRD** and **CMB-S4** aim to improve precision on inflationary observables. Their design sensitivity can probe $r \sim 10^{-3}$, enough to definitively test many geometric inflation models (Table 2).

These missions will help distinguish between low- r models like $f(R)$ and higher- r scenarios such as non-minimally coupled scalar–torsion theories.

Non-Gaussianity and isocurvature constraints

Planck also places stringent bounds on primordial non-Gaussianity, a key discriminator between different classes of inflationary models. The local-type non-Gaussianity parameter is constrained to be:

$$f_{NL}^{local} = -0.9 \pm 5.1 \quad (68\% \text{ CL}), \tag{1}$$

which is fully consistent with zero. This result strongly supports the predictions of single-field, slow-roll inflation, where non-Gaussianities are naturally suppressed due to the near-Gaussian nature of vacuum fluctuations and the approximate shift symmetry of the inflaton.

In contrast, multi-field inflationary scenarios, non-canonical kinetic terms (e.g., DBI inflation), or models involving sharp features in the potential tend to predict larger values of f_{NL} . The absence of detectable non-Gaussianity thus significantly constrains the parameter space of these alternatives, requiring either fine-tuned initial conditions or suppressed interactions between fields.

Additionally, the Planck analysis finds no statistically significant evidence for isocurvature perturbations. The data favor purely adiabatic initial conditions, in which all components of the primordial plasma fluctuate in unison. This poses a challenge to models with multiple uncoupled scalar fields or curvaton-like mechanisms, which generically generate isocurvature modes unless carefully tuned to decay or interact in a way that aligns the perturbations.

Together, these constraints on non-Gaussianity and isocurvature perturbations reinforce the robustness of the standard single-field inflationary paradigm while placing meaningful limits on deviations from this minimal setup.

Model viability summary

To aid comparison, Table 3 summarizes predictions of representative inflationary models within modified gravity frameworks. Each model class exhibits distinct signatures in the scalar spectral index n_s and tensor-to-scalar ratio r , which can be directly compared against the latest cosmological datasets. Starobinsky's $f(R)$ model remains the benchmark for its robust predictions – matching Planck data with $n_s \approx 0.965$ and a very low $r \sim 0.003$. In contrast, simpler power-law extensions of $f(R)$ typically yield too high a tensor amplitude, making them observationally disfavored.

Reconstructed $f(T)$ and $f(Q)$ models offer flexibility through free function choices but require careful tuning to remain within bounds from BICEP/Keck and CMB-S4 forecasts. Scalar–Gauss–Bonnet inflationary models, especially those respecting the gravitational wave speed constraint $c_T = c$, span a range of viable values for r and are actively studied for their natural emergence in string-effective actions.

Chaotic inflation models with quadratic potentials, once dominant in the literature, are now strongly ruled out due to their large predicted tensor modes. This reinforces the preference for plateau-like or asymptotically flat potentials often found in modified gravity models.

Implications for inflationary model building

Current cosmological observations from Planck, BICEP/Keck, and large-scale structure surveys place tight constraints on the inflationary parameter space. The data strongly favor models that yield:

- A scalar spectral index of $n_s \approx 0.96$, consistent with a slight red tilt indicative of scale-dependent suppression of power at small scales,
- A low tensor-to-scalar ratio $r < 0.01$, with an optimal value around $r \sim 10^{-3}$, limiting the energy scale of inflation,
- Negligible non-Gaussianity, supporting nearly Gaussian primordial fluctuations,
- Purely adiabatic initial conditions, with no detectable isocurvature components,
- Tensor perturbations that propagate at luminal speed, $c_T = c$, as required by gravitational wave constraints from GW170817.

These benchmarks act as a filter for theoretical models. Geometrically motivated frameworks such as Starobinsky's $f(R)$ inflation, scalar–torsion reconstructions in teleparallel gravity, and scalar–Gauss–Bonnet couplings that respect the $c_T = 1$ constraint remain highly viable. These models naturally accommodate the observed low values of r , match the

spectral tilt n_s , and avoid exotic features like non-adiabatic or strongly non-Gaussian perturbations.

In contrast, models predicting high tensor amplitudes, such as chaotic inflation with monomial potentials (e.g., $V(\phi) \propto \phi^n$), are increasingly disfavored unless modified or embedded in mechanisms that suppress tensor modes. This motivates ongoing efforts to build inflationary scenarios that balance theoretical naturalness with observational viability – particularly within the extended geometrical frameworks of modified gravity.

3 $f(R)$ gravity and Starobinsky inflation

Modified gravity offers a geometric alternative to conventional scalar-field-driven inflation, broadening the theoretical landscape of early-universe cosmology. Among these frameworks, $f(R)$ gravity stands out as the most minimal and historically influential extension of Einstein's General Relativity. Rather than adding new matter fields, this approach alters the geometric side of Einstein's equations by replacing the Ricci scalar with a more general function. As a result, the theory acquires additional dynamical degrees of freedom, which can naturally drive inflationary expansion in the early universe.

This geometric origin of inflation eliminates the need for exotic scalar potentials or fine-tuned initial conditions. It provides a natural mechanism for both the onset and graceful exit of inflation, often aligning well with observational data. The classic Starobinsky model, a particular case of this framework, remains one of the most successful inflationary scenarios in light of current CMB constraints.

Importantly, modified gravity models like $f(R)$ can also be connected to scalar–tensor theories via suitable mathematical transformations, revealing deeper structural relations. This flexibility allows them to be studied in both the original (Jordan) frame and in a redefined (Einstein) frame where the equations resemble those of standard scalar field inflation. Overall, modified gravity opens up a rich and observationally viable path for understanding cosmic inflation from a fundamentally geometric perspective.

Starobinsky's model: $f(R) = R + \alpha R^2$

The earliest and arguably most successful $f(R)$ inflationary model was proposed by Starobinsky in 1980 [10]. The action is given by:

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \left(R + \alpha R^2 \right), \quad (2)$$

where $\alpha > 0$ is a constant with dimensions of inverse mass squared. At high curvatures ($R \gg 1/\alpha$), the quadratic term dominates, producing a nearly de Sitter phase. As curvature

Table 3 Representative inflationary models and observational status

Model class	n_s	r	Status
Starobinsky $f(R)$	~ 0.965	~ 0.003	Consistent
Power-law $f(R)$	varies	> 0.05	Disfavored
Reconstructed $f(T)$	tunable	tunable	Viable if tuned
Scalar-GB (with $c_T = c$)	0.96–0.97	$10^{-3} - 10^{-2}$	Viable if ghost-free
Chaotic (ϕ^2)	~ 0.96	~ 0.13	Ruled out

decreases, the model smoothly reduces to GR. This built-in transition yields a graceful exit from inflation without additional fields or mechanisms.

The R^2 term generates a scalaron degree of freedom (a massive scalar mode), which slowly rolls along an effective potential. This scalaron is responsible for generating curvature perturbations and tensor fluctuations. Importantly, the dynamics are governed by a single field in the Einstein frame, making the model a minimal and predictive inflationary scenario.

Observationally, Starobinsky inflation predicts a scalar spectral index $n_s \approx 0.965$ and a very low tensor-to-scalar ratio $r \sim 0.003$, both of which align remarkably well with current Planck and BICEP/Keck data. The model also naturally avoids large non-Gaussianities and supports adiabatic initial conditions, satisfying all major CMB constraints.

From a theoretical standpoint, the Starobinsky model is appealing because it arises naturally as the first-order correction to the Einstein–Hilbert action in semiclassical gravity or supergravity frameworks. It has been embedded in higher-derivative supergravity, string-effective actions, and asymptotically safe gravity scenarios, making it a robust candidate for UV completion.

Moreover, reheating in Starobinsky inflation is well understood: the scalaron field oscillates around the GR vacuum and decays gravitationally into light particles, leading to a reheating temperature typically in the range $10^9 - 10^{10}$ GeV. These values are consistent with constraints from thermal leptogenesis and avoid overproduction of unwanted relics.

Overall, Starobinsky’s $f(R)$ inflation model serves as a cornerstone for constructing and evaluating modified gravity models of inflation, providing a benchmark against which alternative scenarios are often compared.

Inflationary dynamics and predictions

Under the slow-roll approximation, the Starobinsky model yields precise inflationary predictions that depend only on the number of e-folds N before the end of inflation. At leading order in $1/N$, the slow-roll parameters and observable

quantities are given by:

$$n_s \simeq 1 - \frac{2}{N},$$

$$r \simeq \frac{12}{N^2},$$

$$\alpha_s \simeq -\frac{2}{N^2}.$$

For a typical range $N = 50 - 60$, corresponding to when the largest observable CMB scales exit the horizon, these expressions yield:

$$n_s \approx 0.96 - 0.967, \quad r \approx 0.003 - 0.005,$$

$$\alpha_s \approx -0.0006 - -0.0004. \tag{3}$$

These values lie well within the observational bounds reported by the Planck 2018 mission and the BICEP/Keck joint analysis [7,8]. In particular, the prediction of a small but nonzero red tilt ($n_s < 1$) and an extremely low tensor-to-scalar ratio r makes Starobinsky inflation one of the best-fitting single-field models to current CMB data.

Moreover, the low value of r is a clear discriminant compared to large-field models such as chaotic inflation, which predict $r \gtrsim 0.1$ and are now strongly disfavored. The absence of significant running (α_s) and non-Gaussianity further strengthens the compatibility of Starobinsky inflation with data, marking it as a benchmark model for inflationary cosmology. Its predictive power, minimal assumptions, and theoretical robustness make it a target scenario for future missions such as LiteBIRD and CMB-S4.

Reheating and exit from inflation

The Starobinsky model incorporates a natural and efficient reheating mechanism. Once inflation ends, the scalaron field – the extra scalar degree of freedom arising from the R^2 term – enters a phase of coherent oscillations around the minimum of its effective potential. During this post-inflationary epoch, the scalaron behaves like a massive scalar field with an effective mass $m \sim 10^{13}$ GeV, and its energy density scales approximately as non-relativistic matter.

These oscillations lead to the production of Standard Model particles via gravitational couplings or through direct,

albeit typically weak, interactions with matter fields. This process gradually converts the scalaron energy into a thermal bath, initiating the radiation-dominated era. Because the mechanism is built into the dynamics of the model and does not require any additional scalar fields or couplings, it provides a minimal and predictive framework for the reheating phase.

Estimates based on perturbative decay calculations yield a typical reheating temperature in the range

$$T_{\text{rh}} \sim 10^9 - 10^{10} \text{ GeV}, \quad (4)$$

which is sufficiently high to accommodate thermal leptogenesis. This ensures that baryon asymmetry generation mechanisms remain viable in the Starobinsky framework. Additionally, such a high reheating scale implies that relic gravitational wave backgrounds could be produced, providing a potential observational probe of the post-inflationary dynamics in upcoming experiments.

Generalizations of $f(R)$ inflation

Building on Starobinsky's foundational work, a wide variety of generalizations within the $f(R)$ gravity framework have been proposed to accommodate broader cosmological dynamics and connect inflation to late-time acceleration:

- **Power-law corrections:** Models of the form $f(R) = R + \mu R^n$, with $n > 2$, introduce steeper effective potentials in the Einstein frame. These typically yield larger tensor-to-scalar ratios r , which are increasingly constrained by CMB data. While such models can reproduce inflationary expansion, they require fine-tuning or additional modifications to remain viable.
- **Exponential models:** A representative example is $f(R) = R - \mu R_s (1 - e^{-R/R_s})$, where μ and R_s are constants. These models exhibit plateau-like potentials similar to Starobinsky inflation but offer greater flexibility in the duration of inflation and reheating dynamics. They often interpolate between early-time de Sitter and late-time cosmic acceleration phases.
- **Quantum-corrected models:** Motivated by effective actions derived from semiclassical gravity or string theory, extensions such as $f(R) = R + \beta R^2 + \gamma \ln R$ incorporate logarithmic or inverse-curvature corrections. These additions can capture quantum-gravitational effects and allow for richer dynamics in both the ultraviolet and infrared regimes.

Such models have been shown to provide a unified description of the universe's evolution, linking inflation to dark energy within a single geometrical framework [12, 16]. Moreover, many viable $f(R)$ models remain consistent with local gravity constraints through screening mechanisms such as

the chameleon effect, which suppress modifications in high-density environments while permitting deviations on cosmological scales.

Einstein frame and scalar–tensor correspondence

The $f(R)$ theory of gravity can be recast into an equivalent scalar–tensor form via a conformal transformation of the metric:

$$\tilde{g}_{\mu\nu} = f'(R) g_{\mu\nu}, \quad (5)$$

where $\tilde{g}_{\mu\nu}$ is the Einstein-frame metric and $f'(R)$ denotes the derivative of the function $f(R)$ with respect to the Ricci scalar. This transformation introduces a scalar degree of freedom, often referred to as the *scalaron*, which encapsulates the dynamical effects of the higher-curvature corrections.

In the Einstein frame, the action corresponds to that of General Relativity minimally coupled to a canonical scalar field ϕ with an effective potential:

$$V(\phi) = \frac{1}{8\alpha} \left(1 - e^{-\sqrt{2/3} \phi/M_P}\right)^2, \quad (6)$$

where α is the coefficient of the R^2 term in the original Jordan-frame action. This potential is exponentially flat in the large-field limit, ensuring a prolonged slow-roll phase and making it an excellent candidate for inflation consistent with current observational bounds.

This conformal mapping not only facilitates analytical treatments using standard inflationary slow-roll techniques but also enables direct comparison with scalar field models derived from other theoretical frameworks. Importantly, physical observables such as the scalar spectral index n_s and tensor-to-scalar ratio r remain invariant under the transformation, ensuring that predictions in the Einstein frame accurately reflect the original modified gravity theory.

Observational viability and status

The Starobinsky model remains one of the most robust inflationary scenarios. Its predictions are fully compatible with current CMB data, and future probes like LiteBIRD and CMB-S4 are expected to measure r at the 10^{-3} level – directly testing this model (Table 4).

$f(R)$ gravity – and particularly the Starobinsky model – offers an elegant and geometrically motivated realization of inflation that avoids trans-Planckian field excursions and fits current data remarkably well. Its scalar–tensor duality provides an intuitive link to canonical inflationary dynamics, while its generalizations allow a wider exploration of early-universe physics. Upcoming experiments will further test its predictions and potentially distinguish between various extensions within this class.

Table 4 Summary of $f(R)$ inflationary models and observational status

Model	n_s	r	Status
Starobinsky $R + \alpha R^2$	0.965	0.003	Consistent
Power-law $R + \mu R^n, n > 2$	< 0.96	> 0.01	Disfavored
Exponential gravity	~ 0.96	~ 0.005	Viable
Log-corrected $f(R)$	Model-dependent	Model-dependent	Viable if tuned

4 Extended $f(R)$ models

While the Starobinsky model is a minimal and observationally successful representative of $f(R)$ gravity, it represents only a specific corner of the theory space. A variety of extended $f(R)$ models have been proposed to generalize inflationary dynamics, improve reheating scenarios, allow better unification with dark energy, and incorporate features inspired by quantum gravity. These models modify the gravitational Lagrangian to include higher-order or non-polynomial curvature terms, often motivated by theoretical considerations or reconstruction from observational data.

Power-law and polynomial models

A straightforward generalization is to consider models of the form:

$$f(R) = R + \alpha R^n, \tag{7}$$

where α is a constant and $n \neq 2$. These models introduce richer inflationary dynamics due to their steeper potentials in the Einstein frame.

For $n > 2$, the inflationary potential becomes increasingly steep, leading to higher values of the tensor-to-scalar ratio r and deviations from the scale-invariant spectrum. Specifically, in the Einstein frame, the scalar field potential associated with $f(R) = R + \alpha R^n$ takes the form:

$$V(\phi) \propto \left(1 - e^{-\sqrt{2/3}\phi/M_P}\right)^{2n/(n-1)}. \tag{8}$$

The steeper the potential, the larger the value of r , often pushing predictions outside the 95% confidence region of Planck+BICEP/Keck data.

Sami and collaborators [17] have shown that such models can yield viable inflation only in narrow regions of parameter space or when combined with additional fields to control reheating and exit. Moreover, for $n < 2$, the effective potential can be too flat, requiring excessive fine-tuning or leading to an eternal inflation regime.

Reheating in these models is typically less efficient than in the Starobinsky case. A smooth transition to the radiation-dominated era may not occur naturally unless extra couplings to matter or auxiliary fields are introduced [12, 18].

Exponential and logarithmic gravity models

To overcome the limitations of polynomial models, various non-polynomial corrections have been considered. Two particularly well-studied examples are:

$$f(R) = R - \mu R_s \left(1 - e^{-R/R_s}\right),$$

$$f(R) = R + \beta R^2 + \gamma \ln(R/\mu^2),$$

where μ, β, γ , and R_s are constants.

These forms are motivated by quantum gravity (e.g., one-loop effective actions) and string-inspired corrections. The exponential gravity model flattens the inflationary potential at large R , producing a slow-roll phase similar to Starobinsky inflation but with improved control over the exit and reheating phases.

Bamba and Odintsov [19, 20] showed that such models can unify inflation and late-time acceleration within a single functional form of $f(R)$. The logarithmic correction arises in effective quantum field theory on curved spacetimes and may improve the UV behavior of the model while also maintaining compatibility with solar system tests.

Reconstruction techniques and model inversion

Instead of postulating a specific form of $f(R)$, another approach is to reconstruct the function from a desired Hubble expansion history $H(t)$ or a scale factor $a(t)$. This inverse technique has been explored extensively by Capozziello, Sebastiani, and Cognola [21, 22].

In this framework, one begins with a target inflationary behavior (e.g., a constant-roll or power-law inflation scenario) and solves for the functional form $f(R)$ that realizes this dynamics. This method enables the design of models that achieve:

- A consistent inflationary phase with slow-roll dynamics,
- A smooth transition to reheating and radiation domination,
- Late-time cosmic acceleration (if desired),
- Compatibility with CMB and large-scale structure data.

The reconstruction approach is particularly useful when observational consistency is required across all epochs, and when one wants to avoid ad hoc assumptions about the form of the action.

Observational viability and theoretical constraints

Extended $f(R)$ models must satisfy both theoretical consistency conditions and empirical constraints:

- **Stability:** $f'(R) > 0, f''(R) > 0$ ensure the absence of ghosts and tachyonic instabilities.
- **Solar system tests:** Compatibility with local gravity experiments often requires the chameleon mechanism or other screening effects.
- **CMB bounds:** $n_s \approx 0.965, r \lesssim 0.01$ must be respected to avoid observational exclusion.
- **Reheating dynamics:** A successful transition to standard Big Bang cosmology is required for completeness (Table 5).

Extended $f(R)$ models constitute a fertile class of inflationary scenarios that go beyond the minimal Starobinsky form. They provide a testing ground for exploring the role of quantum corrections, high-curvature effects, and model reconstruction. While many models are ruled out or strongly constrained by current observations, well-motivated generalizations – especially exponential and reconstructed forms – remain viable and are under active investigation.

With the upcoming sensitivity of LiteBIRD and CMB-S4 to $r \sim 10^{-3}$, these models face critical observational tests in the near future. The ability of extended $f(R)$ theories to unify early- and late-time acceleration makes them particularly appealing for a geometric and minimalistic approach to cosmic evolution.

5 $f(G)$ Gauss–Bonnet inflation

A compelling class of modified gravity models involves the Gauss–Bonnet (GB) invariant:

$$G = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, \tag{9}$$

which arises naturally in string-inspired and higher-dimensional gravitational theories. In four dimensions, G is a topological term and does not contribute dynamically if added linearly to the action. However, nonlinear functions of G , or couplings of G to scalar fields, can introduce nontrivial modifications to the inflationary dynamics while preserving second-order field equations.

Theoretical motivation

The inclusion of $f(G)$ terms in gravitational actions is strongly motivated by low-energy effective string theories, such as heterotic superstring compactifications, where higher-curvature corrections like the GB invariant naturally arise. These terms can cure singularities, provide UV-completion corrections, and affect early-time acceleration.

Bamba et al. [23] and others have shown that such corrections can support successful inflationary evolution without introducing ghost degrees of freedom, provided the function $f(G)$ is chosen appropriately. Moreover, scalar–GB couplings appear in dilaton gravity, moduli stabilization scenarios, and braneworld cosmologies.

General action and scalar–GB couplings

A general class of pure $f(G)$ models is described by:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} + f(G) \right], \tag{10}$$

which modifies gravity directly through a function of the GB term.

A broader and phenomenologically richer framework includes a scalar field ϕ coupled to G via a function $\xi(\phi)$:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \frac{1}{2}(\nabla\phi)^2 - V(\phi) - \xi(\phi)G \right]. \tag{11}$$

This scalar–GB framework appears in the effective low-energy action of string theory, particularly in dilaton and modulus inflation models. The function $\xi(\phi)$ may take polynomial or exponential forms and modifies the slow-roll conditions.

Model examples and inflationary dynamics

Several forms for $f(G)$ or $\xi(\phi)$ have been proposed and studied:

- **Power-law:** $f(G) = \alpha G^n$, which can support quasi-de Sitter inflation for appropriate n and α .
- **Exponential:** $f(G) = \mu e^{\lambda G}$, motivated by non-perturbative string effects.
- **Inverse power:** $f(G) = \beta G^{-n}$, which may unify early- and late-time acceleration.

Capozziello and De Laurentis [24], as well as Bamba and Odintsov [25], analyzed these models and demonstrated that they support slow-roll inflation with graceful exit and viable reheating. Scalar–GB models with exponential coupling $\xi(\phi) \sim e^{-\gamma\phi}$ or monomial $\xi(\phi) \sim \phi^n$ can suppress

Table 5 Inflationary behavior and viability of selected extended $f(R)$ models

Model	n_s	r	Graceful exit
Starobinsky $R + \alpha R^2$	~ 0.965	~ 0.003	Yes
Power-law $R + \alpha R^n$	< 0.96	> 0.01	Not always
Exponential gravity	~ 0.965	~ 0.005	Yes
Log-corrected $f(R)$	Model-dependent	< 0.01	Yes
Reconstructed $f(R)$	Flexible	Flexible	Yes

Table 6 Representative $f(G)$ and scalar–GB inflation models and their predictions

Model	n_s	r
$f(G) = \alpha G^n, n = 1/2$	~ 0.96	~ 0.004
$f(G) = \mu e^{\lambda G}$	~ 0.965	~ 0.002
Scalar–GB $\xi(\phi) \sim e^{-\gamma\phi}$	~ 0.964	< 0.01
Scalar–GB, GW170817-compatible [28]	~ 0.967	~ 0.003

the tensor-to-scalar ratio and stabilize the inflationary trajectory.

Predictions and observational constraints

Inflationary predictions of $f(G)$ and scalar–GB models are consistent with current data for suitable parameter choices. Typically, they yield:

- Scalar spectral index $n_s \sim 0.96–0.967$,
- Tensor-to-scalar ratio $r < 0.01$,
- Negligible or negative running $\alpha_s \sim -10^{-3}$.

These values fall within the 68 or 95% confidence contours from Planck 2018 and BICEP/Keck 2018 [7,8].

Recent studies have also emphasized compatibility with the gravitational wave speed constraint from GW170817. Scalar–GB theories must be tuned such that $c_T = c$, avoiding superluminal propagation. Models satisfying this include those discussed in [26–28] (Table 6).

Reheating and graceful exit

Reheating in these models is generically achievable. Nojiri et al. [11] showed that GB corrections can smoothly terminate inflation and trigger coherent oscillations of the effective scalar degree of freedom, enabling gravitational particle production or coupling-driven reheating.

Scalar–GB theories offer improved reheating dynamics due to the additional scalar degree of freedom. Bamba and collaborators [25,29] found that the reheating temperature

and duration can be controlled via the choice of $\xi(\phi)$, avoiding unwanted relic production.

Gauss–Bonnet inflation, realized either through pure $f(G)$ corrections or scalar–GB couplings, provides a rich framework for embedding inflation within modified gravity. These models are strongly motivated by fundamental theory, remain compatible with current CMB and gravitational wave data, and offer a wide parameter space for observational tests.

In particular, scalar–GB models with carefully tuned couplings can satisfy the GW speed constraint $c_T = c$ while predicting low r and observable non-Gaussian features. Future experiments like LiteBIRD and CMB-S4 will play a crucial role in testing these predictions and constraining the viable form of $f(G)$ and $\xi(\phi)$.

6 $f(R, G)$ mixed models

A rich class of modified gravity models arises when the Ricci scalar R and the Gauss–Bonnet invariant G are combined nonlinearly in the action via a generic function $f(R, G)$. These models generalize both $f(R)$ and $f(G)$ gravity and can support inflation, reheating, radiation, matter-dominated eras, and late-time acceleration in a unified framework. The inclusion of G brings higher-order curvature corrections inspired by string theory, while the R -dependence keeps the formulation grounded in 4D low-energy effective field theory.

Theoretical framework and ghost issues

The general action is given by:

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} f(R, G), \tag{12}$$

where $f(R, G)$ is an arbitrary smooth function. These theories appear naturally in heterotic string compactifications, quantum gravity corrections, and higher-dimensional braneworld scenarios.

However, a major issue with these models is the presence of ghost degrees of freedom due to higher-derivative terms and the nonlocal structure of G . As shown in [30,31], unless the form of $f(R, G)$ is carefully chosen, the equations of

motion propagate ghostlike scalar and tensor modes with negative kinetic energy, leading to quantum instabilities.

Stability conditions to avoid such issues include:

$$f_R > 0, \quad f_{RR} > 0, \quad f_G + 4H^2 f_{GG} > 0, \quad c_T^2 = 1, \quad (13)$$

where c_T is the speed of tensor modes. These ensure stability of both background and perturbations, and are essential for compatibility with GW170817 constraints on gravitational wave speed.

Functional forms and inflationary dynamics

Popular forms of $f(R, G)$ studied in the literature include:

- **Separable forms:** $f(R, G) = f_1(R) + f_2(G)$, allowing a decoupled treatment of curvature and topological corrections.
- **Power-law mixtures:** $f(R, G) = R + \alpha R^2 + \beta G^n$, mimicking Starobinsky inflation plus GB flattening.
- **Product forms:** $f(R, G) = R^m G^n$, which yield richer phase-space behavior and potential non-trivial inflationary attractors.
- **Logarithmic and exponential corrections:** $f(R, G) = R + \mu G \ln(G/G_0)$, inspired by trace anomalies and string corrections.

Sebastiani and Myrzakulov [32] studied exponential-type models and showed that they can reproduce inflation with graceful exit. Capozziello et al. [33] used Noether symmetries to classify viable forms of $f(R, G)$ admitting integrable inflationary solutions. Odintsov and Oikonomou applied dynamical system techniques to map the phase space, showing viable inflationary attractors and trajectories [34,35].

Observational predictions and constraints

The predictions of viable $f(R, G)$ models lie close to the Planck 2018 + BICEP/Keck contours, with characteristic ranges:

- $n_s = 0.960-0.967$,
- $r = 0.005-0.02$,
- $\alpha_s \sim -10^{-3}$ (small running),
- $f_{NL}^{local} \sim \mathcal{O}(1)$, consistent with Planck’s non-Gaussianity limits (Table 7).

Compatibility with GW speed constraints further narrows the parameter space. Some models, such as those with non-minimal scalar–GB terms satisfying $c_T = c$, provide observationally viable extensions (see [36,37]).

Table 7 Inflationary predictions of representative $f(R, G)$ models

Model	n_s	r
$f(R, G) = R + \alpha R^2 + \beta G^{1/2}$	~ 0.964	~ 0.006
$f(R, G) = R^m G^n, m = 1, n = 1/2$	~ 0.961	~ 0.008
$f(R, G) = R + \mu G \ln(G/G_0)$	~ 0.965	$\lesssim 0.01$

Reheating, dynamical features, and phase space

Odintsov et al. [35] demonstrated that $f(R, G)$ models can support successful reheating by dynamically exiting the inflationary slow-roll phase and allowing scalar oscillations or gravitational reheating. Depending on the form of $f(R, G)$, the effective equation of state $w_{\text{eff}}(t)$ transitions smoothly from $w \approx -1$ (inflation) to $w \approx 1/3$ (radiation).

Dynamical system studies reveal critical points corresponding to stable de Sitter inflationary attractors, saddle-point reheating transitions, and matter-domination phases. The works [32,34] show that trajectories remain ghost-free and observationally consistent for broad parameter ranges.

$f(R, G)$ gravity offers a powerful theoretical framework that blends the strengths of $f(R)$ and $f(G)$ inflation while introducing richer dynamics. It accommodates graceful exit, reheating, late-time acceleration, and compatibility with CMB and GW data – all within a geometric, Lagrangian-based formalism. However, care must be taken to avoid ghosts and ensure second-order equations of motion. Models passing these criteria remain strong candidates for inflationary cosmology and provide a promising direction for further exploration.

7 $f(T)$ torsion gravity

While most modified gravity theories rely on curvature-based modifications of General Relativity, an alternative approach is to use torsion as the dynamical origin of gravity. In this framework, called teleparallel gravity, spacetime is endowed with torsion but no curvature. The gravitational interaction is described using the Weitzenböck connection, which leads to the so-called Teleparallel Equivalent of General Relativity (TEGR). Generalizing TEGR to a function $f(T)$ of the torsion scalar introduces new degrees of freedom and provides a promising arena for inflationary model building.

Teleparallel geometry and field equations

In teleparallel gravity, the basic dynamical variables are the tetrad (or vielbein) fields e_μ^A , which form a local orthonormal basis for the tangent space at each spacetime point. The

torsion tensor is defined by:

$$T^{\rho}_{\mu\nu} = \Gamma^{\rho}_{\nu\mu} - \Gamma^{\rho}_{\mu\nu} = e^{\rho}_A \left(\partial_{\mu} e^A_{\nu} - \partial_{\nu} e^A_{\mu} \right), \tag{14}$$

where $\Gamma^{\rho}_{\mu\nu}$ is the Weitzenböck connection. The torsion scalar is then constructed as:

$$T = S^{\mu\nu} T^{\rho}_{\mu\nu}, \tag{15}$$

where $S^{\mu\nu}$ is the superpotential.

The action for $f(T)$ gravity is given by:

$$S = \frac{1}{2\kappa^2} \int d^4x e f(T), \tag{16}$$

where $e = \det(e^A_{\mu})$ is the tetrad determinant, and $f(T)$ is a general function of the torsion scalar. When $f(T) = T$, the theory reduces to TEGR, which is dynamically equivalent to General Relativity.

One of the most attractive features of $f(T)$ gravity is that it leads to second-order field equations, avoiding the Ostrogradsky instabilities that typically arise in higher-order theories like $f(R)$ or $f(R, G)$.

Inflation in $f(T)$ cosmology

Inflation can naturally arise in $f(T)$ gravity without the need for an inflaton field or a scalar potential. The modified Friedmann equations derived from the action admit de Sitter and quasi-de Sitter solutions under appropriate choices of $f(T)$. This allows for a purely geometric realization of inflation, where torsion is the driving agent of accelerated expansion.

A number of specific forms of $f(T)$ have been proposed and studied:

- **Power-law form:** $f(T) = T + \alpha T^n$. For $n < 1$, inflationary solutions with graceful exit can occur.
- **Exponential form:** $f(T) = T + \beta T_0(1 - e^{-T/T_0})$, which allows for a smooth interpolation between early- and late-time acceleration.
- **Logarithmic corrections:** $f(T) = T + \gamma \ln(T/T_0)$, motivated by loop corrections or trace anomalies.

Bamba et al. [38] and Cai et al. [13] have shown that such models yield slow-roll parameters ϵ, η consistent with observational bounds, and can successfully support inflationary evolution with controlled scalar perturbations.

Observational predictions and viability

$f(T)$ gravity is capable of producing observationally viable predictions for the spectral index n_s and the tensor-to-scalar ratio r , often within the tight bounds set by Planck and BICEP/Keck (Table 8):

Table 8 Inflationary predictions for representative $f(T)$ models

Model	n_s	r
$f(T) = T + \alpha T^n, n \sim 0.5$	~ 0.965	$\lesssim 0.01$
$f(T) = T + \beta(1 - e^{-T/T_0})$	~ 0.964	~ 0.005
$f(T) = T + \gamma \ln(T/T_0)$	~ 0.962	< 0.01

The scalar power spectrum is typically close to scale invariance, and the tensor spectrum is suppressed, similar to Starobinsky $f(R)$ inflation. Depending on the model parameters, $f(T)$ inflation may also predict a small running of the spectral index and negligible non-Gaussianity, in agreement with CMB data.

Reheating and exit mechanisms

One of the strengths of $f(T)$ inflation is that exit from inflation and reheating can be achieved geometrically without invoking extra scalar fields. As shown by Myrzakulov et al. and others, the torsion-driven expansion can smoothly decelerate, allowing energy transfer to radiation through gravitational particle production or minimal couplings to matter fields.

Although the reheating temperature is model-dependent, some $f(T)$ models predict values compatible with thermal leptogenesis. This makes them attractive for connecting early-universe inflation with observable particle physics.

Challenges and developments

Despite its strengths, $f(T)$ gravity is not without theoretical issues:

- **Local Lorentz invariance:** In its traditional formulation, $f(T)$ gravity violates local Lorentz invariance due to the dependence on the choice of tetrad. This can lead to unphysical degrees of freedom or frame-dependent results.
- **Covariant formulations:** Krššák and Pereira [39,40] have proposed covariant versions of teleparallel gravity using spin connections that restore full local Lorentz invariance.
- **Tetrad ambiguities:** The choice of tetrad affects the form of the torsion scalar T and the resulting dynamics. Ensuring a proper “good tetrad” is necessary for consistent cosmological evolution.
- **Perturbations and Degrees of Freedom:** The nature of scalar and tensor perturbations in $f(T)$ gravity is subtle, especially in the presence of anisotropies or inhomogeneities. Recent work continues to refine the under-

standing of gravitational wave propagation in these theories.

$f(T)$ gravity presents a promising torsion-based alternative to curvature-driven models of inflation. Its second-order field equations and analytic simplicity make it computationally attractive, while its predictions are competitive with leading models such as Starobinsky inflation. With ongoing developments in covariant formulations and perturbative analysis, $f(T)$ gravity continues to emerge as a serious contender in the landscape of viable inflationary theories. Its unified treatment of inflation and dark energy also makes it appealing for future model-building beyond the standard cosmological paradigm.

8 Scalar field coupled to Ricci scalar

One of the most natural and well-motivated extensions of inflationary cosmology is the introduction of a non-minimal coupling between a scalar field ϕ and the Ricci scalar R . Such a coupling arises inevitably in quantum field theory in curved spacetime due to renormalization effects and is required to maintain renormalizability of scalar field theories at the one-loop level [41]. It also appears in extensions of the Standard Model, including Higgs inflation and induced gravity scenarios.

Action and field equations

The general Jordan-frame action for a non-minimally coupled scalar field is:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} (1 + \xi \phi^2) R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right], \tag{17}$$

where ξ is the non-minimal coupling constant. The case $\xi = 0$ corresponds to minimal coupling, while the conformal coupling is given by $\xi = 1/6$. Large values of $\xi \gg 1$ are often required in inflationary scenarios.

This coupling modifies both the background cosmological dynamics and the evolution of perturbations. The effective Planck mass becomes field-dependent, leading to enhanced friction during inflation and allowing even steep potentials in the Jordan frame to support slow-roll inflation.

Einstein frame formulation and conformal transformation

To analyze inflationary predictions, it is often convenient to perform a conformal transformation:

$$\tilde{g}_{\mu\nu} = \Omega^2(\phi) g_{\mu\nu}, \quad \Omega^2 = 1 + \xi \phi^2, \tag{18}$$

which transforms the action into the Einstein frame where the scalar field couples minimally to gravity but acquires a modified potential and kinetic term. Defining the canonically normalized field χ , the Einstein-frame potential becomes:

$$V_E(\chi) = \frac{V(\phi(\chi))}{(1 + \xi \phi^2)^2}. \tag{19}$$

This transformation flattens the potential at large ϕ , enabling slow-roll inflation even with quartic or quadratic potentials in the Jordan frame.

Higgs inflation and beyond

A notable example is Higgs inflation, proposed by Bezrukov and Shaposhnikov [42], where the Standard Model Higgs doublet is non-minimally coupled to gravity with $\xi \sim 10^4$. In this case, the quartic potential $V(\phi) = \lambda \phi^4/4$ transforms into a Starobinsky-like potential in the Einstein frame:

$$V_E(\chi) \approx \frac{\lambda M_P^4}{4\xi^2} \left(1 - e^{-\sqrt{2/3}\chi/M_P} \right)^2, \tag{20}$$

with predictions:

$$n_s \approx 0.965, \quad r \approx 0.003, \tag{21}$$

in excellent agreement with Planck and BICEP/Keck observations.

Other scalar potentials in this framework include:

- **Quadratic:** $V(\phi) = \frac{1}{2} m^2 \phi^2$, which becomes viable with large ξ ,
- **Hilltop:** $V(\phi) = V_0 \left(1 - \frac{\phi^2}{\mu^2} \right)^2$,
- **Coleman–Weinberg:** Logarithmically corrected potentials arising from radiative corrections.

Bamba, Odintsov, and others have shown that these models yield viable slow-roll parameters in the Einstein frame [20,43].

Inflationary predictions and observables

The observable predictions of non-minimally coupled models depend on the shape of the potential and the value of ξ . For large ξ , even originally steep potentials become flat and produce inflationary observables similar to Starobinsky inflation (Table 9).

These models typically yield negligible running of the spectral index α_s and small local non-Gaussianity $f_{NL} \sim \mathcal{O}(1)$, consistent with Planck constraints.

Table 9 Inflationary predictions of non-minimally coupled scalar field models

Model	n_s	r
Higgs inflation ($\xi \sim 10^4$)	~ 0.965	~ 0.003
Quadratic with $\xi \gg 1$	~ 0.964	$\gtrsim 0.01$
Hilltop with coupling	~ 0.960	$\gtrsim 0.01$

Reheating, unitarity, and UV considerations

Despite their successes, non-minimally coupled models face important theoretical challenges:

- **Unitarity violation:** At large ξ , the unitarity cutoff $\Lambda \sim M_P/\xi$ lies below the Planck scale. This raises concerns about whether inflation occurs in a regime where the effective theory is still valid. Studies by Barbon et al. and Burgess et al. have explored this issue extensively.
- **Reheating dynamics:** The scalar–curvature coupling affects the inflaton’s coupling to matter fields and thereby modifies the reheating process. Depending on the conformal frame and particle content, reheating efficiency and temperature vary significantly.
- **Frame dependence and quantum corrections:** Although physical observables should be frame-independent, quantum corrections may behave differently in Jordan and Einstein frames. Several works have advocated a frame-invariant formalism for consistent treatment at loop level.

Proposed solutions to the unitarity problem include embedding the model in UV-complete frameworks (e.g., asymptotically safe gravity or supersymmetric extensions), or using scale-invariant or induced gravity models that remain weakly coupled throughout inflation [44,45].

Inflation driven by a scalar field non-minimally coupled to the Ricci scalar provides a highly successful and theoretically motivated framework. It unifies ideas from quantum field theory, particle physics, and cosmology. The Higgs inflation model remains a particularly attractive realization, offering minimal extensions to the Standard Model with excellent observational fits.

Nevertheless, open issues such as unitarity bounds, frame dependence, and quantum stability remain under active investigation. As precision data from LiteBIRD and CMB-S4 emerge, the predictions of non-minimal coupling models will be further tested, possibly distinguishing them from curvature- or torsion-driven geometric models.

9 Scalar field coupled to Ricci tensor

Beyond coupling a scalar field to the Ricci scalar R , one may also consider a non-minimal coupling to the Ricci tensor $R_{\mu\nu}$. Such couplings arise naturally in the context of higher-derivative corrections in effective field theory, quantum loop corrections in curved spacetime, and string-theoretic or Kaluza–Klein compactifications. These interactions modify the kinetic structure of the scalar field and can lead to rich inflationary dynamics, including novel friction mechanisms, altered sound speeds, and potentially enhanced non-Gaussianities.

General action and theoretical motivation

The most general second-order scalar–tensor theories allowing Ricci tensor couplings are captured by the Horndeski class and its extensions. A minimal model is given by:

$$S = \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} \xi R^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right], \tag{22}$$

where ξ is a dimensionless coupling constant. The term $\xi R^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$ effectively acts as a curvature-dependent kinetic correction, modifying both the propagation of perturbations and the background scalar dynamics.

Such terms can arise from integrating out heavy modes in quantum gravity or in low-energy expansions of string theory compactifications. They also appear in the generalized Galileon or G-inflation frameworks that extend the standard scalar–tensor paradigm.

G-inflation and enhanced friction mechanism

A prominent realization of scalar–Ricci tensor coupling was proposed by Germani and Kehagias [46], who introduced the G-inflation model:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \frac{1}{2} \left(g^{\mu\nu} - \frac{1}{M^2} R^{\mu\nu} \right) \times \partial_\mu \phi \partial_\nu \phi - V(\phi) \right]. \tag{23}$$

Here, M is a new energy scale controlling the strength of the coupling. In this model, the Ricci tensor coupling increases the effective friction acting on the inflaton, allowing for slow-roll dynamics even with steep potentials and sub-Planckian field excursions. This mechanism is called gravitationally enhanced friction and is of particular interest in string theory constructions where field excursions are limited.

Table 10 Representative predictions from scalar–Ricci tensor inflationary models

Model	n_s	r
G-inflation (Germani–Kehagias)	~ 0.964	$\lesssim 0.01$
Kinetic coupling $\xi R^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$	~ 0.965	~ 0.005
Generalized Galileon models	0.960–0.970	< 0.01

Such models can maintain nearly scale-invariant scalar perturbations while naturally suppressing the tensor-to-scalar ratio r , helping reconcile steep potentials with observational data.

Inflationary dynamics and attractor behavior

The scalar–Ricci tensor coupling modifies the effective Klein–Gordon equation for the scalar field:

$$\square \phi - \xi \nabla_\mu (R^{\mu\nu} \nabla_\nu \phi) + V'(\phi) = 0. \tag{24}$$

During inflation, this extra term acts like a field-dependent Hubble friction, enabling attractor solutions even for otherwise too-steep potentials. Dynamical system studies show that these models admit stable slow-roll attractors for a wide range of initial conditions [47,48].

Observational predictions and parameter space

Scalar–Ricci tensor models generically predict:

- A spectral index $n_s \approx 0.960\text{--}0.970$,
- A tensor-to-scalar ratio $r < 0.01$,
- A running α_s close to zero,
- A nontrivial sound speed $c_s < 1$, which can enhance equilateral-type non-Gaussianity (Table 10).

These predictions align well with Planck 2018 and BICEP/Keck bounds. Importantly, the suppression of tensor modes is a natural feature of these models, which distinguishes them from many large-field canonical inflation models.

Perturbations, stability, and sound speed

The presence of the Ricci tensor coupling modifies the sound speed c_s of scalar perturbations:

$$c_s^2 = \frac{1}{1 + \xi R/M^2}, \tag{25}$$

which implies that even with canonical-looking Lagrangians, the scalar sector may experience nontrivial dispersion relations. This can result in observable equilateral-type non-Gaussianity, quantified by $f_{\text{NL}}^{\text{equil}} \propto (1 - c_s^{-2})$.

However, stability requires:

- No ghost modes (positive-definite kinetic matrix),
- No gradient instabilities ($c_s^2 > 0$),
- Positive energy for all propagating degrees of freedom.

These conditions have been analyzed in detail by Koivisto [49] and Capozziello et al. [48]. Viable models typically require moderate values of ξ and sub-Planckian energy scales M to ensure positivity and bounded Hamiltonians.

UV completion and frame dependence

Being higher-derivative in nature, scalar–Ricci tensor couplings must be embedded carefully within a UV-complete theory. Such terms often arise in low-energy effective actions from string theory or compactified supergravity, but without a UV completion, unitarity and strong coupling issues may arise.

Moreover, these couplings introduce frame-dependent dynamics, and caution is required in defining observable quantities consistently, particularly for loop corrections or reheating analyses. Theories like Horndeski gravity offer a general covariant framework where such terms can be treated consistently without introducing ghosts.

Inflation with a scalar field non-minimally coupled to the Ricci tensor provides a compelling alternative to canonical inflation. These models can accommodate steep potentials, achieve observationally viable predictions for n_s and r , and produce unique signatures in the primordial perturbations – especially in the sound speed and non-Gaussianity sectors.

While further theoretical development is needed to ensure full UV consistency and stability across the cosmological history, scalar–Ricci tensor couplings remain an active area of research in both early-universe cosmology and modified gravity frameworks.

10 Scalar field coupled to torsion tensor

Teleparallel gravity, where gravity is mediated by torsion rather than curvature, provides a rich framework to explore new scalar–geometric couplings. In particular, scalar fields can be coupled not only to the torsion scalar T but also directly to components of the torsion tensor $T^\rho_{\mu\nu}$, leading to novel inflationary mechanisms. These couplings generalize the $f(T)$ paradigm and appear in effective string theory actions, higher-dimensional reductions, and teleparallel scalar–tensor extensions.

Geometric framework and coupling structures

In the teleparallel formulation, the Weitzenböck connection yields torsion but no curvature. The torsion tensor is defined as:

$$T^{\rho}_{\mu\nu} = \Gamma^{\rho}_{\nu\mu} - \Gamma^{\rho}_{\mu\nu}, \tag{26}$$

with the torsion scalar T constructed from contractions:

$$T = S^{\mu\nu}_{\rho} T^{\rho}_{\mu\nu}, \quad \text{where} \\ S^{\mu\nu}_{\rho} = \frac{1}{2}(K^{\mu\nu}_{\rho} + \delta^{\mu}_{\rho} T^{\alpha\nu}_{\alpha} - \delta^{\nu}_{\rho} T^{\alpha\mu}_{\alpha}). \tag{27}$$

Scalar fields can interact with torsion in multiple ways:

- **Non-minimal coupling to torsion scalar:** $\phi^2 T$, analog of $\phi^2 R$,
- **Kinetic coupling to torsion tensor:** $T^{\mu\nu\rho} \partial_{\mu} \phi \partial_{\nu} \phi$,
- **Derivative coupling to vector torsion:** $T^{\mu} \partial_{\mu} \phi$,
- **Axial and trace couplings:** arising in parity-violating or chiral formulations.

These generalizations allow scalar fields to mediate inflation or modify gravity through torsion channels, enriching the landscape of inflationary cosmology.

Non-minimal coupling: $\phi^2 T$

The simplest extension involves a direct coupling between the scalar field and the torsion scalar:

$$S = \int d^4x e \left[\frac{1}{2}(1 + \xi \phi^2) T - \frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - V(\phi) \right], \tag{28}$$

analogous to the scalar–Ricci coupling in curvature-based theories. This model admits de Sitter and quasi–de Sitter phases when ϕ slowly rolls on a flat potential.

Bamba et al. [50] demonstrated that such couplings allow Starobinsky-like inflation with:

$$n_s \approx 0.965, \quad r \approx 0.003, \tag{29}$$

for quartic or exponential potentials, in agreement with Planck 2018 and BICEP/Keck results.

Kinetic couplings to torsion

An even richer class of models includes kinetic couplings between the scalar field and torsion tensors. A general action takes the form:

$$S = \int d^4x e \left[\frac{T}{2\kappa^2} - \frac{1}{2} g^{\mu\nu} \partial_{\mu} \phi \partial_{\nu} \phi + \frac{\eta}{2} T^{\mu\nu\rho} \partial_{\mu} \phi \partial_{\nu} \phi - V(\phi) \right], \tag{30}$$

Table 11 Predictions for scalar–torsion coupled inflationary models

Model	n_s	r
Non-minimal $\phi^2 T$	~ 0.965	~ 0.003
Kinetic torsion coupling	~ 0.964	~ 0.004
Vector torsion–derivative coupling	~ 0.960	$\lesssim 0.01$

where η determines the strength of the kinetic coupling. These models mimic the dynamics of G-inflation by introducing curvature-independent friction terms that damp the evolution of ϕ during inflation. Although the kinetic coupling term $T^{\mu\nu\rho} \partial_{\mu} \phi \partial_{\nu} \phi$ may appear, at first glance, to involve a non-scalar quantity, it is in fact a scalar under general coordinate transformations in the teleparallel framework. This follows because $T^{\mu\nu\rho}$ is the torsion tensor constructed from the Weitzenböck connection, which is a rank-3 tensor, and $\partial_{\mu} \phi$ is a covector. The index contractions over μ and ν ensure that all free indices are saturated, producing a scalar-valued Lagrangian density term. Similarly, for variants involving the torsion vector $T^{\mu} \partial_{\mu} \phi$, T^{μ} is itself a true vector obtained by contraction $T^{\mu} = T^{\nu\mu}_{\nu}$, and the inner product with $\partial_{\mu} \phi$ yields a scalar.¹ We have added this explanation here to make explicit why these terms are permissible in the covariant action. Variants involving couplings to the torsion vector $T^{\mu} = T^{\nu\mu}_{\nu}$, such as $T^{\mu} \partial_{\mu} \phi$, can also lead to dynamical stabilization and modified reheating paths. These have been explored in generalized teleparallel dark energy models and may arise from dimensional reduction of higher-spin theories.

Inflationary dynamics and predictions

Scalar–torsion coupled models admit robust slow-roll inflation, graceful exit, and reheating mechanisms. Depending on the form of the coupling and the potential $V(\phi)$, they produce:

- Spectral index $n_s = 0.960–0.966$,
- Tensor-to-scalar ratio $r \lesssim 0.01$,
- Small running $\alpha_s \sim -10^{-3}$,
- Nearly Gaussian fluctuations in minimal realizations (Table 11).

The ability to naturally suppress tensor modes and maintain nearly scale-invariant scalar spectra places these models among the observationally viable class of geometric inflationary scenarios.

¹ Invariance follows from the tensorial transformation properties of $T^{\mu\nu\rho}$ and T^{μ} under diffeomorphisms in teleparallel geometry. No free indices remain after contraction, and the resulting quantities transform as scalars in the action integral.

Perturbations and stability

Perturbative analysis reveals the following features:

- **Scalar sound speed:** Often remains close to unity, $c_s^2 \approx 1$, suppressing equilateral non-Gaussianity.
- **Ghost avoidance:** Positive coupling constants ($\xi > 0$, $\eta > 0$) prevent ghosts and gradient instabilities.
- **Tensor perturbations:** Propagate similarly to standard $f(T)$ theories, with slight modifications from scalar backreaction.

Bamba et al. [51] emphasized that reheating can proceed via coherent scalar oscillations and gravitational particle production. However, detailed predictions depend sensitively on the chosen tetrad due to the lack of local Lorentz invariance in traditional $f(T)$ models.

To address this, Krššák and Pereira [39,40] developed covariant teleparallel gravity, which introduces a spin connection to restore Lorentz symmetry and allows consistent treatment of perturbations across frames.

Outlook and theoretical significance

Scalar–torsion couplings offer a geometrically rich and observationally consistent alternative to curvature-based inflationary models. They provide:

- Novel mechanisms for inflation and graceful exit,
- Unified treatment of early and late-time acceleration,
- Simplified second-order field equations,
- Unique signatures in CMB polarization and tensor modes.

Ongoing work explores UV completions, quantum corrections, and embeddings in supergravity or string-inspired teleparallel frameworks. With improving observational precision, these models may become testable in the near future, distinguishing torsion-based from curvature-driven inflation via subtle differences in gravitational wave propagation and reheating signatures.

11 Scalar field coupled to Gauss–Bonnet term

Coupling a scalar field to the Gauss–Bonnet (GB) invariant G offers a powerful and theoretically motivated extension of standard inflationary scenarios. These models retain second-order equations of motion, avoid Ostrogradsky instabilities, and provide natural embeddings in low-energy string theory and higher-dimensional gravity. Scalar–Gauss–Bonnet (SGB) inflation can generate viable slow-roll dynamics, suppress tensor modes, and introduce distinct observational

signatures accessible by upcoming CMB polarization experiments.

Theoretical framework and action

The four-dimensional GB term is a topological invariant:

$$G = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}. \quad (31)$$

It does not contribute to the field equations when added linearly to the Einstein–Hilbert action. However, when coupled to a dynamical scalar field ϕ , it introduces nontrivial contributions:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \xi(\phi)G \right]. \quad (32)$$

This coupling appears in heterotic string theory, supergravity compactifications, and loop-corrected effective actions. In particular, it emerges naturally from the dilaton sector in α' -corrected theories. Models with exponential or polynomial forms of $\xi(\phi)$ are commonly derived from such frameworks.

Coupling functions and inflationary dynamics

Several forms of the scalar–GB coupling $\xi(\phi)$ have been investigated:

- **Exponential:** $\xi(\phi) = \lambda e^{-\gamma\phi}$ (string/dilaton origin),
- **Power-law:** $\xi(\phi) = \xi_0 \phi^n$,
- **Logarithmic or reconstructed:** from target inflationary observables.

The GB term effectively modifies the scalar field equation by enhancing the Hubble friction:

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) + \xi'(\phi)G = 0. \quad (33)$$

This allows inflation on steeper potentials by slowing down the field's evolution. Models with otherwise excluded potentials (e.g., quadratic or exponential) become viable under appropriate GB couplings.

Examples and model predictions

Early work by Kanti et al. [52] demonstrated that exponential couplings yield inflation and even bouncing cosmologies. Bouncing models within $f(R)$ and Gauss–Bonnet gravity can provide nonsingular cosmological histories [53].

More recent studies by Bamba, Odintsov, and others [54,55] confirm that a wide variety of $V(\phi)$ and $\xi(\phi)$ combinations lead to successful inflation and exit (Table 12).

These predictions are compatible with Planck 2018 and BICEP/Keck bounds. The tensor-to-scalar ratio is generally

Table 12 Inflationary predictions of scalar–Gauss–Bonnet models

Model	n_s	r
$\xi(\phi) = \lambda\phi^2, V(\phi) = m^2\phi^2/2$	~ 0.963	~ 0.01
$\xi(\phi) = e^{-\gamma\phi}$, exponential $V(\phi)$	~ 0.965	~ 0.005
Plateau $V(\phi)$ with polynomial $\xi(\phi)$	~ 0.964	$\lesssim 0.008$

suppressed, and the scalar spectral index falls within the preferred range $n_s \approx 0.96–0.967$.

Reheating and post-inflation dynamics

A desirable feature of SGB inflation is that inflation can naturally end via oscillations in $\xi(\phi)$ or decay of the GB coupling term. Depending on the sign and slope of $\xi(\phi)$, the scalar field transitions into a coherent oscillation phase, enabling efficient reheating. Models with decaying $\xi(\phi) \propto 1/\phi$ or oscillatory couplings have been studied to ensure a smooth exit and thermal history.

In addition, some constructions connect the GB coupling to late-time acceleration (unification models), though care is needed to avoid violating stability or GW propagation constraints.

Stability, perturbations, and GW speed constraint

Perturbative stability requires that the kinetic term of fluctuations is positive and that no ghost or gradient instabilities arise. For the scalar perturbations, the conditions are:

$$\xi'(\phi)\dot{\phi} < 0, \quad \xi''(\phi) > 0, \tag{34}$$

ensuring positivity of the effective kinetic matrix. For tensor modes, the presence of the GB coupling modifies the tensor propagation speed:

$$c_T^2 = 1 - 4\kappa^2\ddot{\xi}(\phi). \tag{35}$$

This poses a significant constraint due to the observation of GW170817, which showed that $c_T \approx c$ to within parts in 10^{15} . Models like those in [36,55,56] provide GB-coupled inflationary setups that respect the $c_T = 1$ condition by tuning the form of $\xi(\phi)$ or embedding within degenerate higher-order scalar–tensor (DHOST) theories.

Non-Gaussianity and observational prospects

In general, SGB inflation models with standard kinetic terms predict small non-Gaussianities:

$$f_{NL}^{local} \sim \mathcal{O}(1), \tag{36}$$

but models with non-canonical kinetic terms or enhanced couplings may produce observable equilateral-type signa-

tures. Future experiments like LiteBIRD and CMB-S4 will improve constraints on both r and f_{NL} , enabling potential differentiation between SGB and $f(R)/f(T)$ models.

Scalar–Gauss–Bonnet inflation presents a theoretically appealing and observationally viable mechanism for early-universe acceleration. It arises naturally from fundamental theory (e.g., string theory), modifies the inflationary dynamics via geometrically motivated couplings, and produces predictions consistent with CMB data. With ongoing advances in gravitational wave astronomy and precision cosmology, this class of models is poised for critical testing in the near future.

12 Einstein–Cartan models of inflation

Einstein–Cartan (EC) theory is a natural and minimal extension of General Relativity (GR) that incorporates intrinsic spin as a source of torsion. While standard GR assumes a torsion-free connection, EC gravity generalizes the affine structure of spacetime to allow for antisymmetric components of the connection, making spacetime geometry sensitive not only to energy and momentum but also to spin.

Although torsion is generally negligible at low densities, it becomes significant at extremely high densities encountered in the early universe. This renders EC theory particularly relevant for early cosmology, including singularity resolution, inflationary dynamics, and spin-induced bounce scenarios.

Theoretical framework of Einstein–Cartan gravity

In EC theory, the connection $\Gamma^\lambda_{\mu\nu}$ is not symmetric and includes a torsion tensor:

$$T^\lambda_{\mu\nu} = \Gamma^\lambda_{[\mu\nu]} = \frac{1}{2}(\Gamma^\lambda_{\mu\nu} - \Gamma^\lambda_{\nu\mu}), \tag{37}$$

which is algebraically related to the spin density of matter. The full affine connection is decomposed into:

$$\Gamma^\lambda_{\mu\nu} = \tilde{\Gamma}^\lambda_{\mu\nu} + K^\lambda_{\mu\nu}, \tag{38}$$

where $\tilde{\Gamma}^\lambda_{\mu\nu}$ is the Levi-Civita (torsion-free) connection and $K^\lambda_{\mu\nu}$ is the contorsion tensor.

The Einstein–Cartan field equations take the form:

$$G_{\mu\nu} = \kappa^2(T_{\mu\nu} + U_{\mu\nu}), \tag{39}$$

where $T_{\mu\nu}$ is the canonical energy–momentum tensor, and $U_{\mu\nu}$ is a correction term sourced by the spin–torsion interaction. Unlike GR, these equations are derived from the first-order (Palatini) formalism, where the metric and connection are varied independently.

Torsion-induced bounce and singularity resolution

One of the major motivations for EC gravity in cosmology is its ability to avoid the initial singularity. As shown by Hehl et al. [57], at very high matter densities, the torsion–spin interaction induces an effective repulsive force that halts gravitational collapse, resulting in a nonsingular bounce.

Shapiro [58] reviewed this mechanism extensively, showing that torsion can prevent divergences in curvature and energy density. Popławski [59] further demonstrated that fermionic matter with intrinsic spin leads to a cosmic bounce followed by a brief phase of accelerated expansion, even in the absence of an inflaton. These models suggest that:

- The universe undergoes a bounce instead of a Big Bang,
- The spin–torsion coupling generates effective negative pressure,
- Inflation may naturally emerge from spinor dynamics.

Inflation from spin fluids and torsion

In EC theory, fermionic fields act as natural sources of torsion. When modeled as a spin fluid (a statistical ensemble of spin-1/2 particles), the effective equation of state can deviate from that of a perfect fluid, leading to a stiff or even phantom-like behavior.

At early times, the spin–spin interactions mediated by torsion produce a contribution to the energy–momentum tensor that scales faster than radiation. This contribution becomes dominant near the Planck epoch and induces a bounce followed by inflation-like behavior, as studied in both analytical and numerical models [60].

Scalar fields in Einstein–Cartan spacetimes

An alternative EC inflationary scenario includes scalar fields minimally or non-minimally coupled to torsion or contorsion tensors. These models generalize the scalar field dynamics by incorporating torsion-vector couplings such as:

$$S \supset \int d^4x \sqrt{-g} \left[-\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{\xi}{2} S^\mu \phi \partial_\mu \phi - V(\phi) \right], \tag{40}$$

where $S^\mu = T^{\nu\mu}$ is the torsion vector and ξ is a dimensionless coupling constant. This interaction leads to modified slow-roll parameters, effective friction, and new fixed points in the inflationary phase space.

Bamba et al. [61] analyzed scalar–torsion couplings in EC frameworks and showed that such models are consistent with observational constraints, yielding predictions for n_s and r comparable to those in Starobinsky-like inflation.

Table 13 Inflationary behavior in selected Einstein–Cartan models

Model	n_s	r
Spin fluid + EC bounce	~ 0.964	< 0.01
Scalar field in EC background	~ 0.965	~ 0.004
Non-minimal scalar–torsion coupling	~ 0.963	$\lesssim 0.01$

Predictions and observational signatures

The inflationary predictions of EC models vary depending on the mechanism (spinor fluid vs scalar field), but share the following features:

- **Spectral tilt:** $n_s \sim 0.96$, consistent with Planck,
- **Tensor-to-scalar ratio:** $r < 0.01$, often very suppressed,
- **No need for trans-Planckian field values** in scalar models,
- **Natural singularity avoidance** through torsion repulsion,
- **Reheating:** via spinor decay or scalar oscillations post-bounce (Table 13).

Theoretical outlook and connections to quantum gravity

Einstein–Cartan inflation bridges classical gravity with quantum properties of matter, particularly spin. It is closely connected to:

- **Loop quantum gravity (LQG):** where spin networks naturally incorporate torsion,
- **Spin foam models and affine gravity,**
- **Supergravity:** where torsion appears as a supersymmetric auxiliary field,
- **Minimal length scenarios:** torsion helps regulate high-energy divergences.

These features make EC gravity a fertile ground for unifying particle physics, geometry, and cosmology.

Einstein–Cartan cosmology offers a physically motivated and geometrically rich alternative to standard inflation. Whether driven by spin fluids or scalar–torsion couplings, EC models support inflationary expansion, resolve the initial singularity, and yield predictions consistent with current data. They also avoid common theoretical pitfalls such as trans-Planckian excursions or ghost instabilities. As observations sharpen and theoretical models mature, EC inflation remains a compelling path toward a quantum-compatible description of the early universe.

13 Mimetic gravity and inflation

Mimetic gravity is a novel and geometrically motivated modification of General Relativity in which the conformal degree of freedom of the metric is isolated and promoted to a dynamical scalar field. Introduced by Chamseddine and Mukhanov [62], the theory was originally designed to mimic the behavior of cold dark matter, but has since been recognized as a flexible framework for constructing inflationary and bouncing cosmologies without requiring an explicit inflaton field.

The mimetic scalar field arises from a conformal reparametrization of the metric and behaves as a dynamical constraint-enforced degree of freedom. Inflation in mimetic gravity can be realized through the addition of potentials, higher-derivative terms, or couplings to curvature invariants, all of which retain the geometric structure of the theory.

Conformal parametrization and constraint dynamics

The starting point of mimetic gravity is the redefinition of the physical spacetime metric $g_{\mu\nu}$ in terms of an auxiliary metric $\tilde{g}_{\mu\nu}$ and a scalar field ϕ , such that:

$$g_{\mu\nu} = -(\tilde{g}^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi) \tilde{g}_{\mu\nu}. \tag{41}$$

This condition implies the mimetic constraint:

$$g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi = -1, \tag{42}$$

which can be enforced in the action using a Lagrange multiplier λ . The simplest mimetic action becomes:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} + \lambda (g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + 1) \right]. \tag{43}$$

Here, the scalar field ϕ does not propagate via a standard kinetic term, but behaves dynamically due to the constraint. In the absence of other matter fields, this leads to a pressureless fluid behavior, effectively mimicking dark matter.

Extensions for inflationary dynamics

To model inflation, the mimetic framework is extended by introducing additional degrees of freedom:

- **Potential-driven mimetic inflation:** Add a scalar potential $V(\phi)$ to produce slow-roll dynamics.
- **Higher-derivative terms:** Include operators such as $(\square\phi)^2$, $(\nabla_\mu \nabla_\nu \phi)^2$, which arise naturally in Horndeski-like constructions [63].
- **Curvature couplings:** Construct mimetic extensions of $f(R)$, Gauss–Bonnet, or teleparallel gravity (e.g., mimetic $f(G)$, mimetic $f(T)$).
- **Mimetic Horndeski and DHOST models:** Explore higher-order scalar–tensor theories that retain second-order field equations.

Table 14 Inflationary behavior in selected mimetic gravity models

Model	n_s	r
Potential-driven mimetic inflation	~ 0.964	~ 0.003
Mimetic gravity with higher derivatives	~ 0.960	$\lesssim 0.01$
Mimetic $f(R)$ gravity	~ 0.965	~ 0.004
Mimetic Gauss–Bonnet or $f(G)$ models	$0.960\text{--}0.966$	$\lesssim 0.005$

- **LQC-inspired mimetic gravity:** Incorporate loop quantum cosmology corrections into mimetic dynamics.

Nojiri, Odintsov, and collaborators have extensively developed reconstruction techniques within mimetic gravity to generate cosmological histories that support inflation, bounce, or late-time acceleration [64,65].

Inflationary predictions and examples

Mimetic inflationary models are highly flexible and can reproduce a wide range of predictions depending on the structure of the added terms. For suitable choices, they yield:

- Spectral index $n_s \sim 0.960\text{--}0.965$,
- Tensor-to-scalar ratio $r \lesssim 0.01$,
- Modified sound speed $c_s^2 < 1$, potentially enhancing equilateral non-Gaussianity,
- Stability under scalar and tensor perturbations, when higher-derivative operators are handled correctly (Table 14).

Perturbations, stability, and non-Gaussianity

Mimetic models are known to produce stable scalar perturbations in the linear regime, provided the higher-derivative terms are constructed to avoid ghosts and gradient instabilities. In the presence of terms like $(\square\phi)^2$, the sound speed deviates from unity, potentially giving rise to detectable non-Gaussianity:

$$f_{\text{NL}}^{\text{equil}} \propto \frac{1 - c_s^2}{c_s^2}. \tag{44}$$

However, care is required: models with poorly tuned higher-order operators may suffer from instabilities or violate observational bounds. Horndeski-compatible mimetic models have been developed to preserve second-order dynamics and control unwanted degrees of freedom.

Connections to quantum gravity and bouncing cosmologies

Mimetic gravity has also been proposed as a framework for nonsingular bouncing cosmologies, especially in conjunc-

tion with loop quantum cosmology (LQC) corrections. The mimetic scalar can act as an effective anisotropy suppressor and resolve the initial singularity through geometric means.

These models maintain:

- Covariance under diffeomorphisms,
- Constraint-enforced dynamics (Lagrange multiplier formalism),
- Modified Friedmann equations enabling bounce or turnaround phases.

Mimetic gravity offers a geometric mechanism to reproduce inflationary expansion by isolating the conformal degree of freedom of the metric. Through potential-driven or higher-derivative modifications, mimetic models can generate viable inflation consistent with observational constraints. Moreover, their compatibility with Horndeski and quantum gravity frameworks makes mimetic gravity a fertile ground for constructing singularity-free cosmological histories and connecting the early universe to fundamental theory.

As future CMB experiments further tighten bounds on non-Gaussianity and tensor modes, mimetic inflationary models – particularly those with modified sound speed – will face strong tests, potentially distinguishing them from conventional scalar field or curvature-based inflation.

14 Non-local gravity and infinite derivative models

Non-local and infinite-derivative modifications of General Relativity (GR) have gained increasing attention as frameworks capable of addressing the ultraviolet (UV) problems of gravity. These include non-renormalizability, singularity formation, and quantum instabilities. Rooted in string field theory, effective quantum gravity, and higher-spin models, non-local gravity incorporates infinite-order derivative operators, especially in the form of analytic functions of the d’Alembertian \square . When properly constructed, these theories maintain unitarity, preserve causality, and exhibit ghost-free behavior while modifying the structure of spacetime at small scales.

One of the most striking features of such models is that they can support early-universe inflation without requiring an inflaton field or scalar potential. The inflationary dynamics emerge directly from the gravitational sector through non-local curvature operators, offering a minimal and self-contained approach to the origin of cosmological acceleration.

Motivation and effective action framework

Non-local modifications are motivated by both phenomenological and theoretical considerations:

- **String theory:** String field theory and p-adic string models naturally produce actions with non-local operators like e^{\square} ,
- **Asymptotic safety and causal set theory:** These suggest non-localities are needed to ensure UV completeness,
- **Resummation of quantum loops:** Effective actions derived from resummed Feynman diagrams often involve non-local kernels.

A prototypical action for ghost-free non-local gravity takes the form:

$$S = \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R + \frac{1}{2} R \mathcal{F}(\square) R + \mathcal{L}_m \right], \tag{45}$$

where $\mathcal{F}(\square)$ is an analytic form factor (often an entire function) suppressing high-energy divergences. A commonly studied choice is:

$$\mathcal{F}(\square) = \frac{1}{6M_s^2} e^{-\square/M_s^2}, \tag{46}$$

with M_s denoting the scale of non-locality. This structure appears in the UV completion schemes proposed by Biswas, Mazumdar, Koshelev, and Modesto [66–68].

Ghost-free conditions and modified propagator

A key feature of infinite-derivative gravity is that it can be made ghost-free if $\mathcal{F}(\square)$ is chosen to avoid introducing extra poles in the graviton propagator. This demands that $\mathcal{F}(\square)$ be an entire function:

$$\mathcal{F}(\square) = \sum_{n=1}^{\infty} f_n \square^n, \quad \text{with rapid convergence.} \tag{47}$$

The modified graviton propagator in momentum space takes the form:

$$\Pi(k^2) \sim \frac{e^{-k^2/M_s^2}}{k^2}, \tag{48}$$

which suppresses UV contributions and renders loop corrections finite. This allows for a form of perturbative UV completion of gravity that avoids higher-derivative ghosts (unlike generic $f(R, \square R)$ models).

Inflation from purely gravitational non-local terms

Unlike standard scalar-field-driven models, non-local gravity can support inflationary solutions directly from geometric operators. In particular, the exponential form factor in $R\mathcal{F}(\square)R$ generates a de Sitter phase or quasi-de Sitter expansion.

Several classes of inflationary behavior have been found:

Table 15 Inflationary predictions from non-local gravity models

Model	n_s	r
$R + R\mathcal{F}(\square)R$, exponential kernel	~ 0.964	$\lesssim 0.01$
Non-local Starobinsky-like model	~ 0.965	~ 0.003
Bouncing non-local cosmology	~ 0.960	Negligible

- **Exponential inflation:** Driven by the $Re^{-\square}R$ term alone.
- **Bouncing models:** Where the universe undergoes a non-singular contraction–expansion transition.
- **Non-local Starobinsky inflation:** Mimicking $R + R^2$ dynamics with improved UV behavior.

Biswas et al. [66] showed that de Sitter and oscillatory inflationary solutions exist and are stable under perturbations, provided the scale $M_s \sim 10^{14}$ GeV is chosen appropriately (Table 15).

Perturbation theory and observables

Perturbations in non-local gravity are governed by integro-differential equations involving operators like e^{\square} . Nevertheless, linearized analyses around de Sitter or FLRW backgrounds show:

- **Stability:** Scalar and tensor perturbations are stable and ghost-free.
- **Tensor spectrum:** High-energy suppression leads to smaller amplitude and modified tensor tilt.
- **UV finiteness:** Quantum loops are rendered finite, and divergences are smoothed by exponential form factors.

Moreover, the scalar power spectrum remains nearly scale-invariant with $n_s \approx 0.96$, and the tensor-to-scalar ratio is within observable bounds.

Late-time cosmology and unification

Some non-local models can unify inflation with dark energy by introducing other curvature terms such as:

$$R\mathcal{F}_1(\square)R + R_{\mu\nu}\mathcal{F}_2(\square)R^{\mu\nu}. \tag{49}$$

By adjusting the form factors $\mathcal{F}_i(\square)$, a transition from early acceleration to late-time cosmic acceleration can be achieved. This avoids introducing a cosmological constant or quintessence field.

Such scenarios offer a geometric unification of all cosmological epochs within a ghost-free, singularity-free, and UV-complete framework.

Quantum gravity links and theoretical significance

Non-local gravity overlaps with several approaches to quantum gravity:

- **String theory:** Provides natural non-local operators via worldsheet resummations,
- **Asymptotic safety:** Regularizes divergences with exponential UV suppression,
- **Causal set theory and non-commutative geometry:** Support discrete or smeared spacetime structures,
- **Amplitudes and causal perturbation theory:** Benefit from entire-function analyticity.

Non-local operators may also help address trans-Planckian issues in inflation and resolve the black hole information paradox by softening spacetime singularities.

Non-local and infinite-derivative models of gravity offer a compelling route to resolve the ultraviolet problems of General Relativity while simultaneously generating realistic inflationary cosmology. With ghost-free structure, suppressed UV divergences, and predictive power at both early and late times, these models are increasingly recognized as serious candidates for a quantum-consistent and observationally viable theory of the universe’s origin and evolution. Their ability to support inflation without scalar fields, avoid singularities, and unify cosmological epochs underscores their importance in the landscape of modern gravitational theories.

15 Palatini formalism in inflation

The Palatini formalism is an alternative variational principle in gravitational theories in which the metric $g_{\mu\nu}$ and the affine connection $\Gamma^{\lambda}_{\mu\nu}$ are treated as independent variables. Originally proposed by Palatini and later formalized in the context of Einstein’s theory, the Palatini approach leads to the same field equations as the metric formalism in standard General Relativity when applied to the Einstein–Hilbert action. However, this equivalence breaks down for nonlinear extensions of gravity, such as $f(R)$, $f(R, Q)$, and scalar–tensor theories.

In the context of cosmology, this distinction becomes highly nontrivial. Modified gravity models – particularly $f(R)$ models – display markedly different inflationary behavior under the Palatini approach compared to the metric one. In many Palatini models, inflation can be realized without propagating scalar degrees of freedom, and tensor perturbations are naturally suppressed.

Table 16 Inflationary predictions in metric and Palatini $f(R)$ gravity

Model	n_s	r
Starobinsky inflation (metric)	~ 0.965	~ 0.004
Starobinsky inflation (Palatini)	~ 0.965	$\lesssim 0.001$
Palatini $f(R) = R + \alpha R^2 + \beta R^n$	~ 0.964	< 0.01

Metric vs Palatini: a theoretical comparison

In the metric formalism, the connection is fixed to be the Levi-Civita connection, fully determined by the metric. In contrast, the Palatini formalism treats $\Gamma_{\mu\nu}^\lambda$ as independent, and variation with respect to the connection leads to additional equations that constrain the form of $\Gamma_{\mu\nu}^\lambda$.

For an action of the form:

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} f(R) + S_m[g_{\mu\nu}, \psi], \quad (50)$$

where $R = g^{\mu\nu} R_{\mu\nu}(\Gamma)$, variation yields field equations where the scalar curvature becomes an algebraic function of the trace of the energy–momentum tensor T , in contrast to the dynamical behavior in the metric case.

This fundamental distinction implies:

- **Second-order equations** for the Palatini case (vs fourth-order in metric $f(R)$),
- **No extra propagating scalar** degree of freedom in Palatini $f(R)$,
- **Distinct inflationary dynamics**, especially in early-universe cosmology.

Palatini $f(R)$ inflation: Starobinsky and beyond

In the metric formulation, the Starobinsky model $f(R) = R + \alpha R^2$ leads to an effective scalar–tensor theory with a dynamical scalar field and a slow-roll potential in the Einstein frame. However, in the Palatini formulation, the same action results in algebraic dependence of the curvature on matter and does not produce a dynamical scalar field.

Nonetheless, it still leads to inflationary expansion through modified Friedmann equations, yielding:

- A nearly scale-invariant scalar spectrum,
- **Strong suppression of the tensor-to-scalar ratio r** ,
- Observables consistent with Planck 2018 and BICEP/Keck bounds (Table 16).

Because of the algebraic nature of curvature in Palatini $f(R)$, the theory avoids introducing an additional scalar degree of freedom, simplifying the inflationary analysis and reducing fine-tuning concerns.

Scalar–tensor equivalence and Einstein frame interpretation

While metric $f(R)$ gravity can be recast as scalar–tensor theory with a potential derived from $f(R)$, Palatini $f(R)$ models yield a non-dynamical scalar field in the equivalent Einstein frame. The scalar field behaves as an auxiliary variable that modifies the effective gravitational constant and potential shape but lacks independent dynamics.

As a result:

- The slow-roll condition emerges from the structure of $f(R)$,
- The field excursions are sub-Planckian,
- The Einstein-frame potential is flatter than in the metric case.

Reheating and post-inflation evolution

Reheating in Palatini inflation requires special treatment because standard inflaton oscillations may be absent. Proposed mechanisms include:

- Coupling the modified gravity sector to scalar matter fields,
- Using non-oscillatory reheating mechanisms like instant preheating or gravitational reheating,
- Introducing small corrections to $f(R)$ to produce scalar field dynamics post-inflation.

Several studies have explored particle production in Palatini inflation and shown that the universe can successfully reheat, albeit with different energy transfer mechanisms than in scalar-field-driven models.

Extensions and generalizations

Palatini formalism has been extended to include:

- $f(R, Q)$ models, where $Q = R_{\mu\nu} R^{\mu\nu}$ adds further curvature corrections,
- **Palatini–Higgs inflation**, where the Standard Model Higgs is non-minimally coupled in the Palatini frame, reducing r significantly compared to the metric version,
- **Non-minimal curvature–matter couplings**, relevant for unification of early and late-time acceleration,
- **Quantum corrections**, where Palatini quantization offers different renormalization behavior and avoids certain strong coupling issues.

These models maintain second-order field equations and typically exhibit improved stability, reduced tensor modes, and better UV behavior than their metric counterparts.

Observational viability and theoretical advantages

The Palatini formalism offers several attractive features:

- **Suppressed tensor modes** without needing extreme flatness in the potential,
- **Sub-Planckian field values**, avoiding trans-Planckian concerns,
- **Simplified dynamics**, due to absence of new propagating fields,
- **Compatibility with Higgs inflation**, avoiding large radiative corrections,
- **Robust predictions** that lie within 1- σ Planck contours.

The Palatini formalism offers a compelling alternative to the metric approach in modified gravity and inflationary cosmology. Particularly in $f(R)$ models like Starobinsky inflation, it provides distinct predictions – most notably a much lower tensor-to-scalar ratio – without introducing additional degrees of freedom. Its second-order nature, computational simplicity, and ability to realize inflation without fine-tuned potentials make it a strong candidate for constructing viable early-universe scenarios. Ongoing research in Palatini–Higgs inflation, $f(R, Q)$ gravity, and non-minimal matter couplings continues to expand its relevance in both theoretical and observational cosmology.

16 Metric-affine and $f(Q)$ gravity

Metric-affine theories generalize General Relativity (GR) by lifting the assumption that the affine connection is derived from the metric. In such theories, the affine connection $\Gamma^\lambda_{\mu\nu}$ is an independent geometric object that may exhibit torsion $T^\lambda_{\mu\nu}$ and non-metricity $Q_{\lambda\mu\nu}$. Within this broad class, symmetric teleparallel gravity – also known as coincident gravity – focuses on a unique case: curvature and torsion are both set to zero, and gravity is mediated purely through non-metricity.

This construction leads to a new class of theories known as $f(Q)$ gravity, in which the gravitational action depends on the non-metricity scalar Q , analogous to $f(R)$ in curvature-based gravity and $f(T)$ in torsion-based gravity. Due to its second-order nature and ghost-free structure, $f(Q)$ gravity has recently been explored as a viable setting for inflationary cosmology.

Symmetric teleparallelism: geometry of non-metricity

In symmetric teleparallel geometry, the affine connection is chosen such that:

$$R^\lambda_{\mu\nu\rho} = 0, \quad T^\lambda_{\mu\nu} = 0, \tag{51}$$

but non-metricity is non-vanishing:

$$Q_{\lambda\mu\nu} = \nabla_\lambda g_{\mu\nu} \neq 0. \tag{52}$$

The fundamental scalar governing dynamics is the non-metricity scalar Q , defined as:

$$Q = -\frac{1}{4} Q_{\alpha\mu\nu} Q^{\alpha\mu\nu} + \frac{1}{2} Q_{\alpha\mu\nu} Q^{\mu\alpha\nu} + \frac{1}{4} Q_\alpha Q^\alpha - \frac{1}{2} Q_\alpha \tilde{Q}^\alpha, \tag{53}$$

where $Q_\alpha = Q_{\alpha\mu}{}^\mu$ and $\tilde{Q}^\alpha = Q^\mu{}_\mu{}^\alpha$ are traces of non-metricity.

The gravitational action in $f(Q)$ gravity reads:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} f(Q) + \mathcal{L}_m \right], \tag{54}$$

where $f(Q)$ is a general function of the scalar Q , and \mathcal{L}_m is the matter Lagrangian.

Inflation in $f(Q)$ gravity

The inflationary sector of $f(Q)$ gravity has been studied extensively in recent years. Unlike $f(R)$ or $f(T)$, the field equations in $f(Q)$ gravity remain second-order despite nonlinearity in Q , avoiding Ostrogradsky instabilities. This makes the theory attractive for constructing inflationary models without introducing extra degrees of freedom or auxiliary scalar fields.

Heisenberg, Beltrán Jiménez, Lazkoz, and others have shown that:

- Inflation can emerge solely from geometric terms in $f(Q)$,
- One can reconstruct $f(Q)$ functions from a desired scale factor or Hubble evolution,
- The Starobinsky-like expansion and graceful exit are achievable within this framework [69–71].

Additionally, $f(Q)$ inflation can be implemented with or without scalar fields. In the scalar–non-metricity formulation, scalar fields are coupled minimally or non-minimally to Q , enabling further control over the inflationary potential and dynamics.

Inflationary predictions and observables

Viable models yield predictions for scalar perturbations and tensor-to-scalar ratios in agreement with current CMB constraints. The scalar power spectrum is nearly scale-invariant, and the tensor sector is naturally suppressed, depending on the exact form of $f(Q)$ (Table 17).

Models constructed via reconstruction techniques can yield graceful exit, reheating via geometrical mechanisms,

Table 17 Inflationary predictions in selected $f(Q)$ gravity models

Model	n_s	r
Starobinsky-type $f(Q) = Q + \alpha Q^2$	~ 0.965	~ 0.004
Reconstructed $f(Q)$ from Hubble flow parameters	~ 0.964	< 0.01
Power-law $f(Q) = Q^n$	Model-dependent	Model-dependent

and a phase of post-inflation deceleration compatible with Big Bang nucleosynthesis.

Theoretical advantages of $f(Q)$ gravity

The symmetric teleparallel framework provides several notable advantages:

- **Second-order field equations**, avoiding ghosts and pathologies,
- **No need for auxiliary scalar fields**, inflation from geometry alone,
- **Simple matter coupling**, with geodesics determined by the Levi-Civita connection,
- **Compatibility with metric–affine extensions**, including torsion and disformal structures.

Additionally, symmetric teleparallelism permits the construction of scale-invariant and conformal extensions, and naturally admits Lagrangians that unify inflation and dark energy under a single geometric description.

Perturbations, GW speed, and stability

While the background dynamics are well understood, perturbation theory in $f(Q)$ gravity is more subtle. Recent studies have investigated scalar and tensor modes using gauge-invariant formalisms:

- Scalar perturbations are stable for a wide class of $f(Q)$,
- Tensor modes propagate at light speed, ensuring consistency with GW170817,
- The theory avoids extra polarizations or pathologies in the gravitational wave sector.

However, full cosmological perturbation theory – especially nonlinear and second-order calculations – remains an active area of research, including the formulation of consistent initial conditions and treatment of isocurvature modes.

Generalizations and future directions

Extensions of $f(Q)$ gravity include:

- **Scalar–non-metricity theories**, analogs of scalar–torsion and scalar–curvature models,
- **Conformal $f(Q)$ theories**, useful in Higgs inflation and scale-invariant scenarios,
- **Non-minimal couplings**, such as $\xi(\phi)Q$ for scalar field dynamics,
- **Higher-dimensional and braneworld scenarios**, using the metric-affine formalism.

Quantum corrections and attempts to formulate a UV-complete version of symmetric teleparallel gravity are also underway, possibly connecting to affine quantum gravity and background-independent formalisms.

Metric-affine gravity and its subclass of $f(Q)$ theories offer a rich and geometrically novel platform for modeling inflation and cosmic evolution. By constructing gravity from non-metricity rather than curvature or torsion, symmetric teleparallelism retains second-order dynamics, avoids ghost modes, and permits purely geometric mechanisms for early-universe inflation.

Inflation in $f(Q)$ gravity models yields predictions in excellent agreement with CMB observations, and generalizations allow for unified treatment of both early and late acceleration. With increasing interest in non-Riemannian geometries, metric-affine inflation is poised to play an important role in connecting cosmology with fundamental modifications of gravity.

17 Dynamical system approaches to modified gravity inflation

The dynamical systems approach provides a powerful framework for analyzing inflationary dynamics across a wide range of cosmological models, especially in the increasingly complex landscape of modified gravity. Rather than relying on numerical solutions for specific initial conditions, this method re-expresses the cosmological field equations as an autonomous system of first-order differential equations. This enables a systematic exploration of global behavior in phase space, revealing inflationary attractors, stability conditions, and natural exit mechanisms.

Dynamical systems techniques are particularly useful in theories with multiple fields or non-linear geometric terms, such as $f(R)$, $f(T)$, $f(R, T)$, scalar–tensor models, Gauss–Bonnet theories, mimetic gravity, and even Palatini or metric-affine frameworks.

Methodology: autonomous systems and phase space geometry

To construct a dynamical system, one introduces a set of dimensionless variables that capture the relative importance of kinetic energy, potential energy, curvature corrections, and other relevant terms. For example:

$$x = \frac{\dot{\phi}}{\sqrt{6}H}, \quad y = \frac{\sqrt{V(\phi)}}{\sqrt{3}H}, \quad \lambda = -\frac{1}{V} \frac{dV}{d\phi}, \quad z = \frac{F'(R)}{F(R)}, \tag{55}$$

with $N = \ln a$ as the e-folding number.

The cosmological equations are rewritten as an autonomous system:

$$\frac{dx_i}{dN} = f_i(x_j), \tag{56}$$

where the functions f_i depend on the specific modified gravity theory under study. Fixed points $\{x_i^*\}$ correspond to critical cosmological behaviors such as de Sitter inflation, kinetic-dominated regimes, or scaling attractors. Linearization around these points yields the stability matrix $J_{ij} = \partial f_i / \partial x_j$, whose eigenvalues determine local stability.

Applications across modified gravity theories

Dynamical systems methods have been fruitfully applied to a wide array of modified gravity models:

- **$f(R)$ gravity:** Models like Starobinsky inflation exhibit stable de Sitter points and radiation attractors. For $f(R) = R + \alpha R^2$, inflation appears as a saddle or attractor, depending on the parameters [72].
- **$f(T)$ gravity:** The torsion-based formulation leads to qualitatively different fixed points than curvature-based models. Power-law inflation and phantom-like attractors can be identified, with unique reheating trajectories [73].
- **Scalar-tensor and Brans-Dicke models:** Dynamical system analysis reveals the impact of non-minimal couplings and potential shapes on inflationary dynamics and late-time acceleration [74].
- **$f(R, T)$ and mimetic gravity:** These theories show inflationary fixed points even without explicit potentials, enabled by the interaction between matter and geometry [64, 75].
- **Gauss-Bonnet and scalar-GB theories:** Modified friction from coupling functions $\xi(\phi)G$ introduces rich dynamical structures with stable inflationary trajectories [55].
- **Palatini $f(R)$ and metric-affine theories:** These theories often yield simpler, second-order dynamical systems,




Fig. 1 Phase space portrait of $f(R)$ gravity showing key fixed points

Table 18 Representative fixed points in modified gravity inflation

Point	Interpretation	Scale factor behavior	Stability
P_1	de Sitter inflation	$a(t) \propto e^{Ht}$	Stable node
P_2	Power-law inflation	$a(t) \propto t^p$	Saddle point
P_3	Kinetic-dominated	$w = 1$	Unstable source
P_4	Scaling solution	$\rho_\phi / \rho_m = \text{const}$	Depends on potential

but their phase space shows bifurcations associated with the form of $f(R)$.

- **DHOST and beyond-Horndeski models:** Despite complexity, constrained subspaces allow reduced phase-space dynamics where inflation, scaling solutions, and tracker behavior can be identified.

Phase portraits and flow behavior

A crucial feature of this approach is its ability to visualize the dynamical flow of cosmological variables, aiding in the classification of inflationary attractors and trajectories that exit into radiation or matter-dominated eras. Consider a reduced system in the x - y plane for scalar-tensor gravity (Fig. 1):

Classification of fixed points and stability

Typical fixed points are classified by their cosmological significance (Table 18):

The eigenvalue spectrum of the Jacobian matrix at each point determines whether the solution is a node, focus, center, or saddle.

Advantages and limitations

Advantages:

- Captures full asymptotic behavior of cosmological systems,
- Identifies attractors, fixed points, and trajectories without exact solutions,
- Useful for model selection and inflationary viability,
- Enables systematic reconstruction of potentials or coupling functions.

Challenges:

- Complex models may yield high-dimensional phase spaces,
- Non-autonomous systems may arise (e.g., time-dependent parameters),
- Some physically relevant trajectories (e.g., reheating) are transient and may not appear as fixed points,
- Linear analysis may miss subtle non-linear or bifurcation effects.

The dynamical systems approach is an essential tool in the study of inflationary dynamics across broad classes of modified gravity theories. It provides qualitative and quantitative insights into attractor behavior, cosmological stability, and the transitions between inflation and post-inflationary epochs. As the complexity of gravitational models grows, phase-space analysis remains indispensable for organizing the cosmological solution space, identifying observationally viable regimes, and guiding the reconstruction of theoretically motivated inflationary models.

18 Reheating and preheating in modified gravity

The post-inflationary reheating epoch plays a pivotal role in linking the cold, quasi-de Sitter inflationary phase to the hot Big Bang universe. During this transition, the vacuum-like energy of the inflationary sector is converted into relativistic particles, initiating standard thermal history. In some scenarios, this occurs via *preheating*, an explosive non-perturbative particle production mechanism often driven by parametric resonance.

In the context of *modified gravity*, the reheating process acquires new features. The gravitational degrees of freedom are modified or extended, the background dynamics differ from General Relativity (GR), and extra scalar modes or geometrical couplings can impact decay rates, expansion rates, and reheating efficiency.




Fig. 2 Schematic evolution of inflaton and radiation energy densities during preheating in modified gravity

Reheating in $f(R)$ gravity

In $f(R)$ models – especially the Starobinsky model $f(R) = R + \alpha R^2$ – inflation is driven by the scalaron, a dynamical scalar degree of freedom emerging from the R^2 term. After inflation ends, the scalaron oscillates around the minimum of its effective potential $V_{\text{eff}}(\phi) \sim \phi^2$, and reheating occurs through its gravitational or Yukawa-like couplings to other fields.

Key features include:

- Scalaron mass: $M \sim 10^{13}$ GeV,
- Decay via loop-generated interactions or minimal couplings to Standard Model (SM) scalars/fermions,
- Reheating temperature: $T_{\text{reh}} \sim 10^9 - 10^{10}$ GeV,
- Smooth transition into radiation domination.

Perturbative and non-perturbative analyses have been extensively developed by Gorbunov, Starobinsky, and others [76,77].

Preheating: resonant particle production in modified gravity

Preheating offers a more efficient and rapid mechanism for energy transfer than perturbative reheating. When the inflaton or gravitational scalar oscillates after inflation, it can induce time-varying masses for other fields:

$$\ddot{\chi}_k + \left[\frac{k^2}{a^2} + g^2 \phi^2(t) \right] \chi_k = 0, \tag{57}$$

leading to resonance amplification of field modes χ_k . In modified gravity, this mechanism can be triggered by:

- Non-minimal curvature couplings (e.g., $\xi R\phi^2$),
- Scalar–Gauss–Bonnet couplings $\xi(\phi)G$,
- Kinetic torsion couplings in $f(T)$ or Einstein–Cartan frameworks,
- Oscillations of geometric degrees of freedom in $f(Q)$, mimetic, or non-local models (Fig. 2).

Reheating constraints from observables

The reheating phase modifies the total number of e-folds between horizon exit and the end of inflation:

$$N_k = 55 + \frac{1}{4} \ln \left(\frac{V_k^2}{M_P^4 \rho_{\text{end}}} \right) + \frac{1 - 3w_{\text{reh}}}{12(1 + w_{\text{reh}})} \ln \left(\frac{\rho_{\text{reh}}}{\rho_{\text{end}}} \right), \quad (58)$$

where w_{reh} is the average equation of state. The dependence of n_s and r on T_{reh} and w_{reh} allows observational constraints on reheating. In modified gravity:

- Extra geometric degrees of freedom can alter ρ_{end} ,
- Non-trivial effective equations of state arise,
- Entropy production and decay into hidden sectors are model dependent (Table 19).

Examples of modified gravity reheating

Recent developments and open challenges

Recent studies include:

- **Preheating in GB inflation:** Tachyonic amplification from $\xi''(\phi)G$ triggers efficient reheating [78,79].
- **Reheating in $f(Q)$ gravity:** Without extra scalar fields, reheating may occur via gravitational particle production [80].
- **Einstein–Cartan reheating:** Spin-induced torsion can convert into radiation via fermionic interactions, providing a bounce-like reheating scenario [81].
- **Constraints from CMB:** Planck and future missions can indirectly probe reheating via effects on the inflationary consistency relations and gravitational wave background.

Reheating and preheating in modified gravity constitute a crucial but less-explored frontier. Due to their dependence on gravitational geometry and couplings, these phases can significantly alter the post-inflationary universe and provide unique observational signatures. A proper understanding of reheating dynamics is essential for constructing viable inflationary models and ensuring consistency with both early- and late-time cosmological observations.

19 Primordial black holes and gravitational waves in modified gravity

Inflation is not only a mechanism for generating the large-scale structure of the universe, but it also provides fertile ground for the production of *primordial black holes* (PBHs) and stochastic backgrounds of *gravitational waves* (GWs).

These relics encode information about the small-scale features of inflationary dynamics and offer windows into physics beyond General Relativity (GR).

In particular, modified gravity models – such as $f(R)$, scalar–Gauss–Bonnet, mimetic, teleparallel, and non-local gravities – naturally introduce features that amplify curvature perturbations, alter tensor dynamics, or trigger resonant particle production. This makes them powerful frameworks for probing early-universe relics with future astrophysical and gravitational wave observatories.

Enhanced scalar perturbations and PBH formation

Primordial black holes form when enhanced curvature perturbations at small scales reenter the horizon in the early universe and collapse under their own gravity. A scalar perturbation spectrum $\mathcal{P}_\zeta(k)$ exceeding a threshold value (typically $\sim 10^{-2}$) can trigger such collapse:

$$\mathcal{P}_\zeta(k_c) \gtrsim \delta_c^2 \sim 10^{-2}, \quad (59)$$

where k_c corresponds to the comoving scale of PBH formation. Inflationary models with transient periods of ultra-slow-roll, inflection points, or effective sound speed reduction can naturally amplify small-scale perturbations.

In modified gravity theories, PBH formation is facilitated via:

- **Plateau potentials** in $f(R)$ and $f(G)$ models,
- **Scalar–Gauss–Bonnet couplings** $\xi(\phi)G$ enhancing friction,
- **Mimetic constraints** leading to clustering of scalar modes,
- **Non-canonical kinetic terms** or entropy modes in multi-field extensions (Table 20).

Depending on the model, PBHs may contribute to dark matter, seed supermassive black holes, or generate second-order gravitational waves.

Gravitational waves from modified inflation

In addition to scalar perturbations, inflation generically produces tensor modes, which today form a stochastic gravitational wave background (SGWB). In modified gravity:

- The **tensor sound speed** $c_T \neq 1$ can modify the amplitude and shape,
- The **effective Planck mass** $M_{\text{eff}}^2(t)$ may vary during inflation or reheating,
- **Extra degrees of freedom** (e.g., scalar–torsion or scalar–GB couplings) can lead to new tensor sources,
- **Second-order scalar-induced GWs** arise from enhanced $\mathcal{P}_\zeta(k)$ in PBH-generating models.


Table 19 Reheating properties in selected modified gravity models

Model	Scalar degree	T_{reh} (GeV)	Preheating?
Starobinsky $f(R)$	Scaloron	$10^9\text{--}10^{10}$	Efficient
Scalar–GB ($\xi(\phi)G$)	Yes	Model-dependent	Tachyonic instabilities
$f(T)$ gravity	Varies	Typically lower	Ambiguous
Mimetic gravity	Mimetic scalar	$< 10^8$	Yes (for specific $V(\phi)$)
Non-local gravity	No scalar	Gravitational	Unclear
Palatini $f(R)$	No scalar	Indirect	Weak/absent
$f(Q)$ gravity	No scalar	Gravitational	Possible via expansion effects

Table 20 Examples of PBH formation in modified gravity frameworks

Model	Amplification mechanism	PBH mass range
Scalar–Gauss–Bonnet inflation	Ultra-slow-roll phase from $\xi(\phi)$	$10^{18} - 10^{25}$ g
$f(R)$ with inflection potentials	Plateau region in $V_{\text{eff}}(\phi)$	$10^{15} - 10^{22}$ g
Mimetic gravity	Non-canonical kinetic clustering	Broad; model-dependent
$f(Q)$ models	Dynamical suppression of sound speed	$10^{17} - 10^{23}$ g
Einstein–Cartan with spin fluid	Spin-induced enhancements	10^{18} g and higher

Fig. 3 Sample GW energy density spectra in different modified gravity scenarios. Peaks in the LISA, DECIGO, and PTA bands offer testable signatures



Specific predictions include:

- Scalar-induced GWs from $f(R)$ and mimetic PBH models peaking in the mHz range (LISA),
- Direct tensor GWs from scalar–Gauss–Bonnet or teleparallel oscillations,
- Suppression or enhancement of GW amplitude depending on coupling function $\xi(\phi)$ or $f(Q)$ form,
- Resonant amplification of tensor modes during preheating (tachyonic or parametric) (Fig. 3).

Observational prospects and constraints

The synergy between PBH and GW probes makes this sector especially promising for testing high-energy gravity:

- **LISA, DECIGO, BBO:** Sensitive to scalar-induced and tachyonic GW backgrounds in the mHz–Hz regime,
- **PTAs (e.g., SKA, NANOGrav):** Constrain nHz GWs from early phases of modified gravity or domain walls,
- **Microlensing surveys (HSC, OGLE, EROS):** Bound PBH abundances across mass ranges,
- **CMB-S4 and LiteBIRD:** Indirectly constrain the primordial tensor spectrum via r and its scale dependence.

Future multi-messenger constraints on stochastic GW spectra and PBH populations can jointly narrow the allowed inflationary model space.

Modified gravity frameworks not only enrich inflationary dynamics but also profoundly impact the generation of PBHs and gravitational waves. These phenomena serve as probes of ultraviolet physics, geometric couplings, and reheating

mechanisms. As GW detectors and PBH searches improve, they offer one of the most powerful

20 Inflation in braneworld and higher-dimensional theories

The idea that our observable universe is a lower-dimensional hypersurface (brane) embedded in a higher-dimensional spacetime (bulk) provides a compelling extension to standard inflationary cosmology. Inspired by string theory, M-theory, and extra-dimensional gravity, braneworld and higher-dimensional models offer rich mechanisms for generating inflation, modifying Friedmann dynamics, and yielding observational signatures distinct from those of 4D General Relativity.

Braneworld inflation typically arises in models such as the Randall–Sundrum (RS) type II scenario, the Dvali–Gabadadze–Porrati (DGP) model, and their generalizations. In these frameworks, standard matter fields are confined to a 3-brane, while gravity propagates into extra spatial dimensions, altering both the background dynamics and cosmological perturbation theory. In higher-dimensional theories derived from string compactifications, inflation is often driven by moduli fields, brane–antibrane interactions, or the motion of D-branes in warped geometries.

Modified Friedmann equations on the brane

In RS-type braneworlds, the Einstein equations projected onto the 3-brane lead to a modified Friedmann equation of the form:

$$H^2 = \frac{\rho}{3M_p^2} \left(1 + \frac{\rho}{2\lambda}\right) + \frac{\mathcal{E}}{a^4}, \tag{60}$$

where λ is the brane tension and \mathcal{E} is a dark radiation term arising from the bulk Weyl tensor. At high energies ($\rho \gg \lambda$), the quadratic term dominates, modifying the slow-roll conditions and allowing inflation with steeper potentials.

This deviation has several immediate consequences:

- Inflation can occur for potentials that would be too steep in standard 4D GR,
- The number of e-folds is modified as the friction term is enhanced,
- Tensor perturbations are suppressed relative to scalar perturbations due to modified normalization.

The high-energy correction enhances the Hubble friction, slowing down scalar field evolution and extending the inflationary phase even for otherwise marginal potentials. These features make brane inflation scenarios attractive for embedding within supergravity or string-derived landscapes.

Examples of braneworld inflation models

- **RS-II braneworld inflation:** The simplest inflationary scenario in RS models employs standard scalar potentials (e.g., $m^2\phi^2$, $\lambda\phi^4$), but due to the quadratic corrections, the dynamics and spectral predictions change. The scalar spectral index n_s and tensor-to-scalar ratio r become:

$$n_s \simeq 1 - \frac{6\epsilon - 2\eta}{1 + V/\lambda}, \quad r \simeq \frac{16\epsilon}{1 + V/\lambda}, \tag{61}$$

where ϵ, η are the usual slow-roll parameters.

- **DGP braneworld inflation:** In the DGP model, the Friedmann equation includes a term linear in H due to gravity leakage into the bulk:

$$H^2 \pm \frac{H}{r_c} = \frac{\rho}{3M_p^2}, \tag{62}$$

where r_c is the crossover scale between 4D and 5D behavior. The self-accelerating branch (+) can generate late-time acceleration, while the normal branch (−) allows for viable inflationary dynamics when coupled with scalar fields on the brane. However, ghost instabilities have challenged the consistency of the self-accelerating branch.

- **Mirage and DBI inflation:** Derived from the dynamics of D3-branes moving in warped throats (e.g., Klebanov–Strassler geometry), Dirac–Born–Infeld (DBI) inflation introduces non-canonical kinetic terms:

$$\mathcal{L} = -f(\phi)^{-1} \sqrt{1 - f(\phi)(\partial\phi)^2} + f(\phi)^{-1} - V(\phi), \tag{63}$$

leading to inflation with large sound speed suppression. These models can produce detectable non-Gaussianities and may be constrained by bounds on the equilateral shape of the bispectrum.

- **KKLMMT and moduli inflation:** In the KKLT framework, inflation is driven by the motion of D3-branes toward anti-D3-branes in warped throats, with the potential arising from moduli stabilization and supersymmetry breaking. These models typically yield small-field inflation and must overcome the η -problem via tuning or additional mechanisms.

Gravitational perturbations and brane effects

Inflation in higher dimensions significantly modifies the perturbation equations. Key effects include:

- **Suppressed tensor modes:** Due to localization of gravity on the brane and redshift of tensor modes into the bulk,

- **Modified consistency relations:** For example, the standard relation $r = -8n_t$ may not hold in brane models,
- **High-energy corrections:** Affect the amplitude and tilt of scalar and tensor power spectra,
- **KK mode excitations:** Massive graviton modes can leave imprints in the CMB anisotropies and GW spectrum.

The full perturbation analysis in these theories requires solving coupled 4D–5D systems (e.g., the Mukohyama master equation) and often involves numerical simulations or approximations.

Non-Gaussianity and signatures

In several braneworld inflation models – especially DBI-type and multi-brane setups – the non-canonical structure leads to enhanced non-Gaussianity. Observationally:

- The non-linearity parameter $f_{\text{NL}}^{\text{eq}}$ can reach order 10 – 100,
- Shape dependence is often equilateral or orthogonal, distinguishing them from local-type non-Gaussianity,
- Tensor–scalar consistency relations are broken, offering a test of higher-dimensional effects.

The presence of isocurvature modes from extra-dimensional fields or moduli can further affect the perturbation spectrum and reheating history.

Observational constraints and prospects

Observational constraints from Planck, BICEP/Keck, and large-scale structure have already begun constraining these scenarios:

- DBI inflation is constrained by bounds on f_{NL} and the tensor-to-scalar ratio,
- KKLMMT-type models require fine-tuning or additional stabilizing ingredients to be viable,
- RS-II inflation is consistent with data for specific potential shapes and values of λ .

Future experiments such as LiteBIRD and CMB-S4 will tighten constraints on tensor modes and non-Gaussianities, potentially distinguishing brane inflation from standard 4D models. Gravitational wave observatories may also detect relic GWs from brane-localized inflation or KK-graviton leakage.

Inflation in braneworld and higher-dimensional theories provides an elegant extension to early-universe cosmology, embedding inflation into a higher-dimensional or string-theoretic setting. These frameworks modify the Friedmann

dynamics, scalar and tensor spectra, and non-Gaussian signatures in profound ways. The interplay between brane tension, extra-dimensional dynamics, and compactification geometry allows for a wide range of inflationary behaviors.

As observational data becomes increasingly precise, inflation in higher dimensions remains a testable and exciting avenue – offering a bridge between quantum gravity, string theory, and cosmology.

21 Bayesian model selection and machine learning approaches to inflation

Bayesian inference has become an indispensable tool in modern cosmology, providing a rigorous framework for comparing theoretical models against increasingly precise observational data. In the context of inflation, Bayesian methods enable one to quantify how well a given inflationary scenario fits Cosmic Microwave Background (CMB) anisotropy and polarization data, including results from *Planck*, *BICEP/Keck*, and large-scale structure surveys.

In parallel, machine learning (ML) techniques have emerged as powerful tools for accelerating and augmenting cosmological data analysis. Their synergy with Bayesian methods enables efficient navigation of high-dimensional parameter spaces, rapid model discrimination, and even potential-free reconstruction of inflationary dynamics from data.

Bayesian evidence and model comparison

The Bayesian evidence (or marginal likelihood) for a model \mathcal{M}_i is defined as:

$$Z_i = P(D|\mathcal{M}_i) = \int \mathcal{L}(\theta_i)\pi(\theta_i) d\theta_i, \quad (64)$$

where $\mathcal{L}(\theta_i)$ is the likelihood function of the data D , and $\pi(\theta_i)$ is the prior distribution over the model parameters θ_i . Model comparison is then performed using the Bayes factor:

$$B_{ij} = \frac{Z_i}{Z_j}, \quad (65)$$

which quantifies the relative support for model \mathcal{M}_i over \mathcal{M}_j . Interpreting $\ln B_{ij}$ follows the Jeffreys' scale: values > 5 indicate strong preference.

Bayesian inference is typically implemented using Monte Carlo Markov Chain (MCMC) methods or nested sampling algorithms. Notable tools include:

- **CosmoMC:** an MCMC engine coupled with CAMB for Boltzmann integration,

- **MultiNest** and **PolyChord**: efficient nested samplers for high-dimensional and multi-modal posterior distributions,
- **Cobaya**: a modular framework incorporating multiple samplers and cosmological modules.

Applications to modified gravity inflation

Modified gravity inflationary models often introduce extended parameter spaces beyond the minimal Λ CDM model. Bayesian model selection has been employed to evaluate:

- Viability of $f(R) = R + \alpha R^2$ vs. power-law inflation,
- Preference for plateau vs. monomial potentials in $f(T)$, $f(G)$, and $f(Q)$ theories,
- The role of non-minimal couplings in scalar–tensor models,
- The impact of quantum corrections or running kinetic terms in k -inflation or Galileon-type models.

Such analyses often constrain the parameter combinations $(n_s, r, \alpha_s, f_{\text{NL}})$, along with model-specific couplings $\alpha, \beta, \xi, \lambda$, and allow computation of posterior credible intervals.

A representative study by Rinaldi et al. [82] showed that Palatini $f(R)$ inflation yields tighter posterior distributions for r , often favored over chaotic inflation. Recent work by Escudero et al. [83] also performed Bayesian comparison across a wide class of single- and multi-field inflationary potentials using Planck+BAO data.

Machine learning for inflationary cosmology

The growth of cosmological datasets has motivated the application of ML techniques, not only to accelerate parameter estimation but also to explore the structure of theory space. Some key applications include:

- **Reconstruction of the inflaton potential:** Using supervised learning, ML can be trained on simulated CMB observables to learn an inverse mapping from observables (e.g., n_s, r) to potential shapes $V(\phi)$. Gaussian process regression (GPR) and variational autoencoders have been used for this task, providing non-parametric reconstructions with uncertainty quantification.
- **Classification of models:** Neural networks (e.g., convolutional and recurrent architectures) have been trained to distinguish among inflationary models, especially in distinguishing between:
 - Canonical vs. non-canonical kinetic terms,
 - Single-field vs. multi-field scenarios,

– Plateau, hilltop, and chaotic potentials.

- **Emulators and surrogates:** ML surrogates such as neural emulators or random forests can emulate expensive Boltzmann solvers like CLASS or CAMB, reducing inference time from hours to milliseconds. This enables faster scans over modified gravity models with extended parameter spaces.
- **Anomaly detection and forecasting:** Unsupervised learning is used to detect departures from standard inflation (e.g., running, features, oscillations) in future surveys such as LiteBIRD, CMB-S4, and the Simons Observatory.

Synergy of ML and Bayesian tools

ML algorithms can be integrated within Bayesian pipelines in multiple ways:

- Use ML emulators as fast likelihood evaluators within MCMC/nested sampling,
- Employ reinforcement learning to explore rare or fine-tuned inflationary trajectories,
- Apply Bayesian neural networks to generate posterior distributions directly with uncertainty quantification.

This synergy allows cosmologists to explore complex theory spaces that would otherwise be computationally prohibitive.

Challenges and future directions

While promising, ML and Bayesian methods face challenges:

- The choice of priors can heavily influence Bayesian evidence, especially in high-dimensional spaces,
- ML models can overfit or fail to generalize outside trained regimes unless carefully regularized,
- Interpretability of deep learning results remains limited, motivating hybrid analytic-ML frameworks.

Nonetheless, these approaches are expected to play an increasingly central role in the precision cosmology era, especially as datasets from future probes (e.g., CMB-S4, Euclid, SKA) begin to constrain inflationary features with unprecedented accuracy.

Bayesian inference and machine learning together provide a powerful statistical and computational framework for the analysis of inflationary models in modified gravity. They enable efficient exploration, model discrimination, and reconstruction of early-universe dynamics. As cosmological data continue to grow in precision and volume, these tools

will be crucial for identifying viable theories of inflation that go beyond the standard GR framework.

22 Summary of viable inflationary models in modified gravity

Inflationary cosmology embedded in modified gravity offers a rich array of theoretical frameworks, each with its unique geometric motivations, inflationary mechanisms, and observational signatures. In this section, we provide a consolidated comparison of the most promising classes of models, evaluating them based on:

- Compatibility with current observations (CMB, BICEP/Keck, LSS),
- Theoretical consistency and stability (e.g., ghost-freedom, second-order field equations),
- Prospects for ultraviolet (UV) completion (e.g., links to string theory or loop quantum gravity).

Model landscape and comparative table

See Table 21.

Model taxonomy and geometric classification

We provide a conceptual taxonomy of inflationary models in modified gravity, structured according to their geometric origin – whether based on curvature, torsion, non-metricity, or higher-derivative extensions – as well as the nature of their field-theoretic couplings. This classification complements the detailed summary and phenomenological comparison tables in Sects. 21 and 25, offering a broader perspective on the theoretical landscape covered in this review.

We distinguish seven major categories:

- **Curvature-based models:** These include $f(R)$ gravity and scalar fields non-minimally coupled to curvature invariants such as the Ricci scalar R and the Gauss–Bonnet term G . These models typically originate from higher-order corrections to the Einstein–Hilbert action and include Starobinsky-type and scalar– G models.
- **Torsion-based models:** In these scenarios, gravity is mediated by torsion rather than curvature. Examples include $f(T)$ gravity, Einstein–Cartan theory, and scalar–torsion coupling models, which offer second-order field equations and unique reheating mechanisms.
- **Non-metricity-based models:** These arise from the symmetric teleparallel framework, in which gravity is expressed through the non-metricity tensor. The $f(Q)$ class of models maintains second-order field equations

and offers elegant alternatives to curvature- or torsion-based scenarios.

- **Higher-derivative and non-local gravity:** This class introduces infinite-derivative form factors or non-local operators such as $R\mathcal{F}(\square)R$. These models are motivated by quantum gravity and string field theory and can support UV-complete, singularity-free inflationary dynamics.
- **Scalar–geometry couplings:** These models explore scalar fields coupled to curvature tensors such as $R_{\mu\nu}$, torsion vectors T^μ , and non-metricity traces Q_μ . They generalize scalar–tensor theories and can lead to strong friction effects or enhanced preheating.
- **Exotic frameworks:** These include MODified Gravity (MOG or STVG), mimetic gravity, Bekenstein’s varying constant theories, and Carmeli’s cosmological relativity. Though less conventional, these models propose novel geometric or physical mechanisms for inflation.
- **String-inspired and extra-dimensional models:** These are derived from brane dynamics, moduli stabilization, or axionic fields in string theory. They include D-brane inflation, axion monodromy, and scenarios based on warped compactifications or tachyonic DBI actions.

By categorizing these models via their underlying geometry – curvature, torsion, or non-metricity – and their physical mechanisms – scalar couplings, higher derivatives, extra dimensions – we clarify their similarities, differences, and potential observational signatures. This classification also aids in guiding future observational strategies to distinguish between competing inflationary scenarios.

Observational status

Recent observations from *Planck* 2018, BICEP/Keck 2018, and LSS surveys provide stringent bounds on key inflationary parameters such as the scalar spectral index n_s , tensor-to-scalar ratio r , and non-Gaussianity f_{NL} . Notably:

- Models predicting $n_s \sim 0.9649 \pm 0.0042$ and $r < 0.06$ are currently favored,
- Many $f(R)$, scalar–GB, and non-local models lie well within these bounds,
- Some $f(T)$ and mimetic models require parameter tuning to avoid instabilities or inconsistent reheating.

Theoretical robustness

Theoretical consistency is critical for inflationary models. Second-order field equations (as in $f(Q)$ and Palatini $f(R)$) are preferred due to the absence of Ostrogradsky instabilities. Covariant reformulations are important in torsion-based theories like $f(T)$ to preserve local Lorentz invariance. Cou-

Table 21 Viable inflationary models in modified gravity: status and outlook

Model	Obs. viable	Stability	UV completion
$f(R)$ Starobinsky	Yes	Ghost-free; scalar–tensor form	Likely (SUGRA, string)
$f(G)$ Gauss–Bonnet	Yes (tuned)	Possible ghost/gradient issues	Heterotic string-based
$f(T)$ Torsion	Model-dependent	Algebraic; non-covariant	Reformulations ongoing
Scalar–GB coupling	Yes (broad)	Needs $\xi'(\phi)\dot{\phi} < 0$	Natural in string theory
$f(Q)$ non-metricity	Promising	2nd-order; ghost-free	New field; UV unclear
Braneworld (RS/DGP)	Yes with tuning	Ghost-free in some setups	String/M-theory support
Mimetic gravity	Some viable	Gradient instabilities possible	UV unclear; loop ideas exist
Einstein–Cartan	Viable at high density	Algebraic torsion; ghost-free	Possible LQG origin
Non-local gravity	Yes	Ghost-free by design	Linked to string field theory
Palatini $f(R)$	Yes	2nd-order; no scalaron	Minimal UV links; stable

pling functions $\xi(\phi)$ in scalar–GB models must be chosen to avoid ghost and gradient instabilities, while Einstein–Cartan models remain algebraically constrained but singularity-free.

UV completion and embedding in high-energy theory

Some inflationary models naturally emerge from or are compatible with UV-complete frameworks:

- **Starobinsky inflation** appears in supergravity and no-scale models,
- **Scalar–Gauss–Bonnet** models arise in low-energy string compactifications,
- **Braneworld inflation** is supported by M-theory and RS-like warped geometry,
- **Non-local gravity** finds roots in string field theory and p -adic gravity,
- Other frameworks like mimetic, torsion, and non-metricity-based inflation are promising but require further embedding in UV-complete theories.

Role of upcoming observations

Forthcoming missions such as LiteBIRD, CMB-S4, the Simons Observatory, SKA, and gravitational wave detectors (e.g., LISA, DECIGO) will further refine parameter bounds. They are expected to:

- Constrain r to the 10^{-3} level,
- Detect or exclude primordial gravitational waves,
- Probe reheating and PBH production indirectly,
- Discriminate between geometric origins of inflation (e.g., curvature, torsion, non-metricity).

Data-driven and ML-augmented exploration

As discussed in Sect. 26, machine learning and Bayesian inference will continue to play a central role in the model selection process. With vast theoretical landscapes and high-dimensional parameter spaces, data-driven workflows will be essential to:

- Rapidly rule out disfavored models,
- Identify inflationary attractors in extended theories,
- Reconstruct inflaton dynamics or geometric actions directly from observables.

Inflationary cosmology in modified gravity continues to offer fertile ground for exploring fundamental questions about the early universe and quantum gravity. Among the numerous candidate models, $f(R)$ Starobinsky inflation, scalar–Gauss–Bonnet scenarios, non-local gravity, and Palatini extensions remain at the forefront due to their predictive power and compatibility with data. Emerging frameworks like $f(Q)$ gravity and Einstein–Cartan theory offer novel mechanisms worthy of further investigation.

The next decade of precision cosmology will be decisive in narrowing the inflationary paradigm – confronting theory with data and guiding us toward the correct geometric description of the universe’s earliest moments.

23 Tachyon inflation in modified gravity

Tachyon fields arise naturally in the context of string theory, particularly in scenarios involving unstable D-branes. The decay of these D-branes leads to the emergence of a tachyon condensate, described effectively by a non-canonical scalar field with Dirac–Born–Infeld (DBI) dynamics. Remarkably, tachyon fields have been shown to support inflationary phases

in both standard and modified gravity frameworks, yielding rich phenomenological predictions.

The general action for tachyonic inflation in the Einstein frame is given by:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - V(T) \sqrt{1 + g^{\mu\nu} \partial_\mu T \partial_\nu T} \right], \quad (66)$$

where T is the tachyon field and $V(T)$ is the effective potential governing its dynamics. Unlike canonical fields, the tachyon’s kinetic term saturates as $\dot{T}^2 \rightarrow 1$, which naturally slows down the field and supports prolonged inflation.

Motivation and theoretical foundations

Tachyon inflation is deeply rooted in Type IIA/B string theory, where open strings on non-BPS D-branes lead to a rolling tachyon condensate. The action resembles the DBI Lagrangian in brane-world scenarios, and tachyon matter behaves like a pressureless fluid at late times.

Key theoretical motivations include:

- Natural UV embedding from string theory compactifications,
- Stability under higher-derivative corrections,
- Non-trivial connections with DBI inflation, K-inflation, and warm inflation,
- Realization of both inflationary and dark energy epochs within a unified framework.

Inflationary dynamics: background and perturbations

In a spatially flat FRW background, the tachyon field evolves under:

$$H^2 = \frac{\kappa^2}{3} \frac{V(T)}{\sqrt{1 - \dot{T}^2}}, \quad \ddot{T} + (1 - \dot{T}^2) \left(3H\dot{T} + \frac{V'}{V} \right) = 0. \quad (67)$$

The slow-roll approximation is valid when $\dot{T}^2 \ll 1$ and $\ddot{T} \ll 3H\dot{T}$. The effective slow-roll parameters become:

$$\epsilon = \frac{M_P^2}{2} \left(\frac{V'}{V} \right)^2 \frac{1}{V^2}, \quad \eta = M_P^2 \left[\frac{V''}{V^3} - \frac{3}{2} \left(\frac{V'}{V^2} \right)^2 \right]. \quad (68)$$

These expressions differ from canonical cases due to the non-standard kinetic structure, leading to modified predictions for inflationary observables.

Common potentials and predictions

Various functional forms for $V(T)$ have been explored:

- **Inverse square:** $V(T) = V_0/T^2$, motivated by string decays,
- **Exponential:** $V(T) = V_0 e^{-\alpha T}$, supports graceful exit,
- **Gaussian:** $V(T) = V_0 \exp(-\alpha T^2)$, allows PBH production,
- **Hilltop:** $V(T) = V_0(1 - \alpha T^2)$, fine-tuned slow-roll,
- **Padé form:** $V(T) = V_0/(1 + \alpha T^2)^n$, has built-in exit (Table 22).

Modified gravity embeddings

Tachyon fields can be embedded in various modified gravity theories:

- In $f(R)$ gravity, the gravitational sector modifies the effective friction term, altering inflation duration and reheating [84].
- In $f(T)$ teleparallel gravity, torsion couples naturally to tachyon kinetic terms. Geng et al. [85] show that slow-roll inflation and reheating are achievable.
- In scalar–Gauss–Bonnet frameworks, tachyons coupled to curvature invariants generate rich dynamics and potential amplification of gravitational waves.

These embeddings yield distinctive corrections to n_s and r , and can suppress non-Gaussianity or enhance reheating via geometric couplings.

Reheating and preheating

Tachyon models often require a mechanism to reheat the universe after inflation, as the field asymptotically rolls without oscillation. Possible mechanisms include:

- **Coupling to gauge fields:** $f(T)F_{\mu\nu}F^{\mu\nu}$,
- **Gravitational particle production:** effective in low \dot{T} regimes,
- **Hybrid-type models:** additional scalar field triggers waterfall transition.

Recent studies show that even tachyon fields in braneworld scenarios can yield successful reheating via parametric resonance [86].

Dynamical systems analysis and attractor behavior

To analyze global stability, one can define variables:

$$x = \dot{T}, \quad y = \frac{V(T)}{3H^2}, \quad \Omega_T = \frac{V(T)}{\sqrt{1 - x^2} 3H^2}. \quad (69)$$

Setare and Sami [87] demonstrated that:

- Tachyon-dominated attractors exist,

Table 22 Representative tachyon inflation potentials and observational features

Model	$V(T)$	n_s	r
Inverse square	V_0/T^2	~ 0.960	< 0.01
Exponential	$V_0e^{-\alpha T}$	$0.95 - 0.97$	$\lesssim 0.1$
Gaussian	$V_0e^{-\alpha T^2}$	~ 0.964	< 0.01
Hilltop	$V_0(1 - \alpha T^2)$	~ 0.97	Small
Padé-type	$V_0/(1 + \alpha T^2)^n$	Tunable	Tunable

- Late-time scaling solutions are possible,
- Exit from inflation occurs naturally if $V(T) \rightarrow 0$ fast enough.

Observational constraints and non-Gaussianity

Tachyon inflation is compatible with current observational data when the sound speed c_s is near unity. However, in the DBI regime, deviations lead to:

- **Enhanced equilateral-type non-Gaussianity**, quantified by $f_{NL}^{equil} \sim \mathcal{O}(1)$,
- **Suppressed tensor modes** ($r < 0.01$), due to reduced inflaton speed,
- **PBH formation possibilities** via enhanced scalar spectra in flat regions of $V(T)$.

Constraints from Planck 2018 allow modest levels of f_{NL} and favor potentials leading to red-tilted spectra. Future probes like LiteBIRD and CMB-S4 will decisively test these models.

Theoretical challenges and outlook

While tachyon inflation is theoretically appealing, it faces some challenges:

- The field does not oscillate post-inflation, complicating standard reheating,
- Potential singularities in DBI kinetic terms must be avoided,
- The effective field theory cutoff may lie below the inflationary Hubble scale,
- Embedding into consistent superstring compactifications is model-dependent.

Nevertheless, when embedded in extended gravity (e.g., $f(T)$, scalar-GB, non-local), many of these problems are softened. Further work is required to address UV sensitivity and define well-behaved quantum corrections.

Tachyon inflation remains a viable and string-motivated framework for early-universe acceleration, especially when embedded within modified gravity theories. Its non-canonical

dynamics, distinctive reheating mechanisms, and observational predictions make it a fertile ground for theoretical and phenomenological exploration. The next generation of cosmological and gravitational wave surveys will further probe the unique signatures of tachyon-based inflation and its geometric generalizations.

24 Inflation in string-inspired gravity

String theory provides a compelling UV-complete framework that naturally incorporates gravity, gauge interactions, and scalar fields. Inflationary cosmology within this setting benefits from a higher-dimensional perspective, leading to novel mechanisms for driving early-universe acceleration. Unlike traditional scalar field inflation, string-inspired models involve rich structures arising from compactification, D-brane dynamics, flux-induced potentials, and higher-curvature corrections.

The scalar fields responsible for inflation often descend from moduli (e.g., dilaton, volume modulus), axions, or open string modes (brane positions), and their potentials are determined by the geometry and topology of the extra dimensions. The compactification process – typically on Calabi–Yau or toroidal manifolds – generates effective 4D scalar-tensor theories with modified kinetic terms, non-minimal couplings, and multi-field dynamics.

Fundamental classes of string-inspired inflation

String-inspired inflation models can be broadly classified as follows:

- **Brane–antibrane inflation:** A D3-brane moves in a warped throat towards an anti-D3-brane, and their attractive potential serves as the inflaton potential. The prototypical realization is the KKLMNT scenario [88], which includes warping and moduli stabilization.
- **DBI inflation:** The kinetic energy of a brane moving in a warped background is governed by a Dirac–Born–Infeld action, leading to non-canonical inflation with a small sound speed and potentially large non-Gaussianities [89].

- **Axion monodromy inflation:** Arising from the winding of axions around non-trivial cycles, these models produce linear or sinusoidal potentials with periodic modulations, offering controlled large-field inflation [90].
- **Dilaton and moduli inflation:** Fields from compactification, such as the dilaton or volume modulus, can drive inflation under exponential or plateau-like potentials. Loop and flux corrections are used to stabilize the moduli [91,92].
- **String gas and pre-big bang cosmology:** Utilizing T-duality and string thermal dynamics, these models propose loitering or bouncing phases that may seed inflation or structure formation [93,94].

Modified gravity and effective action structure

In the low-energy limit, string-inspired inflation leads to extended gravity models beyond Einstein’s General Relativity. The generic 4D effective action may include:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} R + f(\phi)R + \frac{1}{2}k(\phi)(\nabla\phi)^2 - V(\phi) + \xi(\phi)G + \dots \right], \tag{70}$$

where:

- $f(\phi)$: non-minimal coupling (e.g., dilaton gravity),
- $k(\phi)$: non-canonical kinetic terms (e.g., DBI),
- $\xi(\phi)G$: coupling to the Gauss–Bonnet term from string α' corrections.

These actions support generalized inflationary dynamics, slow-roll attractors, and viable reheating phases while avoiding ghosts and higher-derivative instabilities if constructed carefully.

Phenomenology and observational constraints

Observationally viable string-inspired inflation requires matching Planck data, BICEP/Keck bounds, and future CMB constraints. Key phenomenological features include:

- **Spectral index n_s :** Most models predict $n_s \sim 0.96 - 0.97$,
- **Tensor-to-scalar ratio r :** DBI and monodromy models span a wide range ($r \sim 10^{-3} - 0.1$),
- **Non-Gaussianity:** DBI-type models may produce large equilateral f_{NL} , constrained by Planck ($f_{NL}^{equil} < 50$),
- **Isocurvature modes:** Multi-field inflation (e.g., axion inflation with saxions) introduces potential isocurvature perturbations and trajectory turns (Fig. 4).




Fig. 4 Inter-brane potential in warped throat geometry relevant to DBI and brane–antibrane inflation

Challenges and theoretical consistency

Despite their appeal, several theoretical challenges persist:

- **Moduli stabilization:** Unstabilized moduli (e.g., volume modulus) can disrupt inflationary dynamics or lead to fifth-force constraints.
- **Backreaction:** Brane dynamics or warping may induce significant corrections to the potential and kinetic terms.
- **Swampland conjectures:** The Swampland Distance Conjecture and de Sitter Conjecture restrict the field range and gradient of scalar potentials [95,96], placing pressure on large-field models.
- **Reheating and exit:** Some models, especially DBI inflation, lack well-defined reheating mechanisms without introducing additional fields or couplings.

These challenges have motivated hybrid inflation, two-field models, and multi-throat geometries to circumvent constraints.

Recent advances and extensions

In the past few years, new directions have emerged:

- **Multi-axion inflation (N-flation):** Realistic compactifications yield multiple axions with aligned potentials [97,98].
- **Clockwork axion inflation:** Dynamically generates super-Planckian decay constants [99].
- **Modular inflation:** Uses modular forms (e.g., Eisenstein series) for symmetry-stabilized potentials [100].
- **String–Gauss–Bonnet and non-local stringy inflation:** Incorporates curvature corrections from higher orders in the α' or g_s expansions [101,102] (Fig. 5).

Diagram of axion monodromy

Inflation in string-inspired gravity offers a rich arena where quantum gravity, high-energy field theory, and cosmological observations intersect. While achieving full theoretic-




Fig. 5 Quasi-periodic axion potential from monodromy inflation, supported by wrapped branes and fluxes

cal control remains difficult, these models continue to drive progress in connecting fundamental theory with the inflationary paradigm. Future observations from CMB-S4, Lite-BIRD, gravitational wave probes (LISA, BBO), and large-scale structure surveys will be instrumental in confirming or ruling out stringy inflationary scenarios.

25 Inflation in conformal and Weyl-invariant gravity

Conformal symmetry, or invariance under local rescalings of the metric, is a powerful principle in field theory and gravity. In conformal (or Weyl-invariant) gravity, the action remains invariant under the transformation:

$$g_{\mu\nu}(x) \rightarrow \Omega^2(x)g_{\mu\nu}(x), \tag{71}$$

where $\Omega(x)$ is a smooth, non-zero function of spacetime coordinates. This symmetry naturally eliminates any fixed mass or energy scales from the theory, making it an attractive framework for addressing problems related to quantum gravity, cosmological constant, and scale-free early-universe dynamics.

Theoretical motivation

Conformal gravity models date back to Weyl’s original attempts at unifying gravity and electromagnetism, but modern interest has surged due to their appearance in:

- String theory effective actions,
- Supergravity and higher-derivative gravity,
- Asymptotically safe gravity and anomaly-induced inflation,
- Dynamical scale generation in Higgs inflation and dilaton models.

The conformal symmetry must be spontaneously or anomalously broken to recover a standard cosmology, and the mechanism of this breaking provides the inflationary dynamics.

Generic action and symmetry realization

A typical conformally invariant action includes a scalar field ϕ , often identified with the inflaton or dilaton, coupled non-minimally to the Ricci scalar:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{12} \phi^2 R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{\lambda}{4} \phi^4 \right], \tag{72}$$

which is invariant under:

$$g_{\mu\nu} \rightarrow \Omega^2(x)g_{\mu\nu}, \quad \phi \rightarrow \Omega^{-1}(x)\phi. \tag{73}$$

In this setting, inflation arises from spontaneous breaking of conformal symmetry when the field ϕ develops a vacuum expectation value (VEV), setting the Planck scale.

Anomaly-induced inflation

Quantum corrections to a classically conformal theory break the symmetry through trace anomalies. Starobinsky and others showed that the anomaly-induced effective action includes R^2 and $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$ terms, leading to:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} R + \alpha R^2 + \beta C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} \right], \tag{74}$$

where $C_{\mu\nu\rho\sigma}$ is the Weyl tensor. These models naturally yield Starobinsky-type inflationary dynamics, without requiring a fundamental inflaton.

Conformal attractors and α -models

Conformal attractors are a broad class of models with universal predictions in the n_s - r plane. These models often begin with conformal symmetry and introduce a field-space transformation that maps them to Einstein frame models with plateau-like potentials (Table 23):

$$V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3\alpha}} \phi/M_P} \right)^2. \tag{75}$$

Gauge fixing and Einstein frame equivalence

These theories often use conformal gauge fixing to relate different frames. Choosing the Einstein gauge $\phi = \sqrt{6}M_P$ yields a canonical Einstein–Hilbert action plus a scalar potential. Conversely, choosing a gauge where the metric is fixed allows ϕ to play the role of a varying Planck mass. This duality facilitates model building and numerical analysis.

Table 23 Inflationary predictions from conformal gravity-inspired attractor models

Model	Spectral index n_s	Tensor-to-scalar ratio r
Conformal Starobinsky-like	~ 0.965	~ 0.004
Conformal α -model with $\alpha = 1$	~ 0.965	~ 0.008
Higgs-dilaton inflation	~ 0.964	< 0.01
Anomaly-induced $R^2 + C^2$	~ 0.966	~ 0.003

Table 24 Contrasting conformal gravity inflation with standard scalar field models

Feature	Canonical inflation	Conformal gravity inflation
Symmetry	None or shift symmetry	Local conformal invariance
Mass scales	Introduced by hand	Generated dynamically
Fundamental scalars	Inflaton ϕ	Dilaton or anomaly scalar
UV completion	Often absent	Motivated by SUGRA, strings
Predictions	Model-dependent	Universality classes in n_s, r

Inflation without potentials: conformal gravity proper

A radical class of models based on Weyl gravity uses the action:

$$S = \int d^4x \sqrt{-g} C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma}, \tag{76}$$

where dynamics arise from higher-order geometric operators without any scalar field. Inflation-like expansion emerges via conformal symmetry breaking or induced anomalies. These models are highly constrained by unitarity and observational viability but remain theoretically intriguing (Fig. 24).

Comparison with canonical inflation

See Table 24.

Observational prospects

Conformal models are consistent with CMB observations when symmetry breaking is handled appropriately. Notably:

- Most predict small $r < 0.01$, aligned with Planck and BICEP/Keck.
- Running and non-Gaussianity are typically negligible.
- The UV structure and anomaly-driven inflation offer testable predictions via quantum gravity and holography.

Recent developments and open problems

Active areas of research include:

- Weyl-invariant extensions of supergravity and Horndeski models.
- Conformal inflation in holographic cosmology and AdS/CFT.
- Dynamical conformal symmetry breaking in early-universe phase transitions.
- Compatibility with quantum anomalies and radiative corrections.

Despite theoretical elegance, challenges include:

- Ghost instabilities in pure Weyl models,
- Lack of a standard reheating mechanism,
- Complexity in matching late-time cosmology.

Conformal and Weyl-invariant gravity models offer a symmetry-based pathway to inflation without invoking arbitrary mass scales. By breaking scale invariance dynamically or via quantum effects, they naturally lead to viable inflationary expansion with predictive power and close connections to quantum gravity theories.

26 Phenomenological models of inflation

Phenomenological models of inflation aim to capture the essential features of early-universe acceleration without committing to a full UV-complete theory such as string theory or supergravity. These models are typically constructed using effective field theory (EFT) techniques, often involving a single scalar field ϕ minimally coupled to gravity and evolving under a potential $V(\phi)$. The strength of phenomenological models lies in their simplicity, analytic tractability, and direct connection to observable cosmological parameters such as the scalar spectral index n_s , the tensor-to-scalar ratio r , and non-Gaussianity parameters.

Classification of inflationary potentials

Phenomenological models can be grouped into universality classes based on their asymptotic behavior, shape of the potential, and symmetry structure. The most common categories include:

Table 25 Predictions of key phenomenological inflationary models

Model	$V(\phi)$	n_s	r	Comments
Quadratic chaotic	$\frac{1}{2}m^2\phi^2$	~ 0.966	~ 0.13	Disfavored
Starobinsky	$V_0(1 - e^{-\sqrt{2/3}\phi/M_{Pl}})^2$	~ 0.965	~ 0.004	Excellent fit
Natural inflation	$\Lambda^4[1 + \cos(\phi/f)]$	$0.96 - 0.97$	$0.05 - 0.1$	Viable if $f \gtrsim 5M_{Pl}$
Hilltop ($p = 2$)	$V_0(1 - \phi^2/\mu^2)$	~ 0.96	$\lesssim 0.01$	Viable
α -attractor ($\alpha = 1$)	$V_0 \tanh^2(\phi/\sqrt{6\alpha})$	~ 0.965	~ 0.004	Robust

- **Large-field inflation:** Models such as chaotic inflation [103] with monomial potentials $V(\phi) \propto \phi^n$, where inflation occurs for super-Planckian field values. These predict relatively large values of r and are increasingly constrained by CMB observations.
- **Small-field or hilltop inflation:** The field rolls away from a local maximum near $\phi = 0$, using potentials like $V(\phi) = V_0(1 - \phi^p/\mu^p)$ with $p > 0$. These models typically predict small r , but may face issues with initial conditions.
- **Plateau models:** These include Starobinsky-type potentials and the broader class of α -attractor models [104, 105]. The potential flattens at large ϕ , leading to universal predictions $n_s \approx 1 - 2/N$, $r \approx 12\alpha/N^2$.
- **Hybrid inflation:** Inflation is driven by one field but ends due to an instability in a second field – the waterfall field – allowing controlled reheating and embedding into high-energy models [106].
- **Natural and axion-like inflation:** Models with periodic potentials $V(\phi) = \Lambda^4[1 + \cos(\phi/f)]$ or monodromy-type extensions, inspired by shift symmetries and axion physics [90, 107].
- **Logarithmic, exponential, and non-polynomial forms:** These emerge in quantum-corrected potentials, brane inflation, and holographic dual models [108, 109].

Comparative predictions and constraints

Recent observations from Planck 2018 and BICEP/Keck strongly constrain inflationary models, particularly ruling out large-field monomial potentials with $n \geq 2$. Viable models must produce a spectral tilt $n_s \sim 0.965$, low running $\alpha_s \sim 10^{-3}$, and tensor-to-scalar ratio $r < 0.036$ (95% CL). Table 25 summarizes benchmark potentials and their predictions.

Phenomenological models in modified gravity

Many effective potentials have been embedded into general-relativity frameworks:

- **$f(R)$ gravity:** The Starobinsky model corresponds to $f(R) = R + \alpha R^2$. Other plateau models arise via conformal transformations from Jordan to Einstein frames [110, 111].
- **$f(T)$ gravity:** Power-law and exponential inflation potentials have been constructed via teleparallel modifications with torsion [13, 22].
- **Einstein–Cartan theories:** Hilltop or chaotic inflation with torsion couplings modify the slow-roll parameters and reheating behavior [112].
- **Scalar–Gauss–Bonnet:** Phenomenological potentials coupled to G can realize ultra-slow-roll or exit from inflation, enhancing scalar perturbations and PBH formation [113].

These extensions preserve the empirical success of the original potentials while offering novel dynamics, modified sound speeds, and corrections to the consistency relation $r = -8n_T$.

Bayesian selection and machine learning classifiers

The large landscape of inflationary models necessitates a statistical approach to model comparison:

- **Bayesian evidence \mathcal{Z}** integrates the likelihood over prior parameter space and allows pairwise model comparison via Bayes factors $B_{ij} = \mathcal{Z}_i/\mathcal{Z}_j$ [108].
- **Nested sampling tools** like PolyChord, MultiNest, and CosmoMC have been used to quantify the statistical preference for α -attractors over monomial or hilltop models.
- **Machine learning methods** including neural networks, Gaussian process regression, and support vector machines are increasingly applied to classify viable models, accelerate parameter estimation, and reconstruct the inflationary potential from data [114, 115].

Advantages and limitations

Phenomenological models enjoy the following merits:

- Simplicity and analytical control.
- Direct fit to observables (n_s, r, α_s, f_{NL}).
- Flexibility to test new ideas and coupling schemes.

However, these models also face important limitations:

- Lack of UV completion or embedding in quantum gravity.
- Potentially ad hoc forms for $V(\phi)$, lacking symmetry justification.
- Difficulty in modeling post-inflationary reheating or entropy production.

Phenomenological inflation continues to serve as a key interface between theory and observation. With upcoming surveys (LiteBIRD, CMB-S4, SKA) and better large-scale structure data, many models will be further constrained or ruled out. The synergy between analytical model building, data-driven inference, and computational methods like ML will guide the search for realistic and falsifiable inflationary scenarios.

27 Inflation in Modified Gravity with anisotropic tensor (MOG)

MODified Gravity (MOG), also known as scalar–tensor–vector gravity (STVG), is a relativistic theory introduced by Moffat as a unified framework to explain cosmological observations without invoking dark matter or dark energy [116]. In the context of inflation, MOG introduces extra degrees of freedom beyond General Relativity (GR), including a massive vector field ϕ_μ , a running gravitational coupling $G(x)$, and scalar fields $\mu(x), \omega(x)$ that control the vector field’s mass and coupling strength. These additions fundamentally alter the background evolution, tensor modes, and perturbative behavior during the inflationary era.

Theoretical framework

The action for STVG/MOG is composed of several interacting sectors:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{16\pi G(x)} R + \mathcal{L}_\phi + \mathcal{L}_S + \mathcal{L}_{\text{matter}} \right], \tag{77}$$

where:

- $G(x)$ is promoted to a scalar field, whose dynamics are governed by a separate scalar field Lagrangian \mathcal{L}_S ,

- ϕ_μ is a Proca-type vector field with its own mass term and kinetic structure,
- \mathcal{L}_ϕ includes Maxwell-like and mass terms for ϕ_μ ,
- $\mathcal{L}_{\text{matter}}$ represents matter and radiation fields minimally coupled to the metric.

The full set of field equations derived from this action lead to a highly non-trivial modification of Einstein’s equations, containing contributions from the vector field stress-energy tensor, derivatives of the scalar fields, and effective anisotropic stress.

Cosmological dynamics and inflationary mechanisms

MOG modifies the Friedmann equations via three principal contributions:

1. The time dependence of $G(x)$ introduces a dynamical effective Planck mass $M_{\text{eff}}(t) \sim G^{-1/2}(t)$, thereby altering the strength of Hubble friction and impacting slow-roll conditions.
2. The massive vector field ϕ_μ can generate anisotropic stress during inflation, particularly if it couples non-minimally to the inflaton or contributes a non-negligible energy density.
3. The scalar field $\mu(x)$, governing the mass of ϕ_μ , can modulate the evolution of tensor modes.

Inflation in MOG can proceed in two main scenarios:

- **Minimal coupling scenario:** Standard inflaton ϕ minimally coupled to a MOG background with evolving $G(t)$, leading to enhanced friction and extended inflation even for steep potentials.
- **Vector-driven inflation:** The energy density of ϕ_μ contributes directly to the Hubble parameter, enabling vector-dominated or mixed-sourced inflation, possibly yielding small r and stable anisotropies [117].

Perturbations and tensor anisotropies

A critical feature of MOG inflation is the anisotropic stress induced by the vector field. Although the vector field typically decays during expansion, in MOG it is dynamical and massive, allowing its fluctuations to survive and impact the CMB via:

- Modifying the tensor power spectrum and generating statistical anisotropies.
- Affecting scalar–tensor cross correlations due to coupling with ϕ and $G(x)$.
- Producing distinctive signatures in the B-mode polarization power spectrum.

The evolution equations for scalar and tensor perturbations must be modified to include the effective energy-momentum tensor of ϕ_μ , which includes mass, pressure, and anisotropic components. These corrections can shift the consistency relation $r = -8n_T$ and affect the spectral tilt n_s through corrections to the inflaton slow-roll parameters.

Observational signatures and comparison with data

MOG-based inflationary models yield modified predictions for the scalar spectral index n_s , the tensor-to-scalar ratio r , and the non-Gaussianity f_{NL} , depending on the specific inflationary potential and the dynamics of the additional fields. Key predictions include:

- Lower r values for given monomial potentials due to enhanced Hubble damping,
- Small but detectable anisotropy in the tensor sector for certain ϕ_μ configurations,
- Running of n_s and α_s due to time evolution of $G(x)$ and $\mu(x)$,
- Potential violations of parity symmetry if the vector field has Chern–Simons-type couplings.

Recent analyses suggest that MOG inflation with standard potentials like $V(\phi) = m^2\phi^2/2$ or plateau-like forms can fit Planck 2018 and BICEP2/Keck constraints if the field evolution of $G(x)$ is carefully controlled to avoid instabilities post-inflation [118].

Bayesian evidence and parameter space viability

Although MOG is not as widely tested as other models like $f(R)$ or Gauss–Bonnet inflation, recent Bayesian analyses have begun to explore its likelihood landscape. Using nested sampling tools (PolyChord, MontePython), one can:

- Estimate the posterior distribution of inflationary parameters with evolving $G(t)$,
- Assess the impact of vector anisotropy terms on CMB observables,
- Compare MOG-based models with conventional single-field inflation in terms of Bayesian evidence \mathcal{Z} ,
- Classify the viable potentials via ML methods or model-selection statistics (e.g., Bayes factor B_{ij}).

Preliminary results indicate that for certain parameter ranges (particularly sub-Planckian field excursions), MOG inflation performs comparably to plateau-type models in standard GR frameworks.

Challenges and theoretical issues

Despite its interesting features, inflation in MOG faces several theoretical and phenomenological challenges:

- **Stability of perturbations:** Vector field fluctuations can lead to ghost or gradient instabilities unless carefully tuned.
- **Fine-tuning of scalar field masses:** To match observational constraints, $\mu(x)$ and $G(x)$ must evolve slowly, requiring tuning of the scalar field potentials.
- **Reheating and entropy production:** The end of inflation must allow for efficient conversion of energy into Standard Model particles while ensuring consistency with late-time MOG cosmology.

MOG inflation opens new avenues to test gravity at high energies via its unique coupling structure and dynamical field content. Future directions include:

- Development of full gauge-invariant perturbation theory for MOG inflation,
- Computation of non-Gaussianity and bispectra arising from vector and scalar couplings,
- Embedding MOG in a broader EFT of inflation to assess quantum corrections,
- Cross-checking inflationary predictions with MOG dark matter fits at galactic and cluster scales.

Upcoming data from LiteBIRD, CMB-S4, and large-scale structure surveys could provide key constraints on tensor anisotropies and test the unique signatures of MOG inflation.

28 Inflation in cosmological relativity: Carmeli’s theory

Cosmological relativity (CR), introduced by Moshe Carmeli [119], is a five-dimensional extension of General Relativity (GR) that incorporates a cosmic velocity dimension alongside the usual four spacetime coordinates. In CR, the universe is embedded in a phase space defined by spacetime and Hubble flow velocity. This approach offers a radical departure from traditional scalar-field-based inflationary theories by proposing that the early universe’s rapid expansion is a direct geometric consequence of the structure of space–velocity, rather than being sourced by a vacuum energy or inflaton potential.

Theoretical foundations of Carmeli’s 5D framework

Carmeli’s extension is based on the principle that the universe is best described in the five-dimensional manifold

($x^0 = ct, x^1, x^2, x^3, x^4 = \tau v$), where v represents the radial velocity of cosmological objects and τ is a universal constant with units of time, corresponding to the Hubble time $\tau \approx H_0^{-1}$. The fifth coordinate is not compactified but treated on equal footing with spacetime variables.

The fundamental line element in CR (in the absence of gravitational fields) is given by:

$$ds^2 = \tau^2 dv^2 - dx^2 - dy^2 - dz^2. \quad (78)$$

This geometric structure leads to a set of modified Einstein-like field equations in five dimensions:

$$G_{\mu\nu}^{(5)} + \Lambda^{(5)} g_{\mu\nu}^{(5)} = \kappa T_{\mu\nu}^{(5)}, \quad (79)$$

where the energy–momentum tensor includes velocity contributions, and $\Lambda^{(5)}$ may or may not be needed depending on the cosmological regime.

The remarkable outcome is that cosmic acceleration arises intrinsically due to the geometry of expanding space, eliminating the need for a cosmological constant or dark energy. The Hubble law is recovered naturally, and expansion history equations are modified due to the velocity dimension.

Inflation as a geometric consequence

In CR, inflation is not postulated through a scalar field with a slow-roll potential but is instead encoded in the initial conditions and evolution of the scale–velocity geometry. During the early phase of the universe, the relation between cosmic time and velocity leads to an exponential-like expansion of the form:

$$a(t) \propto \exp\left(\frac{t}{\tau}\right), \quad (80)$$

without requiring vacuum energy. Key inflationary features emerge:

- **Accelerated expansion from first principles:** The geometry of the 5D line element leads directly to a scale factor with de Sitter-like behavior.
- **No scalar field dynamics:** Inflation does not depend on potential tuning, mass terms, or fine-tuned couplings.
- **Graceful exit built-in:** As the universe expands and the relative velocity field saturates, the expansion rate slows, transitioning naturally into a radiation-dominated universe.

Comparison with traditional inflation

Carmeli's approach stands in sharp contrast with scalar field inflation, as shown in Table 26.

While scalar field inflation is flexible and allows a variety of potential shapes, CR inflation is more constrained but achieves acceleration without additional degrees of freedom.

Cosmological implications and observational fits

CR makes several quantitative and qualitative predictions:

- **Flat Universe:** The geometry requires spatial flatness, in agreement with *Planck* results.
- **Type Ia supernovae fit:** Carmeli cosmology fits supernova data without the need for a cosmological constant or dark energy [120, 121].
- **Age of the Universe:** In CR, the age is determined by τ and Ω_m , yielding values consistent with current observations.
- **No need for cold dark matter:** Large-scale structure formation can be explained geometrically, removing the necessity for exotic cold dark matter.

However, it remains unclear how CR models match with the observed tensor-to-scalar ratio or the CMB power spectrum in full detail. A rigorous treatment of linear perturbations in 5D CR is still lacking in the literature.

Summary table of predictions and observations

The entries in Table 27 summarize the key differences between Carmeli relativity (CR) and the standard general relativistic (Λ CDM) cosmology. The first four rows address broad cosmological features: spatial curvature, the need for dark energy, the estimated age of the Universe, and the role of dark matter. These are presented alongside observational benchmarks, such as *Planck* constraints on flatness and Type Ia supernova data. The final row replaces the schematic expansion-rate diagram of the previous version (Fig. 6) with a textual comparison: CR predicts a natural exponential growth of the scale factor without invoking an inflationary scalar field, while GR begins with a radiation-dominated era characterized by a power-law expansion $a(t) \propto t^{1/2}$, followed – under standard inflationary scenarios – by a rapid inflationary phase. By presenting this information in tabular form, we avoid potential ambiguities from graphical plots and provide a concise, quantitative side-by-side reference that is easier to verify against observational results.

Open questions and theoretical challenges

Carmeli's theory, though elegant, faces open issues that must be addressed to be fully competitive with standard inflation:

- **Perturbation theory in 5D:** The generation of density fluctuations and tensor modes must be developed from first principles in the 5D framework.
- **No mechanism for quantum fluctuations:** While classical expansion is geometrically driven, it is unclear how primordial perturbations are seeded.


Table 26 Key differences between scalar field inflation and CR-based inflation

Feature	Scalar field inflation	Carmeli theory (CR)
Inflation mechanism	Potential $V(\phi)$ slow-roll	5D spacetime-velocity geometry
Source of acceleration	Vacuum energy (or R^2)	Velocity expansion geometry
Graceful exit mechanism	Reheating	Saturation of v -expansion
Free parameters	$V(\phi), m_\phi, \xi$ etc.	Only τ and Ω_m
Reheating required	Yes	Not explicitly required
Tensor modes r	Model-dependent	Suppressed naturally
Quantum fluctuations	Sourced by inflaton	Geometry-sourced (needs full treatment)

Table 27 Comparison of Carmeli relativity (CR) predictions with standard GR cosmology

Feature	Carmeli relativity (CR)	Standard GR cosmology
Spatial curvature	Predicts exact spatial flatness without fine-tuning; naturally consistent with <i>Planck</i> data.	Spatial flatness requires inflationary mechanism or fine-tuning of initial conditions.
Supernova type Ia fit	Fits data without invoking dark energy or Λ ; expansion arises from extra-velocity dimension.	Requires Λ or dark energy to fit observed acceleration.
Age of Universe	Age determined by (τ, Ω_m) ; typically in the range 13.7 ± 0.2 Gyr for best-fit values.	Age derived from H_0, Ω_m , and Ω_Λ ; standard Λ CDM gives 13.8 Gyr.
Dark matter requirement	Explains large-scale structure without cold dark matter; baryonic matter suffices in CR framework.	Requires cold dark matter to match galaxy rotation curves and structure formation.
Early-time expansion	Natural exponential expansion in the Carmeli framework without inflationary scalar fields.	Radiation-dominated era with $a(t) \propto t^{1/2}$ followed by inflationary expansion.

Fig. 6 Schematic evolution of the equation of state $w(t)$ during reheating for different modified gravity theories



- **Lack of reheating formalism:** Unlike scalar-field inflation, there is no established method for transferring energy to Standard Model particles in CR.
- **Integration with quantum gravity:** CR is classically formulated and has no known embedding in string theory or loop quantum gravity.

Despite these challenges, the simplicity and predictive nature of CR make it a valuable alternative framework worth further investigation.

Carmeli's cosmological relativity offers a compelling paradigm where inflation is not a phase driven by an exotic scalar field but a natural geometric feature of a five-dimensional spacetime–velocity manifold. It reduces the reliance on arbitrary parameters, avoids fine-tuned potentials, and predicts acceleration from symmetry principles. Further development of its perturbative structure, quantum fluctuation mechanisms, and connections to standard cosmology could make it a serious contender in explaining the earliest moments of the universe.

29 Inflationary insights from Jacob Bekenstein's theoretical frameworks

Jacob Bekenstein is widely recognized for formulating the foundational principles of black hole thermodynamics, particularly the concept of black hole entropy proportional to horizon area. However, his lesser-known but equally profound contributions to scalar field theories with varying fundamental constants offer insightful perspectives for inflationary cosmology. Bekenstein's frameworks bridge gravitational thermodynamics, scalar–tensor dynamics, and cosmological evolution – laying conceptual groundwork that influences modern varying-constant inflation models.

Varying fundamental constants as dynamical fields

In the early 1980s, Bekenstein proposed a model in which the fine-structure constant α could evolve in time and space due to the dynamics of a scalar field ψ [122]. Unlike fixed constants in standard field theory, Bekenstein's framework promoted dimensionless couplings to dynamical variables governed by an action principle.

The Bekenstein–Sandvik–Barrow–Magueijo (BSBM) model later extended this idea to cosmology [123], treating the variation of $\alpha = e^2/(\hbar c)$ via a scalar field coupled to the electromagnetic Lagrangian:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{16\pi G} - \frac{\omega}{2} \partial_\mu \psi \partial^\mu \psi - V(\psi) - \frac{1}{4} B_F(\psi) F_{\mu\nu} F^{\mu\nu} + \mathcal{L}_{\text{matter}} \right], \quad (81)$$

where $B_F(\psi)$ encodes how the electromagnetic field strength depends on the scalar field ψ . This scalar field effectively controls the evolution of α , and through its potential $V(\psi)$, it may induce a period of accelerated expansion akin to inflation.

Inflation via scalar–coupling–driven dynamics

Although Bekenstein did not originally frame his theory in the context of inflation, the scalar field ψ can naturally drive inflation under suitable potentials:

- **Effective inflaton:** The varying α field acts analogously to an inflaton, sourcing energy density via its potential and kinetic terms.
- **Coupling-induced damping:** The interaction between ψ and matter/radiation sectors induces frictional damping that enhances slow-roll conditions.
- **Built-in exit mechanism:** Inflation ends when ψ settles into a vacuum expectation value (VEV), stabilizing α at its current value.

This structure can be naturally embedded in scalar–tensor theories and exhibits features of dilaton inflation, scalar–electromagnetic coupling, and even multifield assisted inflation frameworks. Some models exhibit graceful exit via symmetry restoration or coupling decoupling at late times.

Cosmological predictions and observables

Bekenstein-type inflationary models yield a range of predictions:

- **Scalar spectral index:** Depending on the form of $V(\psi)$, models can match Planck data with $n_s \approx 0.96$.
- **Tensor modes:** Many scenarios suppress tensor modes ($r \lesssim 0.01$) due to sub-Planckian field ranges or modified gravity friction.
- **Running and non-Gaussianity:** Higher-order corrections can arise from scalar–matter couplings and field-space curvature.
- **CMB signatures of α -variation:** Shifts in recombination redshift, baryon–photon sound speed, and fine-structure-dependent spectra offer a complementary test of these models.

Observational searches for spatial and temporal variation in α using quasar absorption lines, Oklo natural reactors, and atomic clock experiments place upper bounds of $|\Delta\alpha/\alpha| \lesssim 10^{-7}$, which constrain the dynamics of ψ today but allow rich evolution in the early universe.

Comparison with canonical models

Bekenstein-inspired inflation models provide an alternative to traditional slow-roll models in GR. They possess the structure of non-minimally coupled or varying coupling models and thus fit within a broader class of extended inflationary frameworks. Table 28 highlights the key differences.

Extensions and embeddings in modified gravity

Bekenstein's framework inspired many generalizations:

- **Varying speed of light (VSL) theories:** Bekenstein's model inspired VSL cosmologies where the dynamics of fundamental constants alter horizon and flatness solutions [124, 125].
- **Scalar–electromagnetic couplings:** Modern quintessence models and varying- α dark energy models adopt similar Lagrangian couplings [126].
- **Einstein–Aether and Horndeski extensions:** Bekenstein's kinetic structure generalizes to higher-derivative and vector-tensor theories with Lorentz-violating backgrounds.
- **String-theoretic and brane scenarios:** In string compactifications, dilaton–gauge field couplings take the Bekenstein form, and moduli stabilization naturally mimics scalar coupling inflation.

Legacy and contemporary relevance

Bekenstein's theories, though not inflationary per se, expanded the conceptual toolkit of early universe cosmology:

- They anticipated the now common idea of scalar fields modifying gravitational or gauge couplings.
- They introduced observational probes of inflationary dynamics through fundamental constants rather than only metric perturbations.
- They offered a physically motivated connection between cosmology, thermodynamics, and quantum field theory in curved spacetime.

Today, models of varying couplings are tested against quasar observations, atomic clock constraints, and CMB data. The philosophy behind Bekenstein's approach – embedding inflationary dynamics within fundamental variations of the constants of nature – remains a viable and compelling frontier in cosmology.

30 Inflation and extra dimensions

The idea that our universe may contain extra spatial dimensions beyond the familiar three has profound implications for both fundamental physics and cosmology. Rooted in Kaluza–Klein theory and refined through string theory and M-theory, higher-dimensional models offer novel mechanisms for realizing inflation, modifying cosmological dynamics, and connecting the early universe with UV-complete quantum gravity frameworks.

Inflation in extra-dimensional theories typically arises either from geometric effects in braneworld scenarios or from dynamical fields emerging through compactification in string theory. These models provide alternative routes to inflation that can yield distinctive predictions for cosmological observables, such as the tensor-to-scalar ratio r , spectral index n_s , and primordial non-Gaussianities.

Braneworld inflation and high-energy corrections

In braneworld cosmology, our observable universe is envisioned as a 3+1-dimensional brane embedded in a higher-dimensional bulk spacetime. The most studied models include the Randall–Sundrum type II scenario, where the bulk is a five-dimensional Anti-de Sitter space (AdS_5) with a single positive-tension brane. The presence of extra dimensions modifies the Einstein equations on the brane, leading to a corrected Friedmann equation of the form:

$$H^2 = \frac{8\pi G}{3} \rho \left(1 + \frac{\rho}{2\lambda} \right) + \frac{\mathcal{C}}{a^4}, \quad (82)$$

where λ is the brane tension and \mathcal{C} represents a dark radiation term from the bulk Weyl tensor [127]. At high energies ($\rho \gg \lambda$), the quadratic correction enhances the effective Hubble friction, allowing inflation to proceed even with steep scalar potentials that would be ruled out in standard 4D cosmology.

This framework supports inflationary models such as chaotic, exponential, and hilltop potentials with modified slow-roll conditions and reheating dynamics. The brane–bulk energy exchange and the radion (interbrane distance) stabilization play crucial roles in ensuring a graceful exit and a transition to standard Big Bang cosmology.

String-theoretic realizations: moduli and brane inflation

In string theory, the existence of 6 or 7 extra compactified dimensions gives rise to a rich set of scalar fields, known as moduli, which control the shape, size, and flux configuration of the compactification manifold (e.g., Calabi–Yau or G_2 holonomy spaces). Inflation can be driven by:

Table 28 Comparison of Bekenstein-type scalar variation models with canonical single-field inflation

Feature	Bekenstein model	Canonical inflation
Driving mechanism	Varying $\alpha(\psi)$ scalar field	Scalar inflaton with $V(\phi)$
Coupling to matter	Yes (via gauge sector)	Minimal or none
Effective potential	Electromagnetically induced	Arbitrary (phenomenological or string-inspired)
Tensor modes r	Typically small	Model-dependent
Exit mechanism	Stabilization of ψ	Violation of slow-roll conditions
Additional signals	Variation in fine-structure constant	None

Table 29 Comparison of higher-dimensional and 4D inflationary scenarios

Feature	Extra-dimensional inflation	4D canonical inflation
Dimensionality	≥ 5 bulk + brane	4D spacetime
Inflation driver	Moduli, brane motion, warped throats	Inflaton potential $V(\phi)$
Hubble friction	Enhanced at high energy densities	GR-derived (standard)
Tensor modes r	Typically suppressed	Depends on slow-roll conditions
Non-Gaussianity	Often large (e.g., DBI)	Small unless specialized mechanisms
Reheating mechanism	Tachyon decay, brane annihilation	Inflaton decay, preheating
Field content	Branes, moduli, axions, KK modes	Single or multifield scalar fields
UV completion	Naturally embedded in string theory	Usually effective-field-theory based

- **Volume modulus inflation:** where the inflaton corresponds to the logarithm of the internal volume [128],
- **Axion inflation:** from stringy generalizations of natural inflation using compactified antisymmetric tensor fields [90],
- **Brane inflation:** such as the KKLMMT scenario, where a mobile D3-brane moves toward a static anti-D3-brane in a warped throat, generating an inflationary potential via the DBI action [88].

Inflation ends with brane annihilation, reheating is triggered via tachyon condensation, and non-Gaussian signatures may be enhanced due to the non-canonical kinetic terms.

Phenomenological features and observables

Extra-dimensional inflationary models often lead to novel observational predictions that can distinguish them from conventional 4D theories:

- **Suppressed tensor-to-scalar ratio:** In RS2-like models, the enhancement of Hubble friction at high energies reduces the amplitude of gravitational waves, leading to $r \ll 0.01$.
- **Modified consistency relations:** Tensor tilt n_T deviates from standard slow-roll predictions, impacting the interpretation of B-mode polarization.
- **Isocurvature and entropy modes:** Multiple moduli and brane fields can source correlated isocurvature fluctuations, depending on post-inflation dynamics.

- **Resonant or oscillatory power spectrum features:** Induced by moduli oscillations, monodromy, or Kaluza–Klein excitations.
- **High non-Gaussianity:** DBI and multifield brane inflation models generically produce equilateral-type non-Gaussianities, constrained by Planck data.

These effects are subject to current and future constraints from surveys such as Planck, CMB-S4, LiteBIRD, and the Simons Observatory (Table 29).

Comparative landscape of inflation models

Challenges and future prospects

Despite their theoretical appeal, extra-dimensional inflationary models face several challenges:

- **Moduli stabilization:** Ensuring the stability of volume and shape moduli during and after inflation.
- **Backreaction and loop corrections:** Especially in warped compactifications and brane inflation.
- **UV sensitivity:** Inflationary dynamics are often sensitive to the details of string compactification, requiring careful top-down construction.

Nonetheless, higher-dimensional models offer fertile ground for unifying quantum gravity with cosmology. Future data on primordial tensor modes, non-Gaussianity, and reheating physics may provide decisive evidence supporting (or falsifying) specific classes of extra-dimensional inflation.

31 Comparative assessment and discriminators among models

To respond to the constructive feedback of both referees, we present a comparative assessment of all the modified gravity models discussed in this review. This section emphasizes the predictive power, observational compatibility, theoretical consistency, and potential discriminators that allow one to distinguish among these frameworks.

Unified inflationary predictions

Inflationary models across modified gravity theories differ in their predictions for the scalar spectral index n_s , tensor-to-scalar ratio r , running α_s , and reheating properties. Table 30 synthesizes key inflationary predictions and theoretical attributes of the primary models.

Model-discriminating observables

Several key observables can serve as discriminators between modified gravity scenarios:

- **Tensor-to-scalar ratio r :** The value of r varies significantly across models. $f(R)$ and $f(Q)$ models predict small r , while braneworld and tachyon models can accommodate larger values depending on the energy scale and brane tension.
- **Non-Gaussianity f_{NL} :** Canonical single-field models predict small f_{NL} , whereas models involving DBI kinetics, higher-derivative mimetic gravity, or multi-field dynamics (e.g., moduli inflation) can produce large and detectable non-Gaussianity.
- **Gravitational waves (GWs):** Modified propagation speed, altered tensor spectrum shape, and amplitude suppression/enhancement due to coupling (e.g., scalar–Gauss–Bonnet or $f(T)$ models) can be probed via CMB B-mode polarization or space-based GW interferometers (LISA, BBO, DECIGO).
- **Reheating signatures:** Different reheating mechanisms (scalaron decay, spin–torsion oscillation, brane collision, or geometric dissipation) leave imprints on post-inflation expansion history and can be tested via spectral distortion or residual particle background.
- **Primordial black holes (PBHs):** Models with enhanced scalar power spectra (e.g., Gauss–Bonnet, mimetic, plateau+feature) can overproduce PBHs. Correlating PBH mass spectrum with inflationary model class offers another discriminator.

Future probes and prospects

Planned and ongoing missions offer critical tests to distinguish among these frameworks:

- **CMB stage-IV (CMB-S4), LiteBIRD:** Precision measurements of n_s , r , and possibly α_s with uncertainties $< 10^{-3}$.
- **Gravitational wave observatories:** LISA, DECIGO, and pulsar timing arrays (PTAs) will test early-universe tensor spectra, scalar-induced GWs, and reheating signals.
- **Large-scale structure and 21-cm surveys:** Can probe isocurvature modes and ultra-slow-roll features specific to modified models.
- **Bayesian inference and machine learning pipelines:** Allow model comparison in high-dimensional parameter space and marginalization over theoretical priors.

Synthesis

From the accumulated evidence, it is clear that:

1. Several modified gravity inflation models remain observationally viable and can fit current Planck+BICEP data.
2. Certain classes, like $f(R)$ Starobinsky, scalar–Gauss–Bonnet, and $f(Q)$ gravity, offer compelling minimalist mechanisms with geometric origin and predictive power.
3. Models predicting larger r (e.g., tachyon, brane) may be increasingly constrained by upcoming B-mode polarization measurements.
4. Theoretical consistency (e.g., avoidance of ghosts, second-order field equations) favors specific geometrically motivated models (e.g., $f(Q)$, non-local, Einstein–Cartan).
5. Future multi-messenger cosmology and precision GW cosmology will be key in eliminating degeneracies among different inflationary mechanisms.

This integrated summary sets the stage for a more informed conclusion, identifying which models stand the best chance of surviving future scrutiny and which are already on the verge of exclusion.

32 Open problems and future directions in inflationary modified gravity

Despite considerable progress in modeling inflation using modified gravity frameworks, several unresolved issues remain. These open questions highlight the current theoretical limitations and guide future research directions at the

Table 30 Inflationary features across modified gravity models

Model	n_s	r	α_s	Reheating	UV completion	Constraints
$f(R)$ (Starobinsky)	0.965	~ 0.004	Small	Scalaron decay	Yes (QG)	Favored by CMB
$f(T)$ gravity	0.962–0.965	< 0.01	Small	Model-dependent	Under study	r, c_s degeneracy
Scalar–GB	0.963–0.966	0.003–0.01	Small	Coupling to G	String-based	GW limits apply
$f(Q)$ gravity	0.964	< 0.01	Negligible	Geometric or scalar	Recent ideas	Needs refinement
Einstein–Cartan	0.963–0.965	< 0.01	Negligible	Spin effects	Partial	Sparse perturbation work
Mimetic gravity	~ 0.964	0.003–0.01	Variable	Scalar/derivative terms	Needs UV	Stability bounds
Non-local gravity	0.964–0.966	$\lesssim 0.01$	Small	Geometry decay	Ghost-free UV	Model-dependent
Braneworld (RS-II)	0.963	< 0.07	Negligible	Tachyon/brane decay	Extra-dim.	λ constraints
Tachyon DBI	0.96–0.97	< 0.01	Possible	Tachyon decay	String-based	n_s, f_{NL} limits
Carmeli CR	Matches Λ CDM	–	–	Built-in geometry	5D relativistic	No CMB predictions
MOG/STVG	~ 0.964	< 0.01	Small	Scalar-vector feedback	Alt. to DM	Vector stability issues

Table 31 Selected theoretical limitations in major modified gravity models

Model	Ghost-free?	Well-posed perturbations?	Ambiguity/challenge
$f(R)$	Yes (in metric)	Yes	Higher-order derivatives
$f(T)$	Often yes	Frame-dependent	Tetrad choice ambiguity
$f(Q)$	Yes	In development	Non-metricity parametrization
Scalar–GB	No (generically)	Potential instabilities	Coupling tuning needed
Einstein–Cartan	Yes	Underexplored	Quantization of torsion

interface of cosmology, quantum gravity, and high-precision observations.

1. Theoretical challenges and internal consistency

While many models fit existing cosmological data, their deeper theoretical consistency often remains incomplete:

- **Ghost and instability issues:** Higher-derivative models ($f(R, G)$, non-local gravity) may introduce Ostrogradsky instabilities unless carefully constructed.
- **Frame ambiguities:** Scalar–tensor theories exhibit differences between the Jordan and Einstein frames, complicating the interpretation of observable quantities.
- **Ambiguity in geometric structure:** In teleparallel ($f(T)$) and symmetric teleparallel ($f(Q)$) gravity, the choice of tetrads or affine connection is not unique, potentially leading to spurious solutions (Table 31).

2. Incomplete understanding of reheating and preheating

One of the most pressing open problems is the absence of a fully self-consistent reheating scenario in many modified gravity models. Current issues include:

- Lack of a unique decay mechanism from geometric degrees of freedom to radiation/matter.

- Model-dependent preheating dynamics in non-minimally coupled or torsion-based theories.
- Sensitivity of CMB predictions to the assumed reheating equation of state (Table 32).

3. Interface with quantum gravity and UV completion

Many inflationary models lack a robust embedding in quantum gravity frameworks. Questions include:

- Which modified gravity models can be derived from string theory or loop quantum gravity?
- Do these models respect swampland constraints (e.g., field range limits, no stable dS vacua)?
- Can UV-complete theories like asymptotically safe gravity reproduce inflationary observables?

4. Perturbations and observables: what needs improvement

In modified gravity, scalar and tensor perturbation theory is often underdeveloped. Some open questions:

- What are the sound speed and stability conditions for scalar and tensor modes in $f(Q)$, $f(T)$, and EC gravity?
- How do entropy and isocurvature modes behave in multi-field or scalar–torsion models?

Table 32 Status of reheating predictions in modified gravity theories

Model	Reheating mechanism known?	Preheating possible?	Predictive T_{reh} ?
$f(R)$	Yes (via scalaron decay)	Yes	$\sim 10^9$ GeV
$f(T)$	Poorly understood	Uncertain	Model-dependent
Mimetic gravity	Partially	Yes (with potential)	$\lesssim 10^8$ GeV
Scalar–GB	Yes (for $\xi(\phi)G$)	Yes	Variable
$f(Q)$	Very limited	Under study	Unknown

- Can we compute B-mode polarization reliably in Gauss–Bonnet and Einstein–Cartan backgrounds?

5. Opportunities with future observations

Upcoming experiments will impose much tighter constraints on modified gravity inflation:

- **LiteBIRD and CMB-S4:** $r < 10^{-3}$ will rule out many high- r inflation models.
- **LISA, DECIGO, and PTA:** Gravitational wave background from inflation or preheating can distinguish torsion- or non-metricity-induced models.
- **21 cm cosmology:** Potential sensitivity to primordial non-Gaussianity and isocurvature (Table 33).

6. Toward a model selection framework

To move beyond individual case studies, the field needs more global model-selection frameworks. Promising directions include:

- Bayesian evidence-based surveys across $f(R)$, $f(G)$, $f(Q)$ and scalar-tensor spaces.
- Machine learning-assisted classification and emulation of Boltzmann codes.
- Unified datasets from CMB, large-scale structure, and gravitational waves.

In sum, modified gravity models of inflation offer a rich arena of ideas, but a full synthesis will require advances in mathematical consistency, quantum gravity embeddings, and high-precision cosmological tests. Addressing these challenges will clarify whether inflation is fundamentally geometric, field-driven, or a hybrid of both.

33 Open questions and future directions

Despite the substantial progress in developing and analyzing inflationary models within modified gravity frameworks, several foundational questions remain open. These questions

are not only critical for theoretical consistency but also offer promising directions for observational discrimination in the upcoming decade. We highlight several key open problems:

1. Baryogenesis and leptogenesis mechanisms:

Can modified gravity frameworks – particularly those involving torsion, non-metricity, or additional scalar/vector fields – naturally support baryogenesis or leptogenesis without the need for ad hoc extensions? For example, can Einstein–Cartan theories or scalar–Gauss–Bonnet couplings generate the CP-violating conditions necessary for matter–antimatter asymmetry?

2. Primordial black hole (PBH) discriminants:

Are the statistical and mass distribution properties of PBHs formed in different model classes (e.g., $f(R)$, $f(T)$, $f(Q)$, mimetic gravity) distinguishable through gravitational wave signatures or microlensing observations? Can scalar-induced GWs from PBH-generating inflation be used as unique fingerprints?

3. UV completion pathways:

What are the ultraviolet completions for torsion-based ($f(T)$, Einstein–Cartan) and non-metricity-based ($f(Q)$) inflationary theories? Can these theories be embedded consistently in a string-theoretic, supergravity, or asymptotically safe quantum gravity framework?

4. Reheating in quantum gravity:

Can the reheating phase – often treated phenomenologically – be embedded within a fully quantized theory of spacetime? For example, how does gravitational reheating operate in loop quantum cosmology, string field theory, or under asymptotic safety constraints?

5. Degeneracy resolution via Next-Gen observations:

How will future missions such as LiteBIRD, CMB-S4, and LISA break degeneracies between different modified gravity inflationary models? Can tensor tilt n_t , isocurvature modes, or precision measurements of r distinguish among models that are currently statistically equivalent?

These open problems highlight the intersection between cosmological model-building, high-energy theoretical physics, and precision astrophysical observations. Their

Table 33 Future observational prospects for constraining modified gravity inflation

Experiment	Observable	Target sensitivity	Relevant models
LiteBIRD	r (tensor modes)	$r < 10^{-3}$	$f(R)$, GB, $f(Q)$
LISA/PTA	Ω_{GW}	10^{-12}	Preheating, PBH GWs
CMB-S4	$n_s, \alpha_s, f_{\text{NL}}$	High-precision	All models
SKA/21cm	Non-Gaussianity, isocurvature	$f_{\text{NL}} < 1$	Multi-field, torsion

resolution may ultimately determine which inflationary paradigm best describes the early universe.

34 Discussion and outlook

The inflationary paradigm has undergone a profound transformation over the last two decades, evolving from a simple scalar-field-driven acceleration to a multi-faceted landscape of geometrical, string-inspired, and effective field theories. Modified gravity frameworks provide a vast extension to the standard lore of inflation, offering alternative geometric origins, enriched particle content, and novel phenomenological signatures. This work has systematically reviewed and extended a broad array of these models – including $f(R)$, $f(T)$, $f(G)$, $f(Q)$, scalar–curvature, scalar–torsion, mimetic, Einstein–Cartan, non-local, and extra-dimensional theories.

Our comparative analysis finds that certain models – such as Starobinsky’s $f(R)$ inflation, scalar–Gauss–Bonnet frameworks, and symmetric teleparallel $f(Q)$ theories – provide predictions in remarkable agreement with current CMB constraints. These models typically yield spectral indices in the range $n_s \sim 0.96$ and tensor-to-scalar ratios $r < 0.01$, compatible with both Planck and BICEP/Keck bounds. The geometric nature of these constructions often reduces the need for trans-Planckian field excursions and minimizes radiative instability problems common in canonical inflation.

Other classes, such as torsion-based theories ($f(T)$, scalar–torsion couplings), Ricci-tensor inflation, and Einstein–Cartan models, offer elegant extensions of spacetime structure. These approaches enable inflation to arise not from an ad hoc inflaton field but from intrinsic geometric properties of spacetime itself, such as torsion, non-metricity, or spin-coupled degrees of freedom. However, they also raise concerns regarding the stability of perturbations, the ambiguity of tetrad choices, or the consistency of their UV behavior.

String-inspired models – including brane–antibrane inflation, axion monodromy, and DBI inflation – leverage the geometry of extra dimensions and flux compactifications to generate inflaton potentials that are naturally flat and radiatively stable. These models, especially those arising from

Type IIB string theory, provide a UV-complete perspective but often require intricate moduli stabilization and control over higher-order corrections. The predictions for non-Gaussianities, reheating mechanisms, and cosmic string remnants make them rich targets for upcoming observations.

Further contributions from less conventional frameworks – such as Carmeli’s cosmological relativity, Moffat’s Scalar–Tensor–Vector Gravity (MOG), and Bekenstein-type varying constant models – suggest that inflationary dynamics can be embedded in broader theories of spacetime, where velocity, coupling constants, or gravitational strength evolve with cosmic time. These theories challenge traditional assumptions and demand new formalisms for perturbations, background solutions, and observational predictions.

From a methodological standpoint, this study has highlighted the importance of dynamical systems, phase-space portraits, and Bayesian model selection in classifying and constraining inflationary theories. Tools such as `CosmoMC`, `PolyChord`, and neural emulators of Boltzmann codes (e.g., for `CAMB` and `CLASS`) are now indispensable for high-dimensional inference. Machine learning algorithms, including decision trees, Gaussian processes, and variational autoencoders, are being increasingly employed to scan large classes of inflationary models, detect degeneracies, and reconstruct potential landscapes from data.

We also stress that while most inflationary models can reproduce the observed n_s and r , they often diverge significantly in other testable quantities such as:

- **Non-Gaussianities** (f_{NL}): Especially in multi-field, non-canonical, and DBI scenarios.
- **Tensor spectral index** (n_t): Modified dispersion relations in non-local and higher-order theories can affect this.
- **Reheating and preheating**: Diverse mechanisms exist across theories – some relying on scalaron decay, others on tachyonic instabilities or geometric transitions.
- **Stochastic gravitational waves**: Second-order sources tied to PBH formation or preheating can help distinguish between models in future experiments like LISA and SKA.

One notable development is the emergence of tachyon inflation as a bridge between effective field theory and string cosmology. Tachyonic DBI-type actions, motivated by D-brane decay, naturally lead to inflationary trajectories without requiring fine-tuned potentials and support a graceful exit to reheating. Their embedding in modified gravity (e.g., $f(T)$, Einstein–Cartan) deserves further exploration, especially in light of their distinctive kinetic structure.

An important epistemological reflection is also warranted: modern inflationary cosmology challenges long-held philosophical and even metaphysical notions of origin. Far from closing doors, it has stimulated renewed dialogues across fields – connecting cosmology with foundational questions about time, causality, and the structure of physical law. In conclusion, three parallel thrusts now define the future of inflation in modified gravity:

1. **Observational precision:** Upcoming missions (Lite-BIRD, CMB-S4, LISA, DECIGO, SKA) will place strong constraints on r , f_{NL} , and gravitational wave backgrounds, thereby excluding large swaths of parameter space.
2. **Computational acceleration:** Bayesian model comparison and machine learning will allow rapid navigation of high-dimensional theory space, enabling real-time filtering of viable inflationary scenarios.
3. **Theoretical innovation:** Continued development of UV-complete models – via string theory, non-local operators, geometric unification, and emergent gravity – will enrich the conceptual landscape.

Whether inflation is fundamentally a manifestation of curvature, torsion, non-metricity, scalar dynamics, or emergent geometry remains an open and compelling question. This work affirms that modified gravity is not merely a theoretical embellishment but a legitimate contender in explaining the early universe, with diverse signatures awaiting discovery. The pursuit of inflationary cosmology in such frameworks promises not only to elucidate the past but also to expand the future frontiers of gravitational physics.

Data Availability Statement This manuscript has no associated data. [Authors' comment: No datasets were generated or analyzed during this study.]

Code Availability Statement This manuscript has no associated code/software. [Authors' comment: No code/software was generated or analyzed during this study.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indi-

cated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.
Funded by SCOAP³.

Appendix

Appendix A: Reheating and preheating in modified gravity

Reheating and preheating constitute a critical phase bridging the end of inflation and the onset of the standard hot Big Bang evolution. In the context of modified gravity theories, this transition can deviate significantly from the canonical scalar field scenario due to modifications in the gravitational action, the presence of extra degrees of freedom, or geometric couplings that affect the inflaton's decay channels and the universe's thermal history.

This appendix expands upon Sect. 18 by providing quantitative estimates, equations-of-state evolution, particle production mechanisms, and comparative analyses of reheating scenarios across a range of modified gravity models.

A.1 Reheating temperature estimates

Table 34 presents typical reheating temperature ranges T_{reh} derived from representative models. These are obtained through semi-analytic or numerical evaluations of inflaton decay widths, oscillation dynamics, and energy transfer efficiency.

A.2 Evolution of the equation of state

The post-inflationary equation of state parameter $w(t) = p/\rho$ governs the expansion history and thermalization efficiency. Modified gravity scenarios often produce non-trivial $w(t)$ profiles:

- **$f(R)$ gravity:** Scalaron oscillations lead to an effective matter-like phase ($w \approx 0$), though deviations may arise from higher-order terms.
- **Scalar–Gauss–Bonnet:** Due to non-minimal couplings, w may transiently exceed 1 before settling.
- **$f(Q)$ gravity:** Non-metricity contributions can induce $w < 0$ transitions or stiff behavior depending on the coupling scheme.

Table 34 Estimated reheating temperatures in selected modified gravity frameworks

Model	Reheating mechanism	T_{reh} [GeV]	References
$f(R)$ Starobinsky	Scalaron decay (gravitational)	$10^9 - 10^{10}$	[76]
Scalar–Gauss–Bonnet	Coupled scalar–curvature terms	$10^7 - 10^9$	[54]
$f(Q)$ gravity	Non-metric-induced reheating	$10^6 - 10^8$	[80]
Mimetic inflation	Lagrange multiplier-mediated decay	$\lesssim 10^8$	[63]
$f(T)$ teleparallel	Direct inflaton–matter couplings	Model-dependent	[129]

A.3 Particle production channels

Inflaton decay governs the particle content and thermalization dynamics. Channels include:

- **Fermionic decay:** $\phi \rightarrow \bar{\psi}\psi$ via Yukawa interactions, $\mathcal{L} \supset y\phi\bar{\psi}\psi$.
- **Bosonic decay:** $\phi \rightarrow \chi\chi$ through $\mathcal{L} \supset g\phi^2\chi^2$ or gravitational particle production.
- **Gravitational reheating:** Particle production from oscillating curvature backgrounds, especially in $f(R)$ or $f(Q)$.
- **Non-minimal couplings:** Scalar–curvature couplings in $f(R, \phi)$, ϕR , or $\xi(\phi)G$ models open new decay modes.

The decay rate and subsequent thermalization depend on coupling constants g, y , the inflaton mass m_ϕ , and the background geometry.

A.4 Gravitational vs direct reheating mechanisms

We now compare the general properties of gravitational reheating (e.g., scalaron decay in $f(R)$) and direct coupling-induced reheating in models like $f(T)$ or Gauss–Bonnet (Table 35):

A.5 Remarks and future directions

The reheating and preheating processes in modified gravity models represent a unique probe into the non-standard inflationary exit and thermalization dynamics. They encode crucial signatures in the CMB and gravitational wave background, which may serve to distinguish models observationally. Future studies should explore:

- Non-perturbative lattice simulations in $f(Q)$ and mimetic scenarios.
- Gravitational wave signatures from preheating in torsion-based models.
- Reheating consistency relations linking T_{reh} to (n_s, r) in non-canonical models.

Appendix B: Conformal transformations in modified gravity

In the study of modified gravity theories – especially $f(R)$, scalar–tensor, and dilaton-based frameworks – conformal transformations provide a powerful mathematical tool to reframe the theory into a more tractable form. These transformations relate the so-called Jordan frame, in which matter is minimally coupled and gravity is modified, to the Einstein frame, in which gravity takes the standard Einstein–Hilbert form but the matter sector acquires additional couplings.

The conformal transformation is defined as:

$$\tilde{g}_{\mu\nu} = \Omega^2(x)g_{\mu\nu}, \tag{83}$$

where $\Omega(x)$ is a smooth, strictly positive function of space-time coordinates – often taken to be a function of a scalar field ϕ . The choice of $\Omega(x)$ is model-dependent and crucial in determining the structure of the resulting Einstein-frame action.

For instance, in $f(R)$ gravity, one typically defines the conformal factor via:

$$\Omega^2(x) = f'(R), \tag{84}$$

where $f'(R) = \frac{df}{dR}$, assuming $f''(R) > 0$ to ensure a ghost-free transformation. By introducing an auxiliary scalar field ϕ , the action can be recast as:

$$S = \int d^4x \sqrt{-\tilde{g}} \left[\frac{1}{2\kappa^2} \tilde{R} - \frac{1}{2} \tilde{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - U(\phi) \right] + S_m[\Omega^{-2} \tilde{g}_{\mu\nu}, \psi], \tag{85}$$

where $U(\phi)$ is the potential derived from the functional form of $f(R)$, and S_m is the matter action, which is now non-minimally coupled to the metric in the Einstein frame. The scalar field ϕ is canonically normalized via:

$$\frac{d\phi}{dR} = \sqrt{\frac{3}{2\kappa^2}} \frac{f''(R)}{f'(R)}. \tag{86}$$

This procedure allows one to interpret $f(R)$ gravity as a scalar–tensor theory with a specific potential and coupling structure. The Einstein frame is particularly useful for analyzing inflationary dynamics, deriving slow-roll parameters, and computing cosmological observables such as n_s and r .

Table 35 Comparison of reheating mechanisms across different gravity theories

Feature	Gravitational reheating	Direct coupling reheating
Universality	Generic (minimal assumptions)	Model-dependent
Efficiency	Generally suppressed	High with strong couplings
Thermalization scale	Delayed	Prompt
Dependence on inflaton potential	Moderate	High
Robustness to UV completion	Strong	May require tuning

Table 36 Comparison between Jordan and Einstein frame in conformally transformed gravity

Feature	Jordan frame	Einstein frame
Gravitational sector	Modified ($f(R)$, R^2 , etc.)	Einstein–Hilbert
Scalar field	Auxiliary or non-canonical	Canonically normalized
Matter coupling	Minimal	Non-minimal (via Ω)
Observables (n_s , r)	Model-dependent	Simplified via potential
Interpretation	Modified gravity	Scalar–tensor theory

Transformation of geometric quantities. Under the conformal transformation:

$$\tilde{g}_{\mu\nu} = \Omega^2 g_{\mu\nu}, \quad \Omega = e^{\sigma(x)}, \tag{87}$$

the Ricci scalar transforms as:

$$R = \Omega^2 \left(\tilde{R} + 6\tilde{\square} \ln \Omega - 6\tilde{g}^{\mu\nu} \partial_\mu \ln \Omega \partial_\nu \ln \Omega \right). \tag{88}$$

This identity ensures that the higher-order terms in $f(R)$ or scalar-curvature actions are absorbed into kinetic and potential terms of a scalar field, thereby simplifying the dynamics.

Physical interpretation. While mathematically equivalent, the Jordan and Einstein frames offer different physical interpretations. In the Jordan frame, gravitational effects are modified directly, and matter follows standard geodesics. In contrast, in the Einstein frame, gravity is described by the Einstein tensor, but matter feels an additional coupling via the scalar field.

Applications in inflation. Conformal transformations are widely used to analyze inflation in Starobinsky models, non-minimally coupled Higgs inflation, scalar–Gauss–Bonnet inflation, and dilaton gravity. In many cases, potentials that appear too steep for inflation in the Jordan frame are flattened in the Einstein frame, enabling slow-roll evolution. For example, in Higgs inflation, the non-minimal coupling $\xi\phi^2 R$ in the Jordan frame becomes a plateau-like potential in the Einstein frame:

$$V_E(\chi) \propto \left(1 - e^{-\sqrt{2/3}\chi/M_P} \right)^2, \tag{89}$$

mimicking the form of Starobinsky inflation.

Limitations and open questions. Although conformal transformations are mathematically robust, debates continue regarding the physical equivalence of the two frames, especially in quantum contexts. Additionally, care must be taken

when transforming perturbations and observables, as these may not be invariant under frame transformations.

Summary table: Einstein vs Jordan frame In conclusion, conformal transformations serve as a unifying bridge between different formulations of modified gravity, allowing for analytic tractability and clearer physical interpretation – particularly in inflationary model-building and cosmological perturbation theory (Table 36).

Appendix C: Perturbation equations in modified gravity

Cosmological perturbation theory plays a central role in connecting inflationary models to observable quantities such as the cosmic microwave background (CMB) anisotropies and the large-scale structure (LSS) of the universe. In modified gravity theories, scalar perturbations are especially important because they dominate the formation of structure and encode information about the early universe dynamics.

Metric perturbations in Newtonian gauge. In the conformal Newtonian gauge, the scalar perturbations to the flat Friedmann–Lemaître–Robertson–Walker (FLRW) metric are given by:

$$ds^2 = -(1 + 2\Phi)dt^2 + a(t)^2(1 - 2\Psi)\delta_{ij}dx^i dx^j, \tag{90}$$

where Φ and Ψ are the Bardeen potentials, which represent the scalar degrees of freedom in the metric. These quantities are gauge-invariant under linear coordinate transformations.

In General Relativity (GR), for minimally coupled matter and no anisotropic stress, one finds $\Phi = \Psi$. However, in modified gravity theories, this equality is often broken, and their evolution must be analyzed separately.

Modified field equations. In a general class of modified gravity models, including scalar–tensor, $f(R)$, $f(T)$, Gauss–Bonnet, and $f(Q)$ theories, the linearized field equa-

tions for scalar perturbations are modified due to additional degrees of freedom or non-standard couplings. The perturbed Einstein equations take the schematic form:

$$\nabla^2\Psi = 4\pi G_{\text{eff}}(k, t) a^2\delta\rho, \tag{91}$$

$$\Phi - \Psi = \Pi_{\text{aniso}}(k, t), \tag{92}$$

where:

- $G_{\text{eff}}(k, t)$ is the effective gravitational coupling, which may depend on scale k and time t ,
- Π_{aniso} is the anisotropic stress potential, often sourced by extra scalar, vector, or tensor fields,
- $\delta\rho$ is the comoving matter density perturbation.

These equations generalize the Poisson equation and gravitational slip in GR, allowing for a scale- and time-dependent description of structure growth.

Model-specific effects.

- In $f(R)$ gravity, the extra scalar degree of freedom (scalaron) mediates an additional force, modifying G_{eff} as:

$$G_{\text{eff}} = \frac{G}{f'(R)} \left(1 + \frac{4k^2 f''(R)/a^2}{1 + 3k^2 f''(R)/a^2} \right), \tag{93}$$

which leads to enhanced growth of structures on sub-horizon scales if $f''(R) > 0$.

- In scalar–tensor theories with Brans–Dicke-type couplings, one finds:

$$G_{\text{eff}} = \frac{1}{8\pi\phi} \left(\frac{2\omega(\phi) + 4}{2\omega(\phi) + 3} \right), \tag{94}$$

where ϕ is the scalar field and $\omega(\phi)$ its kinetic function.

- In $f(T)$ gravity, torsional degrees of freedom contribute to the anisotropic stress and modify the evolution of scalar modes:

$$\Pi_{\text{aniso}} \propto f_{TT} \delta T, \tag{95}$$

with $f_{TT} \equiv d^2 f/dT^2$ and δT the torsion perturbation.

- In mimetic gravity or models with disformal couplings, the relation between Φ and Ψ is governed by the dynamics of a mimetic or hidden scalar mode that can source anisotropic stress even in the absence of anisotropic matter.

Gauge-invariant variables and evolution equations. One commonly introduces the comoving curvature perturbation

\mathcal{R} and the Mukhanov–Sasaki variable v , related via:

$$v = z\mathcal{R}, \quad z = a \frac{\dot{\phi}}{H}, \tag{96}$$

for canonical scalar fields, or appropriately generalized in non-minimally coupled models. The mode equation in Fourier space becomes:

$$v_k'' + \left(k^2 - \frac{z''}{z} \right) v_k = 0, \tag{97}$$

where primes denote derivatives with respect to conformal time η . The power spectrum of scalar perturbations is then:

$$\mathcal{P}_{\mathcal{R}}(k) = \frac{k^3}{2\pi^2} |\mathcal{R}_k|^2. \tag{98}$$

Effective gravitational coupling table

See Table 37.

Observational implications. Modifications in G_{eff} and $\Phi - \Psi$ lead to observable signatures:

- Growth rate of structure: Modified gravity affects the evolution of the density contrast δ and redshift-space distortions.
- Lensing potential: The lensing convergence depends on $\Phi + \Psi$, thus any difference introduces measurable deviations.
- ISW effect: The late-time Integrated Sachs–Wolfe signal is sensitive to time variation in $\Phi + \Psi$.

In summary, scalar perturbation theory in modified gravity offers a versatile diagnostic for probing the inflationary and post-inflationary universe. Models can be distinguished not only by their background evolution but by the scale-dependence of G_{eff} , the presence of anisotropic stress, and the evolution of metric potentials – all of which are observable with current and upcoming cosmological surveys.

Appendix D: Inflationary observables

Inflationary cosmology makes concrete predictions that can be tested via cosmic microwave background (CMB) anisotropies, large-scale structure, and primordial gravitational waves. The key observables include:

- Scalar spectral index n_s
- Tensor-to-scalar ratio r
- Running of the spectral index α_s
- Non-Gaussianity parameters f_{NL}

These quantities are computed either from the slow-roll approximation or directly from the correlation functions of curvature perturbations in a specific inflationary model.

Table 37 Effective gravitational couplings and anisotropic stress in selected models

Model	$G_{\text{eff}}(k, t)$	$\Phi \neq \Psi?$
General Relativity	G	No
$f(R)$ gravity	scale-dependent	Yes
Scalar–Tensor	$\propto 1/\phi$	Yes
$f(T)$ torsion gravity	constant or mild scale-dependence	Yes
Mimetic gravity	variable, via Lagrange multipliers	Yes
Einstein–Cartan theory	G , but extra spin-torsion contributions	Possibly

Slow-roll formalism. In canonical single-field slow-roll inflation, the key slow-roll parameters are defined as:

$$\begin{aligned} \epsilon &\equiv -\frac{\dot{H}}{H^2} = \frac{1}{2}M_P^2 \left(\frac{V'}{V}\right)^2, \\ \eta &\equiv M_P^2 \left(\frac{V''}{V}\right), \\ \xi^2 &\equiv M_P^4 \left(\frac{V'V'''}{V^2}\right), \end{aligned}$$

where $V(\phi)$ is the inflaton potential and primes denote derivatives with respect to the scalar field ϕ .

Scalar spectral index n_s . The spectral index n_s describes the deviation of the scalar power spectrum from exact scale invariance:

$$\mathcal{P}_{\mathcal{R}}(k) = A_s \left(\frac{k}{k_*}\right)^{n_s-1}. \tag{99}$$

In slow-roll approximation:

$$n_s = 1 - 6\epsilon + 2\eta. \tag{100}$$

Tensor-to-scalar ratio r . The ratio r quantifies the amplitude of primordial tensor (gravitational wave) perturbations relative to scalars:

$$r = \frac{\mathcal{P}_T}{\mathcal{P}_{\mathcal{R}}} = 16\epsilon. \tag{101}$$

This directly determines the energy scale of inflation:

$$V^{1/4} \simeq \left(\frac{3\pi^2 A_s r}{2}\right)^{1/4} M_P \sim 10^{16} \text{ GeV} \left(\frac{r}{0.01}\right)^{1/4}. \tag{102}$$

Running of the spectral index α_s . The running is the logarithmic scale dependence of n_s :

$$\alpha_s \equiv \frac{dn_s}{d \ln k} = -24\epsilon^2 + 16\epsilon\eta - 2\xi^2. \tag{103}$$

It can become significant in models with multiple stages of inflation or strong deviations from slow-roll.

Non-Gaussianity f_{NL} . Non-Gaussianities are encoded in the bispectrum of curvature perturbations:

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle = (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B(k_1, k_2, k_3). \tag{104}$$

The amplitude and shape are characterized by f_{NL} :

- $f_{\text{NL}}^{\text{local}}$: peak in squeezed configurations, typical of multi-field models,
- $f_{\text{NL}}^{\text{equil}}$: peak in equilateral triangles, common in DBI/k-inflation,
- $f_{\text{NL}}^{\text{orth}}$: orthogonal shape with mixed-sign contributions.

In standard slow-roll inflation:

$$f_{\text{NL}}^{\text{local}} \sim \mathcal{O}(\epsilon, \eta) \ll 1. \tag{105}$$

Observational constraints (Planck 2018):

See Table 38.

Table 38 Representative inflationary observables from Planck 2018 data

Observable	Planck 2018 value
Amplitude A_s (at $k = 0.05 \text{ Mpc}^{-1}$)	2.1×10^{-9}
Spectral index n_s	0.9649 ± 0.0042
Tensor-to-scalar ratio r (95% CL)	< 0.056
Running α_s	-0.0045 ± 0.0067
$f_{\text{NL}}^{\text{local}}$	-0.9 ± 5.1

Extensions in modified gravity. In theories beyond General Relativity – such as $f(R)$, $f(T)$, $f(Q)$, and scalar–tensor theories – these expressions are modified:

- The effective field may live in the Einstein frame with a different potential and kinetic term,
- Sound speed $c_s < 1$ may suppress tensor modes and enhance $f_{\text{NL}}^{\text{equil}}$,
- Non-minimal couplings alter ϵ , η and their mapping to observables,
- Modified dispersion relations can affect power spectrum tilt.

For example, in $f(R)$ gravity, upon conformal transformation:

$$f(R) = R + \alpha R^2 \rightarrow V(\phi) = \frac{1}{8\alpha} \left(1 - e^{-\sqrt{2/3}\phi/M_P}\right)^2, \tag{106}$$

where ϕ is the scalaron field in the Einstein frame. Inflationary observables are then computed using standard slow-roll but with this potential.

Inflationary observables offer a window into the physics of the early universe. Their derivation – from first principles or from effective field theory – links theory with experiment. Future missions like LiteBIRD, CMB-S4, and LISA will refine constraints on r , α_s , and f_{NL} , helping discriminate between competing modified gravity models.

Appendix E: Dynamical system formulation

In cosmology, the dynamical systems approach provides a powerful method for analyzing the qualitative behavior of cosmological models without solving the full system explicitly. This is especially useful in modified gravity theories where the equations of motion become highly nonlinear.

Autonomous system formulation. We begin by defining suitable dimensionless variables, such as:

$$\begin{aligned} x &= \frac{\dot{\phi}}{\sqrt{6}HM_P}, & y &= \frac{\sqrt{V(\phi)}}{\sqrt{3}HM_P}, & z &= \frac{f'(R)}{f(R)}, \\ \lambda &= -\frac{V'}{V}, & \Gamma &= \frac{VV''}{(V')^2}, \end{aligned} \tag{107}$$

depending on the model under study.

The evolution equations for these variables can then be written as an autonomous system of first-order differential equations:

$$\frac{dx_i}{dN} = f_i(x_j), \tag{108}$$

where $N = \ln a$ is the number of e-folds and x_i are the dynamical variables.

Fixed points and stability. Critical (or fixed) points $x_i^{(c)}$ satisfy:

$$\left. \frac{dx_i}{dN} \right|_{x_j=x_j^{(c)}} = 0 \quad \forall i. \tag{109}$$

To assess stability, we linearize around the critical point:

$$\frac{d\delta x_i}{dN} = \sum_j \left. \frac{\partial f_i}{\partial x_j} \right|_{x^{(c)}} \delta x_j = J_{ij} \delta x_j, \tag{110}$$

where J_{ij} is the Jacobian matrix. The eigenvalues of J determine stability:

- All eigenvalues with $\text{Re}(\lambda_i) < 0$: stable (attractor),

- At least one $\text{Re}(\lambda_i) > 0$: unstable (repeller),
- Mixed signs: saddle point.

Applications to modified gravity. Example 1: $f(R)$ gravity. In $f(R)$ gravity, the scalar degree of freedom introduced by the function $f(R)$ can be cast as a scalar field ϕ , with effective dynamics given by:

$$x = -\frac{f'(R)R - f(R)}{6H^2 f'(R)}, \quad y = \frac{R}{6H^2}. \tag{111}$$

The dynamical system then reads:

$$\begin{aligned} \frac{dx}{dN} &= -4x + y - 3x^2 + (\text{matter terms}), \\ \frac{dy}{dN} &= -2y(1 + q), \end{aligned}$$

where q is the deceleration parameter.

Example 2: teleparallel $f(T)$ gravity. Here, the field equations are second-order, and the autonomous system typically includes:

$$x = \frac{\rho_m}{3H^2}, \quad y = \frac{f(T)}{6H^2}, \quad z = \frac{f_T T}{6H^2}, \tag{112}$$

with modified Friedmann equations rewritten accordingly.

Example 3: symmetric teleparallel $f(Q)$ gravity. For $f(Q)$ models, dimensionless variables might include:

$$x = \frac{f(Q)}{6H^2}, \quad y = \frac{Qf_Q}{6H^2}, \tag{113}$$

yielding a closed dynamical system that respects second-order equations of motion.

Phase space behavior. Each fixed point corresponds to a specific cosmological era:

- de Sitter (inflation): $x \approx 0, y \approx 1$,
- Matter-dominated: $x \approx 1, y \approx 0$,
- Scaling solutions: $x + y = 1$.

Dynamical system techniques allow classification of asymptotic regimes in modified gravity inflation models. They are particularly effective for identifying inflationary attractors, transient reheating dynamics, and transitions to late-time acceleration. These methods complement numerical integration and provide robust tools for phase space exploration.

Appendix F: Auxiliary function reconstructions

Auxiliary function reconstruction techniques offer a versatile method for designing inflationary models in modified gravity. These methods aim to reverse-engineer a gravitational Lagrangian or scalar potential that gives rise to a desired

cosmic evolution, particularly the Hubble parameter $H(t)$ or the scale factor $a(t)$.

Reconstructing scalar potentials. In canonical single-field inflation, if one assumes a desired Hubble parameter as a function of the scalar field $H(\phi)$, one can reconstruct the corresponding scalar potential via the Hamilton–Jacobi formalism. The inflaton’s velocity satisfies:

$$\dot{\phi} = -2M_P^2 H'(\phi), \quad (114)$$

and the potential becomes:

$$V(\phi) = 3M_P^2 H^2(\phi) - 2M_P^4 [H'(\phi)]^2. \quad (115)$$

Alternatively, if $H(t)$ is specified, one first computes $\phi(t)$ from:

$$\dot{\phi} = -2M_P^2 \frac{dH}{dt}, \quad (116)$$

then inverts to find $t(\phi)$ and substitutes back to obtain:

$$V(\phi) = 3M_P^2 H^2(t(\phi)) - \frac{1}{2} \dot{\phi}^2(t(\phi)). \quad (117)$$

Reconstructing $f(R)$ gravity. Given a target scale factor $a(t)$, the Ricci scalar in a flat FLRW universe is:

$$R = 6 \left(2H^2 + \dot{H} \right). \quad (118)$$

Inverting $t(R)$ allows expressing $H(t)$ and hence $a(t)$ as functions of R , yielding a differential equation for $f(R)$:

$$\frac{d^2 f(R)}{dR^2} \ddot{R} + \frac{df(R)}{dR} \dot{R} + \text{terms involving } H, \dot{H}, R = \rho_{\text{eff}}. \quad (119)$$

This reconstruction approach is known as the inverse method and has been used to match late-time acceleration or emulate Starobinsky-like inflation.

Reconstructing $f(T)$ gravity. In teleparallel gravity, the torsion scalar is given by $T = -6H^2$. A similar method applies: choose a scale factor $a(t)$, compute $T(t)$, and solve an algebraic differential equation to obtain $f(T)$ such that the Friedmann equations are satisfied:

$$12H^2 f_T + f = 2\rho. \quad (120)$$

Reconstructing $f(Q)$ gravity. In symmetric teleparallel gravity, where $Q = 6H^2$, one can reconstruct $f(Q)$ functions by assuming a cosmological history $H(t)$ or $a(t)$ and solving the generalized Friedmann equation:

$$f(Q) - 6H^2 f_Q = 2\rho. \quad (121)$$

Reconstruction from observational $H(z)$ data. In observational cosmology, $H(z)$ data from supernovae, BAO, and CMB allow one to numerically reconstruct $V(\phi)$ or $f(R)$:

- Convert $z \rightarrow t(z)$ via $dt = -dz/[(1+z)H(z)]$,
- Use numerical integrals to estimate $\phi(z)$ or $R(z)$,

- Fit a parametric form to obtain $V(\phi)$ or $f(R)$.

Auxiliary function reconstructions provide model-independent methods to study inflationary dynamics consistent with data. They offer flexibility in recovering viable potentials or gravity functions that match CMB and $H(z)$ observations, and are particularly valuable in modified gravity where no fundamental scalar Lagrangian may exist initially.

References

1. A.H. Guth, Inflationary universe: a possible solution to the horizon and flatness problems. *Phys. Rev. D* **23**, 347 (1981)
2. A.D. Linde, A new inflationary universe scenario: a possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems. *Phys. Lett. B* **108**, 389–393 (1982)
3. A. Albrecht, P.J. Steinhardt, Cosmology for grand unified theories with radiatively induced symmetry breaking. *Phys. Rev. Lett.* **48**, 1220 (1982)
4. G.F. Smoot et al., Structure in the COBE differential microwave radiometer first-year maps. *Astrophys. J. Lett.* **396**, L1–L5 (1992)
5. E. Komatsu et al. [WMAP Collaboration], Seven-year Wilkinson microwave anisotropy probe (WMAP) observations: cosmological interpretation. *Astrophys. J. Suppl.* **192**, 18 (2011)
6. N. Aghanim et al. [Planck Collaboration], Planck 2018 results. I. Overview and the cosmological legacy of Planck. *Astron. Astrophys.* **641**, A1 (2020). [arXiv:1807.06205](https://arxiv.org/abs/1807.06205) [astro-ph.CO]
7. Y. Akrami et al. [Planck Collaboration], Planck 2018 results. X. Constraints on inflation. *Astron. Astrophys.* **641**, A10 (2020). [arXiv:1807.06211](https://arxiv.org/abs/1807.06211) [astro-ph.CO]
8. P.A.R. Ade et al. [BICEP/Keck Collaboration], Improved constraints on primordial gravitational waves using Planck, WMAP, and BICEP/Keck observations through the 2018 observing season. *Phys. Rev. Lett.* **127**, 151301 (2021). [arXiv:2110.00483](https://arxiv.org/abs/2110.00483)
9. K. Abazajian et al. [CMB-S4 Collaboration], CMB-S4 Science Book, First Edition. [arXiv:1610.02743](https://arxiv.org/abs/1610.02743) [astro-ph.CO]
10. A.A. Starobinsky, A new type of isotropic cosmological models without singularity. *Phys. Lett. B* **91**, 99–102 (1980)
11. S. Nojiri, S.D. Odintsov, M. Sasaki, Gauss–bonnet dark energy. *Phys. Rev. D* **71**, 123509 (2005). [arXiv:hep-th/0504052](https://arxiv.org/abs/hep-th/0504052)
12. S. Nojiri, S.D. Odintsov, Unified cosmic history in modified gravity: from f(r) theory to lorentz non-invariant models. *Phys. Rep.* **505**, 59–144 (2011). [arXiv:1011.0544](https://arxiv.org/abs/1011.0544)
13. Y.F. Cai, S. Capozziello, M. Laurentis, E.N. Saridakis, F(t) teleparallel gravity and cosmology. *Rep. Prog. Phys.* **79**, 106901 (2016). [arXiv:1511.07586](https://arxiv.org/abs/1511.07586)
14. S.D. Odintsov, V.K. Oikonomou, Bouncing cosmology with future singularity from modified gravity. *Phys. Rev. D* **96**(10), 104049 (2017). [arXiv:1711.02230](https://arxiv.org/abs/1711.02230) [gr-qc]
15. V.K. Oikonomou, A smooth bounce scenario in the presence of dark energy: the case of $f(r)$ gravity. *Mod. Phys. Lett. A* **32**, 1750172 (2017). [arXiv:1712.03107](https://arxiv.org/abs/1712.03107) [gr-qc]
16. L. Sebastiani, G. Cognola, R. Myrzakulov, S.D. Odintsov, S. Zerbini, Nearly starobinsky inflation from modified gravity. *Phys. Rev. D* **89**, 023518 (2014). [arXiv:1311.0744](https://arxiv.org/abs/1311.0744)
17. M. Sami, A. Toporensky, P.V. Tretyakov, S. Tsujikawa, The fate of (phantom) dark energy universe with string curvature corrections. *Phys. Lett. B* **619**, 193–200 (2005). [arXiv:hep-th/0504154](https://arxiv.org/abs/hep-th/0504154)
18. S. Nojiri, S.D. Odintsov, Modified gravity with negative and positive powers of curvature: unification of inflation and cosmic acceleration. *Phys. Rev. D* **68**, 123512 (2003). [arXiv:hep-th/0307288](https://arxiv.org/abs/hep-th/0307288)

19. K. Bamba, C.Q. Geng, S. Nojiri, S.D. Odintsov, Equivalence of modified gravity equation to einstein equation with quantum corrections and its implication for inflation. *Mod. Phys. Lett. A* **25**, 900–910 (2010). [arXiv:0909.4397](#) [hep-th]
20. S. Nojiri, S.D. Odintsov, V.K. Oikonomou, Modified gravity theories on a nutshell: inflation, bounce and late-time evolution. *Phys. Rep.* **692**, 1–104 (2017). [arXiv:1705.11098](#)
21. S. Capozziello, V.F. Cardone, A. Troisi, Reconciling dark energy models with $f(r)$ theories. *Phys. Rev. D* **71**, 043503 (2005). [arXiv:astro-ph/0501426](#)
22. L. Sebastiani, S. Zerbini, Static spherically symmetric solutions in $f(r)$ gravity. *Eur. Phys. J. C* **71**, 1591 (2011). [arXiv:1012.5230](#) [gr-qc]
23. K. Bamba, S. Nojiri, S.D. Odintsov, Inflationary cosmology and the late-time accelerated expansion of the universe in non-minimal yang–mills– $f(r)$ gravity and non-minimal vector–gauss–bonnet gravity. *Phys. Rev. D* **77**, 123532 (2008). [arXiv:0803.3384](#)
24. S. Capozziello, M. Laurentis, Extended theories of gravity. *Phys. Rep.* **509**, 167–321 (2011). [arXiv:1108.6266](#)
25. K. Bamba, C.Q. Geng, S. Nojiri, Inflationary cosmology and the late-time accelerated expansion of the universe in non-minimal yang–mills– $f(r)$ gravity and non-minimal vector–gauss–bonnet gravity. *Phys. Rev. D* **77**, 123532 (2008). [arXiv:0803.3384](#)
26. J.M. Ezquiaga, M. Zumalacárregui, Dark energy after gw170817: dead ends and the road ahead. *Phys. Rev. Lett.* **119**, 251304 (2017). [arXiv:1710.05901](#) [astro-ph.CO]
27. T. Kobayashi, M. Yamaguchi, J. Yokoyama, Generalized g -inflation: inflation with the most general second-order field equations. *Prog. Theor. Phys.* **126**, 511–529 (2011). [arXiv:1105.5723](#) [hep-th]
28. S.D. Odintsov, T. Paul, V.K. Oikonomou, Scalar–Gauss–Bonnet inflation consistent with GW constraints and Planck. *Phys. Rev. D* (2024). [arXiv:2407.12155](#) [gr-qc]
29. K. Bamba, S. Nojiri, S.D. Odintsov, The universe future in modified gravity theories: approaching the finite-time future singularity. *JCAP* **0810**, 045 (2008). [arXiv:0807.2575](#)
30. G. Calcagni, S. Tsujikawa, M. Sami, Dark energy and cosmological solutions in second-order string gravity. *Class. Quantum Gravity* **22**, 3977–4006 (2005). [arXiv:hep-th/0604201](#)
31. S.D. Odintsov, V.K. Oikonomou, L. Sebastiani, Unification of constant-roll inflation and dark energy with logarithmic r^2 correction in the jordan frame. *Nucl. Phys. B* **923**, 608–632 (2017). [arXiv:1807.00335](#) [gr-qc]
32. L. Sebastiani, G. Cognola, R. Myrzakulov, S.D. Odintsov, S. Zerbini, Nearly starobinsky inflation from modified gravity. *Phys. Rev. D* **89**, 023518 (2014). [arXiv:1311.0744](#)
33. S. Capozziello, M. Laurentis, O. Luongo, Connecting early and late universe by $f(r)$ gravity. *Int. J. Mod. Phys. D* **22**, 1350084 (2013). [arXiv:1310.8211](#)
34. S.D. Odintsov, V.K. Oikonomou, Singular inflationary universe from $f(r, g)$ gravity. *Phys. Rev. D* **92**, 124024 (2015). [arXiv:1510.04333](#)
35. S.D. Odintsov, D. Sáez-Gómez, $f(r, g)$ gravity inflation. *Phys. Lett. B* **697**, 255–259 (2011). [arXiv:1010.5375](#)
36. S.D. Odintsov, T. Paul, G.G.L. Nashed, Inflation and dark energy in scalar–Gauss–Bonnet gravity consistent with gravitational wave constraints. [arXiv:2506.08193](#) [gr-qc]
37. S.D. Odintsov, V.K. Oikonomou, T. Paul, Bottom-up reconstruction of viable inflationary potentials in scalar–gauss–bonnet gravity. *Phys. Rev. D* **104**, 083506 (2021). [arXiv:2108.10460](#) [gr-qc]
38. K. Bamba, C.Q. Geng, C.C. Lee, L.W. Luo, Equation of state for dark energy in $f(t)$ gravity. *JCAP* **1101**, 021 (2011). [arXiv:1011.0508](#)
39. M. Krššák, E.N. Saridakis, The covariant formulation of $f(t)$ gravity. *Class. Quantum Gravity* **33**, 115009 (2016). [arXiv:1510.08432](#)
40. M. Krššák, Covariant formulation of teleparallel gravity and its applications. *Universe* **5**, 125 (2019). [arXiv:1811.07586](#) [gr-qc]
41. N.D. Birrell, P.C.W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982)
42. F.L. Bezrukov, M. Shaposhnikov, The standard model higgs boson as the inflaton. *Phys. Lett. B* **659**, 703–706 (2008). [arXiv:0710.3755](#)
43. K. Bamba, S.D. Odintsov, D. Sáez-Gómez, Conformal symmetry and accelerating cosmology in teleparallel gravity. *Phys. Rev. D* **88**, 084042 (2013). [arXiv:1308.5789](#)
44. S.D. Odintsov, V.K. Oikonomou, Inflationary dynamics with a smooth slow-roll to constant-roll era transition. *Phys. Rev. D* **94**, 124026 (2016). [arXiv:1610.02533](#) [gr-qc]
45. K. Bamba, S. Nojiri, S.D. Odintsov, Inflationary cosmology and the late-time accelerated expansion of the universe in non-minimal yang–mills– $f(r)$ gravity and non-minimal vector–gauss–bonnet gravity. *Phys. Rev. D* **94**, 084042 (2016). [arXiv:1608.01730](#) [hep-th]
46. C. Germani, A. Kehagias, New model of inflation with nonminimal derivative coupling of standard model higgs boson to gravity. *Phys. Rev. Lett.* **105**, 011302 (2010). [arXiv:1003.2635](#)
47. K. Bamba, S.D. Odintsov, P.V. Tretyakov, Inflation in a scalar–tensor model with kinetic and gauss–bonnet couplings. *Eur. Phys. J. C* **75**, 344 (2015). [arXiv:1411.3852](#)
48. S. Capozziello, G. Lambiase, A. Stabile, Non-minimal derivative couplings and inflation in generalized theories of gravity. *Class. Quantum Gravity* **31**, 115004 (2014). [arXiv:1304.5440](#)
49. T. Koivisto, Covariant conservation of energy momentum in modified gravities. *Class. Quantum Gravity* **23**, 4289–4296 (2006). [arXiv:gr-qc/0505128](#)
50. K. Bamba, S.D. Odintsov, D. Sáez-Gómez, Conformal symmetry and accelerating cosmology in teleparallel gravity. *Phys. Rev. D* **88**, 084042 (2013). [arXiv:1308.5789](#)
51. K. Bamba, S. Nojiri, S.D. Odintsov, D. Sáez-Gómez, Inflationary cosmology and the late-time accelerated expansion of the universe in teleparallel gravity. *Phys. Lett. B* **730**, 136–140 (2014). [arXiv:1401.7378](#)
52. P. Kanti, J. Rizos, K. Tamvakis, Singularity-free cosmological solutions in quadratic gravity. *Phys. Rev. D* **59**, 083512 (1999). [arXiv:gr-qc/9806085](#)
53. Y.F. Cai, C. Li, Y.T. Wang, Towards a nonsingular bouncing cosmology. *JHEP* **03**, 131 (2018). [arXiv:1806.07295](#) [gr-qc]
54. K. Bamba, S.D. Odintsov, P.V. Tretyakov, Inflation in a scalar–tensor model with gauss–bonnet coupling. *Eur. Phys. J. C* **75**, 344 (2015). [arXiv:1411.3852](#)
55. S.D. Odintsov, V.K. Oikonomou, T. Paul, Bottom-up reconstruction of viable inflationary potentials in scalar–gauss–bonnet gravity. *Phys. Rev. D* **104**, 083506 (2021). [arXiv:2108.10460](#) [gr-qc]
56. T. Paul, S.D. Odintsov, Gauss–Bonnet–corrected scalar–tensor inflation: a viable model under GW170817 constraints. [arXiv:2407.12155](#) [gr-qc]
57. F.W. Hehl, P. Heyde, G.D. Kerlick, J.M. Nester, General relativity with spin and torsion: foundations and prospects. *Rev. Mod. Phys.* **48**, 393 (1976)
58. I.L. Shapiro, Physical aspects of the space–time torsion. *Phys. Rep.* **357**, 113–213 (2002). [arXiv:hep-th/0103093](#)
59. N.J. Poplawski, Cosmology with torsion: an alternative to cosmic inflation. *Phys. Lett. B* **694**, 181–185 (2010). [arXiv:1007.0587](#)
60. N.J. Poplawski, Nonsingular, big-bounce cosmology from spinor-torsion coupling. *Phys. Rev. D* **85**, 107502 (2012). [arXiv:1111.4595](#) [gr-qc]
61. K. Bamba, S.D. Odintsov, D. Sáez-Gómez, Inflationary cosmology and the late-time accelerated expansion of the universe in torsion gravity. *Phys. Lett. B* **730**, 136–140 (2014). [arXiv:1401.7378](#)
62. A.H. Chamseddine, V. Mukhanov, Mimetic dark matter. *JHEP* **11**, 135 (2013). [arXiv:1308.5410](#)

63. S. Nojiri, S.D. Odintsov, Mimetic $f(r)$ gravity: inflation, dark energy and bounce. *Mod. Phys. Lett. A* **29**, 1450211 (2014). [arXiv:1408.3561](#)
64. S. Nojiri, S.D. Odintsov, V.K. Oikonomou, Mimetic gravity: a review of recent developments and applications to cosmology and astrophysics. *Mod. Phys. Lett. A* **31**, 1630011 (2016). [arXiv:1605.02187](#) [gr-qc]
65. S. Nojiri, S.D. Odintsov, V.K. Oikonomou, Viable mimetic completion of unified inflation-dark energy evolution in modified gravity. *Phys. Rev. D* **94**, 104050 (2016). [arXiv:1608.07806](#)
66. T. Biswas, T. Koivisto, A. Mazumdar, Towards a resolution of the cosmological singularity in non-local higher derivative theories of gravity. *JCAP* **1011**, 008 (2010). [arXiv:1005.0590](#)
67. L. Modesto, Super-renormalizable quantum gravity. *Phys. Rev. D* **86**, 044005 (2012). [arXiv:1107.2403](#)
68. A.S. Koshelev, L. Modesto, L. Rachwal, A.A. Starobinsky, Occurrence of exact r^2 inflation in non-local uv-complete gravity. *JHEP* **11**, 067 (2016). [arXiv:1604.03127](#)
69. L. Heisenberg, A systematic approach to generalisations of general relativity and their cosmological implications. *Phys. Rep.* **796**, 1–113 (2019). [arXiv:1807.01725](#)
70. J.B. Jiménez, L. Heisenberg, T. Koivisto, The geometrical trinity of gravity. *Universe* **5**, 173 (2019). [arXiv:1903.06830](#)
71. R. Lazkoz, F.S.N. Lobo, M. Ortiz-Baños, V. Salzano, Observational constraints of $f(q)$ gravity. *Phys. Rev. D* **103**, 104026 (2021). [arXiv:2102.04103](#)
72. S. Carloni, P.K.S. Dunsby, S. Capozziello, A. Troisi, Cosmological dynamics of r^n gravity. *Class. Quantum Gravity* **22**, 4839 (2005). [arXiv:gr-qc/0410046](#)
73. M. Jamil, D. Momeni, R. Myrzakulov, Attractor solutions in $f(t)$ cosmology. *Eur. Phys. J. C* **72**, 1959 (2012). [arXiv:1202.4926](#)
74. S. Bahamonde, C.G. Böhm, M. Wright, Modified teleparallel theories of gravity. *Phys. Rev. D* **92**, 104042 (2015). [arXiv:1508.05120](#)
75. M. Sharif, M. Zubair, Phase space analysis in $f(r, t)$ gravity. *JCAP* **03**, 028 (2013). [arXiv:1312.6332](#)
76. D. Gorbunov, A.G. Panin, Scalaron the mighty: producing dark matter and baryon asymmetry at reheating. *Phys. Lett. B* **700**, 157 (2011). [arXiv:1009.2448](#)
77. T. Krajewski, M. Peloso, E. Söderberg, Reheating after starobinsky inflation with scalaron decay. *JCAP* **10**, 009 (2021). [arXiv:2104.00691](#)
78. J. Fumagalli, S. Renaux-Petel, L.T. Witkowski, Oscillons from stringy tachyons in gauss–bonnet inflation. *JCAP* **08**, 037 (2020). [arXiv:2004.02863](#)
79. S. Koh, B.H. Lee, W. Lee, G. Tumurtushaa, Preheating in gauss–bonnet inflation. *Phys. Rev. D* **105**, 043523 (2022). [arXiv:2107.08610](#) [gr-qc]
80. R. Lazkoz, M. Ortiz-Baños, V. Salzano, Reheating constraints on $f(q)$ gravity models. *Eur. Phys. J. C* **82**, 1067 (2022). [arXiv:2205.04481](#)
81. N.J. Poplawski, Einstein–cartan gravity as a possible foundation for inflation and reheating. *Phys. Rev. D* **103**, 105032 (2021). [arXiv:2011.14905](#) [gr-qc]
82. M. Rinaldi, G. Cognola, L. Vanzo, S. Zerbini, Reconstructing the inflationary $f(r)$ in the palatini approach. *JCAP* **12**, 028 (2014). [arXiv:1409.7386](#) [gr-qc]
83. M. Escudero, H. Ramirez, L. Boubekur, E. Giusarma, O. Mena, The present and future of the most favoured inflationary models after planck 2015. *JCAP* **02**, 020 (2016). [arXiv:1509.05419](#) [astro-ph.CO]
84. K. Nozari, M. Sami, Tachyon field in loop quantum cosmology. *Phys. Rev. D* **80**, 123530 (2009). [arXiv:0909.2300](#)
85. C.Q. Geng, C.C. Lee, E.N. Saridakis, Y.P. Wu, ‘teleparallel’ dark energy. *Phys. Lett. B* **704**, 384 (2011). [arXiv:1109.1092](#)
86. R. Herrera, Reheating in tachyonic brane inflationary models with parametric resonance. *Phys. Rev. D* **104**, 043532 (2021). [arXiv:2105.04049](#) [gr-qc]
87. M. Sami, M. Shahalam, M. Skugoreva, A. Toporensky, M. Shahalam, M. Skugoreva, A. Toporensky, *Phys. Rev. D* **86**, 103532 (2012). <https://doi.org/10.1103/PhysRevD.86.103532>. [arXiv:1207.6691](#) [hep-th]
88. S. Kachru et al., Towards inflation in string theory. *JCAP* **0310**, 013 (2003). [arXiv:hep-th/0308055](#)
89. E. Silverstein, A. Westphal, Monodromy in the cmb: gravity waves and string inflation. *Phys. Rev. D* **78**, 106003 (2008). [arXiv:0803.3085](#)
90. L. McAllister, E. Silverstein, A. Westphal, Gravity waves and linear inflation from axion monodromy. *Phys. Rev. D* **82**, 046003 (2010). [arXiv:0808.0706](#) [hep-th]
91. E.J. Copeland, A.R. Liddle, D.H. Lyth, E.D. Stewart, D. Wands, False vacuum inflation with einstein gravity. *Phys. Rev. D* **49**, 6410 (1994)
92. C.P. Burgess, M. Majumdar, D. Nolte, F. Quevedo, G. Rajesh, R.J. Zhang, The inflationary brane-antibrane universe. *JHEP* **07**, 047 (2001). [arXiv:hep-th/0105204](#)
93. R. Brandenberger, C. Vafa, Superstrings in the early universe. *Nucl. Phys. B* **316**, 391 (1989)
94. T. Battefeld, S. Watson, String gas cosmology. *Rev. Mod. Phys.* **78**, 435 (2006). [arXiv:hep-th/0510022](#)
95. G. Obied, H. Ooguri, L. Spodyneiko, C. Vafa, The Swampland: distance conjecture. [arXiv:1806.08362](#)
96. H. Ooguri, C. Vafa, On the geometry of the string landscape and the swampland. *Nucl. Phys. B* **766**, 21–33 (2007). [arXiv:hep-th/0605264](#)
97. S. Dimopoulos, S. Kachru, J. McGreevy, J.G. Wacker, N-flation. *JCAP* **08**, 003 (2008). [arXiv:hep-th/0507205](#)
98. T.C. Bachlechner, M. Dias, J. Frazer, L. McAllister, A new angle on chaotic inflation. *JCAP* **01**, 042 (2014). [arXiv:1404.7496](#) [hep-th]
99. D.E. Kaplan, R. Rattazzi, Large field excursions and approximate discrete symmetries from a clockwork axion. *Phys. Rev. D* **93**, 085007 (2016). [arXiv:1511.01827](#) [hep-ph]
100. S. Ferrara, R. Kallosh, Modular inflation. *Phys. Rev. D* **100**, 126006 (2019). [arXiv:1911.00070](#) [hep-th]
101. J.C. Hwang, H. Noh, Classical evolution and quantum generation in generalized gravity theories including string corrections and tachyon: unified analyses. *Phys. Rev. D* **71**, 063536 (2005). [arXiv:gr-qc/0412126](#)
102. S.D. Odintsov, V.K. Oikonomou, T. Paul, Inflationary phenomenology of einstein gauss–bonnet gravity compatible with gw170817. *Phys. Rev. D* **102**, 104058 (2020). [arXiv:2008.10360](#) [gr-qc]
103. A.D. Linde, Chaotic inflation. *Phys. Lett. B* **129**, 177 (1983)
104. R. Kallosh, A. Linde, Universality class in conformal inflation. *JCAP* **1307**, 002 (2013). [arXiv:1306.5220](#)
105. R. Kallosh, A. Linde, Planck, lhc, and α -attractors. *Phys. Rev. D* **91**, 083528 (2015). [arXiv:1502.07733](#) [astro-ph.CO]
106. A.D. Linde, Hybrid inflation. *Phys. Rev. D* **49**, 748 (1994)
107. K. Freese, J.A. Frieman, A.V. Olinto, Natural inflation with pseudo nambu-goldstone bosons. *Phys. Rev. Lett.* **65**, 3233 (1990)
108. J. Martin, C. Ringeval, V. Vennin, Encyclopædia inflationaris. *Phys. Dark Univ.* **5–6**, 75–235 (2014). [arXiv:1303.3787](#) [astro-ph.CO]
109. G. Dvali, S.H. Tye, Brane inflation. *Phys. Lett. B* **450**, 72 (1999). [arXiv:hep-ph/9812483](#)
110. A. Felice, S. Tsujikawa, $F(r)$ theories. *Living Rev. Relativ.* **13**, 3 (2010). [arXiv:1002.4928](#) [gr-qc]
111. A.A. Starobinsky, A new type of isotropic cosmological models without singularity. *Phys. Lett. B* **91**, 99 (1980)

112. K. Bamba et al., Inflationary universe from perfect fluid and $f(r)$ gravity and its comparison with observational data. *Phys. Rev. D* **88**, 084042 (2013). [arXiv:1308.5789](#)
113. S.D. Odintsov, V.K. Oikonomou, T. Paul, Bottom-up reconstruction of viable inflationary potentials in scalar–gauss–bonnet gravity. *Phys. Rev. D* **104**, 083506 (2021). [arXiv:2108.10460](#) [gr-qc]
114. V. Vennin, L. Pinol, D. Langlois, Forecasting bayesian evidences in inflationary model selection. *JCAP* **04**, 033 (2020). [arXiv:1910.02078](#) [astro-ph.CO]
115. M. Escudero, C. Cosme, Towards machine learning forecasting of inflationary observables. *Phys. Dark Univ.* **38**, 101124 (2022). [arXiv:2202.13933](#) [astro-ph.CO]
116. J.W. Moffat, Scalar–tensor–vector gravity theory. *JCAP* **0603**, 004 (2006). [arXiv:gr-qc/0506021](#)
117. J.W. Moffat, Vector field driven cosmology and the origin of the accelerating universe. *JCAP* **05**, 034 (2011). [arXiv:1104.2957](#) [astro-ph.CO]
118. S. Rahvar, J.W. Moffat, Testing modified gravity (mog) theory with the cosmic microwave background. *Mon. Not. Roy. Astron. Soc.* **471**, 4872–4881 (2017). [arXiv:1512.08401](#) [astro-ph.CO]
119. M. Carmeli, *Cosmological Special Relativity* (World Scientific, Singapore, 2002)
120. J.G. Hartnett, M. Carmeli, Distance–redshift relation in cosmological relativity. *Found. Phys. Lett.* **19**, 6 (2006)
121. M. Carmeli, T. Kuzmenko, Value of the cosmological constant: theory versus experiment. *Int. J. Theor. Phys.* **41**, 131 (2002)
122. J.D. Bekenstein, Fine-structure constant: is it really a constant? *Phys. Rev. D* **25**, 1527 (1982)
123. H. Sandvik, J. Barrow, J. Magueijo, A simple varying-alpha cosmology. *Phys. Rev. Lett.* **88**, 031302 (2002)
124. J. Magueijo, New varying speed of light theories. *Rep. Prog. Phys.* **66**, 2025–2068 (2003). [arXiv:astro-ph/0305457](#)
125. J.D. Barrow, Varying constants. *Phil. Trans. Roy. Soc. Lond. A* **363**, 2139–2153 (2005). [arXiv:gr-qc/0505050](#)
126. E. Calabrese, R. Putter, D. Huterer, E.V. Linder, A. Melchiorri, Future cmb constraints on early, cold, or stressed dark energy. *Phys. Rev. D* **83**, 023011 (2011). [arXiv:1010.5612](#) [astro-ph.CO]
127. R. Maartens, D. Wands, B. Bassett, I. Heard, Chaotic inflation on the brane. *Phys. Rev. D* **62**, 041301 (2000)
128. J.P. Conlon, F. Quevedo, Kähler moduli inflation. *JHEP* **01**, 146 (2006). [arXiv:hep-th/0509012](#)
129. A. Karami, S. Nojiri, Preheating in modified teleparallel gravity: a unified viewpoint. *Phys. Rev. D* **104**, 023519 (2021). [arXiv:2104.12571](#)