



*universe*

IMPACT  
FACTOR  
**2.5**

CITESCORE  
**4.3**

Article

---

# Scalar Field Kantowski–Sachs Solutions in Teleparallel $F(\mathcal{T})$ Gravity

---

Alexandre Landry

Special Issue

Dark Energy and Dark Matter

Edited by

Prof. Dr. Yan Gong



<https://doi.org/10.3390/universe11010026>

## Article

# Scalar Field Kantowski–Sachs Solutions in Teleparallel $F(T)$ Gravity

Alexandre Landry 

Department of Mathematics and Statistics, Dalhousie University, Halifax, NS B3H 3J5, Canada; a.landry@dal.ca; Tel.: +1-514-503-2051

**Abstract:** In this paper, we investigate time-dependent Kantowski–Sachs spherically symmetric teleparallel  $F(T)$  gravity with a scalar field source. We begin by setting the exact field equations to be solved and solve conservation laws for possible scalar field potential,  $V(\phi)$ , solutions. Then, we find new non-trivial teleparallel  $F(T)$  solutions by using power-law and exponential ansatz for each potential case arising from conservation laws, such as linear, quadratic, or logarithmic, to name a few. We find a general formula allowing us to compute all possible new teleparallel  $F(T)$  solutions applicable for any scalar field potential and ansatz. Then, we apply this formula and find a large number of exact and approximate new teleparallel  $F(T)$  solutions for several types of cases. Some new  $F(T)$  solution classes may be relevant for future cosmological applications, especially concerning dark matter, dark energy quintessence, phantom energy leading to the Big Rip event, and quintom models of physical processes.

**Keywords:** teleparallel gravity; field equations; Kantowski–Sachs spacetimes; scalar field source solutions; frame-based approach; time-dependent spacetimes

## 1. Introduction

Teleparallel  $F(T)$  gravity is an alternative theory to General Relativity (GR) of the frame-based type [1–7]. This theory is very promising and is in great expansion. It is defined in terms of spacetime torsion tensor  $T_{bc}^a$  and torsion scalar  $T$ , and its physical quantities are all defined in terms of the coframe  $h^a$  and the spin-connection  $\omega_{bc}^a$  (and their derivatives). This is in contrast to GR being defined in terms of the metric  $g_{\mu\nu}$  and the spacetime curvature  $R_{b\mu\nu}^a$ . One of the features of teleparallel gravity and its frame-based approach is the new possible spacetime symmetries, especially for non-trivial linear isotropy groups, and its Lorentz-invariant geometries [8–10]. There is thus an approach to determine these symmetries for any independent coframe/spin-connection pairs while treating spacetime curvature and torsion as geometric objects [4,5,11]. From this point, any geometry described by an even coframe/spin-connection whose curvature and non-metricity are both zero ( $R_{b\mu\nu}^a = 0$  and  $Q_{a\mu\nu} = 0$  conditions) is a teleparallel geometry. Covariantly, this type of geometry is defined as being gauge invariant (valid for any  $g_{ab}$ ). In the orthonormal gauge  $g_{ab} = \eta_{ab} = \text{Diag}[-1, 1, 1, 1]$ , the relations to satisfy for such a geometry are [11]:

$$\mathcal{L}_X h^a = \lambda_b^a h^b, \quad (1)$$

$$\mathcal{L}_X \omega_{bc}^a = 0, \quad (2)$$

where  $h^a$  is the orthonormal coframe basis,  $\mathcal{L}_X$  is the Lie derivative in terms of Killing Vectors (KV)  $X$ , and  $\lambda_b^a$  is the generator of the Lorentz transformations  $\Lambda_b^a$ . For a pure



Academic Editor: Yan Gong

Received: 6 December 2024

Revised: 7 January 2025

Accepted: 14 January 2025

Published: 16 January 2025

**Citation:** Landry, A. Scalar Field Kantowski–Sachs Solutions in Teleparallel  $F(T)$  Gravity. *Universe* **2025**, *11*, 26. <https://doi.org/10.3390/universe11010026>

**Copyright:** © 2025 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

teleparallel geometry, one must also satisfy the zero Riemann curvature criterion (see refs. [1–7] and references therein):

$$\begin{aligned} R^a_{b\mu\nu} &= \partial_\mu \omega^a_{b\nu} - \partial_\nu \omega^a_{b\mu} + \omega^a_{e\mu} \omega^e_{b\nu} - \omega^a_{e\nu} \omega^e_{b\mu} = 0, \\ \Rightarrow \omega^a_{b\mu} &= \Lambda^a_c \partial_\mu \Lambda^c_b. \end{aligned} \quad (3)$$

The solution of Equation (3) leads to the teleparallel spin-connection defined in terms of a Lorentz transformation  $\Lambda^a_b$ . Note also that  $\omega^a_{b\mu} = 0$  for all proper frames and that  $\omega^a_{b\mu} \neq 0$  for all non-proper frames. Moreover, a proper frame can be defined in several ways in terms of spin-connections; this can make it difficult to determine the symmetries.

There is a direct equivalent to GR defined under teleparallel gravity: the Teleparallel Equivalent to General Relativity (TEGR or  $F(T) = T$  theory) [1]. However, TEGR easily generalizes to teleparallel  $F(T)$  gravity, with  $F$  as a function of the scalar torsion  $T$  [12–14]. Teleparallel  $F(T)$  gravity is known to be locally invariant under the definition of covariant Lorentz [7]. In addition to the teleparallel  $F(T)$ -type gravity, all the considerations mentioned so far have been adapted for extended theories like New General Relativity (NGR) (see refs. [15–17] and references therein) and symmetric teleparallel  $F(Q)$ -type gravity (see refs. [18–21] and references therein), as well as several intermediate theories like  $F(T, Q)$ -type,  $F(R, Q)$ -type,  $F(R, T)$ -type, and others (see refs. [22–28] and references therein). Therefore, the best approach is to go with the teleparallel  $F(T)$  gravity for the present approach.

There have been several works and papers on spherically symmetric spacetimes and its solutions in teleparallel  $F(T)$  gravity using various approaches and for several types of uses [29–50]. However, solving FEs in the orthonormal gauge has been favored for reasons of determining coframe/spin-connection pairs and also to avoid the extra degrees of freedom (DoF) problem associated with using proper frames to solve FEs (see refs. [44–47] for detailed discussions). However, the symmetric FEs and their  $F(T)$  solutions will be very similar between different gauges (orthonormal or not); this further confirms the gauge invariance of general teleparallel  $F(T)$  gravity FEs. But by opting for the orthonormal gauge, one will solve the non-trivial antisymmetric and symmetric parts of FEs covering all DoFs for a diagonal coframe and a non-trivial spin-connection (see, for example, refs. [44–47]). In the literature, many prefer to use tetrads  $e^a_\mu$  instead of coframes  $h^a_\mu$ , because they are defined in an orthonormal gauge by default [29–43]. However, the tetrad approach prevents the use of frames other than orthonormal, and this is the greatest weakness of the latter approach. It is to avoid this type of limitation, to respect gauge invariance as well as to work on any type of frame that is not necessarily orthonormal, that the coframe notation  $h^a_\mu$  will be used here and has been used in refs. [44–47].

One of the major scopes and aims in the study of teleparallel  $F(T)$  solutions concerns the study of cosmological models involving the different forms of dark energy, such as quintessence, phantom (or negative) energy, quintom, and also dark matter. To achieve this aim, some recent studies in teleparallel  $F(T)$  gravity deal with Kantowski–Sachs (KS) spacetimes and the possible  $F(T)$  solutions by using various ansatz [45–47,51,52]. This type of spacetime is characterized by the radial derivative  $\partial_r$  as the fourth symmetry (KV), leading directly to the pure time dependence for coframes, spin-connections, and FEs [45–47]. Previously, there have been some detailed dynamical studies concerning the evolution of critical points, boundary values, asymptotes, and  $\overset{\circ}{R}$  curvature in KS spacetimes under the GR and/or the  $f(\overset{\circ}{R})$  gravity frameworks [53–55]. Very similar studies using the same types of dynamical methods for KS spacetimes and their generalizations have also been carried out, not only in  $f(\overset{\circ}{R})$  gravity but also in  $F(T, B)$ -type, LRS Bianchi III Universe,  $F(T, R)$ -type, and  $F(Q)$ -type gravity theories [56–61]. Recent works on cosmological so-

lutions in teleparallel KS spacetimes with a quantized scalar field source should also be mentioned [62,63]. All these studies confirm the relevance and the accuracy of cosmological KS spacetimes solutions, especially for models involving different forms of dark energy.

However, most of these previous studies were not carried out in terms of teleparallel  $F(T)$  gravity and do not really provide new  $F(T)$  solutions. In particular, there has been a recent study on perfect fluid KS teleparallel  $F(T)$  solutions, revealing relevant solutions for the dark energy quintessence, but also for the phantom dark energy models [46]. It is briefly mentioned in this study that the different forms of dark energy mentioned above are at the same time explainable by fundamental scalar fields. It is therefore quite logical to study cosmological KS  $F(T)$  solutions for different types of scalar fields as a logical follow-up to this last study. All this will allow one to not only confirm the  $F(T)$  solutions obtained in this ref. [46] but also to better study the  $F(T)$  solutions and the fundamental scalar fields associated with dark matter, dark energy quintessence, phantom energy, and also quintom cosmological models using the teleparallel gravity framework and without the quantization of the scalar field source. Most recently, this type of study has been carried out for Teleparallel Robertson–Walker (TRW) spacetimes in  $F(T)$  and  $F(T, B)$  gravity (spacetime geometry at six KVs instead of four) with various scalar field sources [50,64]. Beyond spacetime and symmetry considerations, this type of study will mainly serve purely physical motivations.

A significant physical motivation for this type of study concerns the dark energy quintessence models [65–69]. This new type of model was first developed by Paul J. Steinhardt et al. in the late 1990s [65–67]. There have been various phenomenological studies performed via the GR frameworks in order to obtain a model explaining the accelerating universe expansion (see refs. [65–72] and several other ones). In these models, a perfect linear fluid of negative pressure is assumed ( $P = \alpha_Q \rho$ , where  $-1 < \alpha_Q < -\frac{1}{3}$  models), supported and induced by a scalar field. From this starting point, there have been various dynamical and other studies on the possibilities of scalar fields and possible potentials that most faithfully describe the dark energy quintessence process and thus the accelerating universe expansion [69,70]. Even recently, there are increasingly advanced studies on this very important subject in cosmology, specifically for observational constraints [71,72]. However, several years before Paul Steinhardt’s works on quintessence processes, there had been attempts at models of accelerated universe expansion through the use of scalar field-based models and their symmetries explaining accelerating universe expansion [73,74]. The same precursor models therefore allowed all of P. Steinhardt’s work on dark energy quintessence, generalizations, and other subsequent studies to emerge and arrive at the current models.

Among the set of recent models are the teleparallel gravity-based dark energy quintessence models and their extensions (in particular,  $F(T)$ -type,  $F(T, B)$ -type, scalar-torsion-based) [75–84]. However, these papers have mainly concentrated on dynamics and especially on stability studies on FEs using simple and predefined  $F(T)$  and  $F(T, B)$  functions as superpositions of power-law and/or logarithm terms. Even if these latter types of function are often used in teleparallel cosmology, this somewhat simplistic approach does not truly and fundamentally solve the FEs of teleparallel  $F(T)$  gravity and/or its extensions. These studies mainly only primarily study the impacts on some physical parameters of teleparallel gravity. At the same time, this type of approach makes the teleparallel  $F(T)$  exact, and correct solutions of FEs that can really model the dark energy quintessence processes go under the radar. There is therefore room to go further by solving the teleparallel  $F(T)$  gravity FEs for scalar field sources, as was the case in refs. [46,47] for teleparallel  $F(T)$  solutions for perfect fluid sources. Most recently, there have been detailed studies of Teleparallel Robertson–Walker (TRW)  $F(T)$  and  $F(T, B)$  solutions with perfect fluids and scalar field sources, allowing a larger number of possible solutions for

teleparallel dark energy quintessence models [49,64]. In this manner, we will find the most fundamental solutions for teleparallel KS spacetimes, allowing us to study the best and most fundamental possible models explaining dark energy quintessence by using the frameworks and FEs of teleparallel  $F(T)$  gravity.

In addition, the first and primarily dark energy quintessence model studies also led to the scenarios of cosmological models involving strong negative pressure dark energy perfect fluids (where  $\alpha_Q < -1$ ) [85,86]. This is also a new physical motivation for studying more extreme scenarios of strong and fast accelerating universe expansion, such as phantom (or negative) dark energy models [87–90]. Even in teleparallel gravity, some studies have been considered in phantom dark energy models using the same approaches as for dark energy quintessence processes studies [78,79,91–93]. These works still focus on dynamic and stability studies using very specific teleparallel  $F(T)$  functions (and/or  $F(T, B)$ ) to study the evolution of some specific physical parameters without really directly solving the FEs of teleparallel gravity. Obviously, there has been less work on this type of scenario, because it is more hypothetical. This type of model often involves nonlinear fluids, but the determining feature is mostly the energy condition violation of such a cosmological system (i.e.,  $P + \rho \not\geq 0$  situation), which makes that hypothetical. Recent works in teleparallel  $F(T)$  and  $F(T, B)$  types of gravity for TRW spacetimes provide a larger amount of material and new solutions to further study the phantom energy models in addition to quintessence processes [49,64]. There is also a way and a reason here to go further by directly solving the teleparallel  $F(T)$  gravity FEs with a scalar field source, especially for the KS spacetimes.

Furthermore, the works mentioned show the existence of at least three possible types of dark energy: quintessence, cosmological constant, phantom energy. There are also the quintom dark energy models defined as a mixture of quintessence dark energy ( $-1 < \alpha_Q < -\frac{1}{3}$ ) and phantom dark energy ( $\alpha_Q < -1$ ), having as the intermediate limit the cosmological constant [94–98]. This type of hybrid model is often defined by spacetime geometry and two scalar fields: the first field can describe quintessence and the second phantom energy or one field for the unified process and the other for coupling between the two types of dark energy [94,95,98]. Some works specifically address this problem through the oscillation between the quintessence and phantom states [96]. However, in teleparallel gravity there have been very few works dealing with this type of more complex physical problem [99]. This is also an additional reason favoring the present approach to solve the teleparallel  $F(T)$  gravity FEs with a scalar field source. The results arising from the current work will provide significantly more materials and tools for better developing the teleparallel quintom dark energy models in the future. There are also anisotropic cosmological models that can be added to the list of possibilities emerging from the present approach [100]. There is no lack of possibilities, but here we focus on the KS teleparallel  $F(T)$  solutions with scalar fields to open up these possibilities for future development.

For this paper, we assume a time-coordinate dependent spherically symmetric teleparallel geometry, in particular Kantowski–Sachs teleparallel geometry, in an orthonormal gauge as defined and used in refs. [45–47]. We will find  $F(T)$  solutions for a power-law defined scalar field and then for an exponential scalar field source. After a summary of the teleparallel FEs and Kantowski–Sachs class of geometries in Section 2, we will find in Section 3 the power-law scalar field exact and approximated  $F(T)$  solutions. We then repeat the exercise in Section 4 with an exponential scalar field source for exact and approximated  $F(T)$  solutions. In both cases, we will use a power-law and an exponential coframe component ansatz for a better comparison between  $F(T)$  solutions. We will finish the investigation by discussing in Section 5 the other possible scalar field sources and solving methods, especially by solving the special case of logarithmic source in Section 5.2. This

paper also has common features, aims, and a similar structure to some recent papers on perfect fluid teleparallel  $F(T)$  solutions (see refs. [44,46,47] for details).

The notation is defined as follows: coordinate indices  $\mu, \nu, \dots$ , tangent space indices  $a, b, \dots$  (see ref. [4]), spacetime coordinates  $x^\mu$ , coframe  $\mathbf{h}_a$  or  $\mathbf{h}^a$ , vierbein  $h_a^\mu$  or  $h_\mu^a$ , metric  $g_{\mu\nu}$ , gauge metric  $g_{ab}$ , spin-connection  $\omega_b^a = \omega_{bc}^a \mathbf{h}^c$ , curvature tensor  $R_{bcd}^a$ , torsion tensor  $T_{bc}^a$ . The derivatives with respect to (w.r.t.)  $t$ ,  $F_t = F'$  (with a prime).

## 2. Summary of Teleparallel Gravity and Scalar Field Kantowski–Sachs Field Equations

### 2.1. Teleparallel $F(T)$ Gravity Theory

As in several recent works, the teleparallel  $F(T)$ -type gravity action integral is [1–5,44–47]:

$$S_{F(T)} = \int d^4x \left[ \frac{h}{2\kappa} F(T) + \mathcal{L}_{Source} \right], \quad (4)$$

where  $h$  is the coframe determinant and  $\kappa$  is the coupling constant. We apply the least-action principle to Equation (4), and we find the general FEs in terms of coframe and spin-connection [1–5]:

$$\begin{aligned} \kappa \Theta_a^\mu = & h^{-1} F_T(T) \partial_\nu (h S_a^{\mu\nu}) + F_{TT}(T) S_a^{\mu\nu} \partial_\mu T + \frac{F(T)}{2} h_a^\mu - F_T(T) T_{av}^b S_b^{\mu\nu} \\ & - F_T(T) \omega_{av}^b S_b^{\mu\nu}, \end{aligned} \quad (5)$$

where  $\Theta_a^\mu$  is the energy–momentum,  $T$  the torsion scalar,  $T_{av}^b$  the torsion tensor,  $h_a^\mu$  the coframe (or tetrad for orthonormal frames),  $\omega_{av}^b$  the spin-connection, and  $S_a^{\mu\nu}$  the superpotential (torsion dependent). We can separate Equation (5) into the symmetric and antisymmetric parts of FEs as [44–47]:

$$\kappa \Theta_{(ab)} = F_T(T) \overset{\circ}{G}_{ab} + F_{TT}(T) S_{(ab)}^\mu \partial_\mu T + \frac{g_{ab}}{2} [F(T) - T F_T(T)], \quad (6)$$

$$0 = F_{TT}(T) S_{[ab]}^\mu \partial_\mu T, \quad (7)$$

with  $\overset{\circ}{G}_{ab}$  being the Einstein tensor and  $g_{ab}$  the gauge metric. Equations (5)–(7) are valid for any type of gauge (they satisfy the gauge invariance principle), coframe, and spin-connection, including the orthonormal frames and gauges [3,5,45]. However, there are some papers where the coframe  $h_a^\mu$  will be replaced for orthonormal frames by the tetrads  $e_\mu^a$  (see refs. [29–43]), but this last notation is not as general as the coframe  $h_\mu^a$  [5,45].

### 2.2. General Conservation Laws

The canonical energy–momentum and its conservation laws are obtained from the  $\mathcal{L}_{Matter}$  term of Equation (4) as [1–3,44–47]:

$$\Theta_a^\mu = \frac{1}{h} \frac{\delta \mathcal{L}_{Source}}{\delta h_\mu^a}, \quad (8)$$

$$\Rightarrow \overset{\circ}{\nabla}_\nu (\Theta^{\mu\nu}) = 0, \quad (9)$$

with  $\overset{\circ}{\nabla}_\nu$  being the covariant derivative and  $\Theta^{\mu\nu}$  the conserved energy–momentum tensor. Equation (9) is also the GR conservation of energy–momentum expression. The Equation (8) antisymmetric and symmetric parts are [44–47]:

$$\Theta_{[ab]} = 0, \quad \Theta_{(ab)} = T_{ab}, \quad (10)$$



where  $T_{ab}$  is the symmetric part of the energy–momentum tensor. Equation (10) is valid only when the matter field interacts with the metric  $g_{\mu\nu}$ , defined from the coframe  $h^a_\mu$  and the gauge  $g_{ab}$ , and is not directly coupled to the  $F(T)$  gravity. Equation (9) also imposes the symmetry of  $\Theta^{\mu\nu}$  and hence the condition stated by Equations (10). All the previous considerations are valid only when the hypermomentum is zero (i.e.,  $\mathfrak{T}^{\mu\nu} = 0$ ), as discussed in refs. [31,44,46,47]. The hypermomentum condition is defined from Equations (6) and (7) as [31]:

$$\mathfrak{T}_{ab} = \kappa \Theta_{ab} - F_T(T) \overset{\circ}{G}_{ab} - F_{TT}(T) S_{ab}{}^\mu \partial_\mu T - \frac{g_{ab}}{2} [F(T) - T F_T(T)] = 0. \quad (11)$$

Therefore, there exists a general non-zero hypermomentum (i.e.,  $\mathfrak{T}^{\mu\nu} \neq 0$ ) conservation law condition for any teleparallel gravity theory which generalizes the Equation (9) condition [31,101–103]. This general hypermomentum conservation law is trivially satisfied for the  $\mathfrak{T}^{\mu\nu} = 0$  situation (the current situation and any GR solution).

### 2.3. Teleparallel Kantowski–Sachs Coframe and Spin-Connection Solutions

The orthonormal time-dependent Kantowski–Sachs resulting vierbein in  $(t, r, \theta, \phi)$  coordinates is [45–47]:

$$h^a_\mu = \text{Diag}[1, A_2(t), A_3(t), A_3(t) \sin(\theta)], \quad (12)$$

where we are able to choose new coordinates such that  $A_1(t) = 1$  without any loss of generality. This will allow us to find cosmological-like solutions.

Another possible orthonormal coframe solution is taking the choice  $A_3(t) = t$  and then solve for  $A_1(t)$  and  $A_2(t)$  [47]:

$$h^a_\mu = \text{Diag}[A_1(t), A_2(t), t, t \sin(\theta)]. \quad (13)$$

In refs. [45,46], we find that the pure vacuum KS coframe solution implies that  $A_3 = t$ . This is another justification for the Equation (13) coframe expression. However, Equation (13) is not as practical and appropriate as Equation (12) for cosmological-like solutions. Equations (12) and (13) can also be expressed in principle in terms of tetrads  $e^a_\mu$  only for orthonormal gauges. But the tetrad notation, as seen in the literature, limits the scope of the gauge choices for physical quantity expression. Because the teleparallel  $F(T)$  gravity is gauge invariant and the FEs are defined for any gauge choice (gauge invariant), it is preferable to express the frame components in terms of  $h^a_\mu$  instead of  $e^a_\mu$  [3,5,45]. By using the  $h^a_\mu$  coframe notation, we are able to make the right gauge choice, and we are only limited to orthonormal gauges.

The spin-connection  $\omega_{abc}$  non-null components for time-dependent spacetimes are expressed using the Equation (12) coframe components as  $\omega_{abc} = \omega_{abc}(\psi(t), \chi(t))$ , where  $\chi$  and  $\psi$  are arbitrary functions and  $a \neq b$  indexes (see refs. [45–47] for detailed discussion and calculations). In refs. [45–47], we found from the antisymmetric part of the  $F(T)$  FEs that the solution is  $\psi = 0$  and  $\chi = \frac{\pi}{2}$  ( $\frac{3\pi}{2}$  is also a solution). In terms of spin-connection, the only non-zero  $\omega_{abc}$  components are [45–47]:

$$\omega_{234} = -\omega_{243} = \delta, \quad \omega_{344} = -\frac{\cos(\theta)}{A_3 \sin(\theta)}, \quad (14)$$

where  $\delta = \pm 1$ . Fundamentally, Equations (12) (or Equation (13)) and (14) are also the solutions of Equations (1) and (2) using the Cartan–Karlhede (CK) algorithm method and also Equation (3) for zero curvature criteria (see refs. [4,5,45] for details and justifications).

#### 2.4. Scalar Field Source Conservation Laws

For the scalar field source action integral, the  $\mathcal{L}_{Source}$  term will be described by the Lagrangian density [2,49,64,76,77,81,82]:

$$\mathcal{L}_{Source} = \frac{h}{2} \overset{\circ}{\nabla}_\nu \phi \overset{\circ}{\nabla}^\nu \phi - h V(\phi). \quad (15)$$

We use, in Equation (16), the covariant derivative  $\overset{\circ}{\nabla}_\nu$  for satisfying the GR conservation law defined by Equation (9).

For a time-dependent  $\phi = \phi(t)$  scalar field source and a cosmological-like spacetime (i.e.,  $A_1 = 1$ ), Equation (15) will be simplified as:

$$\mathcal{L}_{Source} = h \frac{\phi'^2}{2} - h V(\phi), \quad (16)$$

where  $\phi' = \partial_t \phi$ . By applying the least-action principle to Equation (16), we find that the energy–momentum tensor  $T_{ab}$  is [44–47,104–106]:

$$T_{ab} = (P_\phi + \rho_\phi) u_a u_b + g_{ab} P_\phi, \quad (17)$$

where  $u_a = (-1, 0, 0, 0)$  and  $P_\phi$  and  $\rho_\phi$  are, respectively, the pressure and density equivalent for the scalar field defined by [49]:

$$P_\phi = \frac{\phi'^2}{2} - V(\phi) \quad \text{and} \quad \rho_\phi = \frac{\phi'^2}{2} + V(\phi), \quad (18)$$

where  $\phi' = \phi_t$  and  $V = V(\phi)$  is the scalar field potential. Then, the conservation law for scalar fields with density and pressure defined by Equations (18) for time-dependent spacetimes is [45–47]:

$$\phi' \left( \ln(A_2 A_3^2) \right)' + \phi'' + \frac{dV}{d\phi} = 0. \quad (19)$$

Equation (19) is the most general scalar field conservation law equation for any  $A_2$  and  $A_3$  ansatz components with any  $\phi(t)$  scalar field expression. For the coming steps, we will first solve Equation (19) for its possible potential for study in this investigation. Then, we will solve the FEs to find the corresponding classes of teleparallel  $F(T)$  solutions for each potential case satisfying Equation (19). We will do these derivation steps for power-law and exponential scalar field  $\phi(t)$  definitions. Equation (18) pressure and density definitions allow us to use the perfect fluid energy–momentum equivalence, as seen in Equation (17), for solving the FEs with a scalar field source. The Equations (17)–(19) also show that the change from a pure linear perfect fluid of  $P = \alpha \rho$  to a scalar field pressure  $P_\phi$  and density  $\rho_\phi$  is clearly a natural and a straightforward generalization of the physical problem. This transformation will allow us to more easily study the fundamental causes of the dark energy physical processes by only using the mathematical analogy between the two types of physical system.

To complete the discussion on physical implications from Equation (18) and to better make the link with a linear perfect fluid EoS equivalent  $P_\phi = \alpha_Q \rho_\phi$ , we need to find the dark energy coefficient index  $\alpha_Q$  (also called the quintessence coefficient) for making the



parallel between  $\phi$  and the dark energy processes. The  $\alpha_Q$  index will be defined from Equation (18) as [65–67,72]:

$$\alpha_Q = \frac{P_\phi}{\rho_\phi} = \frac{\phi'^2 - 2V(\phi)}{\phi'^2 + 2V(\phi)} = -1 + \frac{2\phi'^2}{\phi'^2 + 2V(\phi)}. \quad (20)$$

We can use this Equation (20) for the dark energy perfect fluid equivalent for every new teleparallel  $F(T)$  solution, any potential  $V(\phi)$ , and any ansatz. But there are physical and especially dark energy state significations for  $\alpha_Q$ , such as:

1. **Quintessence**  $-1 < \alpha_Q < -\frac{1}{3}$ : This is the most relevant and possible dark energy state explaining the controlled universe accelerating expansion by using an associated scalar field  $\phi$ , called the quintessence scalar field [65–67,72].
2. **Phantom Energy**  $\alpha_Q < -1$ : This case leads to phantom dark energy processes characterized by an uncontrolled accelerating universe expansion going to the Big Rip at the end of the physical process [87–89]. There is also a fundamental scalar field  $\phi$  describing the phantom energy processes.
3. **Cosmological constant**  $\alpha_Q = -1$ : This is the first and most fundamental dark energy state. It is also the boundary between quintessence and phantom dark energy states. A constant scalar field leads to the cosmological constant according to Equation (20), a GR solution [64].
4. **Quintom models**: This is a mix of dark energy quintessence and phantom dark energy physical processes and is usually described by a two-scalar-fields model [94–98].

This classification is also to express the new teleparallel  $F(T)$  solutions in terms of dark energy quintessence and phantom and quintom processes.

### 2.5. Symmetric and Unified Field Equations

The torsion scalar and the symmetric FE components for the  $\chi = \frac{\pi}{2}$  ( $\delta = +1$ ) case in Equation (14) are [45–47]:

$$T = 2(\ln(A_3))' \left( (\ln(A_3))' + 2(\ln(A_2))' \right) - \frac{2}{A_3^2}, \quad (21)$$

$$B' = - \left( \ln \left( A_2 A_3^2 \right) \right)' + \frac{\frac{1}{A_3^2} - \left( \ln \left( \frac{A_2}{A_3} \right) \right)''}{\left( \ln \left( \frac{A_2}{A_3} \right) \right)'}, \quad (22)$$

$$\kappa \phi'^2 + 2\kappa V(\phi) = -F(T) + 2 \left( T + \frac{2}{A_3^2} \right) F_T(T), \quad (23)$$

$$-\kappa \phi'^2 = \left[ (\ln(A_3))' \left( B' - \left( \ln \left( \frac{A_2}{A_3} \right) \right)' \right) + (\ln(A_3))'' \right] F_T(T), \quad (24)$$

where  $F_T(T) \neq \text{constant}$  and  $B' = \partial_t(\ln F_T(T))$ . Compared with the Kantowski–Sachs  $F(T)$ -gravity FEs in the literature [51,52], the FEs are different. For the  $\delta = -1$  FEs set, there are some small minor differences for some terms in Equations (22)–(24), mainly some different signs at very specific terms. For the rest, the general form of Equations (22)–(24) remains identical, regardless of  $\delta$ . However, we can simplify Equations (22)–(24) by adding Equations (23) and (24) and then by substituting Equations (21) and (22):

$$2\kappa V(\phi(T)) = -F(T) + \left[ \frac{3}{2} \left( T + \frac{2}{A_3^2} \right) + \frac{(\ln(A_3))'}{\left( \ln \left( \frac{A_2}{A_3} \right) \right)'} \left( \frac{1}{A_3^2} - \left( \ln \left( \frac{A_2}{A_3} \right) \right)'' \right) + (\ln(A_3))'' \right] F_T(T). \quad (25)$$

Equation (25) will be the DE to solve for  $F(T)$ . We will substitute the right ansatz and again use Equation (21) as a characteristic equation, as in ref. [46]. In addition, we will have to calculate for each used ansatz the corresponding  $V(\phi(T))$  by using Equation (19) for conservation laws. The Equation (21) characteristic equation will also be useful for finding the  $\phi(T)$  scalar field and then the  $V(\phi(T))$  functions in terms of  $T$ .

Another method based on Equation (25) consists of setting coframe components, Equation (21) and a  $F(T)$  function ansatz to find the exact scalar field potential  $V(\phi)$ , and then  $\phi(t)$  by using Equation (19). However, this method does not allow one to control the source and/or to take into account the exact homogeneous parts of an  $F(T)$ .

### 3. Power-Law Scalar Field Solutions

In this section, we will set a power-law scalar field  $\phi(t) = p_0 t^p$ , where  $p_0$  is a constant and  $p$  is a real power. The conservation law defined by Equation (19) becomes, with field expression inversion as  $t(\phi) = \left(\frac{\phi}{p_0}\right)^{1/p}$ :

$$0 = p p_0^{1/p} \left( \ln(A_2 A_3^2) \right)' \phi^{1-1/p} + p(p-1) p_0^{2/p} \phi^{1-2/p} + \frac{dV}{d\phi}, \quad (26)$$

where  $(\ln(A_2 A_3^2))'$  is dependent on  $\phi$  and the used ansatz for  $A_2$  and  $A_3$  components. There are a number of possible ansatz:

1. **Power-law:** This is the most simple, used, and well-known ansatz. It is defined as [46]:

$$A_2 = t^b \quad \text{and} \quad A_3 = c_0 t^c, \quad (27)$$

where  $c_0$  is a constant. Then, we find that  $(\ln(A_2 A_3^2))' = \frac{(b+2c)}{t(\phi)} = (b+2c) p_0^{1/p} \phi^{-1/p}$  and Equation (26) becomes:

$$\begin{aligned} 0 &= p p_0^{2/p} (b+2c+p-1) \phi^{1-2/p} + \frac{dV}{d\phi}, \\ \Rightarrow V(\phi) &= \phi_0 - \frac{p^2 p_0^{2/p}}{2(p-1)} (b+2c+p-1) \phi^{2-2/p}. \end{aligned} \quad (28)$$

The  $\alpha_Q$ -index from Equation (20) is, by substituting Equation (28):

$$\alpha_Q = -1 + \frac{p^2 p_0^{2/p} \phi^{2-2/p}}{\phi_0 - \frac{p^2 p_0^{2/p}}{2} \left( \frac{b+2c}{p-1} \right) \phi^{2-2/p}}. \quad (29)$$

There are special cases:

- $p \gg 1$ : Equation (28) can be approximated as:

$$V(\phi) \approx \phi_0 - \frac{p^2}{2} \phi^2. \quad (30)$$

Then, Equation (29) will be, for this case:

$$\alpha_Q \approx -1 + \frac{p^2 \phi^2}{\phi_0 - \frac{p(b+2c)}{2} \phi^2} \approx -1 - \frac{2p}{b+2c}, \quad (31)$$

where we have the physical situations:

- $\frac{1}{3} < \frac{p}{b+2c} < 0$ : dark energy quintessence process. If  $p > 0$ , then  $b+2c < 0$ .
- $\frac{p}{b+2c} > 0$ : phantom energy process. If  $p > 0$ , then  $b+2c > 0$ .

If  $b + 2c$  is not large, Equation (31) will rather be  $\alpha_Q = -1 + \frac{p^2}{\phi_0} \phi^2$ , without any constraint on  $b + 2c$ . This last result can be directly found by using Equation (30) and then the Equation (20)  $\alpha_Q$  definition.

- $p = 1$ : Equation (28) is simplified and then becomes:

$$\begin{aligned} 0 &= p_0^2 (b + 2c) \phi^{-1} + \frac{dV}{d\phi}, \\ \Rightarrow V(\phi) &= \phi_0 - p_0^2 (b + 2c) \ln \phi. \end{aligned} \quad (32)$$

Then, Equation (20) is, by substituting Equation (32):

$$\alpha_Q = -1 + \frac{p_0^2}{\frac{p_0^2}{2} + \phi_0 - p_0^2 (b + 2c) \ln \phi}. \quad (33)$$

2. **Exponential:** This case is defined by [46]:

$$A_2 = e^{bt} \quad \text{and} \quad A_3 = c_0 e^{ct}. \quad (34)$$

The Equation (34) form leads to  $(\ln(A_2 A_3^2))' = (b + 2c)$ , and Equation (26) becomes, in this case:

$$\begin{aligned} 0 &= p p_0^{1/p} (b + 2c) \phi^{1-1/p} + p(p-1) p_0^{2/p} \phi^{1-2/p} + \frac{dV}{d\phi}, \\ \Rightarrow V(\phi) &= \phi_0 - \frac{p^2 (b + 2c)}{(2p-1)} p_0^{1/p} \phi^{2-1/p} - \frac{p^2}{2} p_0^{2/p} \phi^{2-2/p}. \end{aligned} \quad (35)$$

Then, Equation (20) will be:

$$\alpha_Q = -1 + \frac{p^2 p_0^{2/p} \phi^{2-2/p}}{\phi_0 - \frac{p^2 (b+2c)}{(2p-1)} p_0^{1/p} \phi^{2-1/p}}. \quad (36)$$

The previous special cases become:

- $p \gg 1$ : Equations (35) and (36) will be exactly Equations (30) and (31).
- $p = 1$ : Equation (35) yields a linear potential:

$$\begin{aligned} 0 &= p_0 (b + 2c) + \frac{dV}{d\phi}, \\ \Rightarrow V(\phi) &= \phi_0 - p_0 (b + 2c) \phi, \end{aligned} \quad (37)$$

and then Equation (20) becomes, with the Equation (37) potential:

$$\alpha_Q = -1 + \frac{p_0^2}{\frac{p_0^2}{2} + \phi_0 - p_0 (b + 2c) \phi}. \quad (38)$$

We also note the scalar potential  $V(\phi)$  solutions to Equation (26), obtained for a power-law scalar field, confirm those found in one of the first studies on dark energy quintessence [69]. This was just confirmed by the separate use of the Equations (27) and (34) ansatzes, which shows the rightness and relevance of the potential solutions compared to the first models studied in the past literature.

### 3.1. Power-Law Ansatz Solutions

Equation (21), in terms of Equation (27), leads to the characteristic equation (see ref. [46]):

$$0 = 2c(c + 2b)t^{-2} - \frac{2}{c_0^2}t^{-2c} - T, \quad (39)$$

where  $c \neq 0$  ( $c = 0$  subcase leads to GR solutions). Then, Equation (25) becomes, under the Equation (27) ansatz and by replacing  $V(\phi(T)) = \phi_0 + V(T)$ :

$$\begin{aligned} \Lambda_0 + 2\kappa V(T) &= -F(T) + \left[ \frac{3}{2}T + \frac{(3b - 2c)}{c_0^2(b - c)}t^{-2c}(T) \right] F_T(T), \\ &= -F(T) + A(T) F_T(T), \end{aligned} \quad (40)$$

where the function  $A(T)$  is defined as

$$A(T) = \frac{3}{2}T + \frac{(3b - 2c)}{c_0^2(b - c)}t^{-2c}(T), \quad (41)$$

with  $t(T)$  as the Equation (39) solution and  $\Lambda_0 = 2\kappa\phi_0$  as the cosmological constant. From this Equation (39), there are several possible solutions to Equation (40) depending on the values of  $c$ , as in ref. [46]. Each value of  $c$  will lead to specific  $V(T)$  and  $A(T)$  functions, and then we will solve Equation (40) with this form. In general, the Equation (40) solution will be under the form:

$$F(T) = -\Lambda_0 + \exp\left[\int_T \frac{dT'}{A(T')}\right] \left[ F_0 + 2\kappa \int_T dT' \frac{V(T')}{A(T')} \exp\left[-\int_{T'} \frac{dT''}{A(T'')}\right] \right], \quad (42)$$

where  $A(T)$  is Equation (41). Equation (42) is the general formula applicable for any subcases and will be used to find all new teleparallel  $F(T)$  solutions in the current paper.

The subcases are:

1. **c = -2b** ( $b \neq 0$ ): Equation (39) is simplified as [46]:

$$t(T) = \left( \frac{c_0^2}{2}(-T) \right)^{1/4b}. \quad (43)$$

Then, Equation (41) is simplified to  $A(T) = \frac{T}{3}$ . The scalar field will be  $\phi(T) = p_0 \left( \frac{c_0^2}{2}(-T) \right)^{p/4b}$ , and we will use this expression for the following subcases:

- (a) **General:** Equation (28) for scalar field potential is:

$$V(T) = -\frac{V_p}{2}(-T)^{(p-1)/2b}, \quad (44)$$

where  $V_p = \frac{p^2 p_0^2 (p-1-3b)}{2^{(p-1)/2b} (p-1)} c_0^{(p-1)/b}$ . We obtain that  $V_p = 0$  for  $p = 3b + 1$  and  $V(T)$  is undefined for  $p = 1$ . Then, by substituting the simplified  $A(T)$  and Equations (43) and (44) into Equation (42), the teleparallel  $F(T)$  solution will be:

$$F(T) = -\Lambda_0 - \frac{6\kappa b V_p}{(p-1-6b)}(-T)^{(p-1)/2b} + F_0 T^3, \quad (45)$$

where  $F_0$  is an integration constant and  $p \neq 1$  and  $p \neq 1 + 6b$ .

(b)  **$p \gg 1$  case:** By using the same scalar field  $\phi(T)$ , Equation (30) yields as potential:

$$V(T) = -\frac{V_\infty}{2} (-T)^{p/2b}, \quad (46)$$

where  $V_\infty = \frac{p^2 p_0^2 c_0^{p/b}}{2^{p/2b}}$ . Then, by using the same  $A(T)$  and Equations (43) and (46), Equation (42) is:

$$F(T) \approx -\Lambda_0 - \frac{6\kappa b V_\infty}{p} (-T)^{p/2b} + F_0 T^3 \approx -\frac{6\kappa b p p_0^2 c_0^{p/b}}{2^{p/2b}} (-T)^{p/2b}, \quad (47)$$

where  $p \rightarrow \infty$  and  $T \leq 0$  in some situations.

(c)  **$p = 1$  case:** By still using the same scalar field  $\phi(T)$ , Equation (32) becomes:

$$V(T) = \tilde{\phi}_0 - \phi_0 + \frac{3p_0^2}{4} \ln(-T), \quad (48)$$

where  $\tilde{\phi}_0 = \phi_0 + \frac{3p_0^2}{4} \ln\left(\frac{p_0^{4b} c_0^2}{2}\right)$ . Then, Equation (42) becomes, by using the same  $A(T)$  and Equations (43) and (48):

$$F(T) = -\tilde{\Lambda}_0 - \frac{\kappa p_0^2}{2} - \frac{3\kappa p_0^2}{2} \ln(-T) + F_0 T^3, \quad (49)$$

where  $T \leq 0$  and  $\tilde{\Lambda}_0 = 2\kappa \tilde{\phi}_0$  is the modified cosmological constant.

By comparing Equations (45), (47), and (49), we see that the homogeneous parts are very similar and their respective differences are only from the  $V(T)$  parts (particular solution).

2.  **$c = 1$ :** Equation (39) becomes [46]:

$$\begin{aligned} 0 &= \left(2(1+2b) - \frac{2}{c_0^2}\right) t^{-2} - T, \\ \Rightarrow t^{-2}(T) &= \frac{T}{2\left(1+2b - \frac{1}{c_0^2}\right)}. \end{aligned} \quad (50)$$

From Equation (50), we find, as Equation (41):

$$A(T) = \left[ \frac{3}{2} + \frac{(3b-2)}{2c_0^2(b-1)(1+2b-c_0^{-2})} \right] T = C T, \quad (51)$$

where  $C = \frac{3}{2} + \frac{(3b-2)}{2c_0^2(b-1)(1+2b-c_0^{-2})}$ . The scalar field for Equation (50) is:

$$\phi(T) = p_0 2^{p/2} \left(1+2b-c_0^{-2}\right)^{p/2} T^{-p/2}. \quad (52)$$

The potential  $V(T)$  and  $F(T)$  solutions are, for the following subcases:

(a) **General:** Equation (28) becomes:

$$V(T) = -\frac{V_p}{2} T^{1-p}, \quad (53)$$

where  $p \neq 1$  and  $V_p = \frac{p^2 p_0^2 2^{p-1}}{(p-1)} (b+p+1) (1+2b-c_0^{-2})^{p-1}$ . Equation (53) will be constant for  $b = -\frac{1}{2} + \frac{1}{2c_0^2}$  and  $b = -p-1$ . By substituting Equations (51) and (53) into Equation (42), we find, as a solution:

$$F(T) = -\Lambda_0 + \frac{\kappa V_p}{(1+C(p-1))} T^{1-p} + F_0 T^{1/C}, \quad (54)$$

where  $p \neq 1$ .

(b) **Equation (30) potential:** This equation, in terms of Equation (52), is ( $p \gg 1$ ):

$$V(T) = -\frac{V_\infty}{2} T^{-p}, \quad (55)$$

where  $V_\infty = p^2 p_0^2 2^p \left(1+2b-\frac{1}{c_0^2}\right)^p$ . By substituting Equations (51) and (55) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + F_0 T^{1/C} + \frac{\kappa V_\infty}{[1+pC]} T^{-p} \approx -\Lambda_0 + F_0 T^{1/C}. \quad (56)$$

(c) **Equation (32) potential:** This equation, in terms of Equation (52), is ( $p = 1$ ):

$$V(T) = \tilde{\phi}_0 - \phi_0 + \frac{p_0^2}{2} (b+2) \ln(T). \quad (57)$$

where  $b \neq -2$  and  $\tilde{\phi}_0 = \phi_0 - \frac{p_0^2}{2} (b+2) \ln \left[ 2p_0^2 \left(1+2b-\frac{1}{c_0^2}\right) \right]$ . By substituting Equations (51) and (57) into Equation (42), we find as a solution:

$$F(T) = -\tilde{\Lambda}_0 + F_0 T^{1/C} - \kappa p_0^2 (b+2) [\ln(T) + C]. \quad (58)$$

3. **c = -1:** Equation (39) becomes [46]:

$$\begin{aligned} 0 &= t^4 + \frac{c_0^2 T}{2} t^2 - c_0^2 (1-2b), \\ \Rightarrow t^2(T) &= \frac{c_0^2}{4} \left[ -T + \delta_1 \sqrt{T^2 + 16(1-2b)c_0^{-2}} \right], \end{aligned} \quad (59)$$

where  $\delta_1 = \pm 1$ . Equation (41) for Equation (59) is:

$$\begin{aligned} A(T) &= \frac{3}{2} T + \frac{(3b+2)}{4(b+1)} \left[ -T + \delta_1 \sqrt{T^2 + 16(1-2b)c_0^{-2}} \right] \\ &= \left( \frac{3}{2} - C_1 \right) T + \delta_1 C_1 \sqrt{T^2 + C_2}, \end{aligned} \quad (60)$$

where  $C_1 = \frac{(3b+2)}{4(b+1)}$  and  $C_2 = 16(1-2b)c_0^{-2}$ . The scalar field for Equation (59) is:

$$\begin{aligned} \phi(T) &= \frac{p_0 c_0^p}{2^p} \left[ -T + \delta_1 \sqrt{T^2 + 16(1-2b)c_0^{-2}} \right]^{p/2} = \frac{p_0 c_0^p}{2^p} \left[ -T + \delta_1 \sqrt{T^2 + C_2} \right]^{p/2}, \\ &= \frac{p_0 c_0^p}{2^p} [u_\pm(T)]^{p/2}, \end{aligned} \quad (61)$$

where  $u_\pm(T) = -T + \delta_1 \sqrt{T^2 + C_2}$  (+ for  $\delta_1 = +1$  and - for  $\delta_1 = -1$ ). The potential  $V(T)$  and  $F(T)$  solutions are, for the following subcases:



(a) **General:** Equation (28) becomes:

$$V(T) = -\frac{V_p}{2} [u_{\pm}(T)]^{p-1}, \quad (62)$$

where  $p \neq 1$  and  $V_p = \frac{p^2 p_0^2 c_0^{2p-2}}{2^{2p-2}(p-1)} (b + p - 3)$ . Equation (62) will be constant for  $b = 3 - p$ . By substituting Equations (60) and (62) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + [3T + 2C_1 u_{\pm}(T)]^{\frac{2(3-2C_1)}{3(3-4C_1)}} [u_{-}(T)]^{-\frac{4C_1 \delta_1}{3(3-4C_1)}} \left[ F_0 - 2\kappa V_p \right. \\ \left. \times \int_T dT' [u_{\pm}(T')]^{p-1} [3T' + 2C_1 u_{\pm}(T')]^{-\frac{2(3-2C_1)}{3(3-4C_1)}-1} [u_{-}(T')]^{\frac{4C_1 \delta_1}{3(3-4C_1)}} \right]. \quad (63)$$

Equation (63) is difficult to solve under this current form, but there are special case solutions. For example, the  $C_2 = 0$  ( $b = \frac{1}{2}$  and  $C_1 = \frac{7}{12}$ ) and  $\delta_1 = -1$  case where  $u_{-}(T) = -2T$  (and  $u_{+}(T) = 0$  for  $\delta_1 = +1$ ) yields:

$$F(T) = -\Lambda_0 + F_0 T^3 - \frac{3\kappa V_p (-2)^p}{(p-4)} T^{p-1}. \quad (64)$$

The other  $C_2 \neq 0$  possible cases are solved and presented in Appendix A.1.

(b) **Equation (30) potential:** This equation, in terms of Equation (61), is ( $p \gg 1$ ):

$$V(T) = -\frac{V_{\infty}}{2} [u_{\pm}(T)]^p, \quad (65)$$

where  $V_{\infty} = \frac{p^2 p_0^2 c_0^{2p}}{2^{2p}}$ . By substituting Equations (60) and (65) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + [3T + 2C_1 u_{\pm}(T)]^{\frac{2(3-2C_1)}{3(3-4C_1)}} [u_{-}(T)]^{-\frac{4C_1 \delta_1}{3(3-4C_1)}} \left[ F_0 - 2\kappa V_{\infty} \right. \\ \left. \times \int_T dT' [u_{\pm}(T')]^p [3T' + 2C_1 u_{\pm}(T')]^{-\frac{2(3-2C_1)}{3(3-4C_1)}-1} [u_{-}(T')]^{\frac{4C_1 \delta_1}{3(3-4C_1)}} \right]. \quad (66)$$

Once again, Equation (66) does not yield a general solution, but there are specific cases leading to analytical  $F(T)$  solutions. For example, the  $C_2 = 0$  ( $b = \frac{1}{2}$  and  $C_1 = \frac{7}{12}$ ) and  $\delta_1 = -1$  case yields:

$$F(T) = -\Lambda_0 + F_0 T^3 - \frac{3\kappa V_{\infty} (-2)^p}{p} T^p. \quad (67)$$

The other  $C_2 \neq 0$  possible cases are also solved and presented in Appendix A.1.

(c) **Equation (32) potential:** This equation, in terms of Equation (61), is ( $p = 1$ ):

$$V(T) = \tilde{\phi}_0 - \phi_0 - \frac{p_0^2}{2} (b-2) \ln [u_{\pm}(T)], \quad (68)$$

where  $b \neq 2$  and  $\tilde{\phi}_0 = \phi_0 - p_0^2 (b-2) \ln \left[ \frac{p_0 c_0}{2} \right]$ . By substituting Equations (60) and (68) into Equation (42), we find as a solution:

$$F(T) = -\tilde{\Lambda}_0 + [3T + 2C_1 u_{\pm}(T)]^{\frac{2(3-2C_1)}{3(3-4C_1)}} [u_{\pm}(T)]^{-\frac{4C_1 \delta_1}{3(3-4C_1)}} \left[ F_0 - 2\kappa p_0^2 (b-2) \right. \\ \left. \times \int_T dT' \ln[u_{\pm}(T')] [3T + 2C_1 u_{\pm}(T)]^{-\frac{2(3-2C_1)}{3(3-4C_1)}-1} [u_{\pm}(T')]^{\frac{4C_1 \delta_1}{3(3-4C_1)}} \right]. \quad (69)$$

There is no general solution for Equation (69), but only specific solutions. For example, the  $C_2 = 0$  ( $b = \frac{1}{2}$  and  $C_1 = \frac{7}{12}$ ) and  $\delta_1 = -1$  case leads to:

$$F(T) = -\tilde{\Lambda}_0 + F_0 T^{\frac{2}{3-4C_1}} - \frac{\kappa p_0^2 (b-2)}{2} (4C_1 - 3 - 2 \ln(-2T)). \quad (70)$$

All  $C_2 \neq 0$  solvable cases are also presented in Appendix A.1.

4.  $c = 2$ : Equation (39) becomes [46]:

$$0 = t^{-4} - 4c_0^2 (1+b) t^{-2} + \frac{c_0^2 T}{2}, \\ \Rightarrow t^{-2}(T) = 2c_0^2 \left[ (1+b) + \delta_1 \sqrt{(1+b)^2 - \frac{T}{8c_0^2}} \right], \quad (71)$$

where  $\delta_1 = \pm 1$ . Equation (41) for Equation (71) is:

$$A(T) = \frac{3}{2}T + \frac{4(3b-4)(1+b)^2 c_0^2}{(b-2)} \left[ 1 + \delta_1 \sqrt{1 - \frac{T}{8c_0^2(1+b)^2}} \right]^2, \\ = \frac{1}{(b-2)} \left[ -T + C_1 \left( 1 + \delta_1 \sqrt{1 - \frac{T}{C_2}} \right) \right], \\ = \frac{1}{(b-2)} [-T + C_1 w_{\pm}(T)], \quad (72)$$

where  $C_1 = (3b-4)C_2$ ,  $C_2 = 8c_0^2(1+b)^2$ ,  $b \neq 2$ , and  $w_{\pm}(T) = 1 + \delta_1 \sqrt{1 - \frac{T}{C_2}}$  (+ for  $\delta_1 = 1$  and - for  $\delta_1 = -1$ ). The scalar field for Equation (71) is:

$$\phi(T) = \frac{p_0}{2^{p/2} c_0^p (1+b)^{p/2}} [w(T)]^{-p/2}. \quad (73)$$

The potential  $V(T)$  and  $F(T)$  solutions are, for the following subcases:

(a) **General:** Equation (28) becomes:

$$V(T) = -\frac{V_p}{2} [w_{\pm}(T)]^{1-p}, \quad (74)$$

where  $p \neq 1$  and  $V_p = \frac{p^2 p_0^2}{2^{p-1} (p-1) c_0^{2p-2}} (b+p+3)(1+b)^{1-p}$ . Equation (74) will be constant for  $b = -p-3$ . By substituting Equations (72) and (74) into Equation (42), we find as a solution:

$$\begin{aligned}
F(T) = & -\Lambda_0 + [-T + C_1(w_{\pm}(T))]^{(2-b)} [y(T)]^{\frac{(b-2)C_1}{2(C_1-2C_2)}} \\
& \times \left[ \frac{T + (C_1 - 2C_2)(w_{\pm}(T))}{T + (C_1 - 2C_2)(2 - w_{\pm}(T))} \right]^{\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \left[ F_0 C_2^{(2-b)/2} - \kappa V_p (b-2) \right. \\
& \times \int_T dT' [w_{\pm}(T')]^{1-p} [-T' + C_1(w_{\pm}(T'))]^{(b-3)} [y(T')]^{-\frac{(b-2)C_1}{2(C_1-2C_2)}} \\
& \left. \times \left[ \frac{T' + (C_1 - 2C_2)(w_{\pm}(T'))}{T' + (C_1 - 2C_2)(2 - w_{\pm}(T'))} \right]^{-\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \right], \quad (75)
\end{aligned}$$

where  $y(T) = \frac{-C_2 T}{C_1(C_1-2C_2)+C_2 T}$ . There is no general solution for Equation (75), but for the specific cases:

- $C_1 = 0$  and  $\delta_1 = 1$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 + F_0 (-T)^{(2-b)} \\
& - \kappa V_p (2)^{1-p} {}_3F_2 \left( b-2, \frac{p}{2}, \frac{p-1}{2}; p, b-1; \frac{T}{C_2} \right). \quad (76)
\end{aligned}$$

- $C_1 = 0$  and  $\delta_1 = -1$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 + F_0 (-T)^{(2-b)} - \kappa V_p \frac{(b-2)(-2C_2)^{p-1}}{(b-p-1)} (-T)^{1-p} \\
& \times {}_3F_2 \left( b-p-1, \frac{2-p}{2}, \frac{1-p}{2}; b-p, 2-p; \frac{T}{C_2} \right). \quad (77)
\end{aligned}$$

- The  $C_1 = 2C_2$  solutions are presented in Appendix A.2.

(b) **Equation (30) potential:** This equation, in terms of Equation (73), is ( $p \gg 1$ ):

$$V(T) = -\frac{V_{\infty}}{2} [w_{\pm}(T)]^{-p}, \quad (78)$$

where  $V_{\infty} = \frac{p^2 p_0^2 (1+b)^{-p}}{2^p c_0^{2p}}$ . By substituting Equations (71), (72), and (78) into Equation (42), we find as a solution:

$$\begin{aligned}
F(T) = & -\Lambda_0 + [-T + C_1(w_{\pm}(T))]^{(2-b)} [y(T)]^{\frac{(b-2)C_1}{2(C_1-2C_2)}} \\
& \times \left[ \frac{T + (C_1 - 2C_2)(w_{\pm}(T))}{T + (C_1 - 2C_2)(2 - w_{\pm}(T))} \right]^{\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \left[ F_0 C_2^{(2-b)/2} - \kappa V_{\infty} (b-2) \right. \\
& \times \int_T dT' [w_{\pm}(T')]^{-p} [-T' + C_1(w_{\pm}(T'))]^{(b-3)} [y(T')]^{-\frac{(b-2)C_1}{2(C_1-2C_2)}} \\
& \left. \times \left[ \frac{T' + (C_1 - 2C_2)(w_{\pm}(T'))}{T' + (C_1 - 2C_2)(2 - w_{\pm}(T'))} \right]^{-\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \right]. \quad (79)
\end{aligned}$$

There is no general solution for Equation (79). There are solutions for the following cases:

- $C_1 = 0$  and  $\delta_1 = 1$ :

$$F(T) \approx -\Lambda_0 + \left[ F_0 (-T)^{(2-b)} + \frac{\kappa V_\infty}{2^p} {}_3F_2 \left( b-2, \frac{p}{2}, \frac{p}{2}; b-1, p; \frac{T}{C_2} \right) \right]. \quad (80)$$

- $C_1 = 0$  and  $\delta_1 = -1$ :

$$F(T) \approx -\Lambda_0 + \left[ F_0 (-T)^{(2-b)} - \frac{\kappa V_\infty (-2C_2)^p}{p} (b-2) (-T)^{-p} \right. \\ \left. \times {}_3F_2 \left( -p, -\frac{p}{2}, -\frac{p}{2}; -p, p; \frac{T}{C_2} \right) \right]. \quad (81)$$

(c) **Equation (32) potential:** This equation, in terms of Equation (73), is ( $p = 1$ ):

$$V(T) = \tilde{\phi}_0 - \phi_0 + \frac{p_0^2}{2} (b+4) \ln [w_\pm(T)], \quad (82)$$

where  $b \neq -1, -4$  and  $\tilde{\phi}_0 = \phi_0 - p_0^2 (b+4) \ln \left[ \frac{p_0}{2^{1/2} \ell_0 \sqrt{1+b}} \right]$ . By substituting Equations (71), (72), and (82) into Equation (42), we find as a solution:

$$F(T) = -\tilde{\Lambda}_0 + [-T + C_1(w_\pm(T))]^{(2-b)} [y(T)]^{\frac{(b-2)C_1}{2(C_1-2C_2)}} \\ \times \left[ \frac{T + (C_1 - 2C_2)(w_\pm(T))}{T + (C_1 - 2C_2)(2 - w_\pm(T))} \right]^{\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \left[ F_0 C_2^{(2-b)/2} \right. \\ \left. + \frac{\kappa p_0^2 (b+4)(b-2)}{C_1} \int_T dT' \ln [w_\pm(T')] [-T' + C_1(w_\pm(T'))]^{(b-3)} \right. \\ \left. \times [y(T')]^{-\frac{(b-2)C_1}{2(C_1-2C_2)}} \left[ \frac{T' + (C_1 - 2C_2)(w_\pm(T'))}{T' + (C_1 - 2C_2)(2 - w_\pm(T'))} \right]^{-\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \right]. \quad (83)$$

There is no general solution for Equation (83), but only for  $C_1 \neq 0$  cases, as presented in Appendix A.2.

## 5. Late Cosmology $t \rightarrow \infty$ limit:

In this case, there are some subcases according to Equation (39), such as:

(a)  $c > 1$ : The Equation (39)  $t(T)$  relation will be, for the following cases:

$$0 \approx 2c(c+2b)t^{-2} - T, \\ \Rightarrow t^{-2}(T) \approx \frac{T}{2c(c+2b)}. \quad (84)$$

If  $t(T) \rightarrow \infty$ , we find from Equation (84) that  $T \rightarrow 0$ . Then, Equation (41) becomes:

$$A(T) \approx \frac{3}{2}T + \frac{(3b-2c)}{c_0^2(b-c)(2c(c+2b))^c} T^c, \\ \rightarrow \frac{3}{2}T \quad \text{when } T \rightarrow 0. \quad (85)$$

For any  $V(T)$  potential and by inserting Equations (84) and (85) into Equation (42), we find, as the  $F(T)$  solution:

$$F(T) = -\Lambda_0 + T^{\frac{2}{3}} \left[ F_0 + \frac{4\kappa}{3} \int_T dT' V(T') T'^{-\frac{5}{3}} \right]. \quad (86)$$

The Equation (86) solutions for  $\phi(T) = p_0 (2c(c+2b))^{p/2} T^{-p/2}$  are as follows:

- **General:** The  $V(T)$  is exactly as Equation (53), with  $V_p = p^2 p_0^2 \frac{(b+2c+p-1)}{(p-1)} (2c(c+2b))^{p-1}$  as the proportionality constant. Equation (86) will be:

$$- p \neq \frac{1}{3}:$$

$$F(T) = -\Lambda_0 + F_0 T^{\frac{2}{3}} - \frac{4\kappa}{(1-3p)} T^{1-p}. \quad (87)$$

$$- p = \frac{1}{3}:$$

$$F(T) = -\Lambda_0 + T^{\frac{2}{3}} \left[ F_0 - \frac{4\kappa}{3} \ln(T) \right]. \quad (88)$$

- **Equation (30) potential:** The  $V(T)$  is exactly as Equation (55), with  $V_\infty = p^2 p_0^2 (2c(c+2b))^p$  as the proportionality constant. Equation (86) will be:

$$- p \neq -\frac{2}{3}:$$

$$F(T) = -\Lambda_0 + F_0 T^{\frac{2}{3}} + \frac{2\kappa V_p}{(2+3p)} T^{-p}. \quad (89)$$

$$- p = -\frac{2}{3}:$$

$$F(T) = -\Lambda_0 + T^{\frac{2}{3}} \left[ F_0 - \frac{2\kappa V_p}{3} \ln(T) \right]. \quad (90)$$

- **Equation (32) potential:** The  $V(T)$  is exactly  $V(T) = \tilde{\phi}_0 - \phi_0 + \frac{p_0^2}{2}(b+2c) \ln(T)$ , a very similar form to Equation (57). Equation (86) will be ( $p=1$ ):

$$F(T) = -\tilde{\Lambda}_0 + F_0 T^{\frac{2}{3}} - \kappa p_0^2 (b+2c) \left( \ln(T) + \frac{3}{2} \right). \quad (91)$$

- (b)  **$c=1$ :** The  $t(T)$  solution will be described by Equation (50),  $A(T)$  by Equation (51), and  $F(T)$  by Equations (54), (56), and (58) under the  $t \rightarrow \infty$  limit.
- (c)  **$0 < c < 1$  ( $c \neq 0$ ):** Equation (39) leads to:

$$\begin{aligned} 0 &\approx \frac{2}{c_0^2} t^{-2c} + T, \\ \Rightarrow t^{-2c}(T) &\approx \frac{c_0^2}{2} (-T). \end{aligned} \quad (92)$$

If  $t(T) \rightarrow \infty$ , we find that  $T \rightarrow 0$  from Equation (92). Equation (41) becomes:

$$A(T) \approx \frac{c}{2(c-b)} T \quad \text{when } T \rightarrow 0, \quad (93)$$

where  $b \neq c$ . By inserting Equations (92) and (93) into Equation (42), we find as the  $F(T)$  solution for any potential:

$$F(T) = -\Lambda_0 + T^{\frac{2(c-b)}{c}} \left[ F_0 + \frac{4\kappa(c-b)}{c} \int_T dT' V(T') T'^{\frac{2b-3c}{c}} \right]. \quad (94)$$

The Equation (94) solutions for  $\phi(T) = p_0 2^{p/2c} c_0^{-p/c} (-T)^{-p/2c}$  are as follows:

- **General:** Equation (28) becomes:

$$V(T) = -\frac{V_p}{2} (-T)^{\frac{(1-p)}{c}}, \quad (95)$$

where  $V_p = \frac{p^2 p_0^2}{(p-1)} (b+2c+p-1) 2^{\frac{p-1}{c}} c_0^{\frac{2(1-p)}{c}}$ , and Equation (94) will be:

–  $p \neq 1 + 2(b-c)$ :

$$F(T) = -\Lambda_0 + F_0 T^{\frac{2(c-b)}{c}} - \frac{2\kappa(c-b)V_p}{(2b-2c-p+1)} (-T)^{\frac{(1-p)}{c}}. \quad (96)$$

–  $p = 1 + 2(b-c)$ :

$$F(T) = -\Lambda_0 + T^{\frac{2(c-b)}{c}} \left[ F_0 - \frac{2\kappa(c-b)V_p}{c} \ln(T) \right]. \quad (97)$$

- **Equation (30) potential:** This equation, in terms of the  $\phi(T)$ , is:

$$V(T) = -\frac{V_\infty}{2} (-T)^{-p/c}, \quad (98)$$

where  $V_\infty = p^2 p_0^2 2^{p/c} c_0^{-2p/c}$ , and Equation (94) will be:

–  $p \neq 2(b-c)$

$$F(T) = -\Lambda_0 + F_0 T^{\frac{2(c-b)}{c}} - \frac{2\kappa(c-b)V_\infty}{(2b-2c-p)} (-T)^{-\frac{p}{c}}. \quad (99)$$

–  $p = 2(b-c)$

$$F(T) = -\Lambda_0 + T^{\frac{2(c-b)}{c}} \left[ F_0 - \frac{2\kappa(c-b)V_\infty}{c} \ln(T) \right]. \quad (100)$$

- **Equation (32) potential:** This equation, in terms of the  $\phi(T)$ , is ( $p = 1$ ):

$$V(T) = \tilde{\phi}_0 - \phi_0 + \frac{p_0^2}{2c} (b+2c) \ln(-T), \quad (101)$$

where  $\tilde{\phi}_0 = \phi_0 - p_0^2 (b+2c) \ln(p_0 2^{1/2c} c_0^{-1/c})$ , and Equation (94) will be:

$$F(T) = -\tilde{\Lambda}_0 + F_0 T^{\frac{2(c-b)}{c}} - \frac{\kappa p_0^2}{c} (b+2c) \left( \ln(-T) - \frac{c}{2(b-c)} \right). \quad (102)$$

- (d)  **$c < 0$ :** By setting  $c = -|c|$  in Equations (39) and (41), we find Equations (92) and (93) as exactly  $t^{2|c|}(T) \approx \frac{c_0^2}{2} (-T)$  and  $A(T) \approx \frac{|c|}{2(|c|+b)} T$ . We also find that, for



$T \rightarrow -\infty$  and then  $A(T) \rightarrow -\infty$  under the  $t \rightarrow \infty$  limit. In this case, we will find that all  $F(T)$  solutions are under the Equation (94) form, and we will recover the Equations (96)–(102) teleparallel  $F(T)$  solutions by only performing the  $c = -|c|$  transformation.

- (e)  **$c = -2b$ :** We find that  $t(T)$  is Equation (43) and  $A(T) = \frac{T}{3}$ , leading to Equations (45), (47), and (49) as  $F(T)$  solutions under the  $t \rightarrow \infty$  limit. We can summarize the situation as follows:
- $b > 0$ :  $T \rightarrow -\infty$ ,  $A(T) \rightarrow -\infty$  and  $F(T) \rightarrow -\infty$ ,
  - $b < 0$ :  $T \rightarrow 0$ ,  $A(T) \rightarrow 0$  and  $F(T) \rightarrow 0$ .

**6. Early Cosmology  $t \rightarrow 0$  limit:** For this limit, Equation (39) will lead to:

- (a)  **$c > 1$ :**  $t^{-2c}(T)$  and  $A(T)$  are, respectively, Equations (92) and (93), but  $t^{-2c}(T) \rightarrow \infty$  and  $A(T) \rightarrow \infty$  when  $T \rightarrow -\infty$ . We will then recover the same  $F(T)$  solution expressions as Equations (94)–(102), but under the  $T \rightarrow -\infty$  limit.
- (b)  **$0 < c < 1$  and  $c < 0$ :**  $t^{-2}(T)$  and  $A(T)$  are, respectively, Equations (84) and (85), but  $t^{-2}(T) \rightarrow \infty$  and  $A(T) \rightarrow \infty$  when  $T \rightarrow \infty$ . We will then recover the same  $F(T)$  solution expressions as Equations (86)–(91), but under the  $T \rightarrow \infty$  limit.

For other values of  $b$ ,  $c$ , and/or subcases, most Equation (42) integral cases will not lead to a closed and analytic  $F(T)$  solutions, as in ref. [46]. All the previous new teleparallel  $F(T)$  solutions are new results, and some of those are comparable to several  $F(T)$  solutions found in ref. [46] for some perfect fluid sources. Some  $c = 1$  and  $t \rightarrow \infty$   $F(T)$  solutions can also be compared to some TRW  $F(T)$  solutions found in refs. [49,50,64] for perfect fluid and also scalar field sources, a typical case of isotropic cosmological spacetime. This last limit is possible because of additional symmetries (six KVs for TRW spacetime instead of four in a KS spacetime) from the linear isotropy group, as mentioned in Section 1.

### 3.2. Exponential Ansatz Solutions

Equation (21), in terms of Equation (34), leads to the characteristic equation:

$$e^{-2ct(T)} = \frac{c_0^2}{2}(T_0 - T), \quad (103)$$

where  $c \neq 0$  and  $T_0 = 2c(c + 2b)$  ( $c = 0$  subcase leads to GR solutions). By substituting Equation (103) into Equation (25), we find the Equation (40) DE form with

$$A(T) = \frac{c}{2(b-c)} \left[ \left( \frac{3b}{2} - c \right) T_0 - T \right] = \frac{c}{2(b-c)} (T_1 - T), \quad (104)$$

where  $T_1 = c(3b - 2c)(c + 2b)$ , and the solution is also described by Equation (42). The subcases are:

1.  **$c = -2b$ :**  $T_0 = 0$ , and Equation (103) is simplified to:

$$e^{4bt(T)} = \frac{c_0^2}{2}(-T), \quad (105)$$

where  $T \leq 0$ . Then, Equation (104) is simplified to  $A(T) = \frac{T}{3}$ ; the scalar field will be  $\phi(T) = \frac{p_0}{(4b)^p} \left( \ln \left( \frac{c_0^2}{2}(-T) \right) \right)^p$ . The potential  $V(T)$  and  $F(T)$  solutions are, for the following subcases:

(a) **General:** Equation (35) becomes:

$$V(T) = \frac{V_{1p}}{2} \left( \ln \left( \frac{c_0^2}{2} (-T) \right) \right)^{2p-1} - \frac{V_{2p}}{2} \left( \ln \left( \frac{c_0^2}{2} (-T) \right) \right)^{2p-2}, \quad (106)$$

where  $V_{1p} = \frac{3p^2 p_0^2}{2(2p-1)(4b)^{2p-2}}$  and  $V_{2p} = \frac{p^2 p_0^2}{(4b)^{2p-2}}$ . By substituting  $A(T)$  and Equation (106) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + F_0 T^3 + 3\kappa T^3 \left[ V_{1p} \mathcal{N}_{2p-1} \left( -\frac{c_0^2}{2}, T \right) - V_{2p} \mathcal{N}_{2p-2} \left( -\frac{c_0^2}{2}, T \right) \right], \quad (107)$$

where  $F_0$  is an integration constant,  $p \neq \frac{1}{2}$ , and  $\mathcal{N}_k \left( -\frac{c_0^2}{2}, T \right)$  is a new special function class defined as:

$$\mathcal{N}_k(B_1, x) = \int dx \frac{(\ln(B_1 x))^k}{x^4}. \quad (108)$$

Some simple values of Equation (108) are shown in the Table 1.

**Table 1.** Some values of Equation (108)  $\mathcal{N}_k(B_1, x)$  special functions.

$k$	$\mathcal{N}_k(B_1, x)$
0	$-\frac{1}{3x^3}$
1	$-\frac{1}{3x^3} \left[ \ln(Ax) + \frac{1}{3} \right]$
2	$-\frac{1}{3x^3} \left[ \ln^2(Ax) + \frac{2}{3} \ln(Ax) + \frac{2}{9} \right]$
3	$-\frac{1}{3x^3} \left[ \ln^3(Ax) + \ln^2(Ax) + \frac{2}{3} \ln(Ax) + \frac{2}{9} \right]$
4	$-\frac{\ln^4(Ax)}{3x^3} - \frac{4}{9x^3} \left[ \ln^3(Ax) + \ln^2(Ax) + \frac{2}{3} \ln(Ax) + \frac{2}{9} \right]$
5	$-\frac{1}{3x^3} \left[ \ln^5(Ax) + \frac{5}{3} \ln^4(Ax) \right] - \frac{20}{27x^3} \left[ \ln^3(Ax) + \ln^2(Ax) + \frac{2}{3} \ln(Ax) + \frac{2}{9} \right]$
6	$-\frac{1}{3x^3} \left[ \ln^6(Ax) + 2 \ln^5(Ax) + \frac{10}{3} \ln^4(Ax) \right] - \frac{40}{27x^3} \left[ \ln^3(Ax) + \ln^2(Ax) + \frac{2}{3} \ln(Ax) + \frac{2}{9} \right]$

(b) **Equation (30) potential:** This equation, in terms of the Equation (105) scalar field, is ( $p \gg 1$ ):

$$V(T) = -\frac{V_\infty}{2} \left( \ln \left( \frac{c_0^2}{2} (-T) \right) \right)^{2p}, \quad (109)$$

where  $V_\infty = \frac{p^2 p_0^2}{(4b)^{2p}}$ . By substituting  $A(T) = \frac{T}{3}$  and (109) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + F_0 T^3 - 3\kappa V_\infty T^3 \mathcal{N}_{2p} \left( -\frac{c_0^2}{2}, T \right), \quad (110)$$

where  $\mathcal{N}_{2p} \left( -\frac{c_0^2}{2}, T \right)$  is defined by Equation (108).

- (c) **Equation (37) potential:** This equation, in terms of the Equation (105) scalar field, is ( $p = 1$ ):

$$V(T) = \tilde{\phi}_0 - \phi_0 + \frac{3p_0^2}{4} \ln(-T), \quad (111)$$

where  $\tilde{\phi}_0 = \phi_0 + \frac{3p_0^2}{4} \ln\left(\frac{c_0^2}{2}\right)$ . By substituting  $A(T) = \frac{T}{3}$  and (111) into Equation (42), we find as a solution:

$$F(T) = -\tilde{\Lambda}_0 + F_0 T^3 + \frac{9\kappa p_0^2}{2} T^3 \mathcal{N}_1(-1, T), \quad (112)$$

where  $\mathcal{N}_1(-1, T)$  is defined by Equation (108).

2.  $c \neq -2b$  (General case): From Equation (103), we find that the scalar field is:

$$\phi(T) = \frac{p_0}{(-2c)^p} \left[ \ln\left(\frac{c_0^2}{2}(T_0 - T)\right) \right]^p. \quad (113)$$

For potential  $V(T)$  and  $F(T)$  solutions, we find for the following subcases:

- (a) **General:** Equation (35) becomes:

$$V(T) = -\frac{V_{1p}}{2} \left[ \ln\left(-\frac{c_0^2}{2}(T - T_0)\right) \right]^{2p-1} - \frac{V_{2p}}{2} \left[ \ln\left(-\frac{c_0^2}{2}(T - T_0)\right) \right]^{2p-2}, \quad (114)$$

where  $V_{1p} = \frac{2p^2 p_0^2 (b+2c)}{(2p-1)(-2c)^{2p-1}}$  and  $V_{2p} = \frac{p^2 p_0^2}{(-2c)^{2p-2}}$ . By substituting Equations (104) and (114) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + F_0 (T - T_1)^{2(c-b)/c} - \frac{2\kappa(c-b)}{c} (T - T_1)^{2(c-b)/c} \times \left[ V_{1p} \mathcal{Q}_{2p-1}\left(-\frac{c_0^2}{2}, T_0, T_1, T\right) + V_{2p} \mathcal{Q}_{2p-2}\left(-\frac{c_0^2}{2}, T_0, T_1, T\right) \right], \quad (115)$$

where  $\mathcal{Q}_{2p}(B_1, B_2, B_3, x)$  is defined as:

$$\mathcal{Q}_k(B_1, B_2, B_3, x) = \int dx (\ln(B_1(x - B_2)))^k (x - B_3)^{2b/c-3}. \quad (116)$$

Equation (116) is a generalization of the Equation (108) special function class. Some values of Equation (116) are shown in the Table 2.

- (b) **Equation (30) potential:** This equation, in terms of the Equation (113) scalar field, is ( $p \gg 1$ ):

$$V(T) = -\frac{V_\infty}{2} \left[ \ln\left(-\frac{c_0^2}{2}(T - T_0)\right) \right]^{2p}, \quad (117)$$

where  $V_\infty = \frac{p_0^2 p^2}{(-2c)^{2p}}$ . By substituting Equations (104) and (117) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + (T - T_1)^{2(c-b)/c} \left[ F_0 - \frac{2\kappa(c-b)V_\infty}{c} \mathcal{Q}_{2p}\left(-\frac{c_0^2}{2}, T_0, T_1, T\right) \right], \quad (118)$$

where  $\mathcal{Q}_{2p}\left(-\frac{c_0^2}{2}, T_0, T_1, T\right)$  is defined by Equation (116).

**Table 2.** Some values of Equation (116) for  $\mathcal{Q}_k(B_1, B_2, B_3, x)$  special functions.

$k$	$b/c$	$\mathcal{Q}_k(B_1, B_2, B_3, x)$
0	all	$\frac{c}{2(b-c)}(x - B_3)^{2(b-c)/c}$
1	0	$\frac{2}{(B_2 - B_3)^2} \left[ \frac{(B_2 - B_3)}{(x - B_3)} - \ln(B_1(x - B_3)) + \frac{(x - B_2)(B_2 - 2B_3 + x)}{(x - B_3)^2} \ln(B_1(x - B_2)) \right]$
1	1	$dilog\left(\frac{x - B_3}{B_2 - B_3}\right) + \ln(B_1(x - B_2)) \ln\left(\frac{x - B_3}{B_2 - B_3}\right)$
1	$\frac{3}{2}$	$(\ln(B_1(x - B_2)) - 1)(x - B_2)$
1	2	$\frac{(x - B_2)}{2} \left[ (B_2 - 2B_3 + x) \ln(B_1(x - B_2)) - \frac{x}{2} - \frac{3B_2}{2} + 2B_3 \right]$
2	1	$\ln(B_1(x - B_2))^2 \ln\left(\frac{x - B_3}{B_2 - B_3}\right) + 2 \ln(B_1(x - B_2)) \text{polylog}\left(2, \frac{-x + B_2}{B_2 - B_3}\right) - 2 \text{polylog}\left(3, \frac{-x + B_2}{B_2 - B_3}\right)$
2	$\frac{3}{2}$	$\left( \ln(B_1(x - B_2))^2 - 2 \ln(B_1(x - B_2)) + 2 \right) (x - B_2)$
2	2	$\frac{(x - B_2)}{2} \left[ (B_2 - 2B_3 + x) \ln(B_1(x - B_2))^2 + (-x - 3B_2 + 4B_3) \ln(B_1(x - B_2)) \right] + \frac{(x - B_2)}{2} \left[ \frac{x}{2} + \frac{7B_2}{2} - 4B_3 \right]$

- (c) **Equation (37) potential:** This equation, in terms of the Equation (113) scalar field, is ( $p = 1$ ):

$$V(T) = \tilde{\phi}_0 - \phi_0 + \frac{p_0^2(b + 2c)}{2c} \ln(T_0 - T), \quad (119)$$

where  $\tilde{\phi}_0 = \phi_0 + \frac{p_0^2(b + 2c)}{2c} \ln\left(\frac{c_0^2}{2}\right)$ . By substituting Equation (119) into Equation (42), we find as a solution:

$$F(T) = -\tilde{\Lambda}_0 + (T - T_1)^{2(c-b)/c} \left[ F_0 + \frac{2\kappa p_0^2(c - b)(b + 2c)}{c^2} \mathcal{Q}_1(-1, T_0, T_1, T) \right], \quad (120)$$

where  $\mathcal{Q}_1(-1, T_0, T_1, T)$  is defined by Equation (116).

### 3. Late Cosmology $t \rightarrow \infty$ limit:

- (a)  **$c > 0$ :** Equation (103) will lead to  $T = T_0 = \text{constant}$  under this limit, a GR solution similar to a Teleparallel de Sitter (TdS) spacetime ( $c \neq -2b$  general case) [48]. For  $c = -2b$ , we will find that  $T = 0$ , a null torsion scalar spacetime [107]. In the previous situations,  $A(T) = \text{constant}$  according to Equation (104), and we will only find GR solutions.
- (b)  **$c < 0$ :** We will find that  $\exp(2|c|t(T)) \rightarrow \infty$  and then  $T \rightarrow -\infty$  under the same limit with Equation (103):

$$e^{2|c|t(T)} = \frac{c_0^2}{2}(T_0 - T) \approx \frac{c_0^2}{2}(-T) \rightarrow \infty, \quad (121)$$

where  $T_0 = 2|c|(|c| - 2b) \ll |T|$ . Equation (104) becomes:

$$A(T) \approx \frac{|c|}{2(b + |c|)} T \rightarrow -\infty, \quad (122)$$

where  $T_1 = |c| (3b + 2|c|)(|c| - 2b) \ll |T|$ . For any  $V(T)$  potential, Equation (42) will be:

$$F(T) = -\Lambda_0 + T^{\frac{2(b+|c|)}{|c|}} \left[ F_0 + \frac{4\kappa(b+|c|)}{|c|} \int_T dT' V(T') T'^{-\frac{2b+3|c|}{|c|}} \right]. \quad (123)$$

Equation (123) is for  $\phi(T) = \frac{p_0}{(2|c|)^p} \left( \ln \left( \frac{c_0^2}{2} (-T) \right) \right)^p$ :

- **General:** Equation (35) becomes Equation (114), with  $T_1 = T_0 = 0$ ,  $V_{1p} = \frac{2p^2 p_0^2 (b-2|c|)}{(2p-1)(2|c|)^{2p-1}}$  and  $V_{2p} = \frac{p^2 p_0^2}{(2|c|)^{2p-2}}$ . Then, Equation (123) is:

$$F(T) = -\Lambda_0 + T^{\frac{2(b+|c|)}{|c|}} \left[ F_0 - \frac{2\kappa(b+|c|)}{|c|} \left[ V_{1p} \mathcal{Q}_{2p-1} \left( -\frac{c_0^2}{2}, 0, 0, T \right) + V_{2p} \mathcal{Q}_{2p-2} \left( -\frac{c_0^2}{2}, 0, 0, T \right) \right] \right], \quad (124)$$

where  $\mathcal{Q}_{2p}(B_1, 0, 0, x)$  is Equation (116). Equation (124) is also a simplified form of Equation (115).

- **Equation (30) potential:** This equation becomes Equation (117), with  $T_1 = T_0 = 0$  and  $V_\infty = \frac{p^2 p_0^2}{(2|c|)^{2p}}$ . Equation (123) is:

$$F(T) = -\Lambda_0 + T^{\frac{2(b+|c|)}{|c|}} \left[ F_0 - \frac{2\kappa(b+|c|) V_\infty}{|c|} \mathcal{Q}_{2p} \left( -\frac{c_0^2}{2}, 0, 0, T \right) \right], \quad (125)$$

where  $\mathcal{Q}_{2p}(B_1, 0, 0, x)$  is defined by Equation (116). Equation (125) is also a simplified form of Equation (118).

- **Equation (37) potential:** The  $V(T)$  for  $c = -|c|$  will exactly lead to Equation (101), with  $\tilde{\phi}_0 = \phi_0 - \frac{p_0^2 (b-2|c|)}{(2|c|)} \ln \left( \frac{c_0^2}{2} \right)$ , and Equation (123) will be Equation (102) for  $c = -|c|$ . This case proves that some  $F(T)$  solutions will be invariant for any coframe ansatz. In this case, the power-law and exponential ansatzes both yield to the same  $F(T)$  solution.
4. **Early Cosmology  $t \rightarrow 0$  limit:** For any non-zero value of  $c$ , Equation (103) leads to  $T \rightarrow T_0 - \frac{2}{c_0^2} = \text{constant}$  and then  $A(T) = \text{constant}$ , a GR solution. This is a TdS-like case model because of the constant torsion scalar [48].

All the teleparallel  $F(T)$  solutions in this section are new and comparable to those found in ref. [46]. However, there are, in principle, other additional subcases leading to new  $F(T)$  solutions.

#### 4. Exponential Scalar Field Solutions

In this section, we will set an exponential scalar field  $\phi(t) = p_0 \exp(pt)$ , where  $p_0$  is a constant and  $p$  is the exponential coefficient. The conservation law defined by Equation (19) becomes:

$$0 = p \left( \ln(A_2 A_3^2) \right)' \phi + p^2 \phi + \frac{dV}{d\phi}, \quad (126)$$

where  $(\ln(A_2 A_3^2))'$  is still dependent on  $\phi$  and the used ansatz for  $A_2$  and  $A_3$  components, as for Equation (26). Equation (126) is, at first glance, significantly simpler than Equation (26). The possible ansatz are as follows:

1. **Power-law:** By substituting Equation (27) into Equation (126), we find that:

$$0 = p(b+2c) \frac{\phi}{\ln(\phi/p_0)} + p^2 \phi + \frac{dV}{d\phi},$$

$$\Rightarrow V(\phi) = \phi_0 - \frac{p^2}{2} \phi^2 + p(b+2c) p_0^2 \text{Ei}\left(-2 \ln\left(\frac{\phi}{p_0}\right)\right). \quad (127)$$

Equation (127), by the last integral term, is a different kind of potential, and this last can be considered as an additional interaction term. Equation (20) for  $\alpha_Q$  is, by substituting Equation (127):

$$\alpha_Q = -1 + \frac{p^2 \phi^2}{\phi_0 + p(b+2c) p_0^2 \text{Ei}\left(-2 \ln\left(\frac{\phi}{p_0}\right)\right)}. \quad (128)$$

$p \gg 1$ : Equations (127) and (128) become exactly Equations (30) and (31), where the  $p(b+2c) p_0^2 \text{Ei}\left(-2 \ln\left(\frac{\phi}{p_0}\right)\right)$  term is negligible in this case.

2. **Exponential:** This is the most simple and naturally adapted ansatz for a pure exponential scalar field. By substituting Equation (34) into Equation (126), the conservation law becomes:

$$0 = p(b+2c+p) \phi + \frac{dV}{d\phi},$$

$$\Rightarrow V(\phi) = \phi_0 - \frac{p}{2} (b+2c+p) \phi^2, \quad (129)$$

and then Equation (20) will be:

$$\alpha_Q = -1 + \frac{p^2 \phi^2}{\phi_0 - \frac{p}{2} (b+2c) \phi^2}. \quad (130)$$

- $p \gg 1$ : Equations (129) and (130) become Equations (30) and (31), with the same physical process scenarios.
- $p(b+2c+p) < 0$ : Equation (129) is a simple harmonic oscillator (SHO) potential of angular frequency  $\omega_\phi^2 = -p(b+2c+p)$ . In such a case, the scalar field will be:

$$\phi(t) = C_1 \cos(\omega_\phi t) + C_2 \sin(\omega_\phi t). \quad (131)$$

Equation (131) is an oscillating scalar field, but it is not really relevant as a scalar field source for usual cosmological solutions. Beyond that, there are two situations leading to this case:

- $p < 0$  and  $p > -b-2c$ .
- $p > 0$  and  $p < -b-2c$ .

This quadratic  $V(\phi)$  potential case can be easily used for dark energy quintom oscillating models [94–96]. The Equation (131) scalar field definition can only be relevant for this type of physical model.

- $p = 0$  and/or  $p = -b-2c$ : Equation (129) leads to a constant scalar field  $V(\phi) = \phi_0$ .

As for the power-law potentials obtained in Section 3, the Equation (126) solutions are going in the same direction as the first quintessence process studies [69]. This situation is also confirmed separately by the Equations (27) and (34) ansatz approaches in this section.



#### 4.1. Power-Law Ansatz Solutions

By using Equations (39) and (40) with Equation (27) ansatz, we once again obtain the Equation (40) DE form, with  $A(T)$  defined by Equation (41). By using Equation (127) and setting  $V(\phi(T)) = \phi_0 + V(T)$  in the Equation (40) potential, the general solution will be described again by Equation (42). We will find new  $F(T)$  solutions by computing Equation (42) for each  $V(T)$  potential case:

1. **c = -2b**: By using  $A(T) = \frac{T}{3}$  and substituting Equation (43) for the  $t(T)$  solution, the scalar field is  $\phi(T) = p_0 \exp\left(p \left(\frac{c_0^2}{2}(-T)\right)^{1/4b}\right)$ , and Equation (127) for  $V(T)$  potential will be:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp\left(2p \left(\frac{c_0^2}{2}(-T)\right)^{1/4b}\right) - 3pb p_0^2 Ei\left(-2p \left(\frac{c_0^2}{2}(-T)\right)^{1/4b}\right). \quad (132)$$

By substituting  $A(T)$  and Equation (132) into Equation (42), we find as a solution:

$$\begin{aligned} F(T) = & -\Lambda_0 + F_0 T^3 - \frac{\kappa p^2 p_0^2}{(12b-1)} \left[ \left(2p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right)^{6b} \exp\left(-p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right) \right. \\ & \times WhittakerM\left(-6b, -6b + \frac{1}{2}, 2p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right) + (1-12b) \exp\left(-2p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right) \Big] \\ & - \frac{6\kappa p p_0^2}{\left(b - \frac{1}{12}\right)} \left[ 2pb^2 \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}} {}_3F_3\left(1, 1, 1-12b; 2, 2, -12b+2; -2p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right) \right. \\ & \left. - \left(b - \frac{1}{12}\right) \left(b \ln\left(2p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right) + \gamma b + \frac{\ln(T)}{4} + \frac{1}{12}\right) \right], \end{aligned} \quad (133)$$

where  $WhittakerM(B_1, B_2, x) = \exp(-\frac{x}{2}) x^{\frac{1}{2}+B_2} {}_1F_1\left(\frac{1}{2} + B_2 - B_1; 1 + 2B_2; x\right)$  is a special function.

**p ≫ 1**: By using  $A(T) = \frac{T}{3}$  and substituting Equation (43) into the Equation (30) potential,  $V(T)$  becomes:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp\left(2p \left(\frac{c_0^2}{2}(-T)\right)^{1/4b}\right). \quad (134)$$

By substituting Equation (134) into Equation (42), we find as a solution:

$$\begin{aligned} F(T) \approx & -\Lambda_0 + F_0 T^3 - \frac{\kappa p^2 p_0^2}{(12b-1)} \left[ \left(2p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right)^{6b} \exp\left(-p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right) \right. \\ & \left. \times WhittakerM\left(-6b, -6b + \frac{1}{2}, 2p \left(-\frac{c_0^2}{2}T\right)^{\frac{1}{4b}}\right) \right]. \end{aligned} \quad (135)$$

2.  $c = 1$ : By using Equation (51) and substituting Equation (50), we find that:

$$\phi(T) = p_0 \exp \left( \frac{\sqrt{2}p \left( 1 + 2b - \frac{1}{c_0^2} \right)^{1/2}}{\sqrt{T}} \right), \quad (136)$$

and the Equation (127) potential becomes:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp \left( \frac{T_2}{\sqrt{T}} \right) + p(b+2) p_0^2 Ei \left( -\frac{T_2}{\sqrt{T}} \right), \quad (137)$$

where  $T_2 = 2\sqrt{2}p \left( 1 + 2b - \frac{1}{c_0^2} \right)^{1/2}$ . By substituting Equation (137) into Equation (42), we find that:

$$F(T) = -\Lambda_0 + F_0 T^{1/C} - \frac{\kappa p_0^2}{C} \left[ p^2 T^{1/C} \int_T dT' T'^{-1/C-1} \exp \left( \frac{T_2}{\sqrt{T'}} \right) - 2p(b+2) T^{1/C} \int_T dT' T'^{-1/C-1} Ei \left( -\frac{T_2}{\sqrt{T'}} \right) \right]. \quad (138)$$

Equation (138) solutions are possible only for non-zero specific values of  $C$ , such as, for example,  $C = 1$ :

$$F(T) = -\Lambda_0 + F_0 T - \kappa p_0^2 \left[ \frac{2p^2}{T_2^2} \sqrt{T} (\sqrt{T} - T_2) \exp \left( \frac{T_2}{\sqrt{T}} \right) + \frac{2p(b+2)}{T_2^2} \left[ (\sqrt{T}T_2 + T) \exp \left( -\frac{T_2}{\sqrt{T}} \right) + Ei \left( -\frac{T_2}{\sqrt{T}} \right) T_2^2 \right] \right]. \quad (139)$$

There are several possible values of  $C$  yielding to analytical  $F(T)$  solutions, as presented in Appendix B.1.

$p \gg 1$ : By using Equation (51) and substituting Equation (50), we find as the Equation (30) potential:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp \left( \frac{T_2}{\sqrt{T}} \right). \quad (140)$$

By substituting Equation (140) into Equation (42), we find that:

$$F(T) = -\Lambda_0 + T^{1/C} \left[ F_0 - \frac{\kappa p^2 p_0^2}{C} \int_T dT' T'^{-1/C-1} \exp \left( \frac{T_2}{\sqrt{T'}} \right) \right]. \quad (141)$$

As for Equation (138), there is no general solution, but we can solve some cases for  $T_2 \gg 1$ , such as the  $C = 1$  case:

$$F(T) = -\Lambda_0 + F_0 T - 2\kappa p^2 p_0^2 \frac{T}{T_2^2} \left( 1 - \frac{T_2}{\sqrt{T}} \right) \exp \left( \frac{T_2}{\sqrt{T}} \right). \quad (142)$$

There are several other possible  $F(T)$  solutions arising from Equation (141), as presented in Appendix B.1.

3.  $\mathbf{c} = -1$ : By using Equation (60) and substituting Equation (59), we find that:

$$\phi(T) = p_0 \exp\left(\frac{c_0 p}{2} [u_{\pm}(T)]^{1/2}\right), \quad (143)$$

and the Equation (127) potential becomes:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp\left(c_0 p [u_{\pm}(T)]^{1/2}\right) + p(b-2) p_0^2 \operatorname{Ei}\left(-c_0 p [u_{\pm}(T)]^{1/2}\right). \quad (144)$$

By substituting Equation (144) into Equation (42), we find that:

$$\begin{aligned} F(T) = & -\Lambda_0 + [3T + 2C_1 u_{\pm}(T)]^{\frac{2(3-2C_1)}{3(3-4C_1)}} [u_{-}(T)]^{-\frac{4C_1 \delta_1}{3(3-4C_1)}} \left[ F_0 - 2\kappa p_0^2 \right. \\ & \times \int_T dT' \left[ p^2 \exp\left(c_0 p [u_{\pm}(T')]\right)^{1/2} - 2p(b-2) \operatorname{Ei}\left(-c_0 p [u_{\pm}(T')]\right)^{1/2} \right] \\ & \times \left. [3T' + 2C_1 u_{\pm}(T')]^{-\frac{2(3-2C_1)}{3(3-4C_1)}-1} [u_{-}(T')]^{\frac{4C_1 \delta_1}{3(3-4C_1)}} \right]. \end{aligned} \quad (145)$$

As in Section 3.1, there is no general solution for Equation (145). However, there are specific cases yielding to analytical solutions:

- $C_2 = 0$  and  $\delta_1 = -1$ :

$$\begin{aligned} F(T) = & -\Lambda_0 + F_0 T^{\frac{2}{3-4C_1}} \\ & + 2\kappa p_0^2 \left[ \frac{p^2}{(4C_1-3)} T^{\frac{2}{3-4C_1}} \mathcal{R}\left(p, C_1, -2c_0^2 T\right) + p(b-2) \right. \\ & \times \left( \frac{4c_0 p}{(4C_1+1)} \sqrt{-2T} {}_3F_3\left(1, 1, \frac{4C_1+1}{4C_1-3}; 2, 2, \frac{8C_1-2}{4C_1-3}; -c_0 p \sqrt{-2T}\right) \right. \\ & \left. \left. - \left( \Psi\left(\frac{4}{4C_1-3}\right) + \gamma - \Psi\left(\frac{1+4C_1}{4C_1-3}\right) + \ln(c_0 p \sqrt{-2T}) \right) \right] \right], \end{aligned} \quad (146)$$

where  $\Psi(x)$  is the Digamma function and  $\mathcal{R}(p, C_1, x)$  is defined by:

$$\mathcal{R}(p, C_1, x) = \int_0^x dx' x'^{-\frac{2}{3-4C_1}-1} \exp\left(p x'^{1/2}\right), \quad (147)$$

where  $p \neq 0$  and  $C_1 \neq \frac{3}{4}$ . Some simple values of Equation (147) are shown in the Table 3.

- Other values of  $C_1$  and/or  $C_2$ : No analytical and/or closed form of the  $F(T)$  solution.

$\mathbf{p} \gg 1$  : By using Equation (60) and substituting Equation (59), we find as the Equation (30) potential:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp\left(c_0 p [u_{\pm}(T)]^{1/2}\right). \quad (148)$$

By substituting Equation (148) into Equation (42), we find that:

$$\begin{aligned} F(T) = & -\Lambda_0 + [3T + 2C_1 u_{\pm}(T)]^{\frac{2(3-2C_1)}{3(3-4C_1)}} [u_{-}(T)]^{-\frac{4C_1 \delta_1}{3(3-4C_1)}} \left[ F_0 - 2\kappa p^2 p_0^2 \right. \\ & \times \left. \int_T dT' \exp\left(c_0 p [u_{\pm}(T')]\right)^{1/2} [3T' + 2C_1 u_{\pm}(T')]^{-\frac{2(3-2C_1)}{3(3-4C_1)}-1} [u_{-}(T')]^{\frac{4C_1 \delta_1}{3(3-4C_1)}} \right]. \end{aligned} \quad (149)$$

Once again, there is no general solution for Equation (149), except for  $C_2 = 0$  and  $\delta_1 = -1$ :

$$F(T) \approx -\Lambda_0 + T^{\frac{2}{(3-4C_1)}} \left[ F_0 + \frac{2\kappa p^2 p_0^2}{(4C_1 - 3)} \mathcal{R}(p, C_1, -2c_0^2 T) \right], \quad (150)$$

where  $\mathcal{R}(p, C_1, -2c_0^2 T)$  is described by Equation (147).

**Table 3.** Some values of Equation (147) for the  $\mathcal{R}(p, C_1, x)$  special function.

$C_1$	$\mathcal{R}(p, C_1, x)$
0	$\frac{9 \left( e^{p\sqrt{x}} (-p\sqrt{x} - \frac{1}{3}) (-p\sqrt{x})^{\frac{2}{3}} + p^2 x (\Gamma(\frac{2}{3}) - \Gamma(\frac{2}{3}, -p\sqrt{x})) \right)}{2x^{\frac{2}{3}} (-p\sqrt{x})^{\frac{2}{3}}}$
1	$\frac{2e^{p\sqrt{x}} \left( x^{\frac{3}{2}} p^3 - 3p^2 x + 6p\sqrt{x} - 6 \right)}{p^4}$
-1	$\frac{7 \left( -e^{p\sqrt{x}} (-p\sqrt{x})^{\frac{3}{2}} + \sqrt{x} p (\Gamma(\frac{3}{2}) - \Gamma(\frac{3}{2}, -p\sqrt{x})) \right)}{2x^{\frac{2}{2}} (-p\sqrt{x})^{\frac{3}{2}}}$
$\frac{5}{4}$	$\frac{2e^{p\sqrt{x}} (p\sqrt{x} - 1)}{p^2}$
2	$\frac{2x^{\frac{2}{5}} (\Gamma(\frac{4}{5}) - \Gamma(\frac{4}{5}, -p\sqrt{x}))}{(-p\sqrt{x})^{\frac{4}{5}}}$
-2	$\frac{11 \left( -e^{p\sqrt{x}} (-p\sqrt{x})^{\frac{7}{11}} + p\sqrt{x} (\Gamma(\frac{7}{11}) - \Gamma(\frac{7}{11}, -p\sqrt{x})) \right)}{2x^{\frac{2}{11}} (-p\sqrt{x})^{\frac{7}{11}}}$

4.  $c = 2$ : By using Equation (72) and substituting Equation (71), we find that:

$$\phi(T) = p_0 \exp \left( \frac{p(1+b)}{\sqrt{2c_0} [w_{\pm}(T)]^{1/2}} \right), \quad (151)$$

and the Equation (127) potential becomes:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp \left( \frac{2p(1+b)}{\sqrt{2c_0} [w_{\pm}(T)]^{1/2}} \right) + p(b+4) p_0^2 Ei \left( -\frac{2p(1+b)}{\sqrt{2c_0} [w_{\pm}(T)]^{1/2}} \right). \quad (152)$$

By substituting Equation (152) into Equation (42), we find that:

$$\begin{aligned} F(T) = & -\Lambda_0 + [-T + C_1(w_{\pm}(T))]^{(2-b)} [y(T)]^{\frac{(b-2)C_1}{2(C_1-2C_2)}} \left[ \frac{T + (C_1 - 2C_2)(w_{\pm}(T))}{T + (C_1 - 2C_2)(2 - w_{\pm}(T))} \right]^{\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \\ & \times \left[ F_0 + \kappa(b-2) p_0^2 \int_T dT' \left[ -p^2 \exp \left( \frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T')]^{1/2}} \right) + 2p(b+4) \right. \right. \\ & \times Ei \left( -\frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T')]^{1/2}} \right) \left. \left. [-T' + C_1(w_{\pm}(T'))]^{(b-3)} [y(T')]^{-\frac{(b-2)C_1}{2(C_1-2C_2)}} \right. \right. \\ & \times \left. \left. \left[ \frac{T' + (C_1 - 2C_2)(w_{\pm}(T'))}{T' + (C_1 - 2C_2)(2 - w_{\pm}(T'))} \right]^{-\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \right] \right]. \end{aligned} \quad (153)$$

There is no general  $F(T)$  solution for Equation (153). However, there are analytical  $F(T)$  solutions for several subcases, as presented in Appendix B.2.

$p \gg 1$  : By using Equation (72) and substituting Equation (71), we find as the Equation (30) potential:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp\left(\frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T)]^{1/2}}\right). \quad (154)$$

By substituting Equation (154) into Equation (42), we find that:

$$\begin{aligned} F(T) = & -\Lambda_0 + [-T + C_1(w_{\pm}(T))]^{(2-b)} [y(T)]^{\frac{(b-2)C_1}{2(C_1-2C_2)}} \left[ \frac{T + (C_1 - 2C_2)(w_{\pm}(T))}{T + (C_1 - 2C_2)(2 - w_{\pm}(T))} \right]^{\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \\ & \times \left[ F_0 - \kappa p^2 p_0^2 \int_T dT' \exp\left(\frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T')]^{1/2}}\right) [-T' + C_1(w_{\pm}(T'))]^{(b-3)} [y(T')]^{-\frac{(b-2)C_1}{2(C_1-2C_2)}} \right. \\ & \times \left. \left[ \frac{T' + (C_1 - 2C_2)(w_{\pm}(T'))}{T' + (C_1 - 2C_2)(2 - w_{\pm}(T'))} \right]^{-\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \right]. \end{aligned} \quad (155)$$

There is no general  $F(T)$  solution for Equation (155). However, there are analytical  $F(T)$  solution for several subcases, as presented in Appendix B.2.

5. **Late Cosmology  $t \rightarrow \infty$  limit:** We have the same type of scenarios as in Section 3.1, and we can find some  $F(T)$  solutions:

(a)  $c > 1$ : By using Equations (84)–(86), we find for the following potentials with  $\phi(T) = p_0 \exp(p(2c(c+2b))^{1/2} T^{-1/2})$ :

• Equation (127) potential:

$$\begin{aligned} V(T) = & -\frac{p^2 p_0^2}{2} \left[ \exp\left(2p(2c(c+2b))^{1/2} T^{-1/2}\right) \right. \\ & \left. - \frac{2(b+2c)}{p} Ei\left(-2p(2c(c+2b))^{1/2} T^{-1/2}\right) \right], \end{aligned} \quad (156)$$

and then Equation (86):

$$\begin{aligned} F(T) = & -\Lambda_0 + T^{\frac{2}{3}} \left[ F_0 - \frac{2\kappa p^2 p_0^2}{3} \left[ \frac{\sqrt{2}}{2(-p\sqrt{c(c+2b)})^{\frac{4}{3}}} \right. \right. \\ & \times \left[ e^{\frac{2p\sqrt{2}\sqrt{c(c+2b)}}{\sqrt{T}}} \left( -\frac{p\sqrt{c(c+2b)}}{\sqrt{T}} \right)^{\frac{1}{3}} + \frac{\Gamma\left(\frac{1}{3}, -\frac{2p\sqrt{2}\sqrt{c(c+2b)}}{\sqrt{T}}\right)}{3\sqrt{2}} \right] \\ & + \frac{(b+2c)}{4p(p\sqrt{c(c+2b)})^{\frac{4}{3}}} \left[ \Gamma\left(\frac{1}{3}, \frac{2p\sqrt{2}\sqrt{c(c+2b)}}{\sqrt{T}}\right) \right. \\ & \left. \left. - 12 Ei_1\left(\frac{2p\sqrt{2}\sqrt{c(c+2b)}}{\sqrt{T}}\right) \left(p\sqrt{c(c+2b)}\right)^{\frac{4}{3}} T^{-2/3} \right. \right. \\ & \left. \left. + 3\sqrt{2} e^{-\frac{2p\sqrt{2}\sqrt{c(c+2b)}}{\sqrt{T}}} \left( \frac{p\sqrt{c(c+2b)}}{\sqrt{T}} \right)^{\frac{1}{3}} \right] \right] \right]. \end{aligned} \quad (157)$$

- Equation (30) potential:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp\left(2p (2c(c+2b))^{1/2} T^{-1/2}\right), \quad (158)$$

and then Equation (86):

$$F(T) = -\Lambda_0 + T^{\frac{2}{3}} \left[ F_0 - \frac{\sqrt{2}\kappa p^2 p_0^2}{3(-p\sqrt{c(c+2b)})^{\frac{4}{3}}} \right. \\ \left. \times \left[ e^{\frac{2p\sqrt{2}\sqrt{c(c+2b)}}{\sqrt{T}}} \left( -\frac{p\sqrt{c(c+2b)}}{\sqrt{T}} \right)^{\frac{1}{3}} + \frac{\Gamma\left(\frac{1}{3}, -\frac{2p\sqrt{2}\sqrt{c(c+2b)}}{\sqrt{T}}\right)}{3\sqrt{2}} \right] \right]. \quad (159)$$

- (b)  $0 < c < 1$  and  $c < 0$ : By using Equations (92)–(94), we find for the following potentials with  $\phi(T) = p_0 \exp\left(p \left(\frac{2}{c_0^2}\right)^{1/2c} (-T)^{-1/2c}\right)$  the  $F(T)$  solutions:

- Equation (127) potential:

$$V(T) = -\frac{p^2 p_0^2}{2} \left[ \exp\left(2p \left(\frac{2}{c_0^2}\right)^{1/2c} (-T)^{-1/2c}\right) \right. \\ \left. - \frac{2(b+2c)}{p} Ei\left(-2p \left(\frac{2}{c_0^2}\right)^{1/2c} (-T)^{-1/2c}\right) \right], \quad (160)$$

and then Equation (94):

$$F(T) = -\Lambda_0 + T^{\frac{2(c-b)}{c}} \left[ F_0 - 2\kappa p^2 p_0^2 (c-b) \left[ 2 \left( p^{2c} \frac{2^{2c+1}}{c_0^2} \right)^{2(b-c)/c} \right. \right. \\ \times \left( -\Gamma(4(c-b)) + \Gamma\left(4(c-b), -2p \left(\frac{2}{c_0^2}\right)^{\frac{1}{2c}} (-T)^{-\frac{1}{2c}}\right) \right) \\ \left. - \frac{2(b+2c)}{p} T^{\frac{2(b-c)}{c}} \left[ \frac{1}{8(b-c)^2} + \frac{\gamma}{2(b-c)} - \frac{\ln(T)}{4c(b-c)} + \frac{\ln(2)}{2(b-c)} \right. \right. \\ \left. \left. + \frac{\ln\left(p \left(\frac{2}{c_0^2}\right)^{\frac{1}{2c}} (-1)^{-\frac{1}{2c}}\right)}{2(b-c)} + \frac{4p}{1+4(c-b)} (-T)^{-\frac{1}{2c}} \left(\frac{2}{c_0^2}\right)^{\frac{1}{2c}} \right. \right. \\ \left. \left. \times {}_3F_3\left(1, 1, 1+4(c-b); 2, 2, 2+4(c-b); -2p \left(\frac{2}{c_0^2}\right)^{\frac{1}{2c}} (-T)^{-\frac{1}{2c}}\right) \right] \right] \right]. \quad (161)$$

- Equation (30) potential:

$$V(T) = -\frac{p^2 p_0^2}{2} \exp\left(2p \left(\frac{2}{c_0^2}\right)^{1/2c} (-T)^{-1/2c}\right), \quad (162)$$

and then Equation (94):



$$F(T) = -\Lambda_0 + T^{\frac{2(c-b)}{c}} \left[ F_0 - 4\kappa p^2 p_0^2 (c-b) \left( p^{2c} \frac{2^{2c+1}}{c_0^2} \right)^{2(b-c)/c} \right. \\ \left. \times \left( -\Gamma(4(c-b)) + \Gamma\left(4(c-b), -2p \left( \frac{2}{c_0^2} \right)^{\frac{1}{2c}} (-T)^{-\frac{1}{2c}} \right) \right) \right]. \quad (163)$$

#### 6. Early Cosmology $t \rightarrow 0$ limit:

- (a)  $c > 1$ :  $t^{-2c}(T)$  and  $A(T)$  are, respectively, Equations (92) and (93), but satisfying the  $T \rightarrow -\infty$  limit in the Section 3.1 case. We will then recover Equations (161) and (163) under the same limit.
- (b)  $0 < c < 1$  and  $c < 0$ :  $t^{-2}(T)$  and  $A(T)$  are, respectively, Equations (84) and (85), but satisfying the  $T \rightarrow \infty$  limit in the Section 3.1 case. We will recover Equations (157) and (159) under the same limit.

There are several other cases yielding to new analytical  $F(T)$  solutions. All the previous teleparallel  $F(T)$  solutions are new and comparable to those found in ref [46].

#### 4.2. Exponential Ansatz Solutions

By using Equations (103) and (104) with the Equation (34) ansatz, we substitute Equation (129) into Equation (40) for again finding the  $F(T)$  solution defined by the Equation (42) formula. We will compute the Equation (42) results in the following potential cases:

1.  $c = -2b$ : By using  $A(T) = \frac{T}{3}$  and Equation (105), we find that  $\phi(T) = p_0 \left( \frac{c_0^2}{2} \right)^{p/4b} (-T)^{p/4b}$ , and the Equation (129)  $V(T)$  potential is:

$$V(T) = -\frac{p(p-3b)p_0^2}{2} \left( \frac{c_0^2}{2} \right)^{p/2b} (-T)^{p/2b}, \quad (164)$$

where  $p \neq 3b$ . By substituting Equation (164) into Equation (42), the  $F(T)$  solution is:

$$F(T) = -\Lambda_0 + F_0 T^3 - \frac{6\kappa p b (p-3b) p_0^2}{(p-6b)} \left( \frac{c_0^2}{2} \right)^{p/2b} (-T)^{p/2b}. \quad (165)$$

$p \gg 1$ : Under this limit, Equation (30) becomes:

$$V(T) = -\frac{p^2 p_0^2}{2} \left( \frac{c_0^2}{2} \right)^{p/2b} (-T)^{p/2b}. \quad (166)$$

By substituting Equation (166) into Equation (42), we find as a solution:

$$F(T) \approx -6\kappa p b p_0^2 \left( \frac{c_0^2}{2} \right)^{p/2b} (-T)^{p/2b}. \quad (167)$$

2.  $c \neq -2b$  (General case): From Equations (103) and (104), we find that:

$$\phi(T) = p_0 \left( \frac{c_0^2}{2} \right)^{-p/2c} (T_0 - T)^{-p/2c}, \quad (168)$$

and the Equation (129) potential becomes:

$$V(T) = -\frac{p(b+2c+p)p_0^2}{2} \left(\frac{2}{c_0^2}\right)^{p/c} (T_0 - T)^{-p/c}, \quad (169)$$

where  $p \neq -b - 2c$ . By substituting Equation (169) into Equation (42), the  $F(T)$  solution is:

$$F(T) = -\Lambda_0 + (T - T_1)^{2(c-b)/c} \left[ F_0 + \frac{2\kappa p p_0^2 (b-c)(b+2c+p)}{c} \left(-\frac{2}{c_0^2}\right)^{p/c} \right. \\ \left. \times \int_T dT' (T' - T_0)^{-p/c} (T' - T_1)^{(2b-3c)/c} \right]. \quad (170)$$

There is no general solution to Equation (170), but there are  $F(T)$  solutions for the following cases:

- $T_1 = T_0$ :

$$F(T) = -\Lambda_0 + F_0 (T - T_0)^{2-2b/c} + 2\kappa p p_0^2 \frac{(b-c)(b+2c+p)}{(2b-p-2c)} \left(\frac{-2}{c_0^2(T - T_0)}\right)^{p/c}. \quad (171)$$

- $T_1 \neq T_0$ : There are several possible  $F(T)$  solutions only when  $b \geq \frac{3c}{2}$ , as presented in Appendix B.3.

$p \gg 1$ : Under this limit, Equation (30) becomes:

$$V(T) = -\frac{p^2 p_0^2}{2} \left(\frac{2}{c_0^2}\right)^{p/c} (T_0 - T)^{-p/c}. \quad (172)$$

By substituting Equation (169) into Equation (42), we find as a solution:

$$F(T) = -\Lambda_0 + (T - T_1)^{2(c-b)/c} \left[ F_0 - \frac{2\kappa p^2 p_0^2 (c-b)}{c} \left(-\frac{2}{c_0^2}\right)^{p/c} \right. \\ \left. \times \int_T dT' (T' - T_0)^{-p/c} (T' - T_1)^{(2b-3c)/c} \right]. \quad (173)$$

There is no general solution to Equation (173), but there are  $F(T)$  solutions for the following cases:

- $T_1 = T_0$ :

$$F(T) \approx -\Lambda_0 + F_0 (T - T_0)^{2-2b/c} - 2\kappa p p_0^2 (b-c) \left(-\frac{2}{c_0^2}\right)^{p/c} (T - T_0)^{-p/c}. \quad (174)$$

For  $c > 0$  and  $p \rightarrow \infty$ , Equation (174) becomes:

$$F(T) \rightarrow -\Lambda_0 + F_0 (T - T_0)^{2-2b/c}. \quad (175)$$

For  $c < 0$  and  $p \rightarrow \infty$ , Equation (174) becomes:

$$F(T) \rightarrow -2\kappa p p_0^2 (b + |c|) \left(-\frac{c_0^2}{2}\right)^{p/|c|} (T - T_0)^{p/|c|} \rightarrow \infty. \quad (176)$$

- $T_1 \neq T_0$  and  $b \geq \frac{3c}{2}$ : There are several possible  $F(T)$  solutions, as presented in Appendix B.3.
- $T_1 \neq T_0$ ,  $c > 0$  and  $p \rightarrow \infty$  limit: We find that all cases go to the Equation (175) limit. For the  $c < 0$  and  $p \rightarrow \infty$  limit, we obtain that:

$$F(T) \sim (T - T_0)^{\frac{p}{|c|}} (T - T_1)^{-1}. \quad (177)$$

### 3. Late Cosmology $t \rightarrow \infty$ limit:

- $c > 0$ :** Equation (103) leads to  $T = T_0 = \text{constant}$  when  $c \neq -2b$  (TdS-like spacetime),  $T = 0$  (null torsion scalar spacetime) when  $c = -2b$ , and then  $A(T) = \text{constant}$  under this limit, GR solutions [48], as in Section 3.2.
- $c < 0$ :**  $t(T)$  and  $A(T)$  satisfy Equations (121) and (122), and the  $F(T)$  solutions are described by Equation (123) because this is the same situation as in Section 3.2.

By using  $\phi(T) = p_0 \left( \frac{c_0^2}{2} \right)^{\frac{p}{2|c|}} (-T)^{\frac{p}{2|c|}}$ , where  $c = -|c|$ , we find the  $F(T)$  solutions for the following cases:

- **General:** The Equation (129) potential is:

$$V(T) = -\frac{V_p}{2} (-T)^{\frac{p}{|c|}}, \quad (178)$$

where  $V_p = p p_0^2 (b + 2c + p) \left( \frac{c_0^2}{2} \right)^{\frac{p}{|c|}}$ , and then Equation (123) will be:

$$-p \neq 2(b + |c|)$$

$$F(T) = -\Lambda_0 + F_0 T^{\frac{2(b+|c|)}{|c|}} + \frac{2\kappa (b + |c|) V_p}{(2b + 2|c| - p)} (-T)^{\frac{p}{|c|}}. \quad (179)$$

$$-p = 2(b + |c|)$$

$$F(T) = -\Lambda_0 + T^{\frac{2(b+|c|)}{|c|}} \left[ F_0 - \frac{2\kappa (b + |c|) V_p}{|c|} \ln(T) \right]. \quad (180)$$

- **Equation (30) potential:** The  $V(T)$  expression is under the same form as Equation (178), but we replace  $V_p$  with  $V_\infty = p^2 p_0^2 \left( \frac{c_0^2}{2} \right)^{\frac{p}{|c|}}$ . Equation (123) will be under the same forms as Equations (179) and (180), but we replace  $V_p$  with  $V_\infty$  inside these equations. For  $p \rightarrow \infty$ , Equation (179) will become  $F(T) \sim (-T)^{\frac{p}{|c|}}$ , a pure teleparallel cosmological solution.

- Early Cosmology  $t \rightarrow 0$  limit:** For any non-zero value of  $c$ , Equation (103) also leads to  $T \rightarrow T_0 - \frac{2}{c_0^2} = \text{constant}$  and  $A(T) = \text{constant}$ , a GR solution (TdS-like spacetime), as in Section 3.2 [48,107].

All the previous teleparallel  $F(T)$  solutions are new, and most of those are comparable with the solutions found in refs. [46,49,50,64], including the late/early cosmological limit cases. However, there are other possible subcases leading to additional teleparallel  $F(T)$  solutions.

## 5. Other Scalar Field Source Solutions

### 5.1. General Methods of Teleparallel Field Equations Solving

In Sections 3 and 4, we used the power-law and exponential scalar field  $\phi(t)$  sources for finding the main classes of teleparallel  $F(T)$  solutions. These two types of scalar field sources are the most simple and usual in the literature and allow a large number of new analytical  $F(T)$  solutions [50,53,56,57,64]. But there are, in principle, several other possible scalar field source cases, such as logarithmic, oscillating (see Equation (131)), polynomial (superposition of power-law terms), and several others. As previously, we use an appropriate coframe ansatz; we solve for  $t(T)$  by using the torsion scalar expression defined by Equation (21) and then for  $A(T)$ . Then, we will find the scalar field  $\phi(T)$ , the  $V(\phi)$  potential from Equation (19), and the  $V(T)$  expression from the coframe ansatz. Finally, after being sure that we have an Equation (40) DE form, we will compute the  $F(T)$  solutions by using and substituting the previous elements into Equation (42). This is the same process used in Sections 3 and 4 for computing the huge number of new teleparallel  $F(T)$  solutions. The teleparallel  $F(T)$  solution finding approach (and/or the extensions) is not a wall-to-wall and/or an absolute method, and there are some other approaches in the literature. The main advantage of the current approach is that we can control the used scalar field sources and then find the relevant and exact  $F(T)$  solutions for a specific type of source, including the homogeneous part of solutions (fundamental FE  $F(T)$  solutions). This is also why we used this approach in previous and current works [44–50,64].

Therefore, we may also proceed by another manner to solve the same problem, as briefly mentioned in Section 2.5. We can look for the exact scalar field  $\phi(t)$  and  $V(\phi)$  instead of looking for a teleparallel  $F(T)$  solution. We will set the coframe ansatz components and the characteristic equation defined from Equation (21), as done in the current development, and we will study some relevant and simple  $F(T)$  function ansatz. By inserting the coframe ansatz components and the  $F(T)$  function, we will compute the unified FE defined by Equation (25) and find the associated potential  $V(\phi)$ . By using the conservation law defined in Equation (19), we will find the exact expression for the scalar field  $\phi(t)$ , as desired in such a case. We can easily expect to recover the same  $\phi(T)$  solutions by setting similar  $F(T)$  functions to those found in the current paper. However, if we set more complex and/or different  $F(T)$  functions, we will probably find that the  $\phi(T)$  expressions might be slightly more complex and/or different compared to those used in the current paper. However, the scalar field finding approach has the disadvantage of losing control of the type of scalar field source used and to potentially miss the fundamental FEs  $F(T)$  solutions without the source (homogeneous parts of solution). These are the reasons for using the  $F(T)$  solution finding approach, as done in the current paper.

### 5.2. Logarithmic Scalar Field Source

The logarithmic scalar field source  $\phi(t) = p_0 \ln(pt)$  case (or  $t(\phi) = p^{-1} \exp\left(\frac{\phi}{p_0}\right)$ ) is a good example, leading to simple teleparallel  $F(T)$  solutions. By using the conservation law defined by Equation (19), we find as potential  $V(\phi)$  for the ansatz:

1. **Power-law:** By using the Equations (27) ansatz, we find that:

$$\begin{aligned} \frac{dV}{d\phi} &= p_0 p^2 (1 - b - 2c) \exp\left(-\frac{2\phi}{p_0}\right), \\ \Rightarrow V(\phi) &= \phi_0 + \frac{p_0^2 p^2}{2} (b + 2c - 1) \exp\left(-\frac{2\phi}{p_0}\right). \end{aligned} \quad (181)$$

Equation (20) for  $\alpha_Q$  is:

$$\alpha_Q = -1 + \left[ \left( \frac{\phi_0}{p^2 p_0^2} \right) \exp\left(\frac{2\phi}{p_0}\right) + \frac{(b+2c)}{2} \right]^{-1}. \quad (182)$$

There are some examples leading to simple  $F(T)$  solutions:

(a)  $c = 1$ : We will use Equations (50) and (51) to find, as scalar field:

$$\phi(T) = p_0 \ln \left( \sqrt{2} p \left( 1 + 2b - \frac{1}{c_0^2} \right)^{1/2} \right) - \frac{p_0}{2} \ln(T), \quad (183)$$

and then:

$$V(T) = \frac{p_0^2 (b+2c-1)}{4 \left( 1 + 2b - \frac{1}{c_0^2} \right)} T. \quad (184)$$

Equation (42) becomes, for the Equation (181) potential:

$$F(T) = -\Lambda_0 + F_0 T^{1/C} + \frac{2\kappa p_0^2 (b+2c-1)}{4 \left( 1 + 2b - \frac{1}{c_0^2} \right) (C-1)} T. \quad (185)$$

Equation (184) shows that a logarithmic scalar field can lead to a simple  $V(T)$  potential expression and then to a TEGR-like  $F(T)$  source term, as obtained in Equation (185). This term is similar to a case of Teleparallel Robertson–Walker (TRW) spacetime  $F(T)$  solutions for flat cosmological cases, as found in refs. [49,50,64].

(b) **Late Cosmology  $t \rightarrow \infty$  limit:**

- $c > 1$ : From Equation (84), we find that  $\phi(T) = p_0 \ln \left( p \left( \frac{T}{2c(c+2b)} \right)^{-1/2} \right)$  and the  $V(T)$  from Equation (181) becomes:

$$V(T) = \frac{p_0^2 (b+2c-1)}{4c(c+2b)} T. \quad (186)$$

By using Equations (84), (85), and (186), we will find that Equation (86) becomes:

$$F(T) = -\Lambda_0 + F_0 T^{\frac{2}{3}} + \frac{\kappa p_0^2 (b+2c-1)}{c(c+2b)} T. \quad (187)$$

Here again, the Equation (187) source term is a TEGR-like contribution as for Equation (185), as in refs. [49,50,64].

- $c < 1$ : From Equation (92), we find that  $\phi(T) = p_0 \ln \left( p \left( \frac{c_0^2}{2} (-T) \right)^{-1/2c} \right)$  and the  $V(T)$  from Equation (181) becomes:

$$V(T) = \frac{p_0^2}{2} (b+2c-1) \left( \frac{c_0^2}{2} \right)^{1/c} (-T)^{1/c}. \quad (188)$$

By using Equations (92), (93), and (188), we will find that Equation (94) becomes:

$$F(T) = -\Lambda_0 + F_0 T^{\frac{2(c-b)}{c}} + \frac{2\kappa p_0^2 (c-b)(b+2c-1)}{(2b-2c+1)} \left(-\frac{c_0^2}{2}\right)^{1/c} T^{\frac{1}{c}}. \quad (189)$$

Equation (189) is a sum of power-law terms as for the TRW  $F(T)$  spacetimes, as found in refs. [49,50,64].

(c) **Early Cosmology  $t \rightarrow 0$  limit:** For this limit, Equation (39) will lead to the cases satisfying:

- i.  **$c > 1$ :** Equations (92) and (93) for the  $T \rightarrow -\infty$  limit, and we recover Equation (189) under the same limit.
- ii.  **$c < 1$ :** Equations (84) and (85) for the  $T \rightarrow \infty$  limit, and we recover Equation (187) under the same limit.

2. **Exponential:** By using the Equations (34) ansatz, we find that:

$$\begin{aligned} \frac{dV}{d\phi} &= p_0 p^2 \exp\left(-\frac{2\phi}{p_0}\right) - p_0 p (b+2c) \exp\left(-\frac{\phi}{p_0}\right), \\ \Rightarrow V(\phi) &= \phi_0 - \frac{p_0^2 p^2}{2} \exp\left(-\frac{2\phi}{p_0}\right) + p_0^2 p (b+2c) \exp\left(-\frac{\phi}{p_0}\right). \end{aligned} \quad (190)$$

Equation (20) for  $\alpha_Q$  is:

$$\alpha_Q = -1 + \left[ \left( \frac{\phi_0}{p_0^2 p^2} \right) \exp\left(\frac{2\phi}{p_0}\right) + \frac{(b+2c)}{p} \exp\left(\frac{\phi}{p_0}\right) \right]^{-1}. \quad (191)$$

There are some examples leading to simple  $F(T)$  solutions:

- (a)  **$c \neq -2b$ :** By using Equation (103), we find that  $\phi(T) = p_0 \ln\left(-\frac{p}{2c} \ln\left(\frac{c_0^2}{2}(T_0 - T)\right)\right)$ , and then  $V(T)$  becomes, for the Equation (190) potential:

$$V(T) = -2p_0^2 c^2 \left[ \ln\left(\frac{c_0^2}{2}(T_0 - T)\right) \right]^{-2} - 2p_0^2 c (b+2c) \left[ \ln\left(\frac{c_0^2}{2}(T_0 - T)\right) \right]^{-1}. \quad (192)$$

Then, by substituting Equations (103), (104), and (192) into Equation (42), we find as the  $F(T)$  solution:

$$\begin{aligned} F(T) &= -\Lambda_0 + (T - T_1)^{\frac{2(c-b)}{c}} \left[ F_0 + 8\kappa p_0^2 c (b-c) \right. \\ &\quad \times \left[ \int_T dT' \left[ \ln\left(\frac{c_0^2}{2}(T_0 - T')\right) \right]^{-2} (T' - T_1)^{\frac{2b-3c}{c}} \right. \\ &\quad \left. \left. + \frac{(b+2c)}{c} \int_T dT' \left[ \ln\left(\frac{c_0^2}{2}(T_0 - T')\right) \right]^{-1} (T' - T_1)^{\frac{2b-3c}{c}} \right] \right]. \end{aligned} \quad (193)$$

There is no general solution for Equation (193). By setting  $b = \frac{3c}{2}$ , we can find, as a simple  $F(T)$  solution:

$$F(T) = -\Lambda_0 + (T - T_1)^{-1} \left[ F_0 + \frac{4\kappa p_0^2 c^2}{c_0^2} \left[ \frac{c_0^2(T_0 - T)}{-\ln(2) + \ln(-c_0^2(T - T_0))} + 9 \operatorname{Ei}_1(\ln(2) - \ln(-c_0^2(T - T_0))) \right] \right]. \quad (194)$$

There are several possible examples for  $b > \frac{3c}{2}$  and/or  $T_1 \neq T_0$ .

(b) **Late Cosmology  $t \rightarrow \infty$  limit:**

- $c > 0$ : As in Sections 3.2 and 4.2, Equation (103) leads to  $T = T_0 = \text{constant}$  when  $c \neq -2b$  (TdS-like spacetime) and  $T = 0$  (null torsion scalar spacetime) when  $c = -2b$ , GR solutions as for TdS spacetimes [48].
- $c < 0$ : By using Equation (121), we find that  $\phi(T) = p_0 \ln\left(\frac{p}{2|c|} \ln\left(\frac{c_0^2}{2}(-T)\right)\right)$ , and  $V(T)$  becomes, for Equation (190):

$$V(T) = -2c^2 p_0^2 \left[ \ln\left(\frac{c_0^2}{2}(-T)\right) \right]^{-2} + 2p_0^2 |c| (b - 2|c|) \left[ \ln\left(\frac{c_0^2}{2}(-T)\right) \right]^{-1}. \quad (195)$$

Then, by substituting Equations (121) and (195) into Equation (123), we find as the  $F(T)$  solution:

$$F(T) = -\Lambda_0 + T^{\frac{2(b+|c|)}{|c|}} \left[ F_0 - 8\kappa p_0^2 (b + |c|) |c| \times \left[ \int_T dT' \left[ \ln\left(\frac{c_0^2}{2}(-T')\right) \right]^{-2} T'^{-\frac{2b+3|c|}{|c|}} + \frac{(2|c| - b)}{|c|} \int_T dT' \left[ \ln\left(\frac{c_0^2}{2}(-T')\right) \right]^{-1} T'^{-\frac{2b+3|c|}{|c|}} \right] \right]. \quad (196)$$

There is no general solution for Equation (196). By setting  $b = -\frac{3|c|}{2}$ , we can find, as a simple  $F(T)$  solution:

$$F(T) = -\Lambda_0 + T^{-1} \left[ F_0 + \frac{4\kappa p_0^2 |c|^2}{c_0^2} \left[ \frac{T}{\ln(2) - \ln(-c_0^2 T)} + 9 \operatorname{Ei}_1(\ln(2) - \ln(-c_0^2 T)) \right] \right]. \quad (197)$$

The new teleparallel  $F(T)$  solutions found from the Equation (34) ansatz are more complex than in refs. [49,50,64], but they are comparable to the solution classes in ref. [46].

- (c) **Early Cosmology  $t \rightarrow 0$  limit:** As in Sections 3.2 and 4.2, Equation (103) leads to  $T \rightarrow T_0 - \frac{2}{c_0^2} = \text{constant}$ , a GR solution (a TdS-like spacetime) [48,107].

All the previous teleparallel  $F(T)$  solutions are new and comparable to those found in refs. [45,46,49,50,64], especially for the late and/or early cosmological limit scenarios.

## 6. Discussion and Conclusions

The primary aim of this study was to find the teleparallel  $F(T)$  solutions coming from the FEs precisely described by Equations (21)–(24) with two usual types of scalar field  $\phi(t)$ :

power-law and exponential. In addition and to complete the study, we briefly solved the same FEs with the same ansatzes for a logarithmic scalar field source to find more simple  $F(T)$  solutions comparable to the TRW spacetime solutions found in refs. [49,50,64]. In each cases and depending on the values of  $p$  in the scalar field, we solved the laws of conservation and the FEs for various types of potential  $V(T)$ . We first found that the FEs transform by various substitutions into the single Equation (40), for any type of scalar field and ansatz. From this last equation, the possible teleparallel  $F(T)$  solutions all boil down to Equation (42) using the function  $A(T)$  described by Equation (41). All of the new  $F(T)$  solutions obtained in Sections 3.1 through to the end of Section 5.2 were found by applying Equation (42) for various types of scalar potential. For the  $A_2$  and  $A_3$  coframe components, the power-law ansatz has been used in Sections 3.1 and 4.1; the application has been restricted to the values of  $c = -2b, 1, -1$ , and  $2$ . Most of other values of  $c$  do not result in closed analytic functions and/or are not analytically solvable. In addition, we studied and found that the teleparallel  $F(T)$  solutions under the late/early cosmological limit (i.e.,  $t \rightarrow \infty$ ) for the simple cases are close to the TRW solutions, as found in refs. [49,50,64]. In Sections 3.2 and 4.2, the exponential coframe ansatz has been used to study a well-known case of infinite sum of power terms. At the same time, this ansatz is much simpler to treat and only brings out two types of case instead of an infinity: general and  $c = -2b$ . In Section 5.2, we used the same coframe ansatzes, but we restricted the study to  $c = 1$  (power-law only),  $c \neq -2b$  (exponential only), and the late/early cosmological limit cases. For the rest, each type of case treated here leads to its own class of  $F(T)$  solutions, with its own particularities.

For the power-law  $\phi(t)$  treated in Section 3, we first studied in Section 3.1 via the power-law coframe ansatz the cases of potentials described by Equations (28), (30), and (32), i.e., the general case,  $p \gg 1$ , and  $p = 1$ . This was intended to study the effects of different scalar potential types associated with gravitational sources. The  $p \gg 1$  potential cases can be useful for phantom energy model studies involving fast accelerating universe expansion. Often, physical phenomena described by a scalar field are characterized by its associated potential, and teleparallel  $F(T)$ -gravity cosmology is no exception to the rule. In Section 3.1, we find general and exact  $F(T)$  solutions for the  $c = -2b$  and  $c = 1$  cases. For the  $c = -1$  and  $c = 2$  cases, even if there is no real general and analytic large coverage solution, there are analytical  $F(T)$  solutions for specific cases depending on the parameters  $C_1$  and  $C_2$ , specially defined to simply solve these cases. Therefore, larger values of  $c$  subcases can be useful for studies on teleparallel phantom energy and quintom models. For late and early cosmological limit cases, an important number of new  $F(T)$  solutions are comparable to those obtained in refs. [46,50,64], especially with the isotropic TRW solutions. In Section 3.2, via the exponential ansatz coframe and for the potentials described by Equations (35) and (37), we obtain purely analytic  $F(T)$  solutions, but this required the introduction and definition of new special functions depending on the parameters of the scalar field and the ansatz used to achieve them. Even in cosmology and teleparallel gravity, special functions are a necessary evil to achieve analytic solutions. Other types of approach, such as quantum gravity theories or fundamental particle interactions with gravity, regularly require special functions to express certain new solutions (see, for example, refs. [108,109]). It would not be surprising if the special functions were to be used even more in some future work in teleparallel gravity, more specifically for solutions involving more complex and sophisticated source terms to be solved.

By considering exponential  $\phi(t)$  in Section 4, we first have, in Section 4.1, studied the power-law coframe ansatz and the potentials described in Equations (30) and (127). We obtain teleparallel  $F(T)$  solutions that are slightly more complex compared to the results obtained in Section 3.1, but with some clearly visible similarities. Again, the  $c = -2b$  and



$c = 1$  cases lead to analytical  $F(T)$  solutions. However, we only obtain  $F(T)$  solutions for very specific subcases in the  $c = -1$  and  $c = 2$  situations. This is not surprising compared to the results of Section 3.1. For Section 4.2, with the exponential ansatz coframe and the potentials described by Equations (30) and (129), we essentially obtain purely analytical and easily usable  $F(T)$  solutions. These latter solutions confirm, at the same time, some new simple solutions obtained in Section 3. Let us not forget that the exponential ansatz and scalar field are cases fundamentally expressed by an infinite sum of terms in powers, as stated in ref. [46]. This last fact also explains some common points in the newly obtained  $F(T)$  solutions. In comparison, the new simplest  $F(T)$  solutions obtained in this paper and including most of the late cosmological limit  $F(T)$  solutions are also similar to the teleparallel cosmology solutions obtained in the recent literature (see, for example, refs. [46,49,50,75,83]). This finding is also an additional argument in favor of the new solutions' rightness obtained in the present paper.

Beyond the new solutions, this approach aimed to obtain purely analytical teleparallel  $F(T)$  solutions that could be used in the future to study complex cosmological models in more detail using teleparallel  $F(T)$ -gravity tools. One need only think in particular of the quintessence dark energy, the cosmological phantom energy (negative energy), or even cosmological quintom models. In these latter cases, one will be able, in the near future, to study models involving perfect fluids and scalar fields at the same time, in order to represent reality more faithfully. In such a case, we will use the common (or very similar)  $F(T)$  solutions of the present paper and of some recent papers concerning teleparallel  $F(T)$  solutions in perfect fluids [46,47,49]. This will ultimately lead to complete and realistic cosmological models that can fully describe and explain the dark energy quintessence process ( $-1 < \alpha_Q < -\frac{1}{3}$ ), a realistic and most probable scenario according to the recent literature [65–67,72]. After that, one could just as well study with a similar approach some physical models of the cosmological phantom energy type ( $\alpha_Q < -1$ ), leading to the extreme scenario of the acceleration of uncontrolled universe expansion leading to the Big Rip [87–89]. We can also add the quintom dark energy models, because this is, by definition, a mix of the quintessence and phantom models [94–99]. As a safe and right comparison basis, we can also use some scalar field TRW  $F(T)$  solutions found in ref. [50] as a comparison basis for the late/early cosmological limit solutions. Like the fundamental scalar field associated with the dark energy quintessence process, one could obtain, for the different models involving phantom energy, such fundamental scalar fields. Perhaps the cosmological process of uncontrolled universe acceleration leading to the Big Rip would in fact be explainable by a fundamental scalar field. The same questions will also be relevant and useful for future quintom model studies in teleparallel gravity. The questions will then be as follows: What type(s) of  $V(\phi(T))$  exactly? What are the most appropriate teleparallel  $F(T)$  solutions? What are the coframes, spin-connections, and other conditions? All these questions and hypotheses deserve to be answered seriously and tactfully in future works by using the teleparallel gravity framework and toolkit.

**Funding:** A.L. is supported by an Atlantic Association of Research in Mathematical Sciences (AARMS) fellowship.

**Data Availability Statement:** All data materials are included within the paper. No external data have been produced for this paper.

**Acknowledgments:** Thanks to A.A. Coley for useful and constructive comments. Thanks also to R.J. van den Hoogen for relevant discussions on the topic.

**Conflicts of Interest:** The author declares no conflicts of interest.

## Abbreviations

The following abbreviations are used in this manuscript:

AL	Alexandre Landry
CK	Cartan–Karlhede
DE	Differential Equation
EoS	Equation of State
FE	Field Equation
GR	General Relativity
KV	Killing Vector
NGR	New General Relativity
SHO	Simple Harmonic Oscillator
TEGR	Teleparallel Equivalent of General Relativity
TRW	Teleparallel Robertson–Walker
TdS	Teleparallel de Sitter

## Appendix A. Additional Power-Law Scalar Field $F(T)$ Solutions

### Appendix A.1. Power-Law Ansatz $c = -1$ Solutions

Equation (63)  $C_2 \neq 0$  teleparallel  $F(T)$  solutions:

1.  $C_1 = \frac{3}{2}$  ( $b = -\frac{4}{3}$  and  $C_2 = \frac{176}{3c_0^2}$ ): There are three possible solutions:

- $p \neq \frac{1}{3}$ :

$$F(T) = -\Lambda_0 + F_0[-u_-(T)]^{\frac{2\delta_1}{3}} + \frac{2\kappa V_p}{(3p-1)} \delta_1^{p-1} [u_{\pm}(T)]^{p-1}. \quad (A1)$$

- $p = \frac{1}{3}$  and  $\delta_1 = +1$ :

$$F(T) = -\Lambda_0 + [-u_-(T)]^{\frac{2}{3}} \left[ F_0 - \frac{2\kappa V_p}{3C_2^{\frac{2}{3}}} \ln[-u_-(T)] \right]. \quad (A2)$$

- $p = \frac{1}{3}$  and  $\delta_1 = -1$ :

$$F(T) = -\Lambda_0 + [-u_-(T)]^{\frac{2}{3}} \left[ F_0 + \frac{2\kappa V_p}{3(-1)^{\frac{2}{3}}} \ln[-u_-(T)] \right]. \quad (A3)$$

2.  $C_1 \rightarrow \frac{3}{4}$ : By setting  $C_1 = \frac{3}{4} + \epsilon$  where  $\epsilon \ll 1$ , we find:

$$F(T) = -\Lambda_0 + F_0 + \frac{4\kappa V_p}{3C_2(p+1)(p-1)} [u_{\pm}(T)]^p \left[ T + \delta_1 p \sqrt{T^2 + C_2} \right]. \quad (A4)$$

Equation (66)  $C_2 \neq 0$  teleparallel  $F(T)$  solutions ( $p \gg 1$  case):

1.  $C_1 = \frac{3}{2}$  ( $b = -\frac{4}{3}$  and  $C_2 = \frac{176}{3c_0^2}$ ):

$$F(T) = -\Lambda_0 + F_0[-u_-(T)]^{\frac{2\delta_1}{3}} + \frac{2\kappa V_{\infty}}{3p} [u_{\pm}(T)]^p. \quad (A5)$$

2.  $C_1 \rightarrow \frac{3}{4}$ : By setting  $C_1 = \frac{3}{4} + \epsilon$ , where  $\epsilon \ll 1$ , we find:

$$F(T) = -\Lambda_0 + F_0 + \frac{4\kappa V_{\infty}}{3C_2 p} [u_{\pm}(T)]^p (C_2 - T[u_{\pm}(T)]). \quad (A6)$$

Equation (69)  $C_2 \neq 0$  teleparallel  $F(T)$  solutions ( $p = 1$  case):

1.  $C_1 = \frac{3}{2}$  ( $b = -\frac{4}{3}$  and  $C_2 = \frac{176}{3c_0^2}$ ):

$$F(T) = -\tilde{\Lambda}_0 + F_0[-u_-(T)]^{\frac{2\delta_1}{3}} + \kappa p_0^2(b-2) \left( \ln[u_{\pm}(T)] - \frac{3}{2} \right). \quad (\text{A7})$$

2.  $C_1 \rightarrow \frac{3}{4}$ : By setting  $C_1 = \frac{3}{4} + \epsilon$ , where  $\epsilon \ll 1$ , we find:

$$F(T) = -\tilde{\Lambda}_0 + F_0 - \frac{\kappa p_0^2}{3}(b-2) \left[ \delta_1 \ln[-u_-(T)] - \ln^2[u_{\pm}(T)] + \frac{2T}{C_2}[u_{\pm}(T)] \left( \ln[u_{\pm}(T)] - \frac{1}{2} \right) \right]. \quad (\text{A8})$$

#### Appendix A.2. Power-Law Ansatz $c = 2$ Solutions

The general Equation (75) teleparallel  $F(T)$  solution for  $C_1 = 2C_2$  is:

$$F(T) = -\Lambda_0 + \left[ -\frac{T}{C_2} + 2(w_{\pm}(T)) \right]^{(2-b)} \left[ F_0 - \frac{\kappa V_p(b-2)}{C_2} \times \int_T dT' [w_{\pm}(T')]^{1-p} \left[ -\frac{T'}{C_2} + 2(w_{\pm}(T')) \right]^{(b-3)} \right]. \quad (\text{A9})$$

Equation (A9) solutions are possible only for the following subcases:

1.  $b = 3$  and  $\delta_1 = 1$ :

$$F(T) = -\Lambda_0 + \left[ -\frac{T}{C_2} + 2(w_+(T)) \right]^{-1} \left[ F_0 - \frac{\kappa V_p}{C_2} (2)^{1-p} T {}_3F_2 \left( 1, \frac{p}{2}, \frac{p-1}{2}; 2, p; \frac{T}{C_2} \right) \right]. \quad (\text{A10})$$

2.  $b = 3$  and  $\delta_1 = -1$ :

$$F(T) = -\Lambda_0 + \left[ -\frac{T}{C_2} + 2(w_-(T)) \right]^{-1} \left[ F_0 + \frac{2^{p-1}\kappa V_p}{C_2^{2-p}(p-2)} T^{2-p} {}_2F_1 \left( \frac{2-p}{2}, \frac{1-p}{2}; 3-p; \frac{T}{C_2} \right) \right]. \quad (\text{A11})$$

3.  $b = 4$  and  $\delta_1 = 1$ :

$$F(T) = -\Lambda_0 + \left[ -\frac{T}{C_2} + 2(w_+(T)) \right]^{-2} \left[ F_0 - \frac{\kappa V_p}{2^{p-1}} \left[ -\left( \frac{T}{C_2} \right)^2 {}_3F_2 \left( 2, \frac{p}{2}, \frac{p-1}{2}; 3, p; \frac{T}{C_2} \right) + 8 \left( \frac{T}{C_2} \right) {}_3F_2 \left( 1, \frac{p-2}{2}, \frac{p-1}{2}; 2, p-1; \frac{T}{C_2} \right) \right] \right]. \quad (\text{A12})$$

4.  $b = 4$  and  $\delta_1 = -1$ :

$$F(T) = -\Lambda_0 + \left[ -\frac{T}{C_2} + 2(w_-(T)) \right]^{-2} \left[ F_0 - \frac{8\kappa V_p}{(p-3)} \left( \frac{T}{2C_2} \right)^{(3-p)} \times \left[ {}_3F_2 \left( 3-p, \frac{2-p}{2}, \frac{1-p}{2}; 4-p, 2-p; \frac{T}{C_2} \right) + {}_2F_1 \left( \frac{2-p}{2}, \frac{3-p}{2}; 4-p; \frac{T}{C_2} \right) \right] \right]. \quad (\text{A13})$$

The general Equation (79) teleparallel  $F(T)$  solution for  $C_1 = 2C_2$  is ( $p \gg 1$  case):

$$F(T) = -\Lambda_0 + [-T + 2C_2(w_{\pm}(T))]^{(2-b)} \left[ F_0 - \kappa V_{\infty} (b-2) \right. \\ \left. \times \int_T dT' [w_{\pm}(T')]^{-p} [-T' + 2C_2(w_{\pm}(T'))]^{(b-3)} \right]. \quad (\text{A14})$$

Equation (A14) solutions are possible only for the following subcases:

1.  $b = 3$  and  $\delta_1 = 1$ :

$$F(T) \approx -\Lambda_0 + \left[ -\frac{T}{C_2} + 2(w_+(T)) \right]^{-1} \left[ F_0 - \frac{\kappa V_p}{C_2 2^p} T {}_3F_2 \left( 1, \frac{p}{2}, \frac{p}{2}; 2, p; \frac{T}{C_2} \right) \right]. \quad (\text{A15})$$

2.  $b = 3$  and  $\delta_1 = -1$ :

$$F(T) \approx -\Lambda_0 - \left[ \frac{T}{C_2} - 2(w_-(T)) \right]^{-1} \left[ F_0 + \frac{(2C_2)^p \kappa V_p}{p} T^{-p} {}_2F_1 \left( -\frac{p}{2}, -\frac{p}{2}; -p; \frac{T}{C_2} \right) \right]. \quad (\text{A16})$$

3.  $b = 4$  and  $\delta_1 = 1$ :

$$F(T) \approx -\Lambda_0 + \left[ -\frac{T}{C_2} + 2(w_+(T)) \right]^{-2} \left[ F_0 - \frac{\kappa V_p}{2^p} \left[ -\left( \frac{T}{C_2} \right)^2 {}_3F_2 \left( 2, \frac{p}{2}, \frac{p}{2}; 3, p; \frac{T}{C_2} \right) \right. \right. \\ \left. \left. + 8 \left( \frac{T}{C_2} \right) {}_3F_2 \left( 1, \frac{p}{2}, \frac{p}{2}; 2, p; \frac{T}{C_2} \right) \right] \right]. \quad (\text{A17})$$

4.  $b = 4$  and  $\delta_1 = -1$ :

$$F(T) \approx -\Lambda_0 + \left[ \frac{T}{C_2} - 2(w_-(T)) \right]^{-2} \left[ F_0 - \frac{2\kappa V_p}{p} \left( \frac{T}{2C_2} \right)^{-p} {}_2F_1 \left( -\frac{p}{2}, -\frac{p}{2}; -p; \frac{T}{C_2} \right) \right]. \quad (\text{A18})$$

The general Equation (83) teleparallel  $F(T)$  solution for  $C_1 = 2C_2$  is ( $p = 1$  case):

$$F(T) = -\tilde{\Lambda}_0 + [-T + 2C_2(w_{\pm}(T))]^{(2-b)} \left[ F_0 + \frac{\kappa p_0^2 (b+4)(b-2)}{2C_2} \right. \\ \left. \times \int_T dT' \ln [w_{\pm}(T')] [-T' + 2C_2(w_{\pm}(T'))]^{(b-3)} \right]. \quad (\text{A19})$$

The simplest solutions of Equation (83) are:

1.  $b = 3$ :

$$F(T) = -\tilde{\Lambda}_0 + [-T + 2C_2(w_+(T))]^{-1} \left[ F_0 - \frac{7\kappa p_0^2}{2C_2} \left[ \frac{T}{2} + C_2 w_{\pm}(T) - T \ln [w_{\pm}(T)] \right] \right]. \quad (\text{A20})$$

2.  $b = 4$ :

$$F(T) = -\tilde{\Lambda}_0 + [-T + 2C_2(w_+(T))]^{-2} \left[ F_0 + \frac{8\kappa p_0^2}{C_2} \left[ \left[ -\frac{4\delta_1 C_2^2}{3} \left( 1 - \frac{T}{C_2} \right)^{3/2} - 3T^2 + 12C_2 T \right. \right. \right. \\ \left. \left. \left. - 8C_2^2 \right] \ln [w_{\pm}(T)] + \frac{\delta_1 C_2}{18} (2C_2 - 5T) \sqrt{1 - \frac{T}{C_2} + \frac{T^2}{8} - \frac{C_2}{3} T + \frac{C_2^2}{9}} \right] \right]. \quad (\text{A21})$$

## Appendix B. Additional Exponential Scalar Field $F(T)$ Solutions

### Appendix B.1. Power-Law Ansatz $c = 1$ Solutions

Equation (138)  $C_1 \neq 1$  teleparallel  $F(T)$  solutions:

1.  $C = -1$ :

$$F(T) = -\Lambda_0 + F_0 T^{-1} + \kappa p_0^2 \left[ p^2 T^{-1} \left( Ei_1 \left( -\frac{T_2}{\sqrt{T}} \right) T_2^2 + \exp \left( \frac{T_2}{\sqrt{T}} \right) (T_2 \sqrt{T} + T) \right) \right. \\ \left. - p(b+2) \left( \frac{(-T_2 \sqrt{T} + T) \exp \left( -\frac{T_2}{\sqrt{T}} \right)}{T} + \frac{(T_2^2 - 2T) Ei_1 \left( \frac{T_2}{\sqrt{T}} \right)}{T} \right) \right]. \quad (A22)$$

2.  $C = 2$ :

$$F(T) = -\Lambda_0 + F_0 T^{1/2} + \kappa p_0^2 \left[ \left( p^2 - 2p(b+2) \right) \frac{\sqrt{T}}{T_2} \exp \left( \frac{T_2}{\sqrt{T}} \right) - 2p(b+2) Ei \left( -\frac{T_2}{\sqrt{T}} \right) \right]. \quad (A23)$$

3.  $C = -2$ :

$$F(T) = -\Lambda_0 + F_0 T^{-1/2} + \kappa p_0^2 \left[ p^2 \left( \exp \left( \frac{T_2}{\sqrt{T}} \right) + \frac{T_2}{\sqrt{T}} Ei_1 \left( -\frac{T_2}{\sqrt{T}} \right) \right) \right. \\ \left. - 2p(b+2) \left( \exp \left( -\frac{T_2}{\sqrt{T}} \right) - \left( \frac{T_2}{\sqrt{T}} + 1 \right) Ei_1 \left( \frac{T_2}{\sqrt{T}} \right) \right) \right]. \quad (A24)$$

Equation (141)  $C_1 \neq 1$  teleparallel  $F(T)$  solutions ( $p \gg 1$  case):

1.  $C = -1$ :

$$F(T) = -\Lambda_0 + F_0 T^{-1} + \kappa p^2 p_0^2 \left[ \frac{T_2^2}{T} Ei_1 \left( -\frac{T_2}{\sqrt{T}} \right) + \left( \frac{T_2}{\sqrt{T}} + 1 \right) \right]. \quad (A25)$$

2.  $C = 2$ :

$$F(T) = -\Lambda_0 + F_0 T^{1/2} + \kappa p^2 p_0^2 \frac{T^{1/2}}{T_2} \exp \left( \frac{T_2}{\sqrt{T}} \right). \quad (A26)$$

3.  $C = -2$ :

$$F(T) = -\Lambda_0 + F_0 T^{-1/2} + \kappa p^2 p_0^2 \left[ \frac{T_2}{\sqrt{T}} Ei_1 \left( -\frac{T_2}{\sqrt{T}} \right) + \exp \left( \frac{T_2}{\sqrt{T}} \right) \right]. \quad (A27)$$

### Appendix B.2. Power-Law Ansatz $c = 2$ Solutions

Equation (153) teleparallel  $F(T)$  solutions are:

1.  $C_1 = 0$ :

$$F(T) = -\Lambda_0 + (-T)^{(2-b)} \left[ F_0 - \kappa(b-2) p_0^2 \left[ p^2 \int_T dT' (-T')^{(b-3)} \exp \left( \frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T')]^{1/2}} \right) \right. \right. \\ \left. \left. - 2p(b+4) \int_T dT' (-T')^{(b-3)} Ei \left( -\frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T')]^{1/2}} \right) \right] \right]. \quad (A28)$$

There are solutions for specific values of  $b$ , such as:

- $b = 1$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 + (-T) \left[ F_0 - \kappa p_0^2 \left[ -\frac{1}{4\sqrt{w_{\pm}(T)} c_0 C_2 (w_{\pm}(T) - 2)} \right. \right. \\
& \times \left[ -\sqrt{w_{\pm}(T)} p^3 (w_{\pm}(T) - 2) \left( e^{\frac{2p}{c_0}} \text{Ei}_1 \left( -\frac{2p(\sqrt{2} - \sqrt{w_{\pm}(T)})}{c_0 \sqrt{w_{\pm}(T)}} \right) \right. \right. \\
& - e^{-\frac{2p}{c_0}} \text{Ei}_1 \left( -\frac{2p(\sqrt{2} + \sqrt{w_{\pm}(T)})}{c_0 \sqrt{w_{\pm}(T)}} \right) \left. \right] + \left( \left( p^2 + \frac{c_0^2}{2} \right) (w_{\pm}(T) - 1) \right. \\
& + \left. \left( p^2 - \frac{c_0^2}{2} \right) \sqrt{w_{\pm}(T)} - c_0 \sqrt{2} p (w_{\pm}(T) - 2) \right) \exp \left( \frac{2\sqrt{2} p}{c_0 \sqrt{w_{\pm}(T)}} \right) c_0 \left. \right] \\
& + \frac{5}{2(w_{\pm}(T))^{\frac{3}{2}} C_2^2 p (w_{\pm}(T) - 2)} \left[ \left( \frac{T}{2} \text{Ei}_1 \left( \frac{2p(\sqrt{2} - \sqrt{w_{\pm}(T)})}{c_0 \sqrt{w_{\pm}(T)}} \right) e^{-\frac{2p}{c_0}} \right. \right. \\
& + \frac{T}{2} \text{Ei}_1 \left( \frac{2p(\sqrt{2} + \sqrt{w_{\pm}(T)})}{c_0 \sqrt{w_{\pm}(T)}} \right) e^{\frac{2p}{c_0}} - 4C_2 \text{Ei}_1 \left( \frac{2\sqrt{2} p}{c_0 \sqrt{w_{\pm}(T)}} \right) \sqrt{w_{\pm}(T)} p^2 \\
& + \left. \left. \frac{Tc_0}{4} \left( 2\sqrt{2} p + c_0 \sqrt{w_{\pm}(T)} \right) \exp \left( -\frac{2\sqrt{2} p}{c_0 \sqrt{w_{\pm}(T)}} \right) \right] \right].
\end{aligned} \tag{A29}$$

- $b = 3$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 - T^{-1} \left[ F_0 + \kappa p_0^2 \left[ -\frac{p^2}{3c_0^4} \left[ -512 \left( p^2 - \frac{3c_0^2}{8} \right) p^2 C_2 \text{Ei}_1 \left( -\frac{4\sqrt{2} p}{c_0 \sqrt{w_{\pm}(T)}} \right) \right. \right. \right. \\
& + \left( -64pC_2\sqrt{2} \left( \frac{c_0^2}{16} (w_{\pm}(T) - 1) + p^2 - \frac{5c_0^2}{16} \right) \sqrt{w_{\pm}(T)} \right. \\
& + c_0 \left( -16p^2 C_2 (w_{\pm}(T)) + 3Tc_0^2 \right) \left. \right] c_0 \exp \left( \frac{4\sqrt{2} p}{c_0 \sqrt{w_{\pm}(T)}} \right) \left. \right] \\
& + \frac{7p}{3c_0^4} \left[ 2(96C_2 p^2 c_0^2 - 128p^4 C_2 - 3c_0^4 T) \text{Ei}_1 \left( \frac{4\sqrt{2} p}{c_0 \sqrt{w_{\pm}(T)}} \right) \right. \\
& + c_0 \left( 32pC_2\sqrt{2} \left( \frac{c_0^2}{16} (w_{\pm}(T) - 1) + p^2 - \frac{11c_0^2}{16} \right) \sqrt{w_{\pm}(T)} \right. \\
& + \left. \left. \frac{3c_0}{2} \left( 2C_2 \left( -\frac{8p^2}{3} + c_0^2 \right) (w_{\pm}(T) - 1) + (T + 2C_2)c_0^2 - \frac{16p^2 C_2}{3} \right) \right) \right. \\
& \times \left. \left. \exp \left( -\frac{4\sqrt{2} p}{c_0 \sqrt{w_{\pm}(T)}} \right) \right] \right].
\end{aligned} \tag{A30}$$

- $b = 4$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 + (-T)^{-2} \left[ F_0 + 2\kappa p_0^2 \left[ -\frac{p^2}{126c_0^8} \left[ -78125p^4C_2^2 \left( p^4 - \frac{84}{25}p^2c_0^2 + \frac{168}{125}c_0^4 \right) \right. \right. \right. \\
& \times \operatorname{Ei}_1 \left( -\frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) - 63 \left( -\frac{1}{21} \left( 25 \left( \frac{3c_0^2}{5} \left( \left( T - \frac{32C_2}{15} \right) c_0^4 + 20C_2p^2c_0^2 \right. \right. \right. \right. \\
& - \frac{125p^4C_2}{18} \right) (w_{\pm}(T) - 1) - \frac{c_0^6}{25} (39T + 32C_2) + p^2(T - 128C_2)c_0^4 + \frac{2075p^4C_2c_0^2}{6} \\
& - \frac{625p^6C_2}{6} \left. \right) pC_2\sqrt{2}\sqrt{w_{\pm}(T)} \left. \right) + c_0 \left( -\frac{25}{21} \left( \left( T - \frac{76C_2}{5} \right) c_0^4 + 65C_2p^2c_0^2 \right. \right. \\
& - \frac{125p^4C_2}{6} \left. \right) p^2C_2(w_{\pm}(T) - 1) + T^2c_0^6 + \frac{45p^2C_2 \left( T + \frac{76C_2}{27} \right) c_0^4}{7} \\
& - \frac{125p^4C_2(T + 26C_2)c_0^2}{42} + \frac{3125p^6C_2^2}{126} \left. \right) \left. \right] c_0 \exp \left( \frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) \left. \right] \\
& + \frac{p}{63c_0^8} \left[ (-210000p^4c_0^4C_2^2 + 350000c_0^2p^6C_2^2 - 78125C_2^2p^8 + 504T^2c_0^8) \right. \\
& \times \operatorname{Ei}_1 \left( \frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) + 252 \left( \frac{1}{4} \left( -\frac{25pC_2}{21} \sqrt{2} \left( \frac{3}{5}c_0^2 \left( \left( T - \frac{116C_2}{15} \right) c_0^4 \right. \right. \right. \right. \right. \\
& + \frac{250C_2p^2c_0^2}{9} - \frac{125p^4C_2}{18} \right) (w_{\pm}(T) - 1) + \frac{(-67T - 116C_2)c_0^6}{25} \\
& + p^2 \left( T - \frac{790C_2}{3} \right) c_0^4 + \frac{925p^4C_2c_0^2}{2} - \frac{625p^6C_2}{6} \left. \right) \sqrt{w_{\pm}(T)} - c_0 \left( -\frac{25}{21} \right. \\
& \times \left( -\frac{28(T + 2C_2)c_0^6}{25} + p^2 \left( T - \frac{188C_2}{5} \right) c_0^4 + \frac{265p^4C_2c_0^2}{3} - \frac{125p^6C_2}{6} \right) C_2 \\
& \times (w_{\pm}(T) - 1) + \left( T^2 + \frac{8C_2^2}{3} \right) c_0^6 + \frac{205p^2C_2 \left( T + \frac{188C_2}{41} \right) c_0^4}{21} \\
& \left. \left. - \frac{125p^4C_2 \left( T + \frac{106C_2}{3} \right) c_0^2}{42} + \frac{3125p^6C_2^2}{126} \right) \right] \exp \left( -\frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) c_0 \left. \right] \right]. \quad (A31)
\end{aligned}$$

There are other values of  $b$  yielding to analytical  $F(T)$  solutions.

2.  $C_1 = 2C_2$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 + [-T + 2C_2(w_{\pm}(T))]^{(2-b)} \left[ F_0 - \kappa(b-2)p_0^2 \int_T dT' \left[ p^2 \exp \left( \frac{\sqrt{2}p(1+b)}{c_0[w_{\pm}(T')]^{1/2}} \right) \right. \right. \\
& \left. \left. - 2p(b+4) \operatorname{Ei} \left( -\frac{\sqrt{2}p(1+b)}{c_0[w_{\pm}(T')]^{1/2}} \right) \right] [-T' + 2C_2(w_{\pm}(T'))]^{(b-3)} \right]. \quad (A32)
\end{aligned}$$

There are solutions for specific values of  $\delta_1$  and  $b$ :

- $b = 3$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 + [-T + 2C_2(w_{\pm}(T))]^{-1} \left[ F_0 + \kappa p_0^2 \right. \\
& \times \left[ -\frac{p^2}{3c_0^4} \left[ -512p^2 \left( p^2 - \frac{3c_0^2}{8} \right) C_2 \text{Ei}_1 \left( -\frac{4\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) \right. \right. \\
& + \exp \left( \frac{4\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) \left( -64p\sqrt{2}C_2 \left( \frac{c_0^2}{16} (w_{\pm}(T) - 1) + p^2 - \frac{5c_0^2}{16} \right) \right. \\
& \times \sqrt{w_{\pm}(T)} + c_0 (-16p^2C_2(w_{\pm}(T)) + 3Tc_0^2) \Big) c_0 \Big] \\
& + \frac{7p}{3c_0^4} \left[ 2(96C_2c_0^2p^2 - 128C_2p^4 - 3Tc_0^4) \text{Ei}_1 \left( \frac{4\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) \right. \\
& + \left( \left( 32p\sqrt{2}C_2 \left( \frac{c_0^2}{16} (w_{\pm}(T) - 1) + p^2 - \frac{11c_0^2}{16} \right) \sqrt{w_{\pm}(T)} \right. \right. \\
& + \frac{3c_0}{2} \left( 2C_2 \left( -\frac{8p^2}{3} + c_0^2 \right) (w_{\pm}(T) - 1) + (T + 2C_2)c_0^2 - \frac{16p^2C_2}{3} \right) \Big) \\
& \times \exp \left( -\frac{4\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) \Big) c_0 \Big] \Big]. \tag{A33}
\end{aligned}$$

- $b = 4$ :

$$\begin{aligned}
F(T) = & -\Lambda_0 + [-T + 2C_2(w_{\pm}(T))]^{-2} \left[ F_0 + 2\kappa p_0^2 \right. \\
& \times \left[ -\frac{p^2}{126c_0^8} \left[ -78125p^6C_2^2 \left( p^2 - \frac{28c_0^2}{25} \right) \text{Ei}_1 \left( -\frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) \right. \right. \\
& - 63 \exp \left( \frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) c_0 \left( -\frac{25}{21} p\sqrt{2}C_2 \left( \frac{3c_0^2}{5} \left( \left( T - \frac{4C_2}{15} \right) c_0^4 \right. \right. \right. \\
& + \frac{40c_0^2p^2C_2}{9} - \frac{125p^4C_2}{18} \Big) (w_{\pm}(T) - 1) + \frac{(17T - 4C_2)c_0^6}{25} + p^2 \left( T + \frac{8C_2}{3} \right) c_0^4 \\
& + \frac{225p^4C_2c_0^2}{2} - \frac{625p^6C_2}{6} \Big) \sqrt{w_{\pm}(T)} + \left( -\frac{25}{21} C_2 \left( \frac{56(T - C_2)c_0^6}{25} \right. \right. \\
& + p^2 \left( T + \frac{8C_2}{5} \right) c_0^4 + \frac{55p^4C_2c_0^2}{3} - \frac{125p^6C_2}{6} \Big) (w_{\pm}(T) - 1) + \left( T^2 - 4TC_2 + \frac{8}{3}C_2^2 \right) \\
& \times c_0^6 - \frac{5p^2C_2(8C_2 + T)c_0^4}{21} - \frac{125p^4 \left( T + \frac{22C_2}{3} \right) C_2c_0^2}{42} + \frac{3125p^6C_2^2}{126} \Big) c_0 \Big] \\
& + \frac{p}{378c_0^8} \left[ \left( 8064C_2^2c_0^8(w_{\pm}(T) - 1)^3 + \left( 8064C_2^2 - 12096TC_2 + 3024T^2 \right) c_0^8 \right. \right. \\
& + 700000C_2^2p^6c_0^2 - 468750p^8C_2^2 \Big) \text{Ei}_1 \left( \frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) + 1512c_0 \\
& \times \left( \left( -\frac{25p}{84} \left( \frac{3}{5} \left( \left( T + \frac{44C_2}{45} \right) c_0^4 + \frac{190c_0^2p^2C_2}{27} - \frac{125p^4C_2}{18} \right) c_0^2(w_{\pm}(T) - 1) \right. \right. \right. \\
& + \left( \frac{23T}{75} + \frac{44C_2}{75} \right) c_0^6 + p^2 \left( T + \frac{38C_2}{9} \right) c_0^4 + \frac{2725p^4C_2c_0^2}{18} - \frac{625p^6C_2}{6} \Big) \\
& \times \sqrt{2}C_2\sqrt{w_{\pm}(T)} - \frac{c_0}{4} \left( -\frac{25}{21} C_2 \left( \left( \frac{28T}{15} - \frac{56C_2}{75} \right) c_0^6 \right. \right. \\
& + p^2 \left( T + \frac{52C_2}{15} \right) c_0^4 + \frac{235p^4C_2c_0^2}{9} - \frac{125p^6C_2}{6} \Big) (w_{\pm}(T) - 1) \\
& + \left( T^2 - \frac{8}{3}TC_2 + \frac{8}{9}C_2^2 \right) c_0^6 + \frac{55p^2 \left( T - \frac{52C_2}{11} \right) C_2c_0^4}{63} - \frac{125p^4 \left( T + \frac{94C_2}{9} \right) C_2c_0^2}{42} \\
& + \frac{3125p^6C_2^2}{126} \Big) \exp \left( -\frac{5\sqrt{2}p}{c_0\sqrt{w_{\pm}(T)}} \right) \Big) \Big] \Big]. \tag{A34}
\end{aligned}$$

There are other values of  $b$  yielding to analytical  $F(T)$  solutions.

Equation (155)  $C_1 \neq 0$  teleparallel  $F(T)$  solutions ( $p \gg 1$  case):



1.  $C_1 = 0$ :

$$F(T) = -\Lambda_0 + (-T)^{(2-b)} \left[ F_0 - \kappa p^2 p_0^2 \int_T dT' \exp\left(\frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T')]^{1/2}}\right) (-T')^{(b-3)} \right]. \quad (\text{A35})$$

There are solutions for specific values of  $b$ :

•  $b = 1$ :

$$F(T) \approx -\Lambda_0 - T \left[ F_0 - \frac{\kappa p_0^2 p^2}{4 c_0 C_2} \left[ p \exp\left(\frac{2p}{c_0}\right) \text{Ei}_1\left(-\frac{2p(\sqrt{2} - \sqrt{w_{\pm}(T)})}{c_0 \sqrt{w_{\pm}(T)}}\right) + \frac{(w_{\pm}(T))}{(2 - w_{\pm}(T))} \exp\left(\frac{2\sqrt{2}p}{c_0 \sqrt{w_{\pm}(T)}}\right) c_0 \right] \right]. \quad (\text{A36})$$

•  $b = 3$ :

$$F(T) \approx -\Lambda_0 - \left[ \frac{F_0}{T} + \frac{64\kappa p_0^2 p^5 C_2}{3c_0^3} \frac{\sqrt{2}}{T} \sqrt{w_{\pm}(T)} \exp\left(\frac{4\sqrt{2}p}{c_0 \sqrt{w_{\pm}(T)}}\right) \right]. \quad (\text{A37})$$

•  $b = 4$ :

$$F(T) \approx -\Lambda_0 + \left[ \frac{F_0}{T^2} + \frac{15625\kappa p_0^2 p^9 C_2^2}{63\sqrt{2}c_0^7 T^2} \sqrt{w_{\pm}(T)} \exp\left(\frac{5\sqrt{2}p}{c_0 \sqrt{w_{\pm}(T)}}\right) \right]. \quad (\text{A38})$$

2.  $C_1 = 2C_2$ :

$$F(T) = -\Lambda_0 + [-T + C_1(w_{\pm}(T))]^{(2-b)} [y(T)]^{\frac{(b-2)C_1}{2(C_1-2C_2)}} \left[ \frac{T + (C_1 - 2C_2)(w_{\pm}(T))}{T + (C_1 - 2C_2)(2 - w_{\pm}(T))} \right]^{\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \\ \times \left[ F_0 - \kappa p^2 p_0^2 \int_T dT' \exp\left(\frac{\sqrt{2}p(1+b)}{c_0 [w_{\pm}(T')]^{1/2}}\right) [-T' + C_1(w_{\pm}(T'))]^{(b-3)} \right. \\ \left. \times [y(T')]^{-\frac{(b-2)C_1}{2(C_1-2C_2)}} \left[ \frac{T' + (C_1 - 2C_2)(w_{\pm}(T'))}{T' + (C_1 - 2C_2)(2 - w_{\pm}(T'))} \right]^{-\frac{\delta_1(b-2)C_1}{2(C_1-2C_2)}} \right]. \quad (\text{A39})$$

There are solutions for specific values of  $\delta_1$  and  $b$ :

•  $b = 3$ :

$$F(T) \approx -\Lambda_0 + [-T + C_1(w_{\pm}(T))]^{-1} \times \left[ F_0 + \frac{64\kappa p^5 p_0^2}{3c_0^3} \sqrt{2} C_2 \sqrt{w_{\pm}(T)} \exp\left(\frac{4\sqrt{2}p}{c_0 \sqrt{w_{\pm}(T)}}\right) \right]. \quad (\text{A40})$$

•  $b = 4$ :

$$F(T) \approx -\Lambda_0 + [-T + C_1(w_{\pm}(T))]^{-2} \times \left[ F_0 + \frac{15625\kappa p^9 p_0^2}{126\sqrt{2}c_0^7} C_2^2 \sqrt{w_{\pm}(T)} \exp\left(\frac{5\sqrt{2}p}{c_0 \sqrt{w_{\pm}(T)}}\right) \right]. \quad (\text{A41})$$

### Appendix B.3. Exponential Ansatz Solutions

Equation (170)  $T_1 \neq T_0$  and  $b \geq \frac{3c}{2}$  teleparallel  $F(T)$  solutions:

$$1. \quad b = \frac{3c}{2}:$$

$$F(T) = -\Lambda_0 + (T - T_1)^{-1} \left[ F_0 + \kappa p p_0^2 \frac{c(7c + 2p)}{2(c - p)} \left( -\frac{2}{c_0^2} \right)^{p/c} (T - T_0)^{1-p/c} \right]. \quad (\text{A42})$$

$$2. \quad b = 2c:$$

$$F(T) = -\Lambda_0 + (T - T_1)^{-2} \left[ F_0 + 2\kappa p p_0^2 c(4c + p) \left( -\frac{2}{c_0^2} \right)^{p/c} \times \frac{(T - T_0)^{-\frac{p+c}{c}} ((T + T_0 - 2T_1)c - p(T - T_1))}{2c^2 - 3cp + p^2} \right]. \quad (\text{A43})$$

$$3. \quad b = 3c:$$

$$F(T) = -\Lambda_0 + (T - T_1)^{-4} \left[ F_0 + \frac{24\kappa p p_0^2 c(3c + 2c + p)}{(24c^4 - 50c^3p + 35c^2p^2 - 10cp^3 + p^4)} \left( -\frac{2}{c_0^2} \right)^{p/c} \times (T - T_0)^{-\frac{p+c}{c}} \left( (2T_1^2 + (-2T - 2T_0)T_1 + T^2 + T_0^2)(T + T_0 - 2T_1)c^3 \right. \right. \\ \left. \left. - \frac{11}{6} \left( \frac{26T_1^2}{11} + \left( -\frac{31T}{11} - \frac{21T_0}{11} \right) T_1 + T^2 + \frac{9TT_0}{11} + \frac{6T_0^2}{11} \right) p(T - T_1)c^2 \right. \right. \\ \left. \left. + \left( T + \frac{T_0}{2} - \frac{3T_1}{2} \right) p^2(T - T_1)^2c - \frac{p^3(T - T_1)^3}{6} \right) \right]. \quad (\text{A44})$$

Equation (173)  $T_1 \neq T_0$  and  $b \geq \frac{3c}{2}$  teleparallel  $F(T)$  solutions ( $p \gg 1$  case):

$$1. \quad b = \frac{3c}{2}:$$

$$F(T) \approx -\Lambda_0 + F_0 (T - T_1)^{-1} - \kappa p p_0^2 c \left( -\frac{2}{c_0^2} \right)^{p/c} (T - T_0)^{-p/c} (T - T_1)^{-1}. \quad (\text{A45})$$

$$2. \quad b = 2c:$$

$$F(T) \approx -\Lambda_0 + F_0 (T - T_1)^{-2} - 2\kappa p p_0^2 c \left( -\frac{2}{c_0^2} \right)^{p/c} (T - T_0)^{-\frac{p}{c}} (T - T_1)^{-1}. \quad (\text{A46})$$

$$3. \quad b = 3c:$$

$$F(T) \approx -\Lambda_0 + F_0 (T - T_1)^{-4} - 4\kappa p p_0^2 c \left( -\frac{2}{c_0^2} \right)^{p/c} (T - T_0)^{-\frac{p}{c}} (T - T_1)^{-1}. \quad (\text{A47})$$

## References

1. Aldrovandi, R.; Pereira, J.G. *Teleparallel Gravity, An Introduction*; Springer: Berlin/Heidelberg, Germany, 2013.
2. Bahamonde, S.; Dialektopoulos, K.; Escamilla-Rivera, C.; Farrugia, G.; Gakis, V.; Hendry, M.; Hohmann, M.; Said, J.L.; Mifsud, J.; Di Valentino, E. Teleparallel Gravity: From Theory to Cosmology. *Rep. Prog. Phys.* **2023**, *86*, 026901. [[CrossRef](#)]
3. Krssak, M.; van den Hoogen, R.; Pereira, J.; Boehmer, C.; Coley, A. Teleparallel Theories of Gravity: Illuminating a Fully Invariant Approach. *Class. Quantum Gravity* **2019**, *36*, 183001. [[CrossRef](#)]
4. McNutt, D.D.; Coley, A.A.; van den Hoogen, R.J. A frame based approach to computing symmetries with non-trivial isotropy groups. *J. Math. Phys.* **2023**, *64*, 032503. [[CrossRef](#)]
5. Coley, A.A.; van den Hoogen, R.J.; McNutt, D.D. Symmetry and Equivalence in Teleparallel Gravity. *J. Math. Phys.* **2020**, *61*, 072503. [[CrossRef](#)]
6. Lucas, T.G.; Obukhov, Y.; Pereira, J.G. Regularizing role of teleparallelism. *Phys. Rev. D* **2009**, *80*, 064043. [[CrossRef](#)]
7. Krssak, M.; Pereira, J.G. Spin Connection and Renormalization of Teleparallel Action. *Eur. Phys. J. C* **2015**, *75*, 519. [[CrossRef](#)]

8. Chinea, F. Symmetries in tetrad theories. *Class. Quantum Gravity* **1988**, *5*, 135. [\[CrossRef\]](#)
9. Estabrook, F.; Wahlquist, H. Moving frame formulations of 4-geometries having isometries. *Class. Quantum Gravity* **1996**, *13*, 1333. [\[CrossRef\]](#)
10. Papadopoulos, G.; Grammenos, T. Locally homogeneous spaces, induced Killing vector fields and applications to Bianchi prototypes. *J. Math. Phys.* **2012**, *53*, 072502. [\[CrossRef\]](#)
11. Olver, P. *Equivalence, Invariants and Symmetry*; Cambridge University Press: Cambridge, UK, 1995.
12. Linder, E. Einstein's Other Gravity and the Acceleration of the Universe. *Phys. Rev. D* **2010**, *81*, 127301; Erratum in *Phys. Rev. D* **2010**, *82*, 109902. [\[CrossRef\]](#)
13. Ferraro, R.; Fiorini, F. Modified teleparallel gravity: Inflation without an inflation. *Phys. Rev. D* **2007**, *75*, 084031. [\[CrossRef\]](#)
14. Ferraro, R.; Fiorini, F. On Born-Infeld Gravity in Weitzenböck spacetime. *Phys. Rev. D* **2008**, *78*, 124019. [\[CrossRef\]](#)
15. Hayashi, K.; Shirafuji, T. New general relativity. *Phys. Rev. D* **1979**, *19*, 3524. [\[CrossRef\]](#)
16. Jimenez, J.B.; Dialektopoulos, K.F. Non-Linear Obstructions for Consistent New General Relativity. *J. Cosmol. Astropart. Phys.* **2020**, *2020*, 018. [\[CrossRef\]](#)
17. Bahamonde, S.; Blixt, D.; Dialektopoulos, K.F.; Hell, A. Revisiting Stability in New General Relativity. *arXiv* **2024**, arXiv:2404.02972.
18. Heisenberg, L. Review on  $f(Q)$  Gravity. *arXiv* **2023**, arXiv:2309.15958. [\[CrossRef\]](#)
19. Heisenberg, L.; Hohmann, M.; Kuhn, S. Cosmological teleparallel perturbations. *arXiv* **2023**, arXiv:2311.05495. [\[CrossRef\]](#)
20. Flathmann, K.; Hohmann, M. Parametrized post-Newtonian limit of generalized scalar-nonmetricity theories of gravity. *Phys. Rev. D* **2022**, *105*, 044002. [\[CrossRef\]](#)
21. Hohmann, M. General covariant symmetric teleparallel cosmology. *Phys. Rev. D* **2021**, *104*, 124077. [\[CrossRef\]](#)
22. Jimenez, J.B.; Heisenberg, L.; Koivisto, T.S. The Geometrical Trinity of Gravity. *Universe* **2019**, *5*, 173. [\[CrossRef\]](#)
23. Nakayama, Y. Geometrical trinity of unimodular gravity. *Class. Quantum Gravity* **2023**, *40*, 125005. [\[CrossRef\]](#)
24. Xu, Y.; Li, G.; Harko, T.; Liang, S.-D.  $f(Q, T)$  gravity. *Eur. Phys. J. C* **2019**, *79*, 708. [\[CrossRef\]](#)
25. Maurya, D.C.; Yesmakhanova, K.; Myrzakulov, R.; Nugmanova, G. Myrzakulov  $F(T, Q)$  gravity: Cosmological implications and constraints. *arXiv* **2024**, arXiv:2404.09698.
26. Maurya, D.C.; Yesmakhanova, K.; Myrzakulov, R.; Nugmanova, G. Myrzakulov, FLRW Cosmology in Myrzakulov  $F(R, Q)$  Gravity. *arXiv* **2024**, arXiv:2403.11604.
27. Maurya, D.C.; Myrzakulov, R. Exact Cosmology in Myrzakulov Gravity. *arXiv* **2024**, arXiv:2402.02123.
28. Harko, T.; Lobo, F.S.N.; Nojiri, S.; Odintsov, S.D.  $f(R, T)$  gravity. *Phys. Rev. D* **2011**, *84*, 024020. [\[CrossRef\]](#)
29. Golovnev, A.; Guzman, M.-J. Approaches to spherically symmetric solutions in  $f(T)$ -gravity. *Universe* **2021**, *7*, 121. [\[CrossRef\]](#)
30. Golovnev, A. Issues of Lorentz-invariance in  $f(T)$ -gravity and calculations for spherically symmetric solutions. *Class. Quantum Gravity* **2021**, *38*, 197001. [\[CrossRef\]](#)
31. Golovnev, A.; Guzman, M.-J. Bianchi identities in  $f(T)$ -gravity: Paving the way to confrontation with astrophysics. *Phys. Lett. B* **2020**, *810*, 135806. [\[CrossRef\]](#)
32. DeBenedictis, A.; Ilijić, S.; Sossich, M. On spherically symmetric vacuum solutions and horizons in covariant  $f(T)$  gravity theory. *Phys. Rev. D* **2022**, *105*, 084020. [\[CrossRef\]](#)
33. Bahamonde, S.; Camci, U. Exact Spherically Symmetric Solutions in Modified Teleparallel gravity. *Symmetry* **2019**, *11*, 1462. [\[CrossRef\]](#)
34. Awad, A.; Golovnev, A.; Guzman, M.-J.; El Hanafy, W. Revisiting diagonal tetrads: New Black Hole solutions in  $f(T)$ -gravity. *Eur. Phys. J. C* **2022**, *82*, 972. [\[CrossRef\]](#)
35. Bahamonde, S.; Golovnev, A.; Guzmán, M.-J.; Said, J.L.; Pfeifer, C. Black Holes in  $f(T, B)$  Gravity: Exact and Perturbed Solutions. *J. Cosmol. Astropart. Phys.* **2022**, *01*, 037. [\[CrossRef\]](#)
36. Bahamonde, S.; Faraji, S.; Hackmann, E.; Pfeifer, C. Thick accretion disk configurations in the Born-Infeld teleparallel gravity. *Phys. Rev. D* **2022**, *106*, 084046. [\[CrossRef\]](#)
37. Nashed, G.G.L. Quadratic and cubic spherically symmetric black holes in the modified teleparallel equivalent of general relativity: Energy and thermodynamics. *Class. Quantum Gravity* **2021**, *38*, 125004. [\[CrossRef\]](#)
38. Pfeifer, C.; Schuster, S. Static spherically symmetric black holes in weak  $f(T)$ -gravity. *Universe* **2021**, *7*, 153. [\[CrossRef\]](#)
39. El Hanafy, W.; Nashed, G.G.L. Exact Teleparallel Gravity of Binary Black Holes. *Astrophys. Space Sci.* **2016**, *361*, 68. [\[CrossRef\]](#)
40. Aftergood, J.; DeBenedictis, A. Matter Conditions for Regular Black Holes in  $f(T)$  Gravity. *Phys. Rev. D* **2014**, *90*, 124006. [\[CrossRef\]](#)
41. Bahamonde, S.; Doneva, D.D.; Ducobu, L.; Pfeifer, C.; Yazadjiev, S.S. Spontaneous Scalarization of Black Holes in Gauss-Bonnet Teleparallel Gravity. *Phys. Rev. D* **2023**, *107*, 10, 104013. [\[CrossRef\]](#)
42. Bahamonde, S.; Ducobu, L.; Pfeifer, C. Scalarized Black Holes in Teleparallel Gravity. *J. Cosmol. Astropart. Phys.* **2022**, *04*, 018. [\[CrossRef\]](#)
43. Calza, M.; Sebastiani, L. A class of static spherically symmetric solutions in  $f(T)$ -gravity. *Eur. Phys. J. C* **2024**, *84*, 476. [\[CrossRef\]](#)
44. Landry, A. Static spherically symmetric perfect fluid solutions in teleparallel  $F(T)$  gravity. *Axioms* **2024**, *13*, 333. [\[CrossRef\]](#)

45. Coley, A.A.; Landry, A.; van den Hoogen, R.J.; McNutt, D.D. Spherically symmetric teleparallel geometries. *Eur. Phys. J. C* **2024**, *84*, 334. [\[CrossRef\]](#)
46. Landry, A. Kantowski-Sachs spherically symmetric solutions in teleparallel  $F(T)$  gravity. *Symmetry* **2024**, *16*, 953. [\[CrossRef\]](#)
47. van den Hoogen, R.J.; Forance, H. Teleparallel Geometry with Spherical Symmetry: The diagonal and proper frames. *J. Cosmol. Astrophys.* **2024**, *11*, 033. [\[CrossRef\]](#)
48. Coley, A.A.; Landry, A.; van den Hoogen, R.J.; McNutt, D.D. Generalized Teleparallel de Sitter geometries. *Eur. Phys. J. C* **2023**, *83*, 977. [\[CrossRef\]](#)
49. Coley, A.A.; Landry, A.; Gholami, F. Teleparallel Robertson-Walker Geometries and Applications. *Universe* **2023**, *9*, 454. [\[CrossRef\]](#)
50. Landry, A. Scalar field sources Teleparallel Robertson-Walker  $F(T)$ -gravity solutions. *Mathematics* **2025**, in press. [\[CrossRef\]](#)
51. Rodrigues, M. E.; Kpadonou, A.V.; Rahaman, F.; Oliveira, P.J.; Houndjo, M.J.S. Bianchi type-I, type-III and Kantowski-Sachs solutions in  $f(T)$  gravity. *Astrophys. Space Sci.* **2015**, *357*, 129. [\[CrossRef\]](#)
52. Amir, M.J.; Yussouf, M. Kantowski-Sachs Universe Models in  $f(T)$  Theory of Gravity. *Int. J. Theor. Phys.* **2015**, *54*, 2798. [\[CrossRef\]](#)
53. Leon, G.; Roque, A.A. Qualitative analysis of Kantowski-Sachs metric in a generic class of  $f(R)$  models. *J. Cosmol. Astropart. Phys.* **2014**, *05*, 032. [\[CrossRef\]](#)
54. Shaikh, A.A.; Chakraborty, D. Curvature properties of Kantowski-Sachs metric. *J. Geom. Phys.* **2021**, *160*, 103970. [\[CrossRef\]](#)
55. Oliveira-Neto, G.; Canedo, D.L.; Monerat, G.A. An anisotropic Kantowski-Sachs universe with radiation, dust and a phantom fluid. *Braz. J. Phys.* **2022**, *52*, 130. [\[CrossRef\]](#)
56. Leon, G.; Paliathanasis, A. Anisotropic spacetimes in  $f(T, B)$  theory II: Kantowski-Sachs Universe. *Eur. Phys. J. Plus* **2022**, *137*, 855. [\[CrossRef\]](#)
57. Leon, G.; Paliathanasis, A. Anisotropic spacetimes in  $f(T, B)$  theory III: LRS Bianchi III Universe. *Eur. Phys. J. Plus* **2022**, *137*, 927. [\[CrossRef\]](#)
58. Vinutha, T.; Niharika, K.; Sri Kavya, K. The Study of Kantowski-Sachs Perfect Fluid Cosmological Model in Modified Gravity. *Astrophysics* **2023**, *66*, 64. [\[CrossRef\]](#)
59. Samanta, G.C. Kantowski-Sachs Universe Filled with Perfect Fluid in  $f(R, T)$  Theory of Gravity. *Int. J. Theor. Phys.* **2013**, *52*, 2647. [\[CrossRef\]](#)
60. Dimakis, N.; Roumeliotis, M.; Paliathanasis, A.; Christodoulakis, T. Anisotropic Solutions in Symmetric Teleparallel  $f(Q)$ -theory: Kantowski-Sachs and Bianchi III LRS Cosmologies. *Eur. Phys. J. C* **2023**, *83*, 794. [\[CrossRef\]](#)
61. Millano, A.D.; Dialektopoulos, K.; Dimakis, N.; Giacomini, A.; Shababi, H.; Halder, A.; Paliathanasis, A. Kantowski-Sachs and Bianchi III dynamics in  $f(Q)$ -gravity. *Phys. Rev. D* **2024**, *109*, 124044. [\[CrossRef\]](#)
62. Paliathanasis, A. Classical and Quantum Cosmological Solutions in Teleparallel Dark Energy with Anisotropic Background Geometry. *Symmetry* **2022**, *14*, 1974. [\[CrossRef\]](#)
63. Paliathanasis, A. Kantowski-Sachs cosmology in scalar-torsion theory. *Eur. Phys. J. C* **2023**, *83*, 213. [\[CrossRef\]](#)
64. Gholami, F.; Landry, A. Cosmological solutions in teleparallel  $F(T, B)$  gravity. *Symmetry* **2025**, *17*, 060. [\[CrossRef\]](#)
65. Zlatev, I.; Wang, L.; Steinhardt, P. Quintessence, Cosmic Coincidence, and the Cosmological Constant. *Phys. Rev. Lett.* **1999**, *82*, 896. [\[CrossRef\]](#)
66. Steinhardt, P.; Wang, L.; Zlatev, I. Cosmological tracking solutions. *Phys. Rev. D* **1999**, *59*, 123504. [\[CrossRef\]](#)
67. Caldwell, R.R.; Dave, R.; Steinhardt, P. Cosmological Imprint of an Energy Component with General Equation of State. *Phys. Rev. Lett.* **1998**, *80*, 1582. [\[CrossRef\]](#)
68. Carroll, S.M. Quintessence and the Rest of the World. *Phys. Rev. Lett.* **1998**, *81*, 3067. [\[CrossRef\]](#)
69. Doran, M.; Lilley, M.; Schwindt, J.; Wetterich, C. Quintessence and the Separation of CMB Peaks. *Astrophys. J.* **2001**, *559*, 501. [\[CrossRef\]](#)
70. Zeng, X.-X.; Chen, D.-Y.; Li, L.-F. Holographic thermalization and gravitational collapse in the spacetime dominated by quintessence dark energy. *Phys. Rev. D* **2015**, *91*, 046005. [\[CrossRef\]](#)
71. Chakraborty, S.; Mishra, S.; Chakraborty, S. Dynamical system analysis of quintessence dark energy model. *Int. J. Geom. Methods Mod. Phys.* **2025**, *22*, 2450250. [\[CrossRef\]](#)
72. Shlivko, D.; Steinhardt, P.J. Assessing observational constraints on dark energy. *Phys. Lett. B* **2024**, *855*, 138826. [\[CrossRef\]](#)
73. Wetterich, C. Cosmology and the Fate of Dilatation Symmetry. *Nucl. Phys. B* **1988**, *302*, 668. [\[CrossRef\]](#)
74. Ratra, B.; Peebles, P.J.E. Cosmological consequences of a rolling homogeneous scalar field. *Phys. Rev. D* **1988**, *37*, 3406. [\[CrossRef\]](#)
75. Duchaniya, L.K.; Gandhi, K.; Mishra, B. Attractor behavior of  $f(T)$  modified gravity and the cosmic acceleration. *Phys. Dark Universe* **2024**, *44*, 101464. [\[CrossRef\]](#)
76. Hohmann, M. Teleparallel gravity. In *Modified and Quantum Gravity*; Springer: Berlin/Heidelberg, Germany, 2022; p. 145.
77. Hohmann, M. Scalar-torsion theories of gravity III: Analogue of scalar-tensor gravity and conformal invariants. *Phys. Rev. D* **2018**, *98*, 064004. [\[CrossRef\]](#)

78. Zubair, M.; Waheed, S.; Fayyaz, M.A.; Ahmad, I. Energy Constraints and Phenomenon of Cosmic Evolution in  $f(T, B)$  Framework. *Eur. Phys. J. Plus* **2018**, *133*, 452. [\[CrossRef\]](#)
79. Bhattacharjee, S. Constraining  $f(T, B)$  teleparallel gravity from energy conditions. *New Astron.* **2021**, *83*, 101495. [\[CrossRef\]](#)
80. Trivedi, O.; Khlopov, M.; Said, J.L.; Nunes, R. Cosmological singularities in  $f(T, \phi)$  gravity. *Eur. Phys. J. C* **2023**, *83*, 1017. [\[CrossRef\]](#)
81. Paliathanasis, A.; Leon, G.  $f(T, B)$  gravity in a Friedmann–Lemaître–Robertson–Walker universe with nonzero spatial curvature. *Math. Methods Appl. Sci.* **2023**, *46*, 3905. [\[CrossRef\]](#)
82. Paliathanasis, A.; Leon, G. Cosmological evolution in  $f(T, B)$  gravity. *Eur. Phys. J. Plus* **2021**, *136*, 1092. [\[CrossRef\]](#)
83. Kofinas, G.; Leon, G.; Saridakis, E.N. Dynamical behavior in  $f(T, T_G)$  cosmology. *Class. Quantum Gravity* **2014**, *31*, 175011. [\[CrossRef\]](#)
84. Farrugia, G.; Said, J.L.; Finch, A. Gravitoelectromagnetism, Solar System Test and Weak-Field Solutions in  $f(T, B)$  Gravity with Observational Constraints. *Universe* **2020**, *6*, 34. [\[CrossRef\]](#)
85. Chiba, T.; Okabe, T.; Yamaguchi, M. Kinetically Driven Quintessence. *Phys. Rev. D* **2000**, *62*, 023511. [\[CrossRef\]](#)
86. Carroll, S.M.; Hoffman, M.; Trodden, M. Can the dark energy equation-of-state parameter  $w$  be less than  $-1$ ? *Phys. Rev. D* **2003**, *68*, 023509. [\[CrossRef\]](#)
87. Caldwell, R.R. A phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state. *Phys. Lett. B* **2002**, *545*, 23. [\[CrossRef\]](#)
88. Farnes, J.S. A Unifying Theory of Dark Energy and Dark Matter: Negative Masses and Matter Creation within a Modified  $\Lambda$ CDM Framework. *Astron. Astrophys.* **2018**, *620*, A92. [\[CrossRef\]](#)
89. Baum, L.; Frampton, P.H. Turnaround in Cyclic Cosmology. *Phys. Rev. Lett.* **2007**, *98*, 071301. [\[CrossRef\]](#)
90. Hu, W. Crossing the Phantom Divide: Dark Energy Internal Degrees of Freedom. *Phys. Rev. D* **2005**, *71*, 047301. [\[CrossRef\]](#)
91. Karimzadeh, S.; Shojaei, R. Phantom-Like Behavior in Modified Teleparallel Gravity. *Adv. High Energy Phys.* **2019**, *2019*, 4026856. [\[CrossRef\]](#)
92. Pati, L.; Kadam, S.A.; Tripathy, S.K.; Mishra, B. Rip cosmological models in extended symmetric teleparallel gravity. *Phys. Dark Universe* **2022**, *35*, 100925. [\[CrossRef\]](#)
93. Kucukakca, Y.; Akbarieh, A.R.; Ashrafi, S. Exact solutions in teleparallel dark energy model. *Chin. J. Phys.* **2023**, *82*, 47. [\[CrossRef\]](#)
94. Cai, Y.-F.; Saridakis, E.N.; Setare, M.R.; Xia, J.-Q. Quintom Cosmology: Theoretical implications and observations. *Phys. Rep.* **2010**, *493*, 1. [\[CrossRef\]](#)
95. Guo, Z.-K.; Piao, Y.-S.; Zhang, X.; Zhang, Y.-Z. Cosmological evolution of a quintom model of dark energy. *Phys. Lett. B* **2005**, *608*, 177. [\[CrossRef\]](#)
96. Feng, B.; Li, M.; Piao, Y.-S.; Zhang, X. Oscillating quintom and the recurrent universe. *Phys. Lett. B* **2006**, *634*, 101. [\[CrossRef\]](#)
97. Mishra, S.; Chakraborty, S. Dynamical system analysis of quintom dark energy model. *Eur. Phys. J. C* **2018**, *78*, 917. [\[CrossRef\]](#)
98. Tot, J.; Coley, A.A.; Yildirim, B.; Leon, G. The dynamics of scalar-field Quintom cosmological models. *Phys. Dark Universe* **2023**, *39*, 101155. [\[CrossRef\]](#)
99. Bahamonde, S.; Marciu, M.; Rudra, P. Generalised teleparallel quintom dark energy non-minimally coupled with the scalar torsion and a boundary term. *J. Cosmol. Astropart. Phys.* **2018**, *04*, 056. [\[CrossRef\]](#)
100. Tajahmad, B. Exact solutions of an anisotropic universe in a modified teleparallel gravity model via the Noether and B.N.S. approaches. *Int. J. Geom. Methods Mod. Phys.* **2024**, *21*, 2450230. [\[CrossRef\]](#)
101. Iosifidis, D. Cosmological Hyperfluids, Torsion and Non-metricity. *Eur. Phys. J. C* **2020**, *80*, 1042. [\[CrossRef\]](#)
102. Heisenberg, L.; Hohmann, M.; Kuhn, S. Homogeneous and isotropic cosmology in general teleparallel gravity. *Eur. Phys. J. C* **2023**, *83*, 315. [\[CrossRef\]](#)
103. Heisenberg, L.; Hohmann, M. Gauge-invariant cosmological perturbations in general teleparallel gravity. *Eur. Phys. J. C* **2024**, *84*, 462. [\[CrossRef\]](#)
104. Bohmer, C.G.; d'Alfonso del Sordo, A. Cosmological fluids with boundary term couplings. *Gen. Relativ. Gravit.* **2024**, *56*, 75. [\[CrossRef\]](#)
105. Hawking, S.W.; Ellis, G.F.R. *The Large Scale Structure of Space-Time*; Cambridge University Press: Cambridge, UK, 2010.
106. Coley, A.A. *Dynamical Systems and Cosmology*; Kluwer Academic: Dordrecht, The Netherlands, 2003.
107. Landry, A.; van den Hoogen, R.J. Teleparallel Minkowski Spacetime with Perturbative Approach for Teleparallel Gravity on a Proper Frame. *Universe* **2023**, *9*, 232. [\[CrossRef\]](#)
108. Landry, A.; Hammad, F. Landau levels in a gravitational field: The Schwarzschild spacetime case. *Universe* **2021**, *7*, 144. [\[CrossRef\]](#)
109. Hammad, F.; Landry, A. Landau levels in a gravitational field: The Levi-Civita and Kerr spacetimes case. *Eur. Phys. J. Plus* **2020**, *135*, 90. [\[CrossRef\]](#)

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.