

Type II DFT solutions from Poisson–Lie T -duality/plurality

Yuho Sakatani*

Department of Physics, Kyoto Prefectural University of Medicine, Kyoto 606-0823, Japan

*E-mail: yuho@koto.kpu-m.ac.jp

Received April 15, 2019; Accepted May 22, 2019; Published July 16, 2019

.....
String theory has T -duality symmetry when the target space has Abelian isometries. A generalization of T -duality, where the isometry group is non-Abelian, is known as non-Abelian T -duality, which works well as a solution-generating technique in supergravity. In this paper we describe non-Abelian T -duality as a kind of $O(D, D)$ transformation when the isometry group acts without isotropy. We then provide a duality transformation rule for the Ramond–Ramond fields by using the technique of double field theory (DFT). We also study a more general class of solution-generating technique, the Poisson–Lie (PL) T -duality or T -plurality. We describe the PL T -plurality as an $O(n, n)$ transformation and clearly show the covariance of the DFT equations of motion by using the gauged DFT. We further discuss the PL T -plurality with spectator fields, and study an application to the $AdS_5 \times S^5$ solution. The dilaton puzzle known in the context of the PL T -plurality is resolved with the help of DFT.
.....

Subject Index B11, B20, B80

1. Introduction

T -duality was discovered and reported in Ref. [1] as a symmetry of string theory compactified on a circle. The mass spectrum or the partition function of string theory on a D -dimensional torus was studied, for example, in Refs. [2–6], and T -duality was identified as an $O(D, D; \mathbb{Z})$ symmetry. It was further studied from a different approach [7,8], and the transformation rules for the background fields (i.e. metric, the Kalb–Ramond B -field, and the dilaton) under T -duality were determined. In Refs. [9,10], T -duality was understood as an $O(D, D)$ symmetry of the classical equations of motion of string theory. The classical symmetry was clarified in Ref. [11] by using the gauged sigma model, and this approach has proved quite useful, for example when we discuss the global structure of the T -dualized background [12]. The transformation rules for the Ramond–Ramond (R–R) fields and spacetime fermions were determined in Refs. [13–16]. This well-established symmetry of string theory is called Abelian T -duality since it relies on the existence of Killing vectors which commute with each other (see Refs. [17,18] for reviews).

An extension of T -duality to the case of non-commuting Killing vectors was explored in Ref. [19] (see Refs. [20,21] for earlier works), and this is known as non-Abelian T -duality (NATD). Various aspects have been studied in Refs. [12,22–35], but unlike Abelian T -duality, there are still many things to be clarified. For example, the partition function in the dual model may not be the same as that of the original model (see Ref. [36] for a recent study), and NATD may rather be regarded as a map between two string theories. The global structure of the dual geometry is also not clearly understood [12]. However, NATD at least generates many new solutions of supergravity, and it can be utilized as a useful solution-generating technique.

Under NATD the isometries are generally broken, and naively we cannot recover the original model from the dual model. However, this issue was resolved by relaxing the condition for the dualizability [37]. The generalized duality is called the Poisson–Lie (PL) T -duality [38], and it can be performed even in the absence of the usual Killing vectors. The PL T -duality is based on a pair of groups with the same dimension, G and \tilde{G} , that form a larger Lie group known as the Drinfel’d double \mathcal{D} . The PL T -duality is a symmetry that exchanges the role of the subgroups G and \tilde{G} . Conventional NATD can be reproduced as a special case where one of the two groups is an Abelian group. Aspects of the PL T -duality and generalizations have been studied in Refs. [39–47], and concrete applications are given, for example, in Refs. [38,48–51].

Low-dimensional Drinfel’d doubles were classified in Refs. [52–54], and it was stressed that some Drinfel’d double \mathfrak{d} can be decomposed into several different pairs of subalgebras \mathfrak{g} and $\tilde{\mathfrak{g}}$, $(\mathfrak{d}, \mathfrak{g}, \tilde{\mathfrak{g}}) \cong (\mathfrak{d}, \mathfrak{g}', \tilde{\mathfrak{g}}') \cong \dots$. The decomposition is called the Manin triple, and each Manin triple corresponds to a sigma model. The existence of several decompositions suggests that many sigma models are related through a Drinfel’d double. This idea was explicitly realized in Ref. [55], and the classical equivalence of the sigma models was called the PL T -plurality (see Refs. [56–60] for more examples). Various aspects of the PL T -plurality were discussed in refs. [61,62], and in particular quantum aspects of the PL T -duality/plurality were studied in Refs. [55,63–71].

Recent developments in NATD were triggered by Ref. [72], which provided the transformation rule for the R–R fields under NATD. Although the analysis was limited to the case where the isometry group acts freely, that restriction was relaxed in Ref. [73]. By exploiting the techniques, NATD for an $SU(2)$ isometry was extensively studied in Refs. [74–100] (mainly in the context of AdS/CFT correspondence) and many novel solutions were constructed. Subsequently, the transformation rules that can also be applied to the fermionic T -duality were obtained in Ref. [101].

More recently, NATD has received much attention in the context of integrable deformations of string theory, since a class of integrable deformation called the homogeneous Yang–Baxter deformation was shown to be a subclass of NATD [102–105]. Other integrable deformations such as the λ -deformation and the η -deformations can also be understood in the framework of the so-called \mathcal{E} -model [106], which was developed in the PL T -duality [37,39]. Moreover, as discussed in Refs. [106–109], the λ -deformation and the η -deformations are related by a PL T -duality and an analytic continuation. Thus, there is a close relationship between the PL T -duality and integrable deformations (see Refs. [110–113] for recent studies on the \mathcal{E} -model).

Another approach to T -duality has been developed in Refs. [114–130] and is called the double field theory (DFT). This manifests the Abelian $O(D, D)$ T -duality symmetry at the level of supergravity by formally doubling the dimensions of the spacetime. Several formulations of DFT have been proposed, such as the flux formulation (or the gauged DFT) [131–134] and DFT on group manifolds (or DFT_{WZW}) [135–137]. Recently, by applying the idea of DFT_{WZW} , a formulation of DFT which manifests the Poisson–Lie T -duality was proposed in Ref. [138] and the transformation of the R–R fields under the PL T -duality was discussed for the first time. The idea was developed in Ref. [139], and applications to various integrable deformations were studied (see also Ref. [140] for discussion on the PL T -duality, $O(D, D)$ symmetry, and integrable deformations). The covariance of the supergravity equations of motion under the PL T -duality was also shown in Refs. [141,142] using mathematical approaches.

In this paper we revisit the traditional NATD in a general setup where the non-vanishing B -field and the R–R fields are included. By assuming that the isometry group acts freely on the target space, we

describe NATD as a kind of $O(D, D)$ rotation of the supergravity fields. From the obtained $O(D, D)$ matrix, we can easily determine the transformation rule for the R–R fields by using the technique of DFT. Indeed, by using the information of given isometry generators, we provide simple duality transformation rules for bosonic fields.

We then demonstrate the efficiency of the formula by studying some concrete examples. Since many examples have already been studied in the literature, in this paper we will just consider the cases where the isometry group is non-unimodular, $f_{ab}{}^a \neq 0$. This type of NATD is not well studied because the resulting dual geometry does not satisfy the supergravity equations of motion [23,25,28]. However, as pointed out in Refs. [143,144], the dual geometry in fact satisfies the generalized supergravity equations of motion (GSE) [145,146]. When the target space satisfies the GSE, string theory has scale invariance [145,147] and the κ -symmetry [146]. The conformal symmetry may be broken, but recently a local counterterm that cancels out the Weyl anomaly was constructed in Ref. [148] (see also Ref. [149]), and string theory may be consistently defined even in the generalized background. Even if it is not the case, NATD for a non-unimodular algebra still works as a solution-generating technique in supergravity, because an arbitrary GSE solution can be mapped to a solution of the usual supergravity [145,149–151] by performing a (formal) T -duality. Then, combining the NATD with $f_{ab}{}^a \neq 0$ and the formal T -duality, we can generate a new supergravity solution.

We also study the PL T -plurality with the R–R fields. In fact, the PL T -plurality can be regarded as a constant $O(n, n)$ transformation acting on “untwisted fields” $\{\hat{\mathcal{H}}_{AB}, \hat{d}, \hat{\mathcal{F}}\}$. By requiring the untwisted fields to satisfy the dualizability condition or the \mathcal{E} -model condition of Ref. [148], we show that the DFT equations of motion in the original and the transformed background are covariantly related by the $O(n, n)$ transformation. This shows that if the original background satisfies the DFT equations of motion, the transformed background is also a solution of DFT. We also discuss the PL T -plurality with spectator fields. Again, by requiring certain conditions for the untwisted fields, we show that the DFT equations of motion are satisfied in the dual background. By using the proposed duality rules, we study an example of the PL T -plurality with the R–R fields.

In studies of the PL T -plurality the so-called dilaton puzzle has been discussed, for example in Refs. [55–58]. Under a PL T -plurality transformation, a dual-coordinate dependence (i.e. dependence on the coordinates of the dual group \tilde{G}) can appear in the dilaton. When such coordinate dependence appears, the background does not have the usual supergravity interpretation, and we are forced to disallow such transformation. However, in DFT we can treat the dual coordinates and the usual coordinates on an equal footing and we do not need to worry about the dilaton puzzle. As discussed in Refs. [149,151], a DFT solution with a dual-coordinate-dependent dilaton can be regarded as a solution of GSE, and by performing a further formal T -duality, we can obtain a linear dilaton solution of the usual supergravity. In this way the issue of the dilaton puzzle is totally resolved and we can consider an arbitrary PL T -plurality transformation.

This paper is organized as follows. In Sect. 2 we briefly review DFT and GSE. In Sect. 3 we begin with a review of the traditional NATD, and translate the results into the language of DFT. We then provide a general transformation rule for the R–R fields. Examples of NATD without and with the R–R fields are studied in Sects. 4 and 5. In Sect. 6, we study the PL T -plurality in terms of DFT and determine the transformation rules from the DFT equations of motion. As an example of the PL T -plurality, in Sect. 7 we study the PL T -plurality transformation of $\text{AdS}_5 \times S^5$ solution. Section 8 is devoted to conclusions and discussions.

2. A review of DFT and GSE

2.1. Generalized-metric formulation of DFT

There are several equivalent formulations of DFT, but the generalized-metric formulation [120,121, 123,127] may be the most accessible; we thus utilize it as much as possible in this paper. In this formulation, the fundamental fields are a symmetric tensor, called the generalized metric $\mathcal{H}_{MN}(x)$, and a scalar density $e^{-2d(x)}$ called the DFT dilaton. The Lagrangian of DFT is given by

$$\begin{aligned} \mathcal{L}_{\text{DFT}} &= e^{-2d} \mathcal{S}, \\ \mathcal{S} &\equiv \frac{1}{8} \mathcal{H}^{MN} \partial_M \mathcal{H}^{PQ} \partial_N \mathcal{H}_{PQ} - \frac{1}{2} \mathcal{H}^{PQ} \partial_Q \mathcal{H}^{MN} \partial_N \mathcal{H}_{PM} + 4 \partial_M d \partial_N \mathcal{H}^{MN} \\ &\quad - 4 \mathcal{H}^{MN} \partial_M d \partial_N d - \partial_M \partial_N \mathcal{H}^{MN} + 4 \mathcal{H}^{MN} \partial_M \partial_N d. \end{aligned} \tag{2.1}$$

Here, the fields are supposed to depend on the generalized coordinates $(x^M) = (x^m, \tilde{x}_m)$ ($M = 1, \dots, 2D, m = 1, \dots, D$), and we raise or lower the indices M, N by using the $O(D, D)$ -invariant metric η_{MN} and its inverse η^{MN} :

$$\eta_{MN} = \begin{pmatrix} 0 & \delta_m^n \\ \delta_n^m & 0 \end{pmatrix}, \quad \eta^{MN} = \begin{pmatrix} 0 & \delta_n^m \\ \delta_m^n & 0 \end{pmatrix}. \tag{2.2}$$

The generalized metric \mathcal{H}_{MN} is defined to be an $O(D, D)$ matrix,

$$\mathcal{H}_M^P \mathcal{H}_N^Q \eta_{PQ} = \eta_{MN}, \tag{2.3}$$

and this property allows us to define projection operators as

$$P^{MN} \equiv \frac{1}{2} (\eta^{MN} + \mathcal{H}^{MN}), \quad \bar{P}^{MN} \equiv \frac{1}{2} (\eta^{MN} - \mathcal{H}^{MN}), \tag{2.4}$$

which satisfy $P_M^N + \bar{P}_M^N = \delta_M^N$. For consistency, we assume that arbitrary fields or gauge parameters $A(x)$ and $B(x)$ satisfy the so-called section condition,

$$\eta^{MN} \partial_M \partial_N A = 0, \quad \eta^{MN} \partial_M A \partial_N B = 0. \tag{2.5}$$

According to this requirement, none of the fields can depend on more than D coordinates. Under the section condition, the DFT action is invariant under the generalized Lie derivative

$$\begin{aligned} \hat{\mathfrak{L}}_V \mathcal{H}_{MN} &\equiv V^P \partial_P \mathcal{H}_{MN} + (\partial_M V^P - \partial^P V_M) \mathcal{H}_{PN} + (\partial_N V^P - \partial^P V_N) \mathcal{H}_{MP}, \\ \hat{\mathfrak{L}}_V d &\equiv V^M \partial_M d - \frac{1}{2} \partial_M V^M. \end{aligned} \tag{2.6}$$

Namely, the generalized Lie derivative generates the gauge symmetry of DFT, known as the generalized diffeomorphisms. Under the section condition, we can also check that the generalized Lie derivative is closed, $[\hat{\mathfrak{L}}_{V_1}, \hat{\mathfrak{L}}_{V_2}] = \hat{\mathfrak{L}}_{[V_1, V_2]_{\text{C}}}$, by means of the C-bracket,

$$\begin{aligned} [V_1, V_2]_{\text{C}}^M &\equiv \frac{1}{2} (\hat{\mathfrak{L}}_{V_1} V_2^M - \hat{\mathfrak{L}}_{V_2} V_1^M) \\ &= V_1^N \partial_N V_2^M - V_2^N \partial_N V_1^M - V_{[1}^N \partial^M V_{2]N}. \end{aligned} \tag{2.7}$$

In particular, when the gauge parameters V_a^M satisfy $\eta_{MN} V_a^M V_b^N = 2 c_{ab}$ (c_{ab} a constant), we can show that the C-bracket coincides with the generalized Lie derivative,

$$[V_a, V_b]_C = \hat{\mathfrak{L}}_{V_a} V_b^M = -\hat{\mathfrak{L}}_{V_b} V_a^M, \tag{2.8}$$

similar to the case of the usual Lie derivative $\mathfrak{L}_{v_a} v_b^m = [v_a, v_b]^m$.

In fact, the scalar \mathcal{S} in Eq. (2.1) can be understood as the generalized Ricci scalar curvature,

$$\mathcal{S} \equiv \frac{1}{2} (P^{MK} P^{NL} - \bar{P}^{MK} \bar{P}^{NL}) S_{MNKL}, \tag{2.9}$$

where the (semi-covariant) curvature S_{MNPQ} is defined by

$$\begin{aligned} S_{MNPQ} &\equiv R_{MNPQ} + R_{PQMN} - \Gamma_{RMN} \Gamma^R_{PQ}, \\ R_{MNPQ} &\equiv \partial_M \Gamma_{NPQ} - \partial_N \Gamma_{MPQ} + \Gamma_{MPR} \Gamma_N^R Q - \Gamma_{NPR} \Gamma_M^R Q. \end{aligned} \tag{2.10}$$

If we use the curvature \mathcal{S} , the invariance of the DFT action under generalized diffeomorphisms is manifest. Then, the DFT action can be understood as a natural generalization of the Einstein–Hilbert action.

The equations of motion are also summarized in a covariant form as¹

$$\mathcal{S} = 0, \quad \mathcal{S}_{MN} = 0, \tag{2.11}$$

where the generalized Ricci tensor is defined by

$$\mathcal{S}_{MN} \equiv (P_M^P \bar{P}_N^Q + \bar{P}_M^P P_N^Q) S_{RPQ}^R. \tag{2.12}$$

For concrete computation, the following expression may be more useful:

$$\mathcal{S}_{MN} = -2 (P_M^P \bar{P}_N^Q + \bar{P}_M^P P_N^Q) \mathcal{K}_{PQ}, \tag{2.13}$$

$$\begin{aligned} \mathcal{K}_{MN} &\equiv \frac{1}{8} \partial_M \mathcal{H}^{PQ} \partial_N \mathcal{H}_{PQ} - \frac{1}{2} \partial_{(M} \mathcal{H}^{PQ} \partial_P \mathcal{H}_{|N)Q} + 2 \partial_M \partial_N d \\ &\quad + (\partial_P - 2 \partial_P d) \left(\frac{1}{2} \mathcal{H}^{PQ} \partial_{(M} \mathcal{H}_{N)Q} + \frac{1}{2} \mathcal{H}^Q_{(M} \partial_Q \mathcal{H}^P_{|N)} - \frac{1}{4} \mathcal{H}^{PQ} \partial_Q \mathcal{H}_{MN} \right). \end{aligned} \tag{2.14}$$

When we make the connection to conventional supergravity, we suppose $\tilde{\delta}^m = 0$ and parameterize the generalized metric and the DFT dilaton as

$$(\mathcal{H}_{MN}) = \begin{pmatrix} g_{mn} - B_{mp} g^{pq} B_{qn} & B_{mp} g^{pn} \\ -g^{mp} B_{pn} & g^{mn} \end{pmatrix}, \quad e^{-2d} = e^{-2\Phi} \sqrt{|g|}, \tag{2.15}$$

by using the standard NS–NS fields $\{g_{mn}, B_{mn}, \Phi\}$. Then, \mathcal{S} and \mathcal{S}_{MN} reduce to

$$\begin{aligned} \mathcal{S} &= R + 4 D^m \partial_m \Phi - 4 D^m \Phi D_m \Phi - \frac{1}{12} H_{mnp} H^{mnp}, \\ (\mathcal{S}_{MN}) &= \begin{pmatrix} 2 g_{(m|k} s^{[kl]} B_{|n)} - s_{(mn)} - B_{mk} s^{(kl)} B_{ln} & B_{mk} s^{(kn)} - g_{mk} s^{[kn]} \\ s^{[mk]} g_{kn} - s^{(mk)} B_{km} & s^{(mn)} \end{pmatrix}, \end{aligned}$$

¹ They are summarized as $\mathcal{G}_{MN} \equiv \mathcal{S}_{MN} - \frac{1}{2} \mathcal{S} \mathcal{H}_{MN} = 0$. Here, the generalized Einstein tensor \mathcal{G}_{MN} satisfies the Bianchi identity $\nabla^M \mathcal{G}_{MN} = 0$ [152], where ∇_M is the covariant derivative for the connection Γ_{MNP} .

$$s_{mn} \equiv R_{mn} - \frac{1}{4} H_{mpq} H_n{}^{pq} + 2D_m \partial_n \Phi - \frac{1}{2} D^k H_{kmn} + \partial_k \Phi H^k{}_{mn}, \quad (2.16)$$

and the following standard supergravity Lagrangian and the equations of motion are reproduced from the DFT Lagrangian in Eq. (2.1) and the DFT equations of motion in Eq. (2.11):

$$\mathcal{L} = \sqrt{|g|} e^{-2\Phi} \left(R + 4 D^m \partial_m \Phi - 4 D^m \Phi D_m \Phi - \frac{1}{12} H_{mnp} H^{mnp} \right), \quad (2.17)$$

$$R + 4 D^m \partial_m \Phi - 4 D^m \Phi D_m \Phi - \frac{1}{12} H_{mnp} H^{mnp} = 0, \quad s_{(mn)} = 0, \quad s_{[mn]} = 0. \quad (2.18)$$

We can also introduce the R–R fields in a manifestly $O(D, D)$ covariant manner. However, the treatment of the R–R fields is slightly involved, and we will not write out the covariant expression explicitly here (see Appendix B, and also Refs. [149,153] for the detail). In the following, aimed at readers who are not familiar with DFT, we will try to describe the R–R fields as the usual p -form fields as much as possible.

2.2. Gauged DFT

When we manifest the covariance under the PL T -plurality, it is convenient to rewrite the DFT equations of motion in Eq. (2.11) by using the technique of the gauged DFT [131–134].

Suppose that the generalized metric \mathcal{H}_{MN} has the form

$$\mathcal{H}_{MN}(x) = [U(x) \hat{\mathcal{H}} U^\top(x)]_{MN}, \quad U \equiv (U_M{}^A), \quad (2.19)$$

where $\hat{\mathcal{H}}_{AB}$ is a constant matrix, which we call the untwisted metric. In this case it is useful to define \mathcal{F}_{ABC} and \mathcal{F}_A , called the gaugings or the generalized fluxes, as

$$\begin{aligned} \mathcal{F}_{ABC} &\equiv 3 \Omega_{[ABC]}, & \mathcal{F}_A &\equiv \Omega^B{}_{AB} + 2 \mathcal{D}_A \mathcal{D}, \\ \Omega_{ABC} &\equiv -\mathcal{D}_A U_B{}^M U_{MC} = \Omega_{A[BC]} & \mathcal{D}_A &\equiv U_A{}^M \partial_M & U_A{}^M &\equiv (U^{-1})_A{}^M. \end{aligned} \quad (2.20)$$

They behave as scalars under generalized diffeomorphisms.

By using the generalized fluxes, we can show that the DFT equations of motion in Eq. (2.11), under the section condition, are equivalent to

$$\mathcal{R} = 0, \quad \mathcal{G}^{AB} = 0, \quad (2.21)$$

where

$$\begin{aligned} \mathcal{R} &\equiv -2 \bar{P}^{AB} (2 \mathcal{D}_A \mathcal{F}_B - \mathcal{F}_A \mathcal{F}_B) - \frac{1}{3} \bar{P}^{ABCDEF} \mathcal{F}_{ABC} \mathcal{F}_{DEF}, \\ \mathcal{G}^{AB} &\equiv -4 \bar{P}^{C[A} \mathcal{D}^{B]} \mathcal{F}_C + 2 (\mathcal{F}_C - \mathcal{D}_C) \check{\mathcal{F}}^{C[AB]} - 2 \check{\mathcal{F}}^{CD[A} \mathcal{F}_{CD}{}^{B]}. \end{aligned} \quad (2.22)$$

Here, we have defined

$$\begin{aligned} (\eta_{AB}) &\equiv \begin{pmatrix} 0 & \delta_a^b \\ \delta_b^a & 0 \end{pmatrix}, & (\eta^{AB}) &\equiv \begin{pmatrix} 0 & \delta_b^a \\ \delta_a^b & 0 \end{pmatrix}, & \check{\mathcal{F}}^{ABC} &\equiv \bar{P}^{ABCDEF} \mathcal{F}_{DEF}, \\ P_{AB} &\equiv \frac{1}{2} (\eta_{AB} + \hat{\mathcal{H}}_{AB}), & \bar{P}_{AB} &\equiv \frac{1}{2} (\eta_{AB} - \hat{\mathcal{H}}_{AB}), \\ \bar{P}^{ABCDEF} &\equiv \bar{P}^{AD} \bar{P}^{BE} \bar{P}^{CF} + P^{AD} \bar{P}^{BE} \bar{P}^{CF} + \bar{P}^{AD} P^{BE} \bar{P}^{CF} + \bar{P}^{AD} \bar{P}^{BE} P^{CF} \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{4} (\hat{\mathcal{H}}^{AD} \hat{\mathcal{H}}^{BE} \hat{\mathcal{H}}^{CF} - \hat{\mathcal{H}}^{AD} \eta^{BE} \eta^{CF} - \eta^{AD} \hat{\mathcal{H}}^{BE} \eta^{CF} - \eta^{AD} \eta^{BE} \hat{\mathcal{H}}^{CF}) \\
 &\quad + \frac{1}{2} \eta^{AD} \eta^{BE} \eta^{CF}, \tag{2.23}
 \end{aligned}$$

and the indices A, B are raised or lowered with η_{AB} and η^{AB} . Under the section condition we can check that $\mathcal{R} = \mathcal{S}$. The equivalence between $\mathcal{S}_{MN} = 0$ and $\mathcal{G}^{AB} = 0$ is slightly more non-trivial, but it is concisely explained in Ref. [134] (see also Appendix B).

In the flux formulation of DFT [134], we take the untwisted metric $\hat{\mathcal{H}}_{AB}$ as a diagonal Minkowski metric, and then $E_A^M \equiv U_A^M$ is regarded as the generalized vielbein. The fundamental fields are E_M^A and d , and the equations of motion in Eq. (2.21) can be derived from

$$\mathcal{L} = e^{-2d} \mathcal{R}. \tag{2.24}$$

On the other hand, in this paper we rather interpret Eq. (2.19) as a reduction ansatz and the equations of motion in Eq. (2.21) are just rewritings of Eq. (2.14), similar to the gauged DFT [131–133]. For our purpose, it is enough to consider the cases where the generalized fluxes are constant. In that case, the equations of motion are simple algebraic equations,

$$\mathcal{R} = \frac{1}{12} \mathcal{F}_{ABC} \mathcal{F}_{DEF} (3 \hat{\mathcal{H}}^{AD} \eta^{BE} \eta^{CF} - \hat{\mathcal{H}}^{AD} \hat{\mathcal{H}}^{BE} \hat{\mathcal{H}}^{CF}) - \hat{\mathcal{H}}^{AB} \mathcal{F}_A \mathcal{F}_B = 0, \tag{2.25}$$

$$\mathcal{G}^{AB} = \frac{1}{2} (\eta^{CE} \eta^{DF} - \hat{\mathcal{H}}^{CE} \hat{\mathcal{H}}^{DF}) \hat{\mathcal{H}}^{G[A} \mathcal{F}_{CD}{}^{B]} \mathcal{F}_{EFG} + 2 \mathcal{F}_D \check{\mathcal{F}}^{D[AB]} = 0, \tag{2.26}$$

where we have again used the section condition.

In general, the untwisted metric and the DFT dilaton may depend on the coordinates y^μ on the uncompactified external spacetime. In this case, we denote the extended coordinates as $(x^M) = (y^\mu, x^i, \tilde{y}_\mu, \tilde{x}_i)$ and consider

$$\mathcal{H}_{MN} = [U(x^I) \hat{\mathcal{H}}(y^\mu) U^\top(x^I)]_{MN}, \quad d = \hat{d}(y^\mu) + \mathbf{d}(x^I), \tag{2.27}$$

where $(x^I) \equiv (x^i, \tilde{x}_i)$. By following Ref. [133], we assume that $\hat{\mathcal{H}}_{AB}(y)$ and $\hat{d}(y)$ satisfy

$$\mathcal{D}_A \hat{\mathcal{H}}_{BC}(y) = \partial_A \hat{\mathcal{H}}_{BC}(y), \quad \mathcal{D}_A \hat{d}(y) = \partial_A \hat{d}(y) \quad (\partial_A \equiv \delta_A^M \partial_M), \tag{2.28}$$

and then the generalized Ricci scalar (under the section condition) becomes [133] [see Eq. (B.18)]

$$\begin{aligned}
 \mathcal{S} &= \hat{\mathcal{S}} + \frac{1}{12} \mathcal{F}_{ABC} \mathcal{F}_{DEF} (3 \hat{\mathcal{H}}^{AD} \eta^{BE} \eta^{CF} - \hat{\mathcal{H}}^{AD} \hat{\mathcal{H}}^{BE} \hat{\mathcal{H}}^{CF}) - \hat{\mathcal{H}}^{AB} \mathcal{F}_A \mathcal{F}_B \\
 &\quad - \frac{1}{2} \mathcal{F}^A{}_{BC} \hat{\mathcal{H}}^{BD} \hat{\mathcal{H}}^{CE} \mathcal{D}_D \hat{\mathcal{H}}_{AE} + 2 \mathcal{F}_A \mathcal{D}_B \hat{\mathcal{H}}^{AB} - 4 \mathcal{F}_A \hat{\mathcal{H}}^{AB} \mathcal{D}_B \hat{d}, \tag{2.29}
 \end{aligned}$$

where $\hat{\mathcal{S}}$ denotes the generalized Ricci scalar associated with $\{\hat{\mathcal{H}}_{AB}, \hat{d}\}$, and the fluxes \mathcal{F}_A and \mathcal{F}_{ABC} are now made of $\{U_M^A(x), \mathbf{d}(x)\}$. It is important to note that the equation of motion $\mathcal{S} = 0$ is invariant under a constant $O(D, D)$ rotation

$$\hat{\mathcal{H}}_{AB} \rightarrow (C \hat{\mathcal{H}} C^\top)_{AB}, \quad U_A^M \rightarrow C_A^B U_B^M, \tag{2.30}$$

which also transforms the generalized fluxes covariantly. This transformation looks similar to the PL T -plurality discussed in Sect. 6, but they are totally different transformations since Eq. (2.30) does not change \mathcal{H}_{MN} while the PL T -plurality changes \mathcal{H}_{MN} .

2.3. GSE from DFT

As already explained, if we choose a section $\tilde{\delta}^m = 0$, the DFT equations of motion reproduce the usual supergravity equations of motion. On the other hand, we can derive the GSE by choosing another solution of the section condition [149,151],

$$\mathcal{H}_{MN} = \mathcal{H}_{MN}(x^m), \quad d = d_0(x^m) + I^m \tilde{x}_m \quad (I^m \text{ a constant}), \quad (2.31)$$

where the DFT dilaton has a linear dependence on the dual coordinates. In order to satisfy the section condition, we require the vector field I^m to satisfy

$$\hat{\mathfrak{L}}_X \mathcal{H}_{MN} = \hat{\mathfrak{L}}_X d = 0, \quad (\mathbf{X}^M) \equiv \begin{pmatrix} I^m \\ 0 \end{pmatrix}, \quad (2.32)$$

which are equivalent to

$$\mathbf{X}^P \partial_P \mathcal{H}_{MN} = \mathbf{X}^P \partial_P d = \mathbf{X}^P \partial_P d_0 = 0, \quad (2.33)$$

and indeed ensure the section condition,

$$\eta^{MN} \partial_M \mathcal{H}_{PQ} \partial_N d = \mathbf{X}^P \partial_P \mathcal{H}_{MN} = 0, \quad \eta^{MN} \partial_M d \partial_N d = 2 \mathbf{X}^P \partial_P d_0 = 0. \quad (2.34)$$

If we make the ansatz in Eq. (2.31) and parameterize \mathcal{H}_{MN} as usual in terms of $\{g_{mn}, B_{mn}\}$ and d_0 as $e^{-2d_0} = e^{-2\Phi} \sqrt{|g|}$, the DFT equations of motion (without R–R fields) become

$$\begin{aligned} R + 4D^m \partial_m \Phi - 4|\partial\Phi|^2 - \frac{1}{2}|H_3|^2 - 4(I^m I_m + U^m U_m + 2U^m \partial_m \Phi - D_m U^m) &= 0, \\ R_{mn} - \frac{1}{4} H_{mpq} H_n{}^{pq} + 2D_m \partial_n \Phi + D_m U_n + D_n U_m &= 0, \\ -\frac{1}{2} D^p H_{pnm} + \partial_p \Phi H^p{}_{mn} + U^p H_{pnm} + D_m I_n - D_n I_m &= 0, \end{aligned} \quad (2.35)$$

where $U_m \equiv I^n B_{nm}$. They are precisely the GSE studied in Refs. [145–147]. When $I^m = 0$ (where the Killing equations are trivial), they reduce to the usual supergravity equations of motion.

Another way to derive the GSE is to make the modification

$$\partial_M d \rightarrow \partial_M d + \mathbf{X}_M \quad (\mathbf{X}_M \text{ a generalized vector}) \quad (2.36)$$

everywhere in the DFT equations of motion [151]. As long as \mathbf{X}^M satisfies

$$\hat{\mathfrak{L}}_X \mathcal{H}_{MN} = \hat{\mathfrak{L}}_X d = 0, \quad \eta_{MN} \mathbf{X}^M \mathbf{X}^N = 0, \quad (2.37)$$

we can choose a gauge such that \mathbf{X}^M takes the form in Eq. (2.32) [149]. In terms of the generalized flux, obviously this modification corresponds to

$$\mathcal{F}_A \rightarrow \mathcal{F}_A + 2\mathbf{X}_A \quad (\mathbf{X}_A \equiv U_A{}^M \mathbf{X}_M). \quad (2.38)$$

Even in the presence of the R–R fields, this replacement is enough to derive the type II GSE, although we additionally need to require the isometry condition for the R–R fields,

$$\hat{\mathfrak{L}}_I F = 0. \quad (2.39)$$

2.4. A formal T -duality

In generalized backgrounds, where the supergravity fields satisfy the GSE, string theory may not have conformal symmetry. Accordingly, when we obtain a generalized background as a result of NATD, it is usually regarded as a problematic example and such backgrounds have not been considered seriously. However, as discussed in Refs. [145,149–151], by performing a formal T -duality we can always transform a generalized background to a linear-dilaton solution of the usual supergravity. Here, we review what the formal T -duality is.

The DFT equations of motion are covariant under a constant $O(D, D)$ transformation,

$$x^M \rightarrow \Lambda^M_N x^N, \quad \mathcal{H}_{MN} \rightarrow (\Lambda \mathcal{H} \Lambda^\top)_{MN}, \quad \partial_M d \rightarrow \partial_M d. \quad (2.40)$$

In particular, if we consider an $O(D, D)$ matrix,

$$\Lambda = \begin{pmatrix} \mathbf{1} - e_z & e_z \\ e_z & \mathbf{1} - e_z \end{pmatrix}, \quad e_z \equiv \text{diag}(0, \dots, 0, \underbrace{1}_{x^z}, 0, \dots, 0), \quad (2.41)$$

it corresponds to the (factorized) T -duality along the x^z -direction. For a given GSE solution with $d = d_0 + I^z \tilde{x}_z$, the $O(D, D)$ rotation in Eq. (2.40) with Eq. (2.41) exchanges the coordinates x^z and \tilde{x}_z , and the dilaton becomes $d = d_0 + I^z x^z$. According to the Killing equation, the generalized metric and d_0 are independent of x^z , and the dual coordinate \tilde{x}_z does not appear in the resulting background. This means that the GSE background is transformed to a solution of the usual supergravity with a linear dilaton $d = d_0 + I^z x^z$.

The reason we call this $O(D, D)$ transformation a “formal” T -duality is as follows. The usual Abelian T -duality in the presence of D Abelian isometries is an $O(D, D)$ transformation,

$$\mathcal{H}_{MN} \rightarrow \Lambda_M^P \Lambda_N^Q \mathcal{H}_{PQ}, \quad \partial_M d \rightarrow \partial_M d. \quad (2.42)$$

The difference from Eq. (2.40) is whether the coordinates are transformed or not. If we transform the coordinates, Eq. (2.40) is always a symmetry of the DFT equations of motion even without isometries. In the presence of Abelian isometries, due to the coordinate independence, the transformation $x^M \rightarrow \Lambda^M_N x^N$ is trivial and the formal T -duality reduces to the usual T -duality of Eq. (2.42). To stress the difference, when we perform the transformation in Eq. (2.40) with Eq. (2.41) along a non-isometric direction, we call it a formal T -duality.

3. Non-Abelian T -duality

In this section we study the traditional NATD in general curved backgrounds. We begin with a review of NATD for the NS–NS sector. We then describe the duality as a kind of local $O(D, D)$ rotation and provide the general transformation rule for the R–R fields by employing the results of DFT. To provide a closed-form expression for the duality rule, we restrict our discussion to the case where we can take a simple gauge choice, $x^i(\sigma) = \text{const}$.

3.1. NS–NS sector

In the case of the Abelian T -duality, the dual action is obtained with the procedure of Refs. [8,11]. When a target space has a set of Killing vector fields v_a^m that commute with each other, $[v_a, v_b] = 0$, the sigma model has a global symmetry generated by $x^m(\sigma) \rightarrow x^m(\sigma) + \epsilon^a v_a^m(\sigma)$. This global symmetry can be made a local symmetry by introducing gauge fields $A^a(\sigma)$ and replacing $dx^m \rightarrow Dx^m \equiv dx^m - A^a v_a^m$. We also introduce the Lagrange multipliers $\tilde{x}_a(\sigma)$, which constrain the field

strengths to vanish. Then, by integrating out the gauge fields A^a we obtain the dual action, where the Lagrange multiplier \tilde{x}_a becomes the embedding function in the dual geometry. In Ref. [19], this procedure was generalized to the case of non-commuting Killing vectors. It was further developed later, and in the following we review NATD in a general setup as discussed in Refs. [25,35].

We consider a target space with n generalized Killing vectors V_a ($a = 1, \dots, n$) satisfying

$$\hat{\mathfrak{L}}_{V_a} \mathcal{H}_{MN} = 0, \quad [V_a, V_b]_C = f_{ab}{}^c V_c, \quad \eta_{MN} V_a^M V_b^N = 2 c_{ab}, \quad f_{ab}{}^d c_{dc} = 0. \quad (3.1)$$

Here, c_{ab} is a constant symmetric matrix. If we choose a section $\tilde{\delta}^m = 0$ and parameterize the generalized Killing vectors as

$$(V_a^M) \equiv \begin{pmatrix} v_a^m \\ \tilde{v}_{am} \end{pmatrix} \equiv \begin{pmatrix} v_a^m \\ \hat{v}_{am} + B_{mn} v_a^n \end{pmatrix}, \quad (3.2)$$

these conditions reduce to²

$$\begin{aligned} \mathfrak{L}_{v_a} g_{mn} = 0, \quad \iota_{v_a} H_3 + d\hat{v}_a = 0, \quad v_{(a} \cdot \hat{v}_{b)} = c_{ab}, \\ \mathfrak{L}_{v_a} v_b = f_{ab}{}^c v_c, \quad \mathfrak{L}_{v_a} \hat{v}_b = f_{ab}{}^c \hat{v}_c, \quad f_{ab}{}^d c_{dc} = 0, \end{aligned} \quad (3.3)$$

where the dot denotes a contraction of the index m . They are precisely the requirements to perform NATD [25,35] (see Refs. [154,155] for the origin of the conditions).

Under the setup, we consider the gauged action by following the standard procedure [8,11]. Ignoring the dilaton term, the gauged action takes the form [25,35,154,155]

$$\begin{aligned} S \equiv \frac{1}{4\pi\alpha'} \int_{\Sigma} (g_{mn} Dx^m \wedge *Dx^n - 2A^a \wedge \hat{v}_a + B_{ab} A^a \wedge A^b) + \frac{1}{2\pi\alpha'} \int_{\mathcal{B}} H_3 \\ + \frac{1}{4\pi\alpha'} \int_{\Sigma} (2A^a \wedge d\tilde{x}_a + f_{ab}{}^c \tilde{x}_c A^a \wedge A^b) \quad (\partial\mathcal{B} = \Sigma), \end{aligned} \quad (3.4)$$

where we have introduced gauge fields $A^a(\sigma) \equiv A^a_a(\sigma) d\sigma^a$ ($a = 0, 1$) and have defined

$$D_a x^m \equiv \partial_a x^m - A_a^b v_b^m, \quad F^a \equiv dA^a + \frac{1}{2} f_{bc}{}^a A^b \wedge A^c, \quad B_{ab} \equiv \hat{v}_{[a} \cdot v_{b]}. \quad (3.5)$$

Under the conditions in Eq. (3.1), this action is invariant under the local symmetry,

$$\begin{aligned} \delta_{\epsilon} x^m(\sigma) = \epsilon^a(\sigma) v_a^m(x), \quad \delta_{\epsilon} A^a(\sigma) = d\epsilon^a(\sigma) + f_{bc}{}^a A^b(\sigma) \epsilon^c(\sigma), \\ \delta_{\epsilon} \tilde{x}_a(\sigma) = c_{ab} \epsilon^b(\sigma) - f_{ab}{}^c \epsilon^b(\sigma) \tilde{x}_c(\sigma). \end{aligned} \quad (3.6)$$

If we first use the equations of motion for the Lagrange multipliers \tilde{x}_a , the field strengths F^a are constrained to vanish and the gauge fields will become a pure gauge. Then, at least locally, we can choose a gauge $A^a = 0$ and the original theory will be recovered,

$$S_0 = \frac{1}{4\pi\alpha'} \int_{\Sigma} g_{mn} dx^m \wedge *dx^n + \frac{1}{2\pi\alpha'} \int_{\mathcal{B}} H_3. \quad (3.7)$$

² We can easily show that $f_{ca}{}^d c_{db} + f_{cb}{}^d c_{da} = 0$, and then the last condition can be expressed as $f_{c[a}{}^d c_{b]d} = 0$. We can further rewrite the same condition as $\frac{1}{3} \iota_{v_a} \iota_{v_b} \iota_{v_c} H_3 + \iota_{v_{[a}} f_{bc]}{}^d \hat{v}_d = 0$, which was used in Ref. [35].

On the other hand, by using the equations of motion for A^a first, we obtain the dual model. For this purpose, it is convenient to rewrite the action as

$$S = \frac{1}{4\pi\alpha'} \int_{\Sigma} g_{mn} dx^m \wedge * dx^n + \frac{1}{2\pi\alpha'} \int_{\Sigma} B_2 + \frac{1}{4\pi\alpha'} \int_{\Sigma} [2A^a \wedge \nu_a + g_{ab} A^a \wedge *A^b + (B_{ab} + f_{ab}{}^c \tilde{x}_c) A^a \wedge A^b], \quad (3.8)$$

where

$$\nu_a \equiv d\tilde{x}_a - v_a^m g_{mn} * dx^n - \hat{\nu}_a, \quad g_{ab} \equiv g_{mn} v_a^m v_b^n. \quad (3.9)$$

Then, the equations of motion for A^a become³

$$\nu_a = -g_{ab} * A^b - (B_{ab} + f_{ab}{}^c \tilde{x}_c) A^b, \quad (3.10)$$

and this can be solved for A^a as

$$A^a = -N^{(ab)} * \nu_b - N^{[ab]} \nu_b, \quad (3.11)$$

where we have defined

$$(N^{ab}) \equiv (E_{ab} + f_{ab}{}^c \tilde{x}_c)^{-1}. \quad (3.12)$$

After eliminating the gauge fields, the action becomes

$$S = \frac{1}{4\pi\alpha'} \int_{\Sigma} (g_{mn} dx^m \wedge * dx^n + B_{mn} dx^m \wedge dx^n + N^{(ab)} \nu_a \wedge * \nu_b + N^{[ab]} \nu_a \wedge \nu_b) = -\frac{1}{4\pi\alpha'} \int_{\Sigma} d^2\sigma \sqrt{-\gamma} (\gamma^{ab} - \varepsilon^{ab}) (E_{mn} \partial_a x^m \partial_b x^n + N^{ab} \nu_{aa} \nu_{bb}), \quad (3.13)$$

where $E_{mn} \equiv g_{mn} + B_{mn}$. In the above computation, we have assumed that the matrix $(E_{ab} + f_{ab}{}^c \tilde{x}_c)$ is invertible,⁴ but other than that the computation is general.

Now, a major difference from the Abelian case appears. In the Abelian case, by choosing the adapted coordinates $v_a^m = \delta_a^m$ we can always realize a gauge $x^a(\sigma) = 0$. However, in the non-Abelian case, such a gauge choice is not always possible since we cannot realize $v_a^m = \delta_a^m$. In order to provide a closed-form expression for the duality transformation rule, in this paper we assume that the gauge symmetries can be fixed as $x^i(\sigma) = c^i$ (c^i constant) under a suitable decomposition of spacetime coordinates $(x^m) = (y^\mu, x^i)$. This gauge choice removes n coordinates x^i and instead introduces n dual coordinates \tilde{x}_a . Then, the situation is the same as the Abelian case.

Under the gauge choice $x^i(\sigma) = c^i$, the action in Eq. (3.13) reproduces the dual action for the dual coordinates $x'^m = (y^\mu, \tilde{x}_a)$,

$$S = -\frac{1}{4\pi\alpha'} \int_{\Sigma} d^2\sigma \sqrt{-\gamma} (\gamma^{ab} - \varepsilon^{ab}) E'_{mn} \partial_a x'^m \partial_b x'^n, \quad (3.14)$$

³ They can also be expressed as

$$d\tilde{x}_a - (c_{ab} - f_{ab}{}^c \tilde{x}_c) A^b = v_a^m (g_{mn} * Dx^n + B_{mn} Dx^n) + \tilde{\nu}_{am} Dx^m,$$

and reduce to the standard self-duality relation when $\tilde{\nu}_a = 0$ and $f_{ab}{}^c = 0$.

⁴ Note that the invertibility is not ensured even in the Abelian case $f_{ab}{}^c = 0$.

$$(E'_{mn}) \equiv \left(\begin{array}{cc} E_{\mu\nu} - (v_{a\mu} - \hat{v}_{a\mu}) N^{ab} (v_{bv} + \hat{v}_{bv}) & (v_{c\mu} - \hat{v}_{c\mu}) N^{cb} \\ -N^{ac} (v_{cv} + \hat{v}_{cv}) & N^{ab} \end{array} \right) \Big|_{x^i=c^i}. \tag{3.15}$$

Then, the NATD can be understood as a transformation of the target space geometry,

$$E_{mn} \rightarrow E'_{mn}. \tag{3.16}$$

Regarding the transformation rule for the dilaton, we employ the result of Ref. [19],

$$e^{-2\Phi'} = \frac{1}{|\det(N^{ab})|} e^{-2\Phi}. \tag{3.17}$$

3.2. NATD as $O(D, D)$ transformation

In order to show a general transformation rule for the R–R fields, it is convenient to describe NATD as $O(D, D)$ rotations. Starting with the original background,

$$(E_{mn}) = \begin{pmatrix} E_{\mu\nu} & E_{\mu j} \\ E_{i\nu} & E_{ij} \end{pmatrix}, \tag{3.18}$$

we construct the dual background of Eq. (3.15) through the following three steps.

(1) We first perform a $GL(D)$ transformation,

$$E \rightarrow E^{(1)} = \Lambda_\nu E \Lambda_\nu^\top, \quad \Lambda_\nu \equiv \begin{pmatrix} \delta_\mu^\nu & 0 \\ v_a^\nu & v_a^j \end{pmatrix}. \tag{3.19}$$

As we have assumed, we can fix the gauge symmetry $\delta_\epsilon x^i = \epsilon^a v_a^i$ such that $x^i(\sigma) = c^i$ is realized. For this to be possible, $\det(v_a^i) \neq 0$ should be satisfied and the $GL(D)$ matrix Λ_ν is invertible. We then obtain

$$E^{(1)} = \begin{pmatrix} E_{\mu\nu} & E_{\mu n} v_b^n \\ v_a^m E_{m\nu} & v_a^m v_b^n E_{mn} \end{pmatrix} = \begin{pmatrix} E_{\mu\nu} & (v_{b\mu} - \hat{v}_{b\mu}) + \tilde{v}_{b\mu} \\ (v_{a\nu} + \hat{v}_{a\nu}) - \tilde{v}_{a\nu} & E_{ab} + v_{[a} \cdot \tilde{v}_{b]} \end{pmatrix}, \tag{3.20}$$

where we have used

$$v_a^m B_{mv} = \hat{v}_{av} - \tilde{v}_{av}, \quad B_{ab} = \hat{v}_{[a} \cdot v_{b]}. \tag{3.21}$$

(2) We next perform a B -transformation,

$$E^{(1)} \rightarrow E^{(2)} \equiv E^{(1)} + \Lambda_f, \quad \Lambda_f \equiv \begin{pmatrix} 0 & -\tilde{v}_{b\mu} \\ \tilde{v}_{a\nu} & f_{ab}{}^c \tilde{x}_c - v_{[a} \cdot \tilde{v}_{b]} \end{pmatrix}, \tag{3.22}$$

and obtain

$$E^{(2)} = \begin{pmatrix} E_{\mu\nu} & (v_{b\mu} - \hat{v}_{b\mu}) \\ (v_{a\nu} + \hat{v}_{a\nu}) & E_{ab} + f_{ab}{}^c \tilde{x}_c \end{pmatrix}. \tag{3.23}$$

(3) Finally, we perform a T -duality transformation,

$$E^{(2)} \rightarrow E^{(3)} \equiv (\tilde{\Lambda}_T + \Lambda_T E^{(2)}) (\Lambda_T + \tilde{\Lambda}_T E^{(2)})^{-1},$$

$$\Lambda_T \equiv \begin{pmatrix} \mathbf{1}_{d-n} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix}, \quad \tilde{\Lambda}_T \equiv \begin{pmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{1}_n \end{pmatrix}, \tag{3.24}$$

and obtain

$$\begin{aligned}
 E^{(3)} &= \begin{pmatrix} E_{\mu\rho} & (v_{c\mu} - \hat{v}_{c\mu}) \\ 0 & \mathbf{1} \end{pmatrix} \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ (v_{cv} + \hat{v}_{cv}) & E_{cb} + f_{cb}{}^d \tilde{x}_d \end{pmatrix}^{-1} \\
 &= \begin{pmatrix} E_{\mu\nu} - (v_{a\mu} - \hat{v}_{a\mu}) N^{ab} (v_{bv} + \hat{v}_{bv}) & (v_{c\mu} - \hat{v}_{c\mu}) N^{cb} \\ -N^{ac} (v_{cv} + \hat{v}_{cv}) & N^{ab} \end{pmatrix}. \tag{3.25}
 \end{aligned}$$

By choosing the gauge $x^i = c^i$, this precisely reproduces the dual background of Eq. (3.15).

Of course, each step is not a symmetry of supergravity, but this decomposition is useful when we determine the transformation rule of the R–R fields. In terms of the generalized metric \mathcal{H}_{MN} , the above NATD is expressed as a local $O(D, D)$ transformation,

$$\begin{aligned}
 \mathcal{H}_{MN} &\rightarrow \mathcal{H}'_{MN} = (h \mathcal{H} h^T)_{MN} \Big|_{x^i=c^i}, \\
 (h_M{}^N) &\equiv \begin{pmatrix} \Lambda_\top & \tilde{\Lambda}_\top \\ \tilde{\Lambda}_\top & \Lambda_\top \end{pmatrix} \begin{pmatrix} \mathbf{1} & \Lambda_f \\ 0 & \mathbf{1} \end{pmatrix} \begin{pmatrix} \Lambda_\nu & 0 \\ 0 & (\Lambda_\nu)^{-\top} \end{pmatrix}, \tag{3.26}
 \end{aligned}$$

and the $O(D, D)$ matrix $h_M{}^N$ can be straightforwardly constructed from the given set of generalized Killing vectors $V_a = (v_a^m, \tilde{v}_{am})$.

Under a general $O(D, D)$ rotation,

$$\begin{aligned}
 \mathcal{H}_{MN} &\rightarrow \mathcal{H}'_{MN} = (h \mathcal{H} h^T)_{MN}, \quad h_M{}^N \equiv \begin{pmatrix} p_m{}^n & q_{mn} \\ r^{mn} & s^m{}_n \end{pmatrix}, \\
 E_{mn} &\rightarrow E'_{mn} = [(q + p E) (s + r E)^{-1}]_{mn} = [(s^\top - E r^\top)^{-1} (-q^\top + E p^\top)]_{mn}, \tag{3.27}
 \end{aligned}$$

the determinant of the metric transforms as (see, for example, Ref. [156])

$$\sqrt{|g|} \rightarrow \sqrt{|g'|} = |\det(s + r E)|^{-1} \sqrt{|g|}. \tag{3.28}$$

Therefore, under the NATD of Eq. (3.26) we obtain

$$\begin{aligned}
 \sqrt{|g'|} &= |\det(\Lambda_\top + \tilde{\Lambda}_\top E^{(2)})|^{-1} |\det(\Lambda_\nu)| \sqrt{|g|} \Big|_{x^i=c^i} \\
 &= |\det(N^{ab})| |\det(v_a^i)| \sqrt{|g|} \Big|_{x^i=c^i}. \tag{3.29}
 \end{aligned}$$

Combining this with Eq. (3.17), we obtain the transformation rule for the DFT dilaton:

$$e^{-2d'} = |\det(v_a^i)| e^{-2d} \Big|_{x^i=c^i}. \tag{3.30}$$

This shows that the DFT dilaton e^{-2d} transforms covariantly under the $O(D, D)$ rotation.

3.3. R–R sector

Since the NS–NS fields are transformed covariantly under NATD, it is natural to expect that the R–R fields are also transformed covariantly under the same $O(D, D)$ rotation. Indeed, as we see from many examples, under NATD $\mathcal{H}_{MN} \rightarrow \mathcal{H}'_{MN} = (h \mathcal{H} h^T)_{MN} \Big|_{x^i=c^i}$, the generalized Ricci tensors are always transformed covariantly,

$$\mathcal{S}'_{MN} = (h \mathcal{S} h^T)_{MN} \Big|_{x^i=c^i}, \quad \mathcal{S}' = \mathcal{S} \Big|_{x^i=c^i}. \tag{3.31}$$

This shows that the R–R fields should also transform covariantly, in order to satisfy the equations of motion of type II DFT (see Appendix B),

$$\mathcal{S}_{MN} = \mathcal{E}_{MN}, \quad \mathcal{S} = 0, \tag{3.32}$$

where \mathcal{E}_{MN} is an $O(D, D)$ -covariant energy–momentum tensor that contains the R–R fields.

In DFT, there are basically two approaches to describe the R–R fields. One treats the R–R fields as an $O(D, D)$ spinor [125], based on the earlier work in Ref. [157], and the other treats them as an $O(D) \times O(D)$ bi-spinor [128], which is based on the approach of Refs. [158,159].

3.3.1. R–R fields as a polyform

We first explain the former because it is simpler. Since the treatment of the $O(D, D)$ spinor can be rephrased in terms of the differential form, here we treat the R–R field strength as the usual polyform (see Appendices A and B for our convention),

$$F = \sum_{p:\text{even/odd}} \frac{1}{p!} F_{m_1 \dots m_p} dx^{m_1} \wedge \dots \wedge dx^{m_p} \quad (\text{type IIA/IIB}). \tag{3.33}$$

Let us summarize the behavior of an $O(D, D)$ spinor in terms of the polyform.

- (1) Under a $GL(D)$ subgroup of $O(D, D)$ transformation,

$$(h_M^N) = \begin{pmatrix} M & 0 \\ 0 & M^{-\top} \end{pmatrix}, \quad M \in GL(D), \tag{3.34}$$

a polyform F transforms as a $GL(D)$ tensor,

$$F' = F^{(M)} \equiv \sum_p \frac{1}{p!} F_{m_1 \dots m_p}^{(M)} dx^{m_1} \wedge \dots \wedge dx^{m_p},$$

$$F_{m_1 \dots m_p}^{(M)} \equiv M_{m_1}^{n_1} \dots M_{m_p}^{n_p} F_{n_1 \dots n_p}. \tag{3.35}$$

- (2) Under the B -transformation,

$$(h_M^N) = \begin{pmatrix} \mathbf{1}_d & \omega \\ 0 & \mathbf{1}_d \end{pmatrix}, \tag{3.36}$$

a polyform F transforms as

$$F' = e^{\omega \wedge} F \equiv F + \omega \wedge F + \frac{1}{2!} \omega \wedge \omega \wedge F + \dots \tag{3.37}$$

- (3) Under the (factorized) T -duality along the x^m -direction, it transforms as

$$F' = F \cdot \top_{x^m}, \quad F \cdot \top_{x^m} \equiv F \wedge d\tilde{x}_m + F \vee dx^m, \tag{3.38}$$

where \tilde{x}_m is the coordinate dual to x^m , and $\vee dx^m$ denotes the interior product acting from the right.

- (4) An arbitrary $O(D, D)$ transformation can be decomposed into the above three types of transformations, but for later convenience we also show that under the β -transformation,

$$(h_M^N) = \begin{pmatrix} \mathbf{1}_d & 0 \\ \chi & \mathbf{1}_d \end{pmatrix}, \tag{3.39}$$

the transformation rule is given by

$$F' = e^{\chi \vee} F \equiv F + \chi \vee F + \frac{1}{2!} \chi \vee \chi \vee F + \dots, \quad \chi \vee F \equiv \frac{1}{2} \chi^{mn} \iota_m \iota_n. \quad (3.40)$$

By using the rules, the general formula for the R–R fields under the NATD of Eq. (3.26) becomes

$$F' = [e^{\Lambda_f \wedge} F^{(\Lambda_f)}] \cdot \mathbb{T}_{y^1} \cdots \mathbb{T}_{y^n} \Big|_{x^i=c^i}, \quad \Lambda_f \equiv \frac{1}{2} (\Lambda_f)_{mn} dx^m \wedge dx^n, \quad (3.41)$$

where the order of $\mathbb{T}_{y^1} \cdots \mathbb{T}_{y^n}$ is not important since the overall sign flip is a trivial symmetry.

Note that the field strength $F = dA$ is known as the field strength in the A-basis [160] (which is sometimes called the Page form). Another definition, $G \equiv dC + H_3 \wedge C$, is known as the C-basis (see Appendix A). In the dual background, G can be obtained as

$$G' = e^{-B_2 \wedge} F'. \quad (3.42)$$

We also note that the approach of Ref. [80] based on the Fourier–Mukai transformation (see also Ref. [96] for an application) will be closely related to the procedure explained here.

3.3.2. R–R fields as a bi-spinor

Next, let us also explain the treatment of the R–R fields as a bi-spinor \mathcal{G}^α_β . Starting with a polyform G and a vielbein e^m_a associated with g_{mn} , we define the flat components as $G_{a_1 \cdots a_p} = e^{m_1}_{a_1} \cdots e^{m_p}_{a_p} G_{m_1 \cdots m_p}$ and then define the bi-spinor \mathcal{G} as

$$\mathcal{G} = \sum_p \frac{e^\Phi}{p!} G_{a_1 \cdots a_p} \gamma^{a_1 \cdots a_p}, \quad (3.43)$$

where $\gamma^{a_1 \cdots a_p} \equiv \gamma^{[a_1} \cdots \gamma^{a_p]}$ and $(\gamma^a)^\alpha_\beta$ is the usual gamma matrix satisfying $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$. According to Refs. [128, 158, 159] (see also Ref. [153]), under a general $O(D, D)$ rotation

$$\mathcal{H}_{MN} \rightarrow (h \mathcal{H} h^\top)_{MN}, \quad h = \begin{pmatrix} p & q \\ r & s \end{pmatrix}, \quad (3.44)$$

the bi-spinor transforms as

$$\mathcal{G} \rightarrow \mathcal{G} \Omega^{-1}, \quad (3.45)$$

where Ω is a spinor representation of the Lorentz transformation Λ^a_b ,

$$\Omega^{-1} \bar{\gamma}^a \Omega = \Lambda^a_b \bar{\gamma}^b, \quad \Lambda^a_b \equiv [e^\top (s + r E)^{-1} (s - r E^\top) e^{-\top}]^a_b, \quad (3.46)$$

and $\bar{\gamma}^a \equiv \gamma^{11} \gamma^a$. In particular, under a T -duality along a (spatial) x^z -direction, we have

$$\Omega = \Omega^{-1} = \frac{e^a_z \gamma_a}{\sqrt{g_{zz}}}. \quad (3.47)$$

When the vielbein e^a_m has a diagonal form, Ω is just the gamma matrix $\Omega = \gamma_z$. The Ω corresponding to the β -transformation

$$h = \begin{pmatrix} 1 & 0 \\ \chi & 1 \end{pmatrix} \quad (3.48)$$

was obtained in Ref. [153] as

$$\Omega = [\det(\mathcal{E}' \mathcal{E})_e^f]^{-\frac{1}{2}} \mathcal{A}(\frac{1}{2} \beta'^{ab} \gamma_{ab}) \mathcal{A}(-\frac{1}{2} \beta^{ab} \gamma_{ab}), \tag{3.49}$$

where \mathcal{A} is similar to an exponential function defined in Ref. [158],

$$\mathcal{A}(\frac{1}{2} \beta^{ab} \gamma_{ab}) \equiv \sum_{p=0}^5 \frac{1}{2^p p!} \beta^{a_1 a_2} \dots \beta^{a_{2p-1} a_{2p}} \gamma_{a_1 \dots a_{2p}}, \tag{3.50}$$

the position of the indices a, b are changed with η_{ab} , and we have also defined

$$\begin{aligned} \mathcal{E}^{ab} &\equiv e^{am} e^{bn} E_{mn}^\top, & \beta^{ab} &\equiv -\mathcal{E}^{[ab]}, & \tilde{e}_m^a &\equiv e_m^b (\mathcal{E}^\top)_b^a, \\ \mathcal{E}'^{ab} &\equiv \tilde{e}_m^a \tilde{e}_n^b (E^{mn} + \chi^{mn}), & \beta'^{ab} &\equiv -\mathcal{E}'^{[ab]}. \end{aligned} \tag{3.51}$$

Now, let us consider the NATD in Eq. (3.26). Since it is not easy to find a general expression for Ω , let us truncate the B -field and restrict ourselves to a simple background,

$$(E_{mn}) = \begin{pmatrix} g_{\mu\nu} & 0 \\ 0 & e_i^a e_j^b \eta_{ab} \end{pmatrix}. \tag{3.52}$$

We also suppose the generalized Killing vectors have simple forms $V_a = v_a^i \partial_i$ ($v_a^i e_j^b = \delta_a^b$). Then, the vielbein e_m^a has the block-diagonal form

$$(e_m^a) = \begin{pmatrix} \hat{e}_\mu^a & 0 \\ 0 & e_i^a \end{pmatrix}, \tag{3.53}$$

and using this, we define the R–R bi-spinor as

$$\mathcal{G} = \sum_p \frac{e^\Phi}{p!} G_{a_1 \dots a_p} \gamma^{a_1 \dots a_p}, \quad G_{a_1 \dots a_p} \equiv e_{a_1}^{m_1} \dots e_{a_p}^{m_p} G_{m_1 \dots m_p}. \tag{3.54}$$

Under the first $GL(D)$ transformation, \mathcal{G} is invariant while the internal part of the vielbein becomes an identity matrix $e_i^a = \delta_i^a$. We then perform the B -transformation and T -dualities, but it is more useful to perform the T -dualities first, because the vielbein is now just an identity matrix. Namely, we rewrite the B -transformation and T -dualities as T -dualities and the β -transformation with parameter $\chi^{ab} \equiv f_{ab}{}^c \tilde{x}_c$,

$$\begin{pmatrix} \Lambda_\top & \tilde{\Lambda}_\top \\ \tilde{\Lambda}_\top & \Lambda_\top \end{pmatrix} \begin{pmatrix} \mathbf{1} & \Lambda_f \\ 0 & \mathbf{1} \end{pmatrix} = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \Lambda_f & \mathbf{1} \end{pmatrix} \begin{pmatrix} \Lambda_\top & \tilde{\Lambda}_\top \\ \tilde{\Lambda}_\top & \Lambda_\top \end{pmatrix}, \quad \Lambda_f = \begin{pmatrix} 0 & 0 \\ 0 & \chi^{ab} \end{pmatrix}. \tag{3.55}$$

Under the T -dualities and the β -transformation, the bi-spinor is transformed as

$$\mathcal{G} \rightarrow \mathcal{G} \Omega^{-1}, \quad \Omega^{-1} = [\det(\delta_c^d + \chi_c^d)]^{-\frac{1}{2}} \mathcal{A}(\frac{1}{2} \chi^{ab} \gamma_{ab}) \prod_{a=1}^n \gamma_a. \tag{3.56}$$

This appears to be consistent with the formula given in Eq. (3.8) of Ref. [73] up to convention.

If we need to consider the spacetime fermions such as the gravitino and the dilatino, they are also transformed by this Ω , and this approach will be important. However, in order to determine the transformation rule for the R–R fields, the first approach will be more useful.

4. Examples without R–R fields

In this section we study examples of NATD without the R–R fields. In the absence of the R–R fields, our setup is basically the same as the standard one. In order to find new solutions, we consider NATD for non-unimodular algebras $f_{ba}{}^b \neq 0$.

As found in Ref. [23], in non-unimodular cases the dual geometry does not solve the supergravity equations of motion. However, as recently found in Ref. [143], the dual geometry is a solution of GSE. Additional examples were discussed in Ref. [144], and there, by using the result of Ref. [28], it was shown that the Killing vector I in GSE is given by a simple formula,

$$I = f_{ba}{}^b \tilde{\partial}^a. \tag{4.1}$$

As we reviewed in Sect. 2, an arbitrary solution of GSE can be regarded as a solution of DFT with linear dual-coordinate dependence. Then, through a formal T -duality in DFT, the GSE solution can be mapped to a solution of the conventional supergravity. In this section we generate new solutions of supergravity by combining the NATD for a non-unimodular algebra and the formal T -duality.

In fact, by allowing for non-unimodular algebras, we can perform a rich variety of NATD. In order to demonstrate that, we consider several non-Abelian T -dualities of a single solution, the $AdS_3 \times S^3 \times T^4$ background with the H -flux.

4.1. $AdS_3 \times S^3 \times T^4$: Example 1

In the first example, we introduce the coordinates as

$$\begin{aligned} ds^2 &= \frac{2 dx^+ dx^- + dz^2}{z^2} + ds_{S^3}^2 + ds_{T^4}^2 & B_2 &= \frac{dx^+ \wedge dx^-}{z^2} + \omega_2, \\ ds_{S^3}^2 &\equiv \frac{1}{4} [d\theta^2 + \sin^2 \theta d\phi^2 + (d\psi + \cos \theta d\phi)^2], & \omega_2 &\equiv -\frac{1}{4} \cos \theta d\phi \wedge d\psi. \end{aligned} \tag{4.2}$$

We then consider the generalized isometries generated by two generalized Killing vectors,

$$\begin{aligned} V_1 &\equiv (v_1, \tilde{v}_1) \equiv \left(-(x^+)^2 \partial_+ + \frac{z^2}{2} \partial_- - x^+ z \partial_z, dx^+ - \frac{x^+}{z} dz \right), \\ V_2 &\equiv (v_2, \tilde{v}_2) \equiv \left(-x^+ \partial_+ - \frac{z}{2} \partial_z, -\frac{1}{2z} dz \right), \end{aligned} \tag{4.3}$$

which satisfy the algebra $[V_1, V_2]_{\mathcal{C}} = V_1$. The structure constant has the non-vanishing trace $f_{b2}{}^b = f_{12}{}^1 = 1$, and the dual background will be a solution of GSE.

The B -field is not isometric along the v_1 direction, $\mathcal{L}_{v_1} B_2 \neq 0$, and the dual component \tilde{v}_1 is necessary to satisfy the generalized Killing equations $\mathcal{L}_{v_1} B_2 + d\tilde{v}_1 = 0$. Moreover, in order to realize $[V_1, V_2]_{\mathcal{C}} = V_1$, the dual component of V_2 is also necessary. In this case, we find

$$(c_{ab}) = \begin{pmatrix} 0 & 0 \\ 0 & \frac{1}{2} \end{pmatrix} \neq 0, \tag{4.4}$$

but the requirement $f_{ab}{}^d c_{dc} = 0$ in Eq. (3.1) is not violated and we can perform the NATD. The gauge symmetry associated with the generalized Killing vector V_2 ,

$$\delta_{\epsilon^2} x^+(\sigma) = \epsilon^2 v_2^+(x) = -\epsilon^2(\sigma) x^+(\sigma), \tag{4.5}$$

can be fixed by realizing $x^+(\sigma) = 1$. Similarly, the gauge symmetry associated with V_1 ,

$$\delta_{\epsilon^1} z(\sigma) = \epsilon^1 v_1^z(x)|_{x^+=1} = -\epsilon^1(\sigma) z(\sigma), \tag{4.6}$$

can be also fixed as $z(\sigma) = 1$.

The AdS parts of the matrices in Eq. (3.26) (before the gauge fixing) become

$$(\Lambda_V) = \begin{pmatrix} -(x^+)^2 & \frac{z^2}{2} & -x^+ z \\ 0 & 1 & 0 \\ -x^+ & 0 & -\frac{z}{2} \end{pmatrix}, \quad (\Lambda_f) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \tilde{x}_+ - \frac{x^+}{2} \\ -\tilde{x}_+ + \frac{x^+}{2} & 0 & 0 \end{pmatrix}, \tag{4.7}$$

and under the gauge $x^+ = 1$ and $z = 1$, the dual background becomes

$$ds'^2 = \frac{d\tilde{x}_+^2 + 2(1 - 4\tilde{x}_+)d\tilde{x}_+ dx^-}{4\tilde{x}_+^2} + \frac{2 dx^- d\tilde{z}}{\tilde{x}_+} + ds_{S^3 \times T^4}^2, \quad e^{-2\Phi'} = \tilde{x}_+^2, \\ B'_2 = \frac{(1 - 4\tilde{x}_+)d\tilde{x}_+ \wedge dx^-}{4\tilde{x}_+^2} - \frac{(d\tilde{x}_+ + dx^-) \wedge d\tilde{z}}{\tilde{x}_+} + \omega_2. \tag{4.8}$$

As expected, this background does not solve the conventional supergravity equations of motion, but instead satisfies the GSE with the Killing vector

$$I' = f_{ab}{}^a \tilde{\partial}^b = \tilde{\partial}^z. \tag{4.9}$$

Interestingly, this geometry is locally the same as the original $AdS_3 \times S^3$ spacetime. Indeed, by changing coordinates as

$$x'^+ \equiv \tilde{z} - \tilde{x}_+ + \frac{1}{4} \ln \tilde{x}_+, \quad x'^- \equiv x^-, \quad z' \equiv \sqrt{\tilde{x}_+}, \tag{4.10}$$

we obtain the expressions

$$ds^2 = \frac{2 dx'^+ dx'^- + dz'^2}{z'^2} + ds_{S^3 \times T^4}^2, \quad e^{-2\Phi} = z'^4, \\ B_2 = \frac{dx'^+ \wedge dx'^-}{z'^2} + \frac{2 dx'^+ \wedge dz'}{z'} + \omega_2, \quad I = \partial'_+. \tag{4.11}$$

In fact, we can find a two-parameter family of solutions,

$$ds^2 = \frac{2 dx^+ dx^- + dz^2}{z^2} + ds_{S^3 \times T^4}^2, \quad e^{-2\Phi} = z^4 c_0 c_1, \\ B_2 = \frac{dx^+ \wedge dx^-}{z^2} + \frac{2 c_1 dx^+ \wedge dz}{z} + \omega_2, \quad I = c_0 \partial_+, \tag{4.12}$$

and NATD maps the original solution $(c_0, c_1) = (0, 0)$ to the dual solution $(c_0, c_1) = (1, 1)$.

The metric in Eq. (4.11) is the same as the original one in Eq. (4.2), and the B -field is also just shifted by a closed form $B_2 \rightarrow B_2 + 2 dx^+ \wedge d \ln z$. The essential difference from the original background is in the dilaton and I^m . We note that, unlike the case of “trivial solutions” [161], we cannot remove the Killing vector I^m in the dual geometry of Eq. (4.11).⁵

⁵ According to Ref. [162], a solution of GSE is a trivial solution (namely, it also satisfies the supergravity equations of motion with $I = 0$) only when $\tilde{K}^m \equiv I^n B_{np} g^{pm}$ satisfies $\mathfrak{L}_{\tilde{K}} g_{mn} = 0$, $\mathfrak{L}_{\tilde{K}} \Phi + (I + \tilde{K})^2 = 0$, and $dI_1 + \iota_{\tilde{K}} H_3 = 0$ ($I_1 \equiv I^m g_{mn} dx^n$), but they are not satisfied here.

It is natural to consider performing a B -field gauge transformation in order to undo the shift in the B -field. However, in the standard GSE (where the only modification is given by the Killing vector I^m), the gauge symmetry for the B -field is already fixed and we cannot perform a B -field gauge transformation. Indeed, if we remove the closed form in the B -field by hand, we find another solution:

$$\begin{aligned}
 ds^2 &= \frac{2 dx^+ dx^- + dz^2}{z^2} + ds_{S^3 \times T^4}^2, & e^{-2\Phi} &= z^4 c_1, \\
 B_2 &= \frac{dx^+ \wedge dx^-}{z^2} + \omega_2, & I &= c_0 \partial_+,
 \end{aligned}
 \tag{4.13}$$

where c_0 is a free parameter and c_1 can take two values, $c_1 = 0$ or $c_1 = 1$. This is an example of the trivial solution and c_0 can be chosen as $c_0 = 0$. Then, we get two $\text{AdS}_3 \times S^3 \times T^4$ solutions of the supergravity with two different dilatons, $c_1 = 0$ and $c_1 = 1$.

For an arbitrary GSE solution, by taking a coordinate system with $I = I^z \partial_z$ we can regard it as a DFT solution with the DFT dilaton $d = d_0 + I^z \tilde{x}_z$ ($e^{-2d_0} \equiv e^{-2\Phi} \sqrt{|g|}$). Then, if we perform a formal T -duality that exchanges \tilde{x}_z with the physical coordinate x^z , we can get a solution of the conventional supergravity where the DFT dilaton is $d = d_0 + I^z x^z$. In the present example, Eq. (4.12), we perform a formal T -duality along the x^+ -direction, and then the DFT dilaton becomes a function of the physical coordinates,

$$e^{-2d} = e^{-2c_0 x^+} z^{4c_0 c_1} \sqrt{\frac{\sin^2 \theta}{64 z^6}}.
 \tag{4.14}$$

Then, the dual-coordinate dependence disappears from the background fields. However, in this case the AdS part of the dualized generalized metric becomes

$$(\mathcal{H}_{MN}) = \begin{pmatrix} g_{mn} - B_{mp} g^{pq} B_{qn} & B_{mp} g^{pn} \\ -g^{mp} B_{pn} & g^{mn} \end{pmatrix} = \left(\begin{array}{ccc|ccc} 0 & 0 & 0 & 1 & z^2 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & \frac{1}{z^2} & -\frac{2c_1}{z} & -2c_1 z & 0 \\ \hline 1 & 0 & -\frac{2c_1}{z} & 4c_1^2 & 0 & 2c_1 z \\ z^2 & -1 & -2c_1 z & 0 & 0 & 0 \\ 0 & 0 & 0 & 2c_1 z & 0 & z^2 \end{array} \right),
 \tag{4.15}$$

and we cannot extract the supergravity fields $\{g_{mn}, B_{mn}, \Phi\}$ from \mathcal{H}_{MN} due to $\det(g^{mn}) = 0$. This type of (genuinely) DFT solution is called the non-Riemannian background [163], and is studied in detail in Refs. [164–167]. Using a parameterization given in Ref. [165], we find that

$$\begin{aligned}
 (\mathcal{H}_{MN}) &= \begin{pmatrix} \delta_m^p & B_{mp} \\ 0 & \delta_p^m \end{pmatrix} \begin{pmatrix} K_{pq} & X_p^1 Y_1^q - \bar{X}_p^{\bar{1}} \bar{Y}_1^q \\ Y_1^p X_q^1 - \bar{Y}_1^{\bar{p}} \bar{X}_q^{\bar{1}} & H^{pq} \end{pmatrix} \begin{pmatrix} \delta_n^q & 0 \\ -B_{qn} & \delta_q^n \end{pmatrix}, \\
 H &= \begin{pmatrix} 4c_1^2 & 0 & 2c_1 z \\ 0 & 0 & 0 \\ 2c_1 z & 0 & z^2 \end{pmatrix}, & K &= \begin{pmatrix} \frac{1}{4c_1^2} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & B &= \begin{pmatrix} 0 & 0 & -\frac{1}{2c_1 z} \\ 0 & 0 & 0 \\ \frac{1}{2c_1 z} & 0 & 0 \end{pmatrix}, \\
 X^1 &= \begin{pmatrix} -\frac{z}{2} \\ 0 \\ c_1 \end{pmatrix}, & \bar{X}^{\bar{1}} &= \begin{pmatrix} -\frac{z}{2} \\ \frac{1}{z} \\ c_1 \end{pmatrix}, & Y_1 &= \begin{pmatrix} 0 \\ -z \\ \frac{1}{c_1} \end{pmatrix}, & \bar{Y}_1 &= \begin{pmatrix} 0 \\ z \\ 0 \end{pmatrix}.
 \end{aligned}
 \tag{4.16}$$

In the parameterization of Ref. [165], there are in general n pairs of vectors (X^i, Y^i) and \tilde{n} pairs of vectors $(\tilde{X}_{\tilde{i}}, \tilde{Y}_{\tilde{i}})$, and such a non-Riemannian background is called a (n, \tilde{n}) solution. In this classification, this background is a $(1, 1)$ solution.

In this way, in the first example of NATD, the formal T -duality does not produce the usual supergravity solution, and we instead obtain a $(1, 1)$ non-Riemannian background.

4.2. $AdS_3 \times S^3 \times T^4$: Example 2

In the second example, we take the coordinates

$$ds^2 = \frac{-dt^2 + dx^2 + dz^2}{z^2} + ds_{S^3 \times T^4}^2, \quad B_2 = \frac{dt \wedge dx}{z^2} + \omega_2, \quad (4.17)$$

and consider the translation and the dilatation generators as the generalized Killing vectors,

$$V_1 \equiv (v_1, \tilde{v}_1) \equiv (\partial_x, 0), \quad V_2 \equiv (v_2, \tilde{v}_2) \equiv (t \partial_t + x \partial_x + z \partial_z, 0), \quad (4.18)$$

which satisfy $[V_1, V_2]_C = V_1$ and $c_{ab} = 0$. Here, we fix the gauge as $x(\sigma) = 0$ and $z(\sigma) = 1$.

The AdS_3 parts of the transformation matrices are

$$(\Lambda_v) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ t & x & z \end{pmatrix}, \quad (\Lambda_f) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \tilde{x} \\ -\tilde{x} & 0 & 0 \end{pmatrix}, \quad (4.19)$$

and the NATD gives

$$ds'^2 = \frac{-\tilde{x}^2 dt^2 + 2(1 - t\tilde{x}) dt dx + (1 - t^2) d\tilde{x}^2 + d\tilde{z}^2}{1 - 2t\tilde{x} + \tilde{x}^2} + ds_{S^3 \times T^4}^2, \\ B'_2 = \frac{[(t - \tilde{x}) d\tilde{x} - \tilde{x} dt] \wedge d\tilde{z}}{1 - 2t\tilde{x} + \tilde{x}^2} + \omega_2, \quad e^{-2\Phi'} = 1 - 2t\tilde{x} + \tilde{x}^2. \quad (4.20)$$

This satisfies the GSE by introducing the Killing vector as $I' = f_{ab}{}^a \tilde{\partial}^b = \tilde{\partial}^{\tilde{z}}$.

Again, in order to remove the Killing vector I , let us perform a formal T -duality along the \tilde{z} -direction. This yields a simple linear-dilaton solution of the supergravity,

$$ds^2 = 2 dt d\tilde{x} + d\tilde{x}^2 - 2\tilde{x} dt dz + 2(t - \tilde{x}) d\tilde{x} dz + (1 - 2t\tilde{x} + \tilde{x}^2) dz^2 + ds_{S^3 \times T^4}^2, \\ B_2 = \omega_2, \quad \Phi = z, \quad (4.21)$$

where the AdS part of the B -field has disappeared.

4.3. $AdS_3 \times S^3 \times T^4$: Example 3

We next use the Rindler-type coordinates,

$$ds^2 = \frac{-x^2 dt^2 + dx^2 + dz^2}{z^2} + ds_{S^3 \times T^4}^2, \quad B_2 = \frac{x dt \wedge dx}{z^2} + \omega_2, \quad (4.22)$$

and consider the generalized Killing vectors

$$V_1 \equiv (v_1, \tilde{v}_1) \equiv (\partial_t, 0), \quad V_2 \equiv (v_2, \tilde{v}_2) \equiv (e^{-t}(x^{-1} \partial_t + \partial_x), 0), \quad (4.23)$$

which satisfy $[V_1, V_2]_C = -V_2$ and $c_{ab} = 0$. Here, we take a gauge $t(\sigma) = 0$ and $x(\sigma) = 1$.

The AdS parts of the transformation matrices are

$$(\Lambda_v) = \begin{pmatrix} 1 & 0 & 0 \\ e^{-t}x^{-1} & e^{-t} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (\Lambda_f) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -\tilde{x} \\ \tilde{x} & 0 & 0 \end{pmatrix}, \quad (4.24)$$

and the dual background, which satisfies the GSE, becomes

$$ds'^2 = \frac{d\tilde{x}^2 - 2d\tilde{t}d\tilde{x}}{\tilde{x}(2 - \tilde{x}z^2)} + \frac{dz^2}{z^2} + ds_{S^3 \times T^4}^2, \quad e^{-2\Phi'} = \frac{\tilde{x}(\tilde{x}z^2 - 2)}{z^2},$$

$$B'_2 = \frac{1 - \tilde{x}z^2}{\tilde{x}(2 - \tilde{x}z^2)} d\tilde{t} \wedge d\tilde{x} + \omega_2, \quad I' = \tilde{\partial}^t. \quad (4.25)$$

In order to obtain a solution of the supergravity, we again perform a formal T -duality along the \tilde{t} -direction. Again, we find a non-Riemannian background,

$$(\mathcal{H}_{MN}) = \left(\begin{array}{ccc|ccc} \tilde{x}(\tilde{x}z^2 - 2) & 1 - \tilde{x}z^2 & 0 & \tilde{x}z^2 - 1 & \tilde{x}(\tilde{x}z^2 - 2) & 0 \\ 1 - \tilde{x}z^2 & z^2 & 0 & -z^2 & 1 - \tilde{x}z^2 & 0 \\ 0 & 0 & \frac{1}{z^2} & 0 & 0 & 0 \\ \hline \tilde{x}z^2 - 1 & -z^2 & 0 & 0 & 0 & 0 \\ \tilde{x}(\tilde{x}z^2 - 2) & 1 - \tilde{x}z^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & z^2 \end{array} \right), \quad (4.26)$$

where the $S^3 \times T^4$ part of the generalized metric is not displayed. This is also a (1, 1) solution,

$$(\mathcal{H}_{MN}) = \begin{pmatrix} \delta_m^p & B_{mp} \\ 0 & \delta_p^m \end{pmatrix} \begin{pmatrix} K_{pq} & X_p^1 Y_1^q - \bar{X}_p^{\bar{1}} \bar{Y}_1^q \\ Y_1^p X_q^1 - \bar{Y}_1^{\bar{p}} \bar{X}_q^{\bar{1}} & H^{pq} \end{pmatrix} \begin{pmatrix} \delta_n^q & 0 \\ -B_{qn} & \delta_q^n \end{pmatrix},$$

$$H = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & z^2 \end{pmatrix}, \quad K = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \frac{1}{z^2} \end{pmatrix}, \quad B = \begin{pmatrix} 0 & -\frac{1}{2} & 0 \\ \frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$X^1 = \begin{pmatrix} \frac{\tilde{x}z^2}{2} \\ -\frac{z^2}{2} \\ 0 \end{pmatrix}, \quad \bar{X}^{\bar{1}} = \begin{pmatrix} \frac{2}{z^2} - \tilde{x} \\ 1 \\ 0 \end{pmatrix}, \quad Y_1 = \begin{pmatrix} 1 \\ \tilde{x} - \frac{2}{z^2} \\ 0 \end{pmatrix}, \quad \bar{Y}_{\bar{1}} = \begin{pmatrix} \frac{z^2}{2} \\ \frac{\tilde{x}z^2}{2} \\ 0 \end{pmatrix}. \quad (4.27)$$

To briefly summarize, NATD works well as a solution-generating technique of DFT even if the isometry algebra is non-unimodular. If we additionally perform a formal T -duality, we usually obtain the usual supergravity solution. Sometimes, the parameterization of the generalized metric becomes singular and we obtain a non-Riemannian background, which does not have the usual supergravity interpretation. However, they are interesting backgrounds by themselves, as discussed in Refs. [164–167]. Therefore, it is important to study NATD for non-unimodular algebras more seriously.

5. Examples with R–R fields

In this section we consider NATD with non-vanishing R–R fields. After reproducing a known example, we again consider examples for non-unimodular algebras.

For convenience, let us display the summary of the duality rules. Under the setup

$$\hat{\mathfrak{L}}_{V_a} \mathcal{H}_{MN} = 0, \quad [V_a, V_b]_C = f_{ab}{}^c V_c, \quad \eta_{MN} V_a^M V_b^N = 2c_{ab}, \quad f_{ab}{}^d c_{dc} = 0, \quad (5.1)$$

where $(V_a^M) = (v_a^m, \tilde{v}_{am})$, the dual background is given by

$$\begin{aligned} \mathcal{H}'_{MN} &= (h \mathcal{H} h^T)_{MN} \Big|_{x^i=c^i}, & e^{-2d'} &= |\det(v_a^i)| e^{-2d} \Big|_{x^i=c^i}, \\ F' &= [e^{\Lambda_f \wedge} F^{(\Lambda_v)}] \cdot \mathbf{T}_{y^1} \cdots \mathbf{T}_{y^n} \Big|_{x^i=c^i}, & I &= f_{ba}{}^b \tilde{\delta}^a, \end{aligned} \tag{5.2}$$

where

$$\begin{aligned} (h_M^N) &\equiv \begin{pmatrix} \Lambda_\top & \tilde{\Lambda}_\top \\ \tilde{\Lambda}_\top & \Lambda_\top \end{pmatrix} \begin{pmatrix} \mathbf{1} & \Lambda_f \\ 0 & \mathbf{1} \end{pmatrix} \begin{pmatrix} \Lambda_v & 0 \\ 0 & (\Lambda_v)^{-\top} \end{pmatrix}, & \Lambda_f &\equiv \frac{1}{2} (\Lambda_f)_{mn} dx^m \wedge dx^n, \\ \Lambda_v &\equiv \begin{pmatrix} \delta_\mu^\nu & 0 \\ v_a^\nu & v_a^j \end{pmatrix}, & \Lambda_f &\equiv \begin{pmatrix} 0 & -\tilde{v}_{b\mu} \\ \tilde{v}_{av} & f_{ab}{}^c \tilde{x}_c - v_{[a} \cdot \tilde{v}_{b]} \end{pmatrix}, & \Lambda_\top &\equiv \begin{pmatrix} \mathbf{1}_{d-n} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix}, & \tilde{\Lambda}_\top &\equiv \begin{pmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{1}_n \end{pmatrix}, \end{aligned} \tag{5.3}$$

and the coordinates are transformed as $(x^m) = (y^\mu, x^i) \rightarrow (x'^m) = (y^\mu, \tilde{x}_a)$.

5.1. $AdS_3 \times S^3 \times T^4$

As the first example of NATD with the R–R fields, let us review the example of Ref. [72] and demonstrate that our formula gives the same result. The original background is

$$\begin{aligned} ds^2 &= \frac{-dt^2 + dx^2 + dz^2}{\ell^2 z^2} + \frac{1}{4\ell^2} [d\theta^2 + \sin^2 \theta d\phi^2 + (d\psi + \cos \theta d\phi)^2] + ds_{T^4}^2, \\ G_3 &= \frac{2 dt \wedge dx \wedge dz}{\ell^2 z^3} - \frac{\sin \theta}{4\ell^2} d\theta \wedge d\phi \wedge d\psi, \end{aligned} \tag{5.4}$$

where the AdS_3 and S^3 part has the curvature $R = \mp 6 \ell^2$.

We perform NATD associated with three generalized Killing vectors on the S^3 ,

$$\begin{aligned} V_1 &= (\cos \psi \partial_\theta + \frac{\sin \psi}{\sin \theta} \partial_\phi - \frac{\sin \psi}{\tan \theta} \partial_\psi, 0), \\ V_2 &= (-\sin \psi \partial_\theta + \frac{\cos \psi}{\sin \theta} \partial_\phi - \frac{\cos \psi}{\tan \theta} \partial_\psi, 0), & V_3 &= (\partial_\psi, 0), \end{aligned} \tag{5.5}$$

which satisfy

$$[V_1, V_2]_{\mathcal{C}} = V_3, \quad [V_2, V_3]_{\mathcal{C}} = V_1, \quad [V_3, V_1]_{\mathcal{C}} = V_2. \tag{5.6}$$

As is clear from the explicit form of the Killing vectors, we can choose a gauge

$$\theta(\sigma) = \frac{\pi}{2}, \quad \phi(\sigma) = 0, \quad \psi(\sigma) = 0. \tag{5.7}$$

The (θ, ϕ, ψ) parts of the transformation matrices are

$$(\Lambda_v) = \begin{pmatrix} \cos \psi & \frac{\sin \psi}{\sin \theta} & -\frac{\sin \psi}{\tan \theta} \\ -\sin \psi & \frac{\cos \psi}{\sin \theta} & -\frac{\cos \psi}{\tan \theta} \\ 0 & 0 & 1 \end{pmatrix}, \quad (\Lambda_f) = \begin{pmatrix} 0 & \tilde{\psi} & -\tilde{\phi} \\ -\tilde{\psi} & 0 & \tilde{\theta} \\ \tilde{\phi} & -\tilde{\theta} & 0 \end{pmatrix}, \tag{5.8}$$

and the NS–NS fields in the dual background are

$$ds'^2 = \frac{-dt^2 + dx^2 + dz^2}{\ell^2 z^2} + \frac{4\ell^2 (\delta_{ij} + 16\ell^4 u_i u_j) du^i du^j}{1 + 16\ell^4 u_k u^k} + ds_{T^4}^2,$$

$$B'_2 = -\frac{8 \ell^4 \epsilon_{ijk} u^i du^j \wedge du^k}{1 + 16 \ell^4 u_k u^k}, \quad e^{-2\Phi'} = \frac{1 + 16 \ell^4 u_k u^k}{64 \ell^6}, \quad (5.9)$$

where we have denoted $(u^i) \equiv (\tilde{\theta}, \tilde{\phi}, \tilde{\psi})$, $u_i \equiv u^i$, and $\epsilon_{123} = 1$.

Now, let us consider the R–R fields. Under the gauge in Eq. (5.7), the Page form becomes

$$F = \left(\frac{2 dt \wedge dx \wedge dz}{\ell^2 z^3} - \frac{d\theta \wedge d\phi \wedge d\psi}{4 \ell^2} \right) \wedge [1 - \text{vol}(T^4)]. \quad (5.10)$$

The first $GL(D)$ transformation is trivial, $\Lambda_\nu = \mathbf{1}$, under the gauge in Eq. (5.7). We next perform the B -transformation $F \rightarrow e^{\Lambda_f} \wedge F$ where

$$\Lambda_f = u^1 d\phi \wedge d\psi + u^2 d\psi \wedge d\theta + u^3 d\theta \wedge d\phi. \quad (5.11)$$

Finally, by performing T -dualities along the (θ, ϕ, ψ) -directions, we obtain

$$\begin{aligned} F' &= (F + \Lambda_f \wedge F) (\wedge du^1 + \vee d\theta) (\wedge du^2 + \vee d\phi) (\wedge du^3 + \vee d\psi) \\ &= \left[\frac{1}{4 \ell^2} - \frac{2 dt \wedge dx \wedge dz \wedge (u_i du^i - du^1 \wedge du^2 \wedge du^3)}{\ell^2 z^3} \right] \wedge [1 - \text{vol}(T^4)]. \end{aligned} \quad (5.12)$$

From this Page form we get the R–R field strengths in the C-basis as

$$\begin{aligned} G'_0 &= \frac{1}{4 \ell^2}, & G'_2 &= \frac{2 \ell^2 \epsilon_{ijk} u^i du^j \wedge du^k}{1 + 16 \ell^4 u_i u^i}, \\ G'_4 &= -\frac{2 dt \wedge dx \wedge dz \wedge u_i du^i}{\ell^2 z^3} - \frac{\text{vol}(T^4)}{4 \ell^2}. \end{aligned} \quad (5.13)$$

These are precisely the solution of the massive type IIA supergravity obtained in Ref. [72].

Since the R–R potential also behaves as an $O(D, D)$ spinor in DFT, let us also explain how to determine the R–R potential in the dual background. Due to the gauge fixing of Eq. (5.7), the Page form takes the form in Eq. (5.10). Then the R–R potential in the A-basis is

$$A = -\left(\frac{dt \wedge dx}{\ell^2 z^2} + \frac{\theta d\phi \wedge d\psi}{4 \ell^2} \right) \wedge [1 - \text{vol}(T^4)], \quad (5.14)$$

where θ should not be set to $\theta = \pi/2$ in order to realize $F = dA$. Similar to the field strength, $GL(D)$ transformation is trivial, and the B -transformation $A \rightarrow e^{\Lambda_f} \wedge A$ and T -dualities along the (θ, ϕ, ψ) -directions give

$$A' = \left[\frac{\tilde{u}_1 du^1}{4 \ell^2} + \frac{dt \wedge dx \wedge (u_i du^i - du^1 \wedge du^2 \wedge du^3)}{\ell^2 z^2} \right] \wedge [1 - \text{vol}(T^4)], \quad (5.15)$$

where we have denoted $\tilde{u}_1 \equiv \theta$ as it is dual to $u^1 = \tilde{\theta}$. Since A depends on the dual coordinate explicitly, the relation between F and A is generalized as [see Eq. (B.38)]

$$F = dA, \quad d \equiv dx^m \wedge \partial_m + \iota_m \tilde{\partial}^m, \quad (5.16)$$

and the A' in Eq. (5.15) correctly reproduces the F' obtained in Eq. (5.12). This result is consistent with Ref. [126], where the massive type IIA supergravity was reproduced from DFT by introducing a linear dual-coordinate dependence into the R–R one-form potential. The potential in the C-basis can also be obtained by computing $C' = e^{-B'_2} \wedge A'$.

5.2. $AdS_5 \times S^5$

As the second example, let us consider a NATD of the $AdS_5 \times S^5$ background associated with a non-unimodular algebra. The original $AdS_5 \times S^5$ background is

$$ds^2 = \frac{\eta_{\mu\nu} dx^\mu dx^\nu + dz^2}{z^2} + ds_{S^5}^2, \quad (\eta_{\mu\nu}) \equiv \text{diag}(-1, 1, 1, 1),$$

$$G = 4(-dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3 \wedge dz + \omega_5), \tag{5.17}$$

where

$$ds_{S^5}^2 \equiv dr^2 + \sin^2 r d\xi^2 + \sin^2 r \cos^2 \xi d\phi_1^2 + \sin^2 r \sin^2 \xi d\phi_2^2 + \cos^2 r d\phi_3^2,$$

$$\omega_5 \equiv \sin^3 r \cos r \sin \xi \cos \xi dr \wedge d\xi \wedge d\phi_1 \wedge d\phi_2 \wedge d\phi_3. \tag{5.18}$$

We consider a NATD associated with two Killing vectors,

$$V_1^M = (z \partial_z + x^\mu \partial_\mu, 0), \quad V_2^M = (\partial_1, 0), \tag{5.19}$$

which satisfy $[V_1, V_2]_C = -V_2$. The gauge symmetry can be fixed as $z(\sigma) = 1$ and $x^1(\sigma) = 0$, and the AdS parts of the transformation matrices are

$$(\Lambda_\nu) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ x^0 & x^1 & x^2 & x^3 & z \end{pmatrix}, \quad (\Lambda_f) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \tilde{x}_1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & -\tilde{x}_1 & 0 & 0 & 0 \end{pmatrix}. \tag{5.20}$$

For simplicity, we denote $(u^\mu) \equiv (x^0, \tilde{x}_1, x^2, x^3)$; then the dual background becomes

$$ds'^2 = \frac{d\tilde{z}^2 + a_{\mu\nu} du^\mu du^\nu}{1 + \eta_{\rho\sigma} u^\rho u^\sigma} + \eta_{\mu\nu} du^\mu du^\nu + ds_{S^5}^2, \quad e^{-2\Phi'} = 1 + \eta_{\mu\nu} u^\mu u^\nu,$$

$$B'_2 = \frac{(-u^0 du^0 - u^1 du^1 + u^2 du^2 + u^3 du^3) \wedge d\tilde{z}}{1 + \eta_{\rho\sigma} u^\rho u^\sigma}, \tag{5.21}$$

where

$$(a_{\mu\nu}) = \begin{pmatrix} -u^0 u^0 & -u^0 u^1 & u^0 u^2 & u^0 u^3 \\ -u^1 u^0 & -u^1 u^1 & u^1 u^2 & u^1 u^3 \\ u^2 u^0 & u^2 u^1 & -u^2 u^2 & -u^2 u^3 \\ u^3 u^0 & u^3 u^1 & -u^3 u^2 & -u^3 u^3 \end{pmatrix}. \tag{5.22}$$

Regarding the R–R fields, the first $GL(D)$ transformation does not change the Page form and the next B -transformation gives

$$F = 4(-du^0 \wedge du^1 \wedge du^2 \wedge du^3 \wedge dz + \omega_5) + 4u^1 \omega_5 \wedge du^1 \wedge dz. \tag{5.23}$$

The Abelian T -dualities along the z and x^1 directions give

$$F' = -4 du^0 \wedge du^2 \wedge du^3 + 4 \omega_5 \wedge d\tilde{z} \wedge du^1 + 4 u^1 \omega_5. \tag{5.24}$$

From this Page form, we find that

$$G'_3 = -4 du^0 \wedge du^2 \wedge du^3, \quad G'_5 = -\frac{4 u^1 du^0 \wedge du^1 \wedge du^2 \wedge du^3 \wedge d\tilde{z}}{1 + \eta_{\mu\nu} u^\mu u^\nu} + u^1 d\omega_4. \quad (5.25)$$

Then, by introducing $I = f_{ba}{}^b \tilde{\partial}^a = \tilde{\partial}^z$, they satisfy the type IIB GSE.

In order to obtain a solution of the usual supergravity, we perform a formal T -duality along the \tilde{z} -direction. By using the T -duality rule in Eq. (A.14), we obtain a simple type IIA solution:

$$ds^2 = (1 + \eta_{\mu\nu} u^\mu u^\nu) dz^2 + 2(-u^0 du^0 - u^1 du^1 + u^2 du^2 + u^3 du^3) dz + \eta_{\mu\nu} du^\mu du^\nu + ds_{S^5}^2, \quad \Phi = z, \quad G_4 = 4e^{-z} dz \wedge du^0 \wedge du^2 \wedge du^3. \quad (5.26)$$

5.3. $AdS_3 \times S^3 \times T^4$ with NS–NS and R–R fluxes

In order to demonstrate the efficiency of our formula, let us consider a more involved example. We start with the $AdS_3 \times S^3 \times T^4$ solution with the NS–NS and the R–R fluxes,

$$ds^2 = \frac{-dt^2 + dx^2 + dz^2}{z^2} + \frac{1}{4} [d\theta^2 + \sin^2 \theta d\phi^2 + (d\psi + \cos \theta d\phi)^2] + ds_{T^4}^2, \\ B_2 = p \left(\frac{dt \wedge dx}{z^2} - \frac{\cos \theta d\phi \wedge d\psi}{4} \right), \quad G_3 = q \left(\frac{2 dt \wedge dx \wedge dz}{z^3} - \frac{\sin \theta d\theta \wedge d\phi \wedge d\psi}{4} \right), \quad (5.27)$$

where p and q are constants satisfying $p^2 + q^2 = 1$. The Page form is

$$F = G_3 + F_5 - (G_3 + F_5) \wedge \text{vol}_{T^4}, \quad F_5 \equiv d \left(\frac{pq \cos \theta}{4z^2} \right) \wedge dt \wedge dx \wedge d\phi \wedge d\psi. \quad (5.28)$$

Then, we consider two generalized Killing vectors,

$$V_1 \equiv (v_1, \tilde{v}_1) \equiv (t \partial_t + x \partial_x + z \partial_z, 0), \\ V_2 \equiv (v_2, \tilde{v}_2) \equiv (-2tx \partial_t + (-t^2 - x^2 + z^2) \partial_x - 2xz \partial_z, 2p dt - \frac{2pt}{z} dz), \quad (5.29)$$

which satisfy $[V_1, V_2]_C = V_2$ and $c_{ab} = 0$. The B -field is isometric along the dilatation generator $\mathcal{L}_{v_1} B_2 = 0$, but it is not isometric along the special-conformal generator $\mathcal{L}_{v_2} B_2 \neq 0$ and the dual component \tilde{v}_2 is important. Here, we choose the gauge as $t(\sigma) = 1$ and $x(\sigma) = 1$.

The AdS parts of the transformation matrices are

$$(\Lambda_v) = \begin{pmatrix} t & x & z \\ -2tx & -t^2 - x^2 + z^2 & -2xz \\ 0 & 0 & 1 \end{pmatrix}, \quad (\Lambda_f) = \begin{pmatrix} 0 & \tilde{x} & 0 \\ -\tilde{x} & 0 & -\frac{2p}{z} \\ \frac{2p}{z} & 0 & 0 \end{pmatrix}, \quad (5.30)$$

and the NS–NS fields and the Killing vector take the form

$$ds'^2 = \frac{z^2 d\tilde{t}^2 + 2 d\tilde{t} d\tilde{x} + d\tilde{x}^2 + \frac{(\tilde{x}-p)^2-1}{z^2} dz^2 - \frac{2}{z} [2\tilde{x} d\tilde{t} + (\tilde{x}-p) d\tilde{x}] dz}{z^2 + (\tilde{x}+p)^2 - 1} + ds_{S^3}^2 + ds_{T^4}^2, \\ B'_2 = \frac{-z(\tilde{x}+p) d\tilde{t} \wedge d\tilde{x} - [z^2 + 2p(\tilde{x}+p) - 2] d\tilde{t} \wedge dz + d\tilde{x} \wedge dz}{z[z^2 + (\tilde{x}+p)^2 - 1]} - \frac{p \cos \theta d\phi \wedge d\psi}{4}, \\ e^{-2\Phi'} = z^2 + (\tilde{x}+p)^2 - 1, \quad I' = -\tilde{\partial}^t. \quad (5.31)$$

For the R–R fields, the first GL(D) transformation makes the replacement

$$dt \wedge dx \wedge dz \rightarrow z^2 dt \wedge dx \wedge dz \tag{5.32}$$

in the Page form of Eq. (5.28), and by further acting $e^{A_f \wedge}$ and $T_t \cdot T_x$, we obtain the Page form in the dual background,

$$\begin{aligned} F'_1 &= -\frac{2q dz}{z}, & F'_3 &= \frac{2q}{z} \left[p dz \wedge \frac{\cos \theta d\phi \wedge d\psi}{4} + z(\tilde{x} + p) \omega_{S^3} \right], \\ F'_5 &= \frac{2q}{z} \left[dz \wedge \text{vol}_{T^4} - (z d\tilde{t} \wedge d\tilde{x} + 2p d\tilde{t} \wedge dz) \wedge \omega_{S^3} \right], \\ F'_7 &= -\frac{2q}{z} \left[p dz \wedge \frac{\cos \theta d\phi \wedge d\psi}{4} + z(\tilde{x} + p) \omega_{S^3} \right] \wedge \text{vol}_{T^4}, \\ F'_9 &= \frac{2q}{z} (z d\tilde{t} \wedge d\tilde{x} + 2p d\tilde{t} \wedge dz) \wedge \omega_{S^3} \wedge \text{vol}_{T^4}, \end{aligned} \tag{5.33}$$

where $\omega_{S^3} \equiv \frac{1}{8} \sin \theta d\theta \wedge d\phi \wedge d\psi$. Finally, the field strength $G' = e^{-B'_2 \wedge} F'$ becomes

$$\begin{aligned} G'_1 &= -\frac{2q dz}{z}, & G'_3 &= 2q(\tilde{x} + p) \left[-\frac{z^{-1} d\tilde{t} \wedge d\tilde{x} \wedge dz}{(\tilde{x} + p)^2 + z^2 - 1} + \omega_{S^3} \right], \\ G'_5 &= 2q \frac{[\tilde{x}(z^2 - 2) - pz^2] d\tilde{t} \wedge dz - (\tilde{x} + p) d\tilde{x} \wedge dz - z(z^2 - 1) d\tilde{t} \wedge d\tilde{x}}{z[(\tilde{x} + p)^2 + z^2 - 1]} \wedge \omega_{S^3} \\ &\quad + \frac{2q dz \wedge \text{vol}_{T^4}}{z}. \end{aligned} \tag{5.34}$$

These satisfy type IIB GSE under the original constraint $p^2 + q^2 = 1$.

By performing a formal T -duality along the \tilde{t} -direction, we obtain

$$\begin{aligned} ds^2 &= \frac{(z^2 + 4p^2 - 4) dz^2}{z^4} + \frac{2[(z^2 + 2p\tilde{x} + 2p^2 - 2) dt + 2p dx] dz}{z^3} \\ &\quad + \frac{(z^2 + \tilde{x}^2 + 2p\tilde{x} + p^2 - 1) dt^2 + 2(\tilde{x} + p) dt d\tilde{x} + d\tilde{x}^2}{z^2} + ds_{S^3}^2 + ds_{T^4}^2, \\ B_2 &= -\frac{dt \wedge d\tilde{x}}{z^2} + \frac{2(x dt + d\tilde{x}) \wedge dz}{z^3} - \frac{p \cos \theta d\phi \wedge d\psi}{4}, & e^{-2\Phi} &= z^2 e^{2t}, \\ G_2 &= \frac{2q e^t dt \wedge dz}{z}, & G_4 &= -\frac{2q e^t [z(\tilde{x} + p) dt + z d\tilde{x} + 2p dz] \wedge \omega_{S^3}}{z}, \end{aligned} \tag{5.35}$$

which is a solution of type IIA supergravity.

5.4. Extremal black D3-brane background

In order to show that the AdS factor is not important, let us consider an extremal black D3-brane background. To manifest the Bianchi type V symmetry we employ a non-standard coordinate system,

$$\begin{aligned} ds^2 &= H^{\frac{1}{2}}(r) \left\{ -dt^2 + t^2 [dx_1^2 + e^{2x^1} (dx_2^2 + dx_3^2)] \right\} + \frac{dr^2}{H^2(r)} \\ &\quad + r^2 (d\theta^2 + \sin^2 \theta d\xi^2 + \sin^2 \theta \cos^2 \xi d\phi_1^2 + \sin^2 \theta \sin^2 \xi d\phi_2^2 + \cos^2 \theta d\phi_3^2), \end{aligned}$$

$$G_5 = -\frac{4r_+^4 t^3 e^{2x^1} dt \wedge dx^1 \wedge dx^2 \wedge dx^3 \wedge dr}{r^5} + 4r_+^4 \sin^3 \theta \cos \theta \sin \xi \cos \xi d\theta \wedge d\xi \wedge d\phi_1 \wedge d\phi_2 \wedge d\phi_3, \tag{5.36}$$

where $H(r) \equiv 1 - (r_+/r)^4$ and the four-dimensional metric inside the brackets $\{\dots\}$ is flat. We consider the following three Killing vectors,

$$V_1 \equiv (\partial_1 + x^2 \partial_2 + x^3 \partial_3, 0), \quad V_2 \equiv (\partial_2, 0), \quad V_3 \equiv (\partial_3, 0), \tag{5.37}$$

that satisfy the algebra

$$[V_1, V_2]_{\text{C}} = -V_2, \quad [V_1, V_3]_{\text{C}} = -V_3, \quad [V_2, V_3]_{\text{C}} = 0. \tag{5.38}$$

The (x^1, x^2, x^3) parts of the matrices are

$$(\Lambda_v) = \begin{pmatrix} 1 & x^2 & x^3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (\Lambda_f) = \begin{pmatrix} 0 & -\tilde{x}_2 & -\tilde{x}_3 \\ \tilde{x}_2 & 0 & 0 \\ \tilde{x}_3 & 0 & 0 \end{pmatrix}, \tag{5.39}$$

and the gauge symmetry is fixed as $x^i(\sigma) = 0$ ($i = 1, 2, 3$). The dual background becomes

$$ds'^2 = -H^{\frac{1}{2}} dt^2 + \frac{t^4 H (d\tilde{x}_1^2 + d\tilde{x}_2^2 + d\tilde{x}_3^2) + \tilde{x}_3^2 d\tilde{x}_2^2 - 2\tilde{x}_2 \tilde{x}_3 d\tilde{x}_2 d\tilde{x}_3 + \tilde{x}_2^2 d\tilde{x}_3^2}{t^2 H^{\frac{1}{2}} (H t^4 + \tilde{x}_2^2 + \tilde{x}_3^2)} + \frac{dr^2}{H^2} + r^2 (d\theta^2 + \sin^2 \theta d\xi^2 + \sin^2 \theta \cos^2 \xi d\phi_1^2 + \sin^2 \theta \sin^2 \xi d\phi_2^2 + \cos^2 \theta d\phi_3^2),$$

$$B'_2 = \frac{d\tilde{x}_1 \wedge (\tilde{x}_2 d\tilde{x}_2 + \tilde{x}_3 d\tilde{x}_3)}{H t^4 + \tilde{x}_2^2 + \tilde{x}_3^2}, \quad e^{-2\Phi'} = t^2 H^{\frac{1}{2}} (H t^4 + \tilde{x}_2^2 + \tilde{x}_3^2), \quad I' = 2\tilde{\partial}^1,$$

$$G'_2 = -\frac{4r_+^4 t^3 dt \wedge dr}{r^5}, \quad G'_4 = -\frac{4r_+^4 t^3 dt \wedge d\tilde{x}_1 \wedge (\tilde{x}_2 d\tilde{x}_2 + \tilde{x}_3 d\tilde{x}_3) \wedge dr}{r^5 (H t^4 + \tilde{x}_2^2 + \tilde{x}_3^2)}, \tag{5.40}$$

and this is a solution of type IIA GSE.

Again, by performing a formal T -duality along the \tilde{x}_1 -direction we obtain a solution of type IIB supergravity,

$$ds^2 = H^{\frac{1}{2}} (-dt^2 + t^2 dx_1^2) + \frac{(d\tilde{x}_2 - \tilde{x}_2 dx^1)^2 + (d\tilde{x}_3 - \tilde{x}_3 dx^1)^2}{H^{\frac{1}{2}} t^2} + \frac{dr^2}{H^2} + r^2 (d\theta^2 + \sin^2 \theta d\xi^2 + \sin^2 \theta \cos^2 \xi d\phi_1^2 + \sin^2 \theta \sin^2 \xi d\phi_2^2 + \cos^2 \theta d\phi_3^2),$$

$$e^{-2\Phi} = t^4 e^{-4x^1} H(r), \quad G_3 = e^{-2x^1} \frac{4r_+^4 t^3 dt \wedge dx^1 \wedge dr}{r^5}. \tag{5.41}$$

We note that, as discussed in Ref. [148], some supergravity solutions obtained by a combination of NATD and a formal T -duality can also be obtained from another route, a combination of diffeomorphisms and Abelian T -dualities. Similarly, the solutions obtained in this paper may also be realized from such procedure.

6. Poisson–Lie T -duality/plurality

Here we study a more general class of T -duality known as the Poisson–Lie T -duality [37,38] or T -plurality [55]. We can perform the PL T -duality/plurality when the target space has a set of vectors

v_a satisfying the dualizability conditions [37]

$$[v_a, v_b] = f_{ab}{}^c v_c, \quad \mathfrak{L}_{v_a} E_{mn} = -\tilde{f}^{bc}{}_a E_{mp} v_b^p v_c^q E_{qn}. \tag{6.1}$$

The traditional NATD (with $\tilde{v}_a = 0$) can be regarded as a special case, $\tilde{f}^{bc}{}_a = 0$. We begin with a brief review of the idea and techniques, and show the covariance of the DFT equations of motion under the PL T -plurality. Namely, we show that if we start with a DFT solution, the PL T -dualized background is also a DFT solution. In some examples the Killing vector I^m appears, and the dualized DFT solutions are regarded as GSE solutions. However, through a formal T -duality the GSE solutions can always be transformed into linear-dilaton solutions of the conventional supergravity.

6.1. Review of PL T -duality

We review the PL T -duality as a symmetry of the classical equations of motion of the string sigma model. To make the discussion transparent, we first ignore spectator fields $y^\mu(\sigma)$, which are invariant under the PL T -duality. As studied in Refs. [37,38] it is straightforward to introduce spectators, and their treatment is discussed in Sect. 6.2.4.

Let us consider a sigma model with a target space M , on which a group G acts transitively and freely (i.e. M itself can be regarded as a group manifold),

$$S = \frac{1}{4\pi\alpha'} \int_{\Sigma} E_{mn}(x) (dx^m \wedge * dx^n + dx^m \wedge dx^n). \tag{6.2}$$

Under an infinitesimal right action of a group G , the coordinates x^m are shifted as

$$g(x) \rightarrow g(x) (1 + \epsilon^a T_a) \equiv g(x + \delta x), \quad \delta x^m = \epsilon^a(\sigma) v_a^m(x), \tag{6.3}$$

where T_a ($a = 1, \dots, n$) are the generators of the algebra \mathfrak{g} satisfying

$$[T_a, T_b] = f_{ab}{}^c T_c, \tag{6.4}$$

and v_a^m are the left-invariant vector fields satisfying

$$[v_a, v_b] = f_{ab}{}^c v_c, \quad v_a^m \ell_m^b = \delta_a^b, \quad \ell \equiv \ell^a T_a \equiv g^{-1} dg. \tag{6.5}$$

In general, the variation of the action becomes

$$\delta_\epsilon S = \frac{1}{2\pi\alpha'} \int_{\Sigma} \left\{ -\epsilon^a \left[dJ_a - \frac{1}{2} \mathfrak{L}_{v_a} E_{mn} (dx^m \wedge * dx^n + dx^m \wedge dx^n) \right] + d(\epsilon^a J_a) \right\}, \tag{6.6}$$

where

$$J_a \equiv v_a^m (g_{mn} * dx^n + B_{mn} dx^n). \tag{6.7}$$

If the v_a^m satisfy the Killing equation $\mathfrak{L}_{v_a} E_{mn} = 0$, equations of motion for x^m can be written as

$$dJ_a = 0. \tag{6.8}$$

In particular, if v_a^m further satisfy $[v_a, v_b] = 0$ we can find a coordinate system where $v_a^m = \delta_a^m$ is realized. Then, the Abelian T -duality can be realized as the exchange of $x^m(\sigma)$ with the dual coordinates $\tilde{x}_a(\sigma)$, which are defined as

$$d\tilde{x}_a \equiv J_a. \tag{6.9}$$

The Bianchi identity $d^2\tilde{x}_a = 0$ corresponds to the equations of motion in the original theory.

The PL T -duality is a generalization of this duality when the vector fields v_a satisfy

$$\mathfrak{L}_{v_a} E_{mn} = -\tilde{f}^{bc}{}_a E_{mp} v_b^p v_c^q E_{qn}. \tag{6.10}$$

In this case, the variation becomes

$$\delta S = \frac{1}{2\pi\alpha'} \int_{\Sigma} \left[-\epsilon^a \left(dJ_a - \frac{1}{2} \tilde{f}_a{}^{bc} J_b \wedge J_c \right) + d(\epsilon^a J_a) \right], \tag{6.11}$$

and the equations of motion for x^m become the Maurer–Cartan equation,

$$dJ_a - \frac{1}{2} \tilde{f}_a{}^{bc} J_b \wedge J_c = 0. \tag{6.12}$$

This suggests introducing the dual coordinates $\tilde{x}_m(\sigma)$ through a non-Abelian generalization of Eq. (6.9), namely,

$$\tilde{r}_a \equiv J_a \quad (\tilde{r} \equiv \tilde{r}_a \tilde{T}^a \equiv d\tilde{g} \tilde{g}^{-1}, \quad \tilde{g} \equiv \tilde{g}(\tilde{x}) \in \tilde{G}), \tag{6.13}$$

where \tilde{T}^a are the generators of the dual algebra $\tilde{\mathfrak{g}}$ (associated with a dual group \tilde{G}) satisfying

$$[\tilde{T}^a, \tilde{T}^b] = \tilde{f}^{ab}{}_c \tilde{T}^c. \tag{6.14}$$

Then, under the equations of motion, the physical coordinates $x^m(\sigma)$ describe the motion of the string on the group G while the dual coordinates $\tilde{x}_m(\sigma)$ describe the motion of the string on the dual group \tilde{G} .

It is important to note that the condition in Eq. (6.10) and the identity

$$[\mathfrak{L}_{v_a}, \mathfrak{L}_{v_b}] E_{mn} = \mathfrak{L}_{[v_a, v_b]} E_{mn} \tag{6.15}$$

show the relation

$$f_{ae}{}^c \tilde{f}^{ed}{}_b + f_{ae}{}^d \tilde{f}^{ce}{}_b - f_{be}{}^c \tilde{f}^{ed}{}_a - f_{be}{}^d \tilde{f}^{ce}{}_a = f_{ab}{}^e \tilde{f}^{cd}{}_e. \tag{6.16}$$

By considering the vector space $\tilde{\mathfrak{g}}$ as the dual space of \mathfrak{g} , $\langle T_a, \tilde{T}^b \rangle = \delta_a^b$, the relation gives the structure of the Lie bialgebra. By further introducing an ad -invariant bilinear form as

$$\langle T_A, T_B \rangle = \eta_{AB}, \quad (\eta_{AB}) = \begin{pmatrix} 0 & \delta_a^b \\ \delta_a^b & 0 \end{pmatrix}, \quad (T_A) \equiv (T_a, \tilde{T}^a), \tag{6.17}$$

the commutation relations on a direct sum $\mathfrak{d} \equiv \mathfrak{g} \oplus \tilde{\mathfrak{g}}$ are determined as

$$[T_a, T_b] = f_{ab}{}^c T_c, \quad [T_a, \tilde{T}^b] = \tilde{f}^{bc}{}_a T_c - f_{ac}{}^b \tilde{T}^c, \quad [\tilde{T}^a, \tilde{T}^b] = \tilde{f}^{ab}{}_c \tilde{T}^c, \tag{6.18}$$

and the pair of algebras can be regarded as that of the Drinfel'd double \mathfrak{D} . Given the structure of the Drinfel'd double, the differential equation in Eq. (6.10) can be integrated [37,38] as

$$\mathbf{E}_{ab} \equiv v_a^m v_b^n E_{mn} = [a^{-1} \hat{E} (a^\top + b^\top \hat{E})^{-1}]_{ab}, \tag{6.19}$$

where the matrices a and b are defined by

$$g^{-1} T_A g = (\text{Ad}_{g^{-1}})_A{}^B T_B, \quad \text{Ad}_{g^{-1}} = \begin{pmatrix} a_a{}^b & 0 \\ b^{ab} & (a^{-\top})^a{}_b \end{pmatrix}, \tag{6.20}$$

and \hat{E}_{ab} is an arbitrary constant matrix (that corresponds to $E_{ab}(x)$ at $g = 1$). We can check that the E_{mn} given by Eq. (6.19) indeed satisfy Eq. (6.10).⁶

Now, we rewrite the relation in Eq. (6.13), namely

$$\tilde{r}_a = J_a = g_{ab} * \ell^b + B_{ab} \ell^b \quad (g_{ab} \equiv E_{(ab)}, \quad B_{ab} \equiv E_{[ab]}), \quad (6.21)$$

into two equivalent expressions (by following the standard trick [10] in the Abelian case),

$$\begin{aligned} \ell^a &= -(g^{-1} B)^a_b * \ell^b + g^{ab} * \tilde{r}_b, \\ \tilde{r}_a &= (g - B g^{-1} B)_{ab} * \ell^b + (B g)_a^b * \tilde{r}_b. \end{aligned} \quad (6.22)$$

They can be neatly expressed as a self-duality relation,

$$\begin{aligned} P^A &= H^A_B(x) * P^B, \quad (P^A) \equiv \begin{pmatrix} \ell^a \\ \tilde{r}_a \end{pmatrix}, \\ (H_{AB}) &\equiv \begin{pmatrix} (g - B g^{-1} B)_{ab} & B_{ac} g^{cb} \\ -g^{ac} B_{cb} & g^{ab} \end{pmatrix}, \end{aligned} \quad (6.23)$$

where the indices A, B, \dots are raised or lowered with η_{AB} and its inverse η^{AB} . In terms of the metric H_{AB} , the relation in Eq. (6.19) can be expressed as

$$H_{AB}(x) = (\text{Ad}_g)_A^C (\text{Ad}_g)_B^D \hat{\mathcal{H}}_{CD}, \quad (\hat{\mathcal{H}}_{AB}) \equiv \begin{pmatrix} (\hat{g} - \hat{B} \hat{g}^{-1} \hat{B})_{ab} & \hat{B}_{ac} \hat{g}^{cb} \\ -\hat{g}^{ac} \hat{B}_{cb} & \hat{g}^{ab} \end{pmatrix}, \quad (6.24)$$

where $\hat{g}_{ab} \equiv \hat{E}_{(ab)}$, $\hat{B}_{ab} \equiv \hat{E}_{[ab]}$. Then, Eq. (6.23) gives the important relation

$$\hat{P}^A = \hat{\mathcal{H}}^A_B * \hat{P}^B, \quad \hat{P}(\sigma) \equiv \hat{P}^A T_A \equiv dl l^{-1}, \quad l \equiv g \tilde{g}, \quad (6.25)$$

where we have used⁷

$$\hat{P}(\sigma) \equiv dl l^{-1} = g (\ell^a T_a + \tilde{r}_a \tilde{T}^a) g^{-1} = P^B (\text{Ad}_g)_B^A T_A. \quad (6.26)$$

Expressed in this form, the equations of motion are given in terms of the Drinfel'd double \mathcal{D} ; the decomposition $l = g \tilde{g}$ is no longer important.

Similar to the Abelian T -duality, we can recover the same equations of motion from the dual model by exchanging the role of g and \tilde{g} . Starting with the dual background \tilde{E}_{mn} , which has a set of vector fields \tilde{v}^a satisfying

$$[\tilde{v}^a, \tilde{v}^b] = \tilde{f}^{ab}_c \tilde{v}^c, \quad \mathfrak{L}_{\tilde{v}^a} E_{mn} = -f_{bc}{}^a \tilde{E}_{mp} \tilde{v}^{bp} \tilde{v}^{cq} \tilde{E}_{qn}, \quad (6.27)$$

⁶ For example, when E_{mn} is invertible, we can easily check an equivalent expression $\mathfrak{L}_{v_a} E^{mn} = \tilde{f}^{bc}{}_a v_b^m v_c^n$ by using the rewriting of Eq. (6.35) and $v_c^m \partial_m \Pi^{ab} = -(a^{-\tau})^a_d (a^{-\tau})^b_e \tilde{f}^{dc}{}_e$, which can be derived from Eq. (6.20) (see Ref. [44]).

⁷ If we expand the right-invariant form as $\hat{P} = \mathcal{P}^A_M dx^M T_A$, we find that \mathcal{P}^A_M is not an $O(n, n)$ matrix:

$$(\mathcal{P}^A_M) = \begin{pmatrix} r_m^a & \Pi^{ab} a_b^c \tilde{r}_c^m \\ 0 & a_a^b \tilde{r}_b^m \end{pmatrix}.$$

the equations of motion can be expressed as

$$\hat{\mathcal{P}}_A = \tilde{\mathcal{H}}_A^B * \hat{\mathcal{P}}_B, \quad \hat{\mathcal{P}}_A T^A \equiv d\tilde{l}\tilde{l}^{-1}, \quad \tilde{l} \equiv \tilde{h} h \quad (h \in \mathfrak{g}, \quad \tilde{h} \in \tilde{\mathfrak{g}}) \quad (6.28)$$

by using a constant matrix $\tilde{\mathcal{H}}_A^B$. For the duality equivalence, we demand that Eqs. (6.25) and (6.28) are equivalent. This leads to the identifications

$$\hat{\mathcal{H}}_{AB} = \tilde{\mathcal{H}}_{AB}, \quad g \tilde{g} = l = \tilde{l} \equiv \tilde{h} h. \quad (6.29)$$

After this identification, string theory defined on the original background E_{mn} and the dual background E'_{mn} give the same equations of motion, and are classically equivalent.

In summary, in PL T -dualizable backgrounds the generalized metric $\mathcal{H}_{MN}(x)$ is always related to a constant matrix $\hat{\mathcal{H}}_{AB}$ as

$$\mathcal{H}_{MN} = (U \hat{\mathcal{H}} U^\top)_{MN}, \quad (6.30)$$

where the matrix U is defined as

$$U_M^A \equiv L_M^B (\text{Ad}_g)_B^A, \quad (L_M^A) \equiv \begin{pmatrix} \ell_m^a & 0 \\ 0 & v_a^m \end{pmatrix}. \quad (6.31)$$

By comparing this with Eq. (2.19), we call the matrix U the twist matrix and call the constant matrix $\hat{\mathcal{H}}_{AB}$ the untwisted metric. The dual geometry also has the same structure, where the twist matrix is $\tilde{U}_{MA} \equiv \tilde{L}_{MB} (\text{Ad}_{\tilde{g}})^B_A$. The relation between the original and the dual background becomes

$$\tilde{\mathcal{H}}_{MN} = (h \mathcal{H} h^\top)_{MN}, \quad h_M^N \equiv \tilde{U}_{MA} \eta^{AB} U_B^N. \quad (6.32)$$

For later convenience, we rewrite the twist matrix as

$$U = L \text{Ad}_g = R \Pi, \quad (6.33)$$

where we have defined

$$(R_M^A) \equiv \begin{pmatrix} r_m^a & 0 \\ 0 & e_a^m \end{pmatrix}, \quad (\Pi_A^B) \equiv \begin{pmatrix} \delta_a^b & 0 \\ -\Pi^{ab} & \delta_b^a \end{pmatrix}, \\ r \equiv r^a T_a \equiv dg g^{-1}, \quad r_m^a e_b^m = \delta_b^a, \quad \Pi^{ab} \equiv (b a^{-1})^{ab} = -(a^{-\top} b^\top)^{ab}, \quad (6.34)$$

and used $r^a = (a^{-\top})^a_b \ell^b$. Then, in terms of $E_{mn}(x)$, Eq. (6.30) can be expressed as

$$E_{mn}(x) = [(\hat{E}^{-1} - \Pi)^{-1}]_{ab} r_m^a r_n^b, \quad (6.35)$$

and, similarly, the dual background is

$$\tilde{E}_{mn}(\tilde{x}) = [(\hat{E} - \tilde{\Pi})^{-1}]^{ab} \tilde{r}_{am} \tilde{r}_{bn}. \quad (6.36)$$

In the special case where $\tilde{f}^{ab}_c = 0$, by parameterizing $\tilde{g} = e^{\tilde{x}_a \tilde{T}^a}$ we obtain $\tilde{r} = d\tilde{x}_a \tilde{T}^a$, $\Pi^{ab} = 0$, and $\tilde{\Pi}_{ab} = -f_{ab}^c \tilde{x}_c$. This is precisely the case of NATD. In the dualized background, in general the isometries are broken, and in the traditional NATD we cannot recover the original model. However, the dual background has the form

$$\tilde{E}^{mn} = (\hat{E} - \tilde{\Pi})_{ab} \tilde{e}^{am} \tilde{e}^{bn} = (\hat{E}_a + f_{ab}^c \tilde{x}_c) \tilde{v}^{am} \tilde{v}^{bn}, \quad (6.37)$$

where $\tilde{e}^a = \tilde{v}^a = \tilde{\partial}^a$, and we find that the dual background is T -dualizable,

$$\mathfrak{L}_{\tilde{v}^a} E^{mn} = \tilde{\partial}^a E^{mn} = f_{bc}{}^a \tilde{v}^{bm} \tilde{v}^{cn}. \tag{6.38}$$

Thus, through the PL T -duality we can recover the original background $E_{mn} = E_{ab} r_m^a r_n^b$.

As a side remark, we note that in the case of the Abelian $O(D, D)$ T -duality, the covariant equations of motion of string $dx^M = \hat{\mathcal{H}}^M{}_N * dx^N$ [10] can be derived from the double sigma model (DSM) [163,168–172]. The correspondent of the DSM for the PL T -duality has been studied in Refs. [40, 41,64,173–175], and this approach will be useful to manifest the PL T -duality.

6.2. PL T -plurality

The Lie algebra \mathfrak{d} of the Drinfel'd double \mathfrak{D} can be constructed as a direct sum of two algebras, \mathfrak{g} and $\tilde{\mathfrak{g}}$, which are maximally isotropic with respect to the bilinear form $\langle \cdot, \cdot \rangle$, and $(\mathfrak{d}, \mathfrak{g}, \tilde{\mathfrak{g}})$ is called the Manin triple. In general, a Drinfel'd double has several decompositions into Manin triples, and this leads to the notion of the PL T -plurality [55]. More concretely, let us consider a redefinition of the generators T_A of \mathfrak{d} ,

$$T'_A \equiv C_A{}^B T_B, \tag{6.39}$$

such that the new generators also satisfy the algebra of the Drinfel'd double,

$$[T'_a, T'_b] = f'{}_{ab}{}^c T'_c, \quad [T'_a, \tilde{T}'^b] = \tilde{f}'{}^{bc}{}_a T'_c - f'{}^b{}_ac \tilde{T}'^c, \quad [\tilde{T}'^a, \tilde{T}'^b] = \tilde{f}'{}^{ab}{}_c \tilde{T}'^c, \tag{6.40}$$

and the bilinear form is preserved,

$$\langle T'_A, T'_B \rangle = \eta_{AB}. \tag{6.41}$$

The latter condition shows that the matrix $C_A{}^B$ should be a certain $O(n, n)$ matrix. Since the rescaling of the generators is trivial, we choose $C_A{}^B$ as a “volume-preserving” $O(n, n)$ transformation that does not change the DFT dilaton.

The transformation of the background fields under the $O(n, n)$ transformation can be found in the same manner as the PL T -duality. Starting with a background E'_{mn} satisfying

$$[v'_a, v'_b] = f'{}_{ab}{}^c v'_c, \quad \mathfrak{L}_{v'_a} E'_{mn} = -\tilde{f}'{}^{bc}{}_a E'_{mp} v'_b{}^p v'_c{}^q E'_{qn}, \tag{6.42}$$

we again obtain the same equations of motion,

$$\mathcal{P}'^A = \hat{\mathcal{H}}'^A{}_B * \mathcal{P}'^B, \quad \mathcal{P}'^A T'_A \equiv dl' l'^{-1}, \quad l' \equiv g' \tilde{g}'. \tag{6.43}$$

From the identification $l = l'$ we obtain

$$\hat{\mathcal{P}}^A T_A = dl l^{-1} = dl' l'^{-1} = \hat{\mathcal{P}}'^A T'_A = \hat{\mathcal{P}}'^A C_A{}^B T_B, \tag{6.44}$$

and the relation between the untwisted metrics becomes

$$\hat{\mathcal{H}}'_{AB} = (C \hat{\mathcal{H}} C^\top)_{AB}. \tag{6.45}$$

The generalized metric in the transformed frame has the form

$$\hat{\mathcal{H}}'_{MN} = (U' \hat{\mathcal{H}}' U'^\top)_{MN}, \tag{6.46}$$

and the relation between the original and the dual generalized metric is

$$\mathcal{H}'_{MN} = (h \mathcal{H} h^T)_{MN}, \quad (h_M^N) \equiv U' C U^{-1}. \tag{6.47}$$

In terms of $E_{mn}(x)$, the original background is

$$E_{mn}(x) = [(\hat{E}^{-1} - \Pi)^{-1}]_{ab} r_m^a r_n^b, \tag{6.48}$$

while the dual background is

$$E'_{mn}(x') = [(\hat{E}'^{-1} - \Pi')^{-1}]_{ab} r_m'^a r_n'^b, \quad E'_{mn} = [(\mathbf{q} + \mathbf{p} \hat{E})(s + \mathbf{r} \hat{E})^{-1}]_{mn}, \tag{6.49}$$

where we parameterized the $O(n, n)$ matrix C as

$$C = \begin{pmatrix} \mathbf{p}_m^n & \mathbf{q}_{mn} \\ \mathbf{r}^m_n & \mathbf{s}^m_n \end{pmatrix}. \tag{6.50}$$

Note that the PL T -duality is a special case of the T -plurality where

$$C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \tag{6.51}$$

and the original background corresponds to the trivial choice $C = 1$.

6.2.1. Duality rule for the dilaton

The transformation rule for the dilaton was studied in Ref. [64] in the context of the PL T -duality. This was improved in Ref. [55] in the study of the PL T -plurality. In our convention, the result is

$$e^{-2\Phi'} = e^{-2\bar{\Phi}} \frac{|\det(\mathbf{q} + \mathbf{p} \hat{E})|}{|\det(E'_{ab})| |\det a'^{-1}|} \quad (E'_{ab} \equiv e_a^m e_b^n E'_{mn}), \tag{6.52}$$

where $\bar{\Phi}(x)$ is an arbitrary function. By using the formula in Eq. (3.28), we obtain

$$\begin{aligned} \sqrt{|g'|} &= |\det(r_m'^a)| |\det(\mathbf{1} - \Pi' \hat{E}')|^{-1} |\det(s + \mathbf{r} \hat{E})|^{-1} \sqrt{|\hat{g}|} \\ &= |\det(r_m'^a)| |\det E'_{ab}| |\det(\mathbf{q} + \mathbf{p} \hat{E})|^{-1} \sqrt{|\hat{g}|}, \end{aligned} \tag{6.53}$$

and the DFT dilaton in the dual background becomes

$$e^{-2d'} = e^{-2\bar{d}} |\det(r_m'^a)| |\det a'| = e^{-2\bar{d}} |\det(\ell_m'^a)|, \quad e^{-2\bar{d}} \equiv e^{-2\bar{\Phi}} \sqrt{|\hat{g}|}. \tag{6.54}$$

Namely, the duality rule for the DFT dilaton is

$$|\det(v_a^m)| e^{-2(d' - \bar{d})} = 1 = |\det(v_a^m)| e^{-2(d - \bar{d})}. \tag{6.55}$$

If \bar{d} (or equivalently $\bar{\Phi}$) is constant, this duality rule coincides with the recent proposal of Ref. [139], where the PL T -duality was studied by utilizing “the DFT on a Drinfel’d double” proposed in Ref. [138]. There, it was shown that the dilaton transformation rule is also consistent with Ref. [141]. Moreover, when the dual algebra is Abelian, $\tilde{f}^{ab}_c = 0$, we have $|\det(v_a^m)| = 1$ and the result in Eq. (3.30) known in NATD is also reproduced as a particular case.

In fact, as demonstrated in Ref. [55], the PL T -plurality works even if \bar{d} has a coordinate dependence. A subtle point is that when $e^{-2\bar{d}}$ depends on the original coordinates x^m it is not clear how to

understand the x^m -dependence in the dual model. A prescription proposed in Ref. [55] is as follows. We first identify the relation between coordinates $(x^M) = (x^m, \tilde{x}_m)$ and $(x'^M) = (x'^m, \tilde{x}'_m)$ through the identification

$$g'(x') \tilde{g}'(\tilde{x}') = l = g(x) \tilde{g}(\tilde{x}). \tag{6.56}$$

We next substitute the relation $x^M = x^M(x')$ into $e^{-2\bar{d}(x)}$ as $e^{-2\bar{d}(x)} = e^{-2\bar{d}(x(x'))} \equiv e^{-2\bar{d}(x')}$. Then, the relation in Eq. (6.55) can be understood on both sides:

$$|\det(v_a^m)| e^{-2[d' - \bar{d}(x')]} = 1 = |\det(v_a^m)| e^{-2[d - \bar{d}(x)]}. \tag{6.57}$$

In general, $\bar{d}(x')$ may depend on the dual coordinates \tilde{x}'_m , and the background does not have the usual supergravity description. However, in our examples the DFT dilaton has at most a linear dependence on the dual coordinates, and it can be absorbed into the Killing vector I^m in the GSE.

6.2.2. Covariance of equations of motion

In the approach of Refs. [138,139], the PL T -duality was realized as a manifest symmetry of DFT. We discuss here the covariance under a more general PL T -plurality by using the gauged DFT. The approach may be slightly different from Refs. [138,139] but the essence will be the same.

In PL T -dualizable backgrounds, the generalized metric always has the simple form

$$\mathcal{H}_{MN} = [U(x) \hat{\mathcal{H}} U^T(x)]_{MN}. \tag{6.58}$$

Since the twist matrix U is explicitly determined, we can compute the generalized fluxes \mathcal{F}_{ABC} and \mathcal{F}_A defined in Eq. (2.20). In fact, as shown in Ref. [139], in PL T -dualizable backgrounds the three-index flux is precisely the structure constant of the Drinfel'd double,

$$\mathcal{F}_{abc} = 0, \quad \mathcal{F}_{ab}{}^c = f_{ab}{}^c, \quad \mathcal{F}{}^{ab}{}_c = \tilde{f}{}^{ab}{}_c, \quad \mathcal{F}{}^{abc} = 0. \tag{6.59}$$

We can check this by using the explicit form of the twist matrix and its inverse,

$$(U_M{}^A) = \begin{pmatrix} r_m^a & 0 \\ -e_b^m \Pi^{ba} & e_a^m \end{pmatrix}, \quad (U^A{}_M) = \begin{pmatrix} e_a^m & 0 \\ \Pi^{ab} e_b^m & r_m^a \end{pmatrix}, \tag{6.60}$$

and the relations $\mathbb{E}_{e_a} e_b = -f_{ab}{}^c e_c$, $\mathbb{E}_{e_a} r^b = f_{ac}{}^b r^c$, $\partial_m \Pi^{ab} = -(a^{-T})^a{}_d (a^{-T})^b{}_e \tilde{f}{}^{de}{}_f a^f{}_c r_m^c$, and $\tilde{f}{}^{ab}{}_c = (a^{-T})^a{}_d (a^{-T})^b{}_e a_c{}^f \tilde{f}{}^{de}{}_f - 2f_{ce}{}^{[a} \Pi^{b]e}$ (see Ref. [44] for useful identities).

We can also compute the single-index flux as

$$\mathcal{F}_A = \begin{pmatrix} 2e_a^m \partial_m d + e_c^n \partial_n r_m^c e_a^m \\ -(a^{-T})^a{}_b \tilde{f}{}^{cb}{}_c + \Pi^{ab} (2e_b^m \partial_m d + e_c^n \partial_n r_m^c e_b^m) + 2r_m^a \tilde{\partial}^m d \end{pmatrix}. \tag{6.61}$$

By using the expression for the DFT dilaton in Eq. (6.54), $e^{-2d} = e^{-2\bar{d}} |\det(r_m^a)| |\det a|$, we find

$$\mathcal{F}_A = U_A{}^M \mathcal{F}_M, \quad \mathcal{F}_M \equiv 2\partial_M \bar{d} + \begin{pmatrix} 0 \\ -\tilde{f}{}^{ba}{}_b v_a^m \end{pmatrix}, \tag{6.62}$$

where we have used $a_b{}^e a_c{}^f f_{ef}{}^a = -f_{cb}{}^e a_e{}^a$ and $\partial_m a_a{}^b = a_a{}^c f_{cd}{}^b \ell_m^d$.

As we discuss below, for the covariance of the equation of motion under the PL T -plurality, \mathcal{F}_A needs to transform covariantly. However, even in the particular case $\bar{d} = 0$, for example, we find

that \mathcal{F}_A does not transform covariantly. Indeed, we have $\mathcal{F}_A = 0$ in a duality frame where $\tilde{f}^{ab}_a = 0$, while \mathcal{F}_A appears in a frame where $\tilde{f}^{ab}_a \neq 0$. Therefore, in order to transform \mathcal{F}_A covariantly, we eliminate the non-covariant term by adding a vector field X_M as

$$\partial_M d \rightarrow \partial_M d + X_M, \quad (X_M) \equiv \begin{pmatrix} 0 \\ I^m \end{pmatrix}, \quad I^m = \frac{1}{2} \tilde{f}^{ba}_b v_a^m, \quad (6.63)$$

which was suggested in Ref. [139]. This shift is a bit artificial, but without this procedure we need to abandon all Manin triples with non-unimodular dual algebra. In fact, this shift is precisely the modification of DFT equations of motion Eq. (2.36) that reproduces the GSE after removing the dual-coordinate dependence. After this prescription, we obtain the simple flux

$$\mathcal{F}_A = 2 \mathcal{D}_A \bar{d}. \quad (6.64)$$

In fact, as we see later, $\mathcal{F}_A = 2 \mathcal{D}_A \bar{d}$ are covariantly transformed under the PL T -plurality $\mathcal{F}'_A = C_A^B \mathcal{F}_B$,⁸ and the prescription in Eq. (6.63) works well in our examples.

Now, let us discuss the covariance of the equations of motion. Since the derivative \mathcal{D}_A generally does not transform covariantly, we assume that $\mathcal{F}_A = 2 \mathcal{D}_A \bar{d}$ is constant. Since \mathcal{F}_{ABC} is also constant in PL T -dualizable backgrounds, the DFT equations of motion become simple algebraic equations, Eqs. (2.25) and (2.26).

Under the PL T -plurality $T'_A = C_A^B T_B$, the generalized fluxes are mapped as

$$\mathcal{F}'_{ABC} = C_A^D C_B^E C_C^F \mathcal{F}_{DEF}, \quad \mathcal{F}'_A = C_A^B \mathcal{F}_B \quad (6.65)$$

by introducing X_M when the dual algebra is non-unimodular. According to Eq. (6.45), the untwisted metric $\hat{\mathcal{H}}_{AB}$ is also related covariantly,

$$\hat{\mathcal{H}}'_{AB} = (C \hat{\mathcal{H}} C^T)_{AB}. \quad (6.66)$$

Then, we find that the equations of motion in the original and the dual background are covariantly related by the $O(n, n)$ transformation C . Thus, as long as the original configuration is a DFT solution, the dual background also satisfies the DFT equations of motion.

We note that this $O(n, n)$ transformation is totally different from the transformation in Eq. (2.30), which is just a redefinition of U , and the generalized metric \mathcal{H}_{MN} is invariant. On the other hand, in the case of PL T -plurality, $U(x)$ in the original model and $U'(x')$ in the dual model are defined on a different manifold and there is no clear connection between $U(x)$ and $U'(x')$. Only the constant fluxes made out of $U(x)$ and $U'(x')$ are related by a constant $O(n, n)$ transformation, and this non-trivial relation connects the two equations of motion in a covariant manner.

Before moving on to the R–R sector, we make a brief comment on the vector field I^m . In order to reproduce the (generalized) supergravity from (modified) DFT, we need to choose the standard section $\tilde{\partial}^m = 0$. Therefore, when \bar{d} has a dual-coordinate dependence, we need to make an additional field redefinition. Supposing that \bar{d} only has a linear dual-coordinate dependence $\bar{d} = \bar{d}_0(x^m) + d^m \tilde{x}_m$, we make the field redefinition

$$\bar{d} \rightarrow \bar{d}' = \bar{d}_0(x^m), \quad I^m \rightarrow I'^m = \frac{1}{2} \tilde{f}^{ba}_b v_a^m + d^m. \quad (6.67)$$

⁸ This is non-trivial, because in general the derivative \mathcal{D}_A does not transform covariantly, $\mathcal{D}'_A \neq C_A^B \mathcal{D}_B$, which can be checked by performing the coordinate transformation $x'^M = x'^M(x)$ through Eq. (6.56). Therefore, at the present time, the covariance of \mathcal{F}_A needs to be checked on a case-by-case basis. Of course, when \bar{d} is constant, the covariance is manifest because $\mathcal{F}_A = 0$ and $\mathcal{F}'_A = 0$.

Then, the dual-coordinate dependence disappears from the background. Note that this is different from the shift in Eq. (6.63) and is just a field redefinition. In the following, when we display a (generalized) supergravity solution we always make this redefinition.

Let us also make a brief comment on the Killing vector I^m . In the case of NATD, the Killing vector I^m is given by Eq. (4.1), but Eq. (4.1) is apparently different from the formula in Eq. (6.63) by the factor 2. Here, we will roughly sketch how to resolve the discrepancy by using the redefinition in Eq. (6.67). In the case of NATD, $\partial_m |\det(v_a^m)| = 0$ and $\partial_m \Phi = 0$ are usually satisfied in the original background (under the gauge fixing $x^m = c^m$). Then, we have

$$\begin{aligned} \partial_m \bar{d} &= \partial_m d = -\frac{1}{2} \partial_m \ln \sqrt{|g|} = -\frac{1}{2} \partial_m \ln |\det(r_m^a)| \\ &= \frac{1}{2} \partial_m \ln |\det a| = \frac{1}{2} f_{ba}{}^b \ell_m^a. \end{aligned} \tag{6.68}$$

Namely, \bar{d} has a linear coordinate dependence along the v_a^m direction,

$$v_a^m \partial_m \bar{d} = \frac{1}{2} f_{ba}{}^b. \tag{6.69}$$

After performing NATD, this gives a dual-coordinate dependence of \bar{d} in the dual theory,

$$\bar{d} = \frac{1}{2} \tilde{f}^{ba}{}_{\tilde{b}} \tilde{x}_a, \tag{6.70}$$

where the dual structure constants $\tilde{f}^{ab}{}_c$ correspond to $f_{ab}{}^c$ in the original frame. Then, the modified I^m in Eq. (6.67) recovers the formula in Eq. (4.1),

$$I^m = \frac{1}{2} \tilde{f}^{ba}{}_{\tilde{b}} v_a^m + d^m = \tilde{f}^{ba}{}_{\tilde{b}}, \tag{6.71}$$

where we have used $v_a^m = \delta_a^m$ in the dual theory. In a general setup Eq. (4.1) does not work correctly, and we use the results discussed in this section.

6.2.3. Duality rule for R–R fields

Now, let us determine the duality rule for the R–R fields. We will first find the duality rule from a heuristic approach, and then clarify the result in terms of the gauged DFT.

In the presence of the R–R fields, the equations of motion for \mathcal{H}_{MN} and d are

$$\mathcal{S}_{MN} = \mathcal{E}_{MN}, \quad \mathcal{S} = 0, \tag{6.72}$$

and since \mathcal{S}_{MN} is transformed covariantly under the PL T -duality, the energy–momentum tensor \mathcal{E}_{MN} should also transform covariantly,

$$\mathcal{E}'_{MN} = (h \mathcal{E} h^T)_{MN}. \tag{6.73}$$

The energy–momentum tensor \mathcal{E}_{MN} is a bilinear form of the combination $\mathcal{F} \equiv e^d F$ and it does not contain a derivative of \mathcal{F} . Therefore, we can covariantly transform \mathcal{E}_{MN} simply by rotating the combination \mathcal{F} covariantly, and this gives the transformation rule for the R–R fields.

Under a PL T -plurality, $\mathcal{H}'_{MN} = (h \mathcal{H} h^T)_{MN}$ with $h = U' C U^{-1}$, the $O(n, n)$ -covariant transformation rule for a scalar density e^{-2d} is

$$e^{-2d^{(h)}} = \frac{|\det(e_a^m)|}{|\det(e_a^m)|} e^{-2d}. \tag{6.74}$$

Indeed, the twist matrix has the form $U = R \mathbf{\Pi}$, and the scalar density is invariant under the β -transformation $\mathbf{\Pi}$ while it is multiplied by $|\det(e_a^m)|^{-1}$ under the twist R . Moreover, the scalar density is invariant under the $O(n, n)$ transformation C by the definition of C_A^B . Thus, $e^{-2d^{(h)}}$ in Eq. (6.74) is the covariantly transformed DFT dilaton.

On the other hand, let us denote the covariantly transformed R–R polyform as $\mathcal{F}^{(h)}$. By denoting the action of an $O(n, n)$ transformation h on the polyform as $F \rightarrow \mathbb{S}_h F$,⁹ we have

$$F^{(h)} = \mathbb{S}'_U \mathbb{S}_C \mathbb{S}_U^{-1} F. \tag{6.75}$$

Then, the energy–momentum tensor made of the combination $e^{d^{(h)}} F^{(h)}$ is the expected \mathcal{E}'_{MN} . However, importantly, the actual DFT dilaton is given by

$$|\det(a'^{-1})| |\det(e_a^m)| e^{-2[d' - \bar{d}(x')]} = |\det(a^{-1})| |\det(e_a^m)| e^{-2[d - \bar{d}(x)]}, \tag{6.76}$$

and $e^{d'}$ is related to the covariant one $e^{d^{(h)}}$ as

$$e^{d'} = \frac{\sqrt{|\det a|} e^{-\bar{d}(x)}}{\sqrt{|\det a'|} e^{-\bar{d}(x')}} e^{d^{(h)}}. \tag{6.77}$$

Therefore, if we identify the dual R–R polyform as

$$F' \equiv \frac{\sqrt{|\det a'|} e^{-\bar{d}(x')}}{\sqrt{|\det a|} e^{-\bar{d}(x)}} F^{(h)}, \tag{6.78}$$

the energy–momentum tensor made from $e^{d'} F' = e^{d^{(h)}} F^{(h)}$ is \mathcal{E}'_{MN} . Namely, Eq. (6.78) is the rule for the R–R fields.

Now, as \mathcal{E}_{MN} is transformed covariantly, it is already clear that the equations of motion for \mathcal{H}_{MN} and d are satisfied in the dual background. However, the equation of motion for the R–R fields is still not clear. To clarify the covariance, let us rewrite Eq. (6.78) as

$$\hat{\mathcal{F}}' = \mathbb{S}_C \hat{\mathcal{F}}, \tag{6.79}$$

where we have defined

$$\hat{\mathcal{F}} \equiv \frac{e^{\bar{d}}}{\sqrt{|\det a|}} \mathbb{S}_{U^{-1}} F = \frac{e^d}{\sqrt{|\det(e_a^m)|}} \mathbb{S}_{U^{-1}} F. \tag{6.80}$$

Then, we find that the $\hat{\mathcal{F}}$ is precisely the R–R field strength appearing in the gauged DFT or the flux formulation of DFT [see Eq. (B.56)],

$$|\mathcal{F}\rangle = \sum_p \frac{1}{p!} \hat{\mathcal{F}}_{a_1 \dots a_p} \Gamma^{a_1 \dots a_p} |0\rangle. \tag{6.81}$$

Here, $\Gamma^{a_1 \dots a_p} \equiv \Gamma^{[a_1 \dots a_p]}$ and $(\Gamma^A) \equiv (\Gamma^a, \Gamma_a)$ satisfy the algebra

$$\{\Gamma^A, \Gamma^B\} = \eta^{AB}, \tag{6.82}$$

⁹ An explicit form of the operation \mathbb{S}_h is given in Appendix B.

and the so-called Clifford vacuum $|0\rangle$ is defined by $\Gamma_a |0\rangle = 0$. By using a nilpotent operator,

$$\nabla = \partial - \frac{1}{2} \Gamma^A \mathcal{F}_A + \frac{1}{3!} \Gamma^{ABC} \mathcal{F}_{ABC} \quad (\partial \equiv \Gamma^A \mathcal{D}_A), \tag{6.83}$$

the Bianchi identity can be expressed as (see Appendix B)

$$\nabla |\mathcal{F}\rangle = \left(\partial - \frac{1}{2} \Gamma^A \mathcal{F}_A + \frac{1}{3!} \Gamma^{ABC} \mathcal{F}_{ABC} \right) |\mathcal{F}\rangle = 0. \tag{6.84}$$

As is well known in the democratic formulation [157,160], the Bianchi identity is equivalent to the equations of motion when the self-duality relation $G_p = (-1)^{\frac{p(p-1)}{2}} * G_{10-p}$ is satisfied.

Now, we require the dualizability condition for the R–R fields,

$$\partial |\mathcal{F}\rangle = 0, \tag{6.85}$$

which will be the same as the proposal of Ref. [139]. Then, the Bianchi identity or the equation of motion for the R–R fields becomes an algebraic equation:

$$\left(\frac{1}{3!} \Gamma^{ABC} \mathcal{F}_{ABC} - \frac{1}{2} \Gamma^A \mathcal{F}_A \right) |\mathcal{F}\rangle = 0. \tag{6.86}$$

Note that when the dual algebra is non-unimodular, \mathcal{F}_A should be modified as $\mathcal{F}_A + 2 U_A^M \mathbf{X}_M$ as we explained in the discussion of the NS–NS fields. By denoting the spinor representative of the $O(n, n)$ transformation by S_C , the duality relation of Eq. (6.79) becomes simply

$$|\mathcal{F}'\rangle = S_C |\mathcal{F}\rangle. \tag{6.87}$$

Then, the equation of motion in Eq. (6.86) after the $O(n, n)$ PL T -plurality transformation is

$$\left(\frac{1}{3!} \Gamma^{ABC} C_A^D C_B^E C_C^F \mathcal{F}_{ABC} - \frac{1}{2} \Gamma^A C_A^B \mathcal{F}_A \right) S_C |\hat{\mathcal{F}}\rangle = 0, \tag{6.88}$$

but from the relations $S_C^{-1} \Gamma_A S_C = C_A^B \Gamma_B$ and $C_A^C C_B^D \eta_{CD} = \eta_{AB}$ this is equivalent to the equation of motion in the original background, Eq. (6.86). In this manner, the equation of motion for the R–R fields in Eq. (6.86) is also covariantly transformed.

We call the object $\hat{\mathcal{F}}$ the untwisted R–R fields, and once $\hat{\mathcal{F}}'_{a_1 \dots a_p}$ in the dual background is determined from Eq. (6.79), we can construct the Page form in the dual background as

$$F' = e^{-\bar{d}(x')} \sqrt{|\det a'|} \mathbb{S}_{U'}, \hat{\mathcal{F}}' = e^{-\bar{d}(x')} \sqrt{|\det a'|} e^{-\Pi' \vee} \left(\sum_p \frac{1}{p!} \hat{\mathcal{F}}'_{a_1 \dots a_p} r'^{a_1} \wedge \dots \wedge r'^{a_p} \right), \tag{6.89}$$

where $\Pi' \vee \equiv \frac{1}{2} \Pi'^{ab} \iota_{e'_a} \iota_{e'_b}$.

6.2.4. Spectator fields

In the following, we consider more general cases where spectator fields are also included. Namely, we suppose that the original model takes the form

$$S = -\frac{1}{4\pi \alpha'} \int_{\Sigma} d^2 \sigma \sqrt{-\gamma} (\gamma^{ab} - \varepsilon^{ab}) \left(\partial_a y^\mu \quad r_i^a \partial_a x^i \right) \begin{pmatrix} E_{\mu\nu} & E_{\mu b} \\ E_{a\nu} & E_{ab} \end{pmatrix} \begin{pmatrix} \partial_b y^\nu \\ r_j^b \partial_b x^j \end{pmatrix}. \tag{6.90}$$

Here, we denote the coordinates as $(x^m) = (y^\mu, x^i)$ ($i = 1, \dots, n$). By assuming that the background field $(E_{mn}) = \begin{pmatrix} E_{\mu\nu} & E_{\mu b} \\ E_{a\nu} & E_{ab} \end{pmatrix}$ satisfies the condition

$$\mathfrak{L}_{\nu a} E_{mn} = -\tilde{f}^{bc}{}_a E_{mp} v_b^p v_c^q E_{qn}, \tag{6.91}$$

we can again determine E_{mn} as [37,38]

$$\begin{aligned} E_{\mu\nu} &= \hat{E}_{\mu\nu} + \hat{E}_{\mu c} \hat{E}^{cd} N_{de} \Pi^{ef} \hat{E}_{fv}, & E_{\mu b} &= \hat{E}_{\mu c} \hat{E}^{cd} N_{db}, \\ E_{a\nu} &= N_{ad} \hat{E}^{de} \hat{E}_{e\nu}, & E_{ab} &= N_{ab}, \end{aligned} \tag{6.92}$$

where $(N_{ab}) \equiv (\hat{E}^{ab} - \Pi^{ab})^{-1}$. This reduces to Eq. (6.48) when there is no spectator field. Now, an important difference is that \hat{E}_{mn} is not necessarily constant, but can depend on the spectator fields y^μ , $\hat{E}_{mn} = \hat{E}_{mn}(y)$. The dependence should be determined from the DFT equations of motion and is independent of the structure of the Drinfel'd double.

In terms of the generalized metric \mathcal{H}_{MN} , we can clearly see that the relation in Eq. (6.92) is a straightforward generalization of Eq. (6.30),

$$\begin{aligned} \mathcal{H}_{MN} &= [U(x) \hat{\mathcal{H}}(y) U^\top(x)]_{MN}, & U(x) &\equiv R \Pi, \\ (R_M{}^B) &\equiv \begin{pmatrix} \delta_\mu^\beta & 0 & 0 & 0 \\ 0 & r_i^b & 0 & 0 \\ 0 & 0 & \delta_\beta^\mu & 0 \\ 0 & 0 & 0 & e_b^i \end{pmatrix}, & (\Pi_A{}^B) &\equiv \begin{pmatrix} \delta_\alpha^\beta & 0 & 0 & 0 \\ 0 & \delta_a^b & 0 & 0 \\ 0 & 0 & \delta_\beta^\alpha & 0 \\ 0 & -\Pi^{ab} & 0 & \delta_b^a \end{pmatrix}, \end{aligned} \tag{6.93}$$

where $(x^M) = (y^\mu, x^i, \tilde{y}_\mu, \tilde{x}_i)$. The T -plurality transformation of Eq. (6.45) is also generalized, in a natural manner, as an $O(n, n)$ transformation,

$$\hat{\mathcal{H}}'_{AB} = (C \hat{\mathcal{H}} C^\top)_{AB} \quad (C_A{}^B) = \begin{pmatrix} \delta_\alpha^\beta & 0 & 0 & 0 \\ 0 & p_a^b & 0 & q_{ab} \\ 0 & 0 & \delta_\beta^\alpha & 0 \\ 0 & r^{ab} & 0 & s^a_b \end{pmatrix}. \tag{6.94}$$

The dilaton can also have an additional dependence on the spectators similar to Eq. (2.27),

$$e^{-2d} = e^{-2\hat{d}(y)} e^{-2\mathbf{d}(x)}, \quad e^{-2\mathbf{d}(x)} \equiv e^{-2\bar{d}(x)} |\det(\ell_i^a)|. \tag{6.95}$$

We also suppose that the untwisted R–R fields can depend on the spectator fields $\hat{\mathcal{F}} = \hat{\mathcal{F}}(y)$.

Then, by defining the fluxes \mathcal{F}_{ABC} and \mathcal{F}_A from $U_M{}^A(x)$ and $\mathbf{d}(x)$, we again obtain

$$\begin{aligned} \mathcal{F}_{ab}{}^c &= f_{ab}{}^c, & \mathcal{F}^{ab}{}_c &= \tilde{f}^{ab}{}_c, & \mathcal{F}_{abc} &= \mathcal{F}^{abc} = \mathcal{F}_{\alpha BC} = \mathcal{F}^\alpha{}_{BC} = 0, \\ (\mathcal{F}_A) &= (\mathcal{F}_\alpha, \mathcal{F}_a, \mathcal{F}^\alpha, \mathcal{F}^a) = (0, 2\mathcal{D}_a \mathbf{d}, 0, 2\mathcal{D}^a \mathbf{d}). \end{aligned} \tag{6.96}$$

Here, we again need to perform the shift $\partial^M d \rightarrow \partial^M d + \mathbf{X}^M$, Eq. (6.63), when the dual algebra is non-unimodular.

The requirement in Eq. (2.28) is automatically satisfied with our twist matrix, and by using Eq. (2.29) the dilaton equation of motion becomes

$$\hat{S} + \frac{1}{12} \mathcal{F}_{ABC} \mathcal{F}_{DEF} (3 \hat{\mathcal{H}}^{AD} \eta^{BE} \eta^{CF} - \hat{\mathcal{H}}^{AD} \hat{\mathcal{H}}^{BE} \hat{\mathcal{H}}^{CF}) - \hat{\mathcal{H}}^{AB} \mathcal{F}_A \mathcal{F}_B$$

$$-\frac{1}{2} \mathcal{F}^A{}_{BC} \hat{\mathcal{H}}^{BD} \hat{\mathcal{H}}^{CE} \mathcal{D}_D \hat{\mathcal{H}}_{AE} + 2 \mathcal{F}_A \mathcal{D}_B \hat{\mathcal{H}}^{AB} - 4 \mathcal{F}_A \hat{\mathcal{H}}^{AB} \mathcal{D}_B \hat{d} = 0. \tag{6.97}$$

By requiring that the untwisted fields $\{\hat{\mathcal{H}}_{AB}(y), \hat{d}(y), \hat{\mathcal{F}}(y)\}$ in the original and the dual background are covariantly related by the $O(n, n)$ transformation,

$$\hat{\mathcal{H}}_{AB} = (C \hat{\mathcal{H}} C^T)_{AB}, \quad \hat{d}' = \hat{d}, \quad \hat{\mathcal{F}}' = \mathbb{S}_C \hat{\mathcal{F}}, \tag{6.98}$$

we can easily see that $\mathcal{D}_C \hat{\mathcal{H}}_{AB} = \partial_C \hat{\mathcal{H}}_{AB}$ and $\mathcal{D}_A \hat{d} = \partial_A \hat{d}$ are also transformed covariantly,

$$\begin{aligned} \mathcal{D}'_C \hat{\mathcal{H}}'_{AB}(y) &= C_A{}^D C_A{}^E \partial_C \hat{\mathcal{H}}_{DE}(y) = C_C{}^F C_A{}^D C_A{}^E \mathcal{D}_F \hat{\mathcal{H}}_{DE}(y), \\ \mathcal{D}'_A \hat{d}(y) &= \partial_A \hat{d}(y) = C_A{}^B \mathcal{D}_B \hat{d}(y), \quad \mathcal{D}'_A \hat{\mathcal{F}}(y) = \partial_A \hat{\mathcal{F}}(y) = C_A{}^B \mathcal{D}_B \hat{\mathcal{F}}(y). \end{aligned} \tag{6.99}$$

Then, the dilaton equation of motion in Eq. (6.97) is satisfied in the dualized background if it is satisfied in the original background. As long as the untwisted R–R field satisfies “the Bianchi identity” $\partial|\mathcal{F}\rangle = 0$, which is equivalent to $d\hat{\mathcal{F}}(y) = 0$, the equation of motion for the R–R fields is again a simple algebraic equation,

$$\left(\frac{1}{3!} \Gamma^{ABC} \mathcal{F}_{ABC} - \frac{1}{2} \Gamma^A \mathcal{F}_A \right) |\mathcal{F}(y)\rangle = 0, \tag{6.100}$$

and its covariance is manifest. The covariance of the equations of motion for the generalized metric $S_{MN} = \mathcal{E}_{MN}$ can also be shown in a similar manner. Since the computation is a little complicated, the details are discussed in Appendix B.

7. PL T -plurality for $\text{AdS}_5 \times S^5$

In this section we show an example of the Poisson–Lie T -plurality. As already mentioned, the Lie algebra \mathfrak{d} of the Drinfel’d doubles can be realized as a direct sum of two maximally isotropic algebras \mathfrak{g} and $\tilde{\mathfrak{g}}$, and $(\mathfrak{d}, \mathfrak{g}, \tilde{\mathfrak{g}})$ is called the Manin triple. Following Ref. [54], we denote the pair simply as $(\mathfrak{g}|\tilde{\mathfrak{g}})$. The classification of six-dimensional real Drinfel’d doubles was worked out in Ref. [54], where the following series of Manin triples corresponding to a single Drinfel’d double \mathfrak{d} was found:

$$\begin{aligned} (5|1) &\cong (6_0|1) \cong (5|2.i) \cong (6_0|5.ii) \\ &\cong (1|5) \cong (1|6_0) \cong (2.i|5) \cong (5.ii|6_0). \end{aligned} \tag{7.1}$$

Here, the characters in each slot denote the Bianchi type of the three-dimensional Lie algebra,

$$\begin{aligned} \mathbf{1} : & \quad [X_1, X_2] = 0 & \quad [X_2, X_3] = 0, & \quad [X_3, X_1] = 0, \\ \mathbf{2.i} : & \quad [X_1, X_2] = 0, & \quad [X_2, X_3] = X_1, & \quad [X_3, X_1] = 0, \\ \mathbf{5} : & \quad [X_1, X_2] = -X_2, & \quad [X_2, X_3] = 0, & \quad [X_3, X_1] = X_3, \\ \mathbf{5.ii} : & \quad [X_1, X_2] = -X_1 + X_2, & \quad [X_2, X_3] = X_3 & \quad [X_3, X_1] = -X_3, \\ \mathbf{6_0} : & \quad [X_1, X_2] = 0, & \quad [X_2, X_3] = X_1, & \quad [X_3, X_1] = -X_2. \end{aligned} \tag{7.2}$$




Fig. 1. The PL T -plurality procedure.

Using an $O(3, 3)$ transformation $T'_A = C_A{}^B T_B$,¹⁰ the PL T -plurality for this chain of Manin triples was studied in Ref. [55]. However, in Ref. [55], since the initial background is the flat space (or the Bianchi type V universe) the R–R fields were absent in any of the dual backgrounds. Moreover, there has been an issue in the treatment of the dual-coordinate dependence of the dilaton, known as the dilaton puzzle (see also Refs. [56–58] for detailed discussion of the issue). Accordingly, the only three backgrounds discussed in Ref. [55] were

$$(5|1) \cong (6_0|1) \cong (5|2.i). \tag{7.3}$$

In this section we identify the $\text{AdS}_5 \times S^5$ solution as a background with the $(5|1)$ symmetry, and write down all of the eight backgrounds associated with the Manin triples given in Eq. (7.1).

For convenience, we summarize the procedure of the PL T -plurality in Fig. 1. We first prepare the untwisted fields $\{\hat{\mathcal{H}}_{AB}(y), \hat{d}(y), \hat{\mathcal{F}}(y)\}$ that satisfy

$$\mathcal{D}_A \hat{\mathcal{H}}_{BC}(y) = \partial_A \hat{\mathcal{H}}_{BC}(y), \quad \mathcal{D}_A \hat{d}(y) = \partial_A \hat{d}(y), \quad \mathcal{D}_A \hat{\mathcal{F}}(y) = \partial_A \hat{\mathcal{F}}(y). \tag{7.4}$$

They are independent of the structure of the Drinfel'd double and can be chosen freely. Under the $O(n, n)$ PL T -plurality they are transformed covariantly,

$$\hat{\mathcal{H}}_{AB} \rightarrow (C \hat{\mathcal{H}} C^\top)_{AB}, \quad \hat{d} \rightarrow \hat{d}, \quad \hat{\mathcal{F}} \rightarrow \mathbb{S}_C \hat{\mathcal{F}}. \tag{7.5}$$

By using the generators T_A in each frame, we construct the twist matrix U as

$$\begin{aligned}
 U(x) &\equiv R \Pi, & \Pi \vee &\equiv \frac{1}{2} \Pi^{ab} \iota_{e_a} \iota_{e_b}, & \text{Ad}_{g^{-1}} &= \begin{pmatrix} \delta_a^c & 0 \\ \Pi^{ac} & \delta_c^a \end{pmatrix} \begin{pmatrix} a_c^b & 0 \\ 0 & (a^{-\top})^c_b \end{pmatrix}, \\
 (R_M{}^B) &\equiv \begin{pmatrix} \delta_\mu^\beta & 0 & 0 & 0 \\ 0 & r_i^b & 0 & 0 \\ 0 & 0 & \delta_\beta^\mu & 0 \\ 0 & 0 & 0 & e_b^i \end{pmatrix}, & (\Pi_A{}^B) &\equiv \begin{pmatrix} \delta_\alpha^\beta & 0 & 0 & 0 \\ 0 & \delta_a^b & 0 & 0 \\ 0 & 0 & \delta_\beta^\alpha & 0 \\ 0 & -\Pi^{ab} & 0 & \delta_b^a \end{pmatrix}.
 \end{aligned} \tag{7.6}$$

¹⁰ As pointed out in Ref. [62], the matrix C which connects two Manin triples may not be unique, and a different choice of C may give a different background. We will use the matrices C that are given in Ref. [54].

Originally, the indices A, B in T_A and $C_A{}^B$ run from 1 to $2n$ ($n = 3$ here), but we extend the matrix $C_A{}^B$ as in Eq. (6.94); T_A should then be understood as $(T_A) = (T_\alpha, T_a, \tilde{T}^\alpha, \tilde{T}^a) = (0, T_a, 0, \tilde{T}^a)$.

Then, by twisting the untwisted fields, we construct the DFT fields as

$$\begin{aligned} \mathcal{H}_{MN} &= [U(x) \hat{\mathcal{H}}(y) U^T(x)]_{MN}, \\ e^{-2d} &= e^{-2\hat{d}(y)} e^{-2\bar{d}(x)} |\det(\ell_i^a)|, \quad (\mathbf{X}^M) = \begin{pmatrix} \frac{1}{2} \tilde{f}^{ba} v_a^i \delta_i^m \\ 0 \end{pmatrix}, \\ F &= e^{-\bar{d}(x)} \sqrt{|\det a|} e^{-\Pi(x) \vee} \left[\sum_p \frac{1}{p!} \hat{\mathcal{F}}_{a_1 \dots a_p}(y) r^{a_1} \wedge \dots \wedge r^{a_p} \right]. \end{aligned} \tag{7.7}$$

The function $\bar{d}(x)$ is given in the initial configuration, and after the PL T -plurality it is rewritten in the new coordinates determined through $g(x^i) \tilde{g}(\tilde{x}_i) = l = g'(x'^i) \tilde{g}'(\tilde{x}'_i)$. When $\bar{d}(x)$ has a linear dual-coordinate dependence $d^i \tilde{x}_i$, we make a redefinition and absorb the dependence into the Killing vector, $I^i = \frac{1}{2} \tilde{f}^{ba} v_a^i + d^i$.

7.1. **(5|1):** $AdS_5 \times S^5$

We start with the $AdS_5 \times S^5$ background (in a non-standard coordinate system):

$$\begin{aligned} ds^2 &= \frac{-dt^2 + t^2 [dx_1^2 + e^{-2x^1} (dx_2^2 + dx_3^2)] + dz^2}{z^2} + ds_{S^5}^2, \\ G_5 &= \frac{-4e^{-2x^1} t^3 dt \wedge dx^1 \wedge dx^2 \wedge dx^3 \wedge dz}{z^5} + 4\omega_5, \end{aligned} \tag{7.8}$$

where

$$\begin{aligned} ds_{S^5}^2 &\equiv dr^2 + \sin^2 r d\xi^2 + \cos^2 \xi \sin^2 r d\phi_1^2 + \sin^2 r \sin^2 \xi d\phi_2^2 + \cos^2 r d\phi_3^2, \\ \omega_5 &\equiv \sin^3 r \cos r \sin \xi \cos \xi dr \wedge d\xi \wedge d\phi_1 \wedge d\phi_2 \wedge d\phi_3. \end{aligned} \tag{7.9}$$

This background has Killing vectors

$$v_1 \equiv \partial_1 + x^2 \partial_2 + x^3 \partial_3, \quad v_2 \equiv \partial_2, \quad v_3 \equiv \partial_3 \tag{7.10}$$

satisfying the **(5|1)** algebra,

$$[v_a, v_b] = f_{ab}{}^c v_c, \quad f_{12}{}^2 = f_{13}{}^3 = -1, \quad \mathfrak{L}_{v_a} E_{mn} = 0. \tag{7.11}$$

We can reconstruct this background by providing the parameterization

$$l = g \tilde{g}, \quad g = e^{x^1 T_1} e^{x^2 T_2} e^{x^3 T_3}, \quad \tilde{g} = e^{\tilde{x}_1 \tilde{T}^1} e^{\tilde{x}_2 \tilde{T}^2} e^{\tilde{x}_3 \tilde{T}^3}, \tag{7.12}$$

where $(T_A) = (T_a, \tilde{T}^a)$ are generators of the Manin triple **(5|1)**. We obtain

$$\begin{aligned} \ell &= dx^1 T_1 + (dx^2 - x^2 dx^1) T_2 + (dx^3 - x^3 dx^1) T_3, \\ r &= dx^1 T_1 + e^{-x^1} (dx^2 T_2 + dx^3 T_3), \end{aligned} \tag{7.13}$$

$$a = \begin{pmatrix} 1 & -x^2 & -x^3 \\ 0 & e^{x^1} & 0 \\ 0 & 0 & e^{x^1} \end{pmatrix}, \quad \Pi^{ab} = 0, \tag{7.14}$$

and they give the twist matrix U_M^A . We can easily determine the untwisted metric from the relation $\hat{\mathcal{H}}_{MN} = (U^{-1} \mathcal{H} U^{-T})_{MN}$, and the result is

$$(\hat{E}_{mn}) = \text{diag}\left(-\frac{1}{z^2}, \frac{t^2}{z^2}, \frac{t^2}{z^2}, \frac{t^2}{z^2}, \frac{1}{z^2}, 1, \sin^2 r, \sin^2 r \cos^2 \xi, \sin^2 r \sin^2 \xi, \cos^2 r\right), \quad (7.15)$$

in the coordinate system $(x^m) = (t, x^1, x^2, x^3, z, r, \xi, \phi_1, \phi_2, \phi_3)$. Since the dilaton is absent, $\Phi = 0$, the DFT dilaton becomes

$$e^{-2d} = \sqrt{|g|} = \frac{t^3 e^{-2x^1} \sin^3 r \cos r \sin \xi \cos \xi}{z^5}. \quad (7.16)$$

We also have $|\det(\ell_m^a)| = 1$, and we can identify $\hat{d}(y)$ and $\bar{d}(x)$ as

$$e^{-2d} = e^{-2\hat{d}(y)} e^{-2\bar{d}(x)}, \quad e^{-2\hat{d}(y)} \equiv \frac{t^3 \sin^3 r \cos r \sin \xi \cos \xi}{z^5}, \quad e^{-2\bar{d}(x)} \equiv e^{-2x^1}. \quad (7.17)$$

From this, we obtain the $(x^1, x^2, x^3, \tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$ -components of the single-index flux as

$$\mathcal{F}_A = (2, 0, 0, 0, 0, 0) \equiv \mathcal{F}_A^{(5|1)}. \quad (7.18)$$

In addition, from $e^{-\bar{d}(x)} \sqrt{|\det a|} = 1$, the untwisted R–R fields become

$$\hat{\mathcal{F}} \equiv \sum_p \frac{1}{p!} \hat{\mathcal{F}}_{a_1 \dots a_p} dx^{a_1} \wedge \dots \wedge dx^{a_p} = -\frac{4t^3 dt \wedge dx^1 \wedge dx^2 \wedge dx^3 \wedge dz}{z^5} + 4\omega_5, \quad (7.19)$$

which is a function of the spectator fields $(y^\mu) = (t, z, r, \xi, \phi_1, \phi_2, \phi_3)$, as expected.

Note that if we choose the untwisted fields as

$$(\hat{E}_{mn}) = \text{diag}(-1, t^2, t^2, t^2, 1, 1, 1, 1, 1, 1), \quad e^{-2\hat{d}} = t^3, \quad \hat{\mathcal{F}} = 0, \quad (7.20)$$

the purely NS–NS solutions studied in Ref. [55] can be recovered.

7.2. (1|5): Type IIA GSE

In order to consider the NATD background, we perform a redefinition of generators,

$$T'_A = C_A^B T_B^{(5|1)}, \quad C = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad (7.21)$$

and give a parameterization,

$$l = g' \tilde{g}', \quad g' = e^{x'^1} T'_1 e^{x'^2} T'_2 e^{x'^3} T'_3, \quad \tilde{g}' = e^{\tilde{x}'_1 \tilde{T}'_1} e^{\tilde{x}'_2 \tilde{T}'_2} e^{\tilde{x}'_3 \tilde{T}'_3}. \quad (7.22)$$

Then, from the identification with the original background,

$$g(x) \tilde{g}(\tilde{x}) = l = g'(x') \tilde{g}'(\tilde{x}'), \quad (7.23)$$

we find the following relation between the coordinates:

$$\begin{aligned} x^1 &= \tilde{x}'_1, & x^2 &= \tilde{x}'_2, & x^3 &= \tilde{x}'_3, \\ \tilde{x}_1 &= x'^1 + x'^2 e^{-\tilde{x}'_1 \tilde{x}'_2} + x'^3 e^{-\tilde{x}'_1 \tilde{x}'_3}, & \tilde{x}_2 &= e^{-\tilde{x}'_1} x'^2, & \tilde{x}_3 &= e^{-\tilde{x}'_1} x'^3. \end{aligned} \tag{7.24}$$

From this relation, we can identify \bar{d} as

$$e^{-2\bar{d}} = e^{-2x^1} = e^{-2\tilde{x}'_1}. \tag{7.25}$$

For notational simplicity, in the following we drop the prime.

The untwisted fields in this frame become

$$\begin{aligned} (\hat{E}_{mn}) &= \text{diag}\left(-\frac{1}{z^2}, \frac{z^2}{t^2}, \frac{z^2}{t^2}, \frac{z^2}{t^2}, \frac{1}{z^2}, 1, \sin^2 r, \sin^2 r \cos^2 \xi, \sin^2 r \sin^2 \xi, \cos^2 r\right), \\ e^{-2\hat{d}} &= \frac{t^3 \sin^3 r \cos r \sin \xi \cos \xi}{z^5}, & \hat{\mathcal{F}} &= -\frac{4t^3 dt \wedge dz}{z^5} + 4\omega_5 \wedge dx^1 \wedge dx^2 \wedge dx^3, \end{aligned} \tag{7.26}$$

and we twist them by using the quantities

$$\begin{aligned} \ell &= dx^1 T_1 + dx^2 T_2 + dx^3 T_3, & r &= dx^1 T_1 + dx^2 T_2 + dx^3 T_3, \\ v_1 &= \partial_1, & v_2 &= \partial_2, & v_3 &= \partial_3, \end{aligned} \tag{7.27}$$

$$a = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \Pi^{ab} = -\tilde{f}^{ab}{}_c x^c. \tag{7.28}$$

The resulting metric and the B -field are

$$\begin{aligned} ds^2 &= \frac{-dt^2 + dz^2}{z^2} + \frac{z^2 [t^4 (dx_1^2 + dx_2^2 + dx_3^2) + z^4 (x^3 dx^2 - x^2 dx^3)^2]}{t^2 [t^4 + (x_2^2 + x_3^2) z^4]} + ds_{S^5}^2, \\ B_2 &= \frac{z^4 dx^1 \wedge (x^2 dx^2 + x^3 dx^3)}{t^4 + (x_2^2 + x_3^2) z^4}. \end{aligned} \tag{7.29}$$

Since the dual algebra $\mathfrak{5}$ is non-unimodular, we need to introduce the Killing vector

$$I = \frac{1}{2} \tilde{f}^{ba}{}_b v_a^i \partial_i = \partial_1. \tag{7.30}$$

We can check that the flux \mathcal{F}_A is transformed covariantly from the original one, $\mathcal{F}_B^{(51)}$,

$$\mathcal{F}_A = (0, 0, 0, 2, 0, 0) = C_A{}^B \mathcal{F}_B^{(51)}, \tag{7.31}$$

which ensures that the equations of motion are transformed covariantly. In order to make the background a solution of GSE we make the redefinition in Eq. (6.67), which gives

$$\bar{d} = 0, \quad I = \left(\frac{1}{2} \tilde{f}^{ba}{}_b v_a^i + \tilde{\partial}^i \bar{d}\right) \partial_i = 2 \partial_1. \tag{7.32}$$

After this redefinition, the dual geometry becomes

$$ds^2 = \frac{-dt^2 + dz^2}{z^2} + \frac{z^2 [t^4 (dx_1^2 + dx_2^2 + dx_3^2) + z^4 (x^3 dx^2 - x^2 dx^3)^2]}{t^2 [t^4 + (x_2^2 + x_3^2) z^4]} + ds_{S^5}^2,$$

$$\begin{aligned}
 e^{-2\Phi} &= \frac{t^2 [t^4 + (x_2^2 + x_3^2) z^4]}{z^6}, & B_2 &= \frac{z^4 dx^1 \wedge (x^2 dx^2 + x^3 dx^3)}{t^4 + (x_2^2 + x_3^2) z^4}, & (7.33) \\
 G_2 &= -\frac{4t^3 dt \wedge dz}{z^5}, & G_4 &= -\frac{4t^3 dt \wedge dx^1 \wedge (x^2 dx^2 + x^3 dx^3) \wedge dz}{[t^4 + (x_2^2 + x_3^2) z^4] z}, & I &= 2 \partial_1,
 \end{aligned}$$

which is a solution of type IIA GSE. We can explicitly check that this background has the (1|5) symmetry,

$$[v_a, v_b] = f_{ab}{}^c v_c = 0, \quad \mathfrak{L}_{v_a} E^{mn} = \tilde{f}^{bc}{}_a v_b^m v_c^n. \quad (7.34)$$

A formal T -duality along the x^1 -direction gives a simple solution of type IIB supergravity,

$$\begin{aligned}
 ds^2 &= \frac{-dt^2 + dz^2 + t^2 dx_1^2}{z^2} + \frac{z^2 [(dx^2 - x^2 dx^1)^2 + (dx^3 - x^3 dx^1)^2]}{t^2} + ds_{S^5}^2, \\
 \Phi &= \ln\left(\frac{z^2}{t^2}\right) + 2x^1, & G_3 &= \frac{4t^3 e^{-2x^1} dt \wedge dx^1 \wedge dz}{z^5}. & (7.35)
 \end{aligned}$$

7.3. (6₀|1): Type IIA SUGRA

We next perform the following redefinition of the original (5|1) generators:

$$T'_A = C_A{}^B T_B^{(5|1)}, \quad C = \begin{pmatrix} 0 & 0 & -\frac{1}{2} & 0 & 1 & 0 \\ 0 & 0 & \frac{1}{2} & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & -1 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}. \quad (7.36)$$

This time, we provide the parameterization

$$l = g' \tilde{g}', \quad g' = e^{-x^3} T'_3 e^{x^2} T'_2 e^{x^1} T'_1, \quad \tilde{g}' = e^{\tilde{x}'_1 \tilde{T}'^1} e^{\tilde{x}'_2 \tilde{T}'^2} e^{-\tilde{x}'_3 \tilde{T}'^3}, \quad (7.37)$$

and the coordinates are related to the original ones as

$$\begin{aligned}
 x^1 &= x'^3, & x^2 &= \frac{\tilde{x}'_1 + \tilde{x}'_2}{2}, & x^3 &= \frac{x'^2 - x'^1}{2}, \\
 \tilde{x}_1 &= \tilde{x}'_3 + \frac{(x'^1 + x'^2)(\tilde{x}'_1 + \tilde{x}'_2)}{2}, & \tilde{x}_2 &= x'^1 + x'^2, & \tilde{x}_3 &= \tilde{x}'_2 - \tilde{x}'_1. & (7.38)
 \end{aligned}$$

Then, in this frame, \bar{d} becomes

$$e^{-2\bar{d}} = e^{-2x^1} = e^{-2x'^3}. \quad (7.39)$$

Again we remove the prime, and then the (t, x^1, x^2, x^3, z) -part of the untwisted metric becomes

$$(\hat{E}_{mn}) = \begin{pmatrix} -\frac{1}{z^2} & 0 & 0 & 0 & 0 \\ 0 & \frac{t^2}{4z^2} + \frac{z^2}{t^2} & \frac{z^2}{t^2} - \frac{t^2}{4z^2} & 0 & 0 \\ 0 & \frac{z^2}{t^2} - \frac{t^2}{4z^2} & \frac{t^2}{4z^2} + \frac{z^2}{t^2} & 0 & 0 \\ 0 & 0 & 0 & \frac{t^2}{z^2} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{z^2} \end{pmatrix}. \quad (7.40)$$

In order to obtain the untwisted R–R fields, it may be useful to decompose the matrix C into products of $GL(D)$ transformation, B -transformation, T -dualities, and β -transformation. In this case, for example, we can use the decomposition

$$C = \begin{pmatrix} 0 & 1 & -\frac{1}{2} & 0 & 0 & 0 \\ 0 & 1 & \frac{1}{2} & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{2} & -1 \\ 0 & 0 & 0 & 0 & \frac{1}{2} & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{7.41}$$

Then, the T -duality along the x^2 -direction and the $GL(3)$ transformation give

$$\hat{\mathcal{F}} = \frac{2t^3 dt \wedge (dx^1 - dx^2) \wedge dx^3 \wedge dz}{z^5} - 4(dx^1 + dx^2) \wedge \omega_5. \tag{7.42}$$

In order to obtain the twist matrix, we compute

$$\begin{aligned} \ell &= (dx^1 + x^2 dx^3) T_1 + (dx^2 + x^2 dx^3) T_2 - dx^3 T_3, \\ r &= (\cosh x^3 dx^1 + \sinh x^3 dx^2) T_1 + (\sinh x^3 dx^1 + \cosh x^3 dx^2) T_2 - dx^3 T_3, \\ v_1 &= \partial_1, \quad v_2 = \partial_2, \quad v_3 = x^2 \partial_1 + x^1 \partial_2 - \partial_3, \end{aligned} \tag{7.43}$$

$$a = \begin{pmatrix} \cosh x^3 & -\sinh x^3 & 0 \\ -\sinh x^3 & \cosh x^3 & 0 \\ -x^2 & -x^1 & 1 \end{pmatrix}, \quad \Pi^{ab} = 0. \tag{7.44}$$

Again, the flux \mathcal{F}_A is transformed covariantly,

$$\mathcal{F}_A = (0, 0, -2, 0, 0, 0) = C_A{}^B \mathcal{F}_B^{(5|1)}. \tag{7.45}$$

The background fields are determined as

$$\begin{aligned} ds^2 &= \frac{-dt^2 + t^2 dx_3^2 + dz^2}{z^2} + \frac{e^{-2x^3} t^2 (dx^1 - dx^2)^2}{4z^2} + \frac{e^{2x^3} z^2 (dx^1 + dx^2)^2}{t^2} + ds_{S^5}^2, \\ e^{-2\Phi} &= \frac{e^{-2x^3} t^2}{z^2}, \quad G_4 = \frac{2e^{-2x^3} t^3 (dx^1 - dx^2) \wedge dt \wedge dx^3 \wedge dz}{z^5}, \end{aligned} \tag{7.46}$$

and this is a solution of type IIA supergravity.

7.4. $(\mathbf{1}|\mathbf{6}_0)$: Type IIB GSE

The NATD of the $(\mathbf{6}_0|\mathbf{1})$ background, namely $(\mathbf{1}|\mathbf{6}_0)$, can be realized by

$$T'_A = C_A{}^B T_B^{(5|1)}, \quad C = \begin{pmatrix} 0 & \frac{1}{2} & 0 & 0 & 0 & -1 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & 1 & 0 \\ 0 & 0 & \frac{1}{2} & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{7.47}$$

We give the parameterization

$$l = g' \tilde{g}', \quad g' = e^{x^1 T'_1} e^{x^2 T'_2} e^{-x^3 T'_3}, \quad \tilde{g}' = e^{-\tilde{x}'_3 \tilde{T}'^3} e^{\tilde{x}'_2 \tilde{T}'^2} e^{\tilde{x}'_1 \tilde{T}'^1}. \quad (7.48)$$

In order to determine \bar{d} it is enough to identify the coordinate x^1 , and we find

$$e^{-2\bar{d}} = e^{-2x^1} = e^{-2\tilde{x}'_3}. \quad (7.49)$$

Note that the appearance of the dual-coordinate dependence was discussed in Ref. [55], but at that time DFT had not been developed and the interpretation was not clear.

We can construct the twist matrix U from

$$\begin{aligned} \ell &= dx^1 T_1 + dx^2 T_2 - dx^3 T_3, & r &= dx^1 T_1 + dx^2 T_2 - dx^3 T_3, \\ v_1 &= \partial_1, & v_2 &= \partial_2, & v_3 &= -\partial_3. \end{aligned} \quad (7.50)$$

$$a = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (\Pi^{ab}) = \begin{pmatrix} 0 & 0 & -x^2 \\ 0 & 0 & -x^1 \\ x^2 & x^1 & 0 \end{pmatrix}, \quad (7.51)$$

and the flux \mathcal{F}_A becomes

$$\mathcal{F}_A = (0, 0, 0, 0, 0, -2) = C_A{}^B \mathcal{F}_B^{(5|1)}. \quad (7.52)$$

Thus, the DFT equations of motion are covariantly transformed.

Although the dual algebra is unimodular, in order to absorb the dual coordinate dependence in \bar{d} we make a field redefinition, Eq. (6.67), and obtain

$$e^{-2\bar{d}} = 1, \quad I = \partial_3. \quad (7.53)$$

After the redefinition we obtain a solution of type IIB GSE,

$$\begin{aligned} ds^2 &= \frac{-dt^2 + dz^2}{z^2} + ds_{S^5}^2 \\ &+ \frac{t^6 (dx^1 + dx^2)^2 + 4t^2 z^4 [(dx^1 - dx^2)^2 + (x^1 dx^1 - x^2 dx^2)^2 + dx_3^2]}{t^4 z^2 [(x^1 + x^2)^2 + 4] + 4z^6 (x^1 - x^2)^2}, \\ B_2 &= \frac{t^4 (x^1 + x^2) (dx^1 + dx^2) - 4z^4 (x^1 - x^2) (dx^1 - dx^2)}{t^4 [(x^1 + x^2)^2 + 4] + 4z^4 (x^1 - x^2)^2} \wedge dx^3, \quad I = \partial_3, \\ e^{-2\Phi} &= \frac{t^4 [(x^1 + x^2)^2 + 4] + 4z^4 (x^1 - x^2)^2}{4z^4}, \quad G_3 = \frac{2t^3 (dx^1 + dx^2) \wedge dt \wedge dz}{z^5}, \\ G_5 &= 2(x^1 - x^2) \left[\frac{8t^3 z^{-1} dt \wedge dx^1 \wedge dx^2 \wedge dx^3 \wedge dz}{t^4 [(x^1 + x^2)^2 + 4] + 4z^4 (x^1 - x^2)^2} - 2\omega_{S^5} \right]. \end{aligned} \quad (7.54)$$

It is important to note that the duality $(\mathbf{6}_0|\mathbf{1}) \rightarrow (\mathbf{1}|\mathbf{6}_0)$ is a NATD for traceless structure constants. In the literature, it has been discussed that if the structure constants are traceless then the NATD background satisfies the supergravity equations of motion, but here we obtained a solution of GSE. The consistency is to be clarified in a future study. Of course, the existence of the R–R fields is not

important here. As already mentioned, we can obtain a purely NS–NS solution by starting with the untwisted fields, Eq. (7.20). The $(\mathbf{6}_0|\mathbf{1})$ background is

$$\begin{aligned}
 ds^2 &= -dt^2 + t^2 dx_3^2 + \frac{1}{4}e^{-2x^3}t^2(dx^1 - dx^2)^2 + e^{2x^3}t^{-2}(dx^1 + dx^2)^2 + ds_{T^6}^2, \\
 e^{-2\Phi} &= e^{-2x^3}t^2,
 \end{aligned}
 \tag{7.55}$$

while its NATD, namely the $(\mathbf{1}|\mathbf{6}_0)$ background, is a GSE solution,

$$\begin{aligned}
 ds^2 &= -dt^2 + ds_{T^6}^2 \\
 &\quad + \frac{t^6(dx^1 + dx^2)^2 + 4t^2[(dx^1 - dx^2)^2 + (x^1 dx^1 - x^2 dx^2)^2 + dx_3^2]}{t^4[(x^1 + x^2)^2 + 4] + 4(x^1 - x^2)^2}, \\
 B_2 &= \frac{t^4(x^1 + x^2)(dx^1 + dx^2) - 4(x^1 - x^2)(dx^1 - dx^2)}{t^4[(x^1 + x^2)^2 + 4] + 4(x^1 - x^2)^2} \wedge dx^3, \quad I = \partial_3, \\
 e^{-2\Phi} &= \frac{t^4[(x^1 + x^2)^2 + 4] + 4(x^1 - x^2)^2}{4}.
 \end{aligned}
 \tag{7.56}$$

It will be interesting to study string theory on these backgrounds in detail.

We also note that in the $(\mathbf{1}|\mathbf{6}_0)$ background, Eq. (7.54), if we perform a formal T -duality along the x^3 -direction we obtain a solution of type IIA supergravity,

$$\begin{aligned}
 ds^2 &= \frac{-dt^2 + dz^2}{z^2} + \frac{z^2[dx^1 - dx^2 - (x^1 - x^2)dx^3]^2}{t^2} \\
 &\quad + \frac{t^2[(dx^1 + dx^2)^2 + 2(x^1 + x^2)(dx^1 + dx^2)dx^3 + [(x^1 + x^2)^2 + 4]dx_3^2]}{4z^2} + ds_{S^5}^2, \\
 e^{-2\Phi} &= \frac{e^{-2x^3}t^2}{z^2}, \quad G_4 = -\frac{2e^{-x^3}t^3(dx^1 + dx^2) \wedge dt \wedge dx^3 \wedge dz}{z^5}.
 \end{aligned}
 \tag{7.57}$$

7.5. $(\mathbf{5}|\mathbf{2.i})$: Type IIB SUGRA

In order to obtain the Manin triple $(\mathbf{5}|\mathbf{2.i})$, we perform the redefinition

$$T'_A = C_A{}^B T_B^{(\mathbf{5}|\mathbf{1})}, \quad C = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & -\frac{1}{2} \\ 0 & 0 & 1 & 0 & \frac{1}{2} & 0 \end{pmatrix}.
 \tag{7.58}$$

Again we consider the parameterization

$$l = g' \tilde{g}', \quad g' = e^{x^1} T'_1 e^{x^2} T'_2 e^{-x^3} T'_3, \quad \tilde{g}' = e^{\tilde{x}'_1} \tilde{T}'^1 e^{\tilde{x}'_2} \tilde{T}'^2 e^{-\tilde{x}'_3} \tilde{T}'^3,
 \tag{7.59}$$

and from the coordinate transformation we obtain

$$e^{-2\bar{d}} = e^{-2x^1} = e^{2x^1}.
 \tag{7.60}$$

The necessary quantities are obtained as

$$\begin{aligned} \ell &= dx^1 T_1 + (dx^2 - x^2 dx^1) T_2 - (dx^3 - x^3 dx^1) T_3, \\ r &= dx^1 T_1 + e^{-x^1} (dx^2 T_2 - dx^3 T_3), \\ v_1 &= \partial_1 + x^2 \partial_2 + x^3 \partial_3, \quad v_2 = \partial_2, \quad v_3 = -\partial_3, \end{aligned} \tag{7.61}$$

$$a = \begin{pmatrix} 1 & -x^2 & x^3 \\ 0 & e^{x^1} & 0 \\ 0 & 0 & e^{x^1} \end{pmatrix}, \quad (\Pi^{ab}) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -e^{-x^1} \sinh x^1 \\ 0 & e^{-x^1} \sinh x^1 & 0 \end{pmatrix}, \tag{7.62}$$

and again the flux \mathcal{F}_A is covariantly transformed,

$$\mathcal{F}_A = (-2, 0, 0, 0, 0, 0) = C_A{}^B \mathcal{F}_B^{(5|1)}. \tag{7.63}$$

A straightforward computation gives

$$\begin{aligned} ds^2 &= \frac{-dt^2 + t^2 dx_1^2 + dz^2}{z^2} + \frac{4e^{2x^1} t^2 z^2 (dx_2^2 + dx_3^2)}{4e^{4x^1} t^4 + z^4} + ds_{S^5}^2, \\ B_2 &= -\frac{2z^4 dx^2 \wedge dx^3}{4e^{4x^1} t^4 + z^4}, \quad e^{-2\Phi} = \frac{4e^{4x^1} t^4 + z^4}{4z^4}, \\ G_3 &= -e^{2x^1} \frac{4t^3 dt \wedge dx^1 \wedge dz}{z^5}, \\ G_5 &= -\frac{8e^{2x^1} t^3}{z} \frac{dt \wedge dx^1 \wedge dx^2 \wedge dx^3 \wedge dz}{4e^{4x^1} t^4 + z^4} + 2\omega_{S^5}, \end{aligned} \tag{7.64}$$

and this is a solution of type IIB supergravity.

7.6. (2.i|5): Type IIA SUGRA

We next consider the transformation

$$T'_A = C_A{}^B T_B^{(5|1)}, \quad C = \begin{pmatrix} 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & -\frac{1}{2} \\ 0 & 0 & 1 & 0 & \frac{1}{2} & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \tag{7.65}$$

and provide the parameterization

$$l = g' \tilde{g}', \quad g' = e^{x^1} T'_1 e^{x^2} T'_2 e^{-x^3} T'_3, \quad \tilde{g}' = e^{\tilde{x}^1} \tilde{T}'^1 e^{\tilde{x}^2} \tilde{T}'^2 e^{-\tilde{x}^3} \tilde{T}'^3. \tag{7.66}$$

The coordinate transformation gives

$$e^{-2\tilde{d}} = e^{-2x^1} = e^{2\tilde{x}^1}. \tag{7.67}$$

Again, we compute

$$\begin{aligned} \ell &= (dx^1 - x^3 dx^2) T_1 + dx^2 T_2 - dx^3 T_3, \\ r &= (dx^1 - x^2 dx^3) T_1 + dx^2 T_2 - dx^3 T_3, \end{aligned}$$

$$v_1 = \partial_1, \quad v_2 = x^3 \partial_1 + \partial_2, \quad v_3 = -\partial_3, \tag{7.68}$$

$$a = \begin{pmatrix} 1 & 0 & 0 \\ -x^3 & 1 & 0 \\ -x^2 & 0 & 1 \end{pmatrix}, \quad (\Pi^{ab}) = \begin{pmatrix} 0 & x^2 & -x^3 \\ -x^2 & 0 & 0 \\ x^3 & 0 & 0 \end{pmatrix}, \tag{7.69}$$

and we can check that the flux is covariantly transformed,

$$\mathcal{F}_A = (0, 0, 0, -2, 0, 0) = C_A^B \mathcal{F}_B^{(5|1)}. \tag{7.70}$$

Since the dual algebra **5** is non-unimodular, we have

$$I = \frac{1}{2} \tilde{f}^{ba} v_a^m \partial_m = \partial_1. \tag{7.71}$$

We thus expect that this background is a solution of the GSE. However, according to the field redefinition in Eq. (6.67), we obtain

$$e^{-2\bar{d}} = 1, \quad I = \partial_1 - \partial_1 = 0. \tag{7.72}$$

As the result, we obtain a solution of the conventional type IIA supergravity,

$$\begin{aligned} ds^2 &= \frac{-dt^2 + dz^2}{z^2} + \frac{z^2 [4 dx^1 (dx^1 - x^3 dx^2 - x^2 dx^3) + (x^3 dx^2 + x^2 dx^3)^2]}{4 t^2 (1 + x_2^2 + x_3^2)} \\ &\quad + \frac{t^2 [dx_2^2 + dx_3^2 + (x^3 dx^2 - x^2 dx^3)^2]}{z^2 (1 + x_2^2 + x_3^2)} + ds_{S^5}^2, \\ B_2 &= \frac{dx^1 \wedge (x^2 dx^2 + x^3 dx^3)}{1 + x_2^2 + x_3^2} + \frac{(1 + 2x_2^2) dx^2 \wedge dx^3}{2 (1 + x_2^2 + x_3^2)}, \\ e^{-2\Phi} &= \frac{t^2 (1 + x_2^2 + x_3^2)}{z^2}, \quad G_4 = -\frac{4 t^3 dt \wedge dx^2 \wedge dx^3 \wedge dz}{z^5}. \end{aligned} \tag{7.73}$$

Namely, even if the dual algebra is non-unimodular, the background can satisfy the usual supergravity equations of motion. This is a remarkable example of such unusual cases.

7.7. (5.ii|6₀): Type IIB GSE

We next consider

$$T'_A = C_A^B T_B^{(5|1)}, \quad C = \begin{pmatrix} 1 & 0 & 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & -1 & 0 & 0 & 0 & \frac{1}{2} \\ 0 & 1 & 0 & 1 & 0 & \frac{1}{2} \\ -1 & 0 & 1 & 0 & \frac{1}{2} & 0 \end{pmatrix}, \tag{7.74}$$

and give the parameterization

$$l = g' \tilde{g}', \quad g' = e^{x^1 T'_1} e^{(x^2 - x^1) T'_2} e^{x^3 T'_3}, \quad \tilde{g}' = e^{\tilde{x}_3 \tilde{T}'_3} e^{\tilde{x}_2 \tilde{T}'_2} e^{(\tilde{x}'_1 + \tilde{x}'_2) \tilde{T}'_1}. \tag{7.75}$$

We then obtain \bar{d} as

$$e^{-2\bar{d}} = e^{-2x^1} = e^{-2(x^2 - \tilde{x}'_3)}. \tag{7.76}$$

From a straightforward computation,

$$\begin{aligned} \ell &= e^{x^1-x^2} dx^1 T_1 + (dx^2 - e^{x^1-x^2} dx^1) T_2 + (dx^3 + x^3 dx^2) T_3, \\ r &= [e^{x^1} dx^1 + (1 - e^{x^1}) dx^2] T_1 + e^{x^1} (dx^2 - dx^1) T_2 + e^{x^2} dx^3 T_3, \\ v_1 &= e^{x^2-x^1} \partial_1 + \partial_2 - x^3 \partial_3, \quad v_2 = \partial_2 - x^3 \partial_3, \quad v_3 = \partial_3, \end{aligned} \tag{7.77}$$

$$\begin{aligned} a &= \begin{pmatrix} e^{x^1-x^2} & 1 - e^{x^1-x^2} & x^3 \\ e^{-x^2}(e^{x^1} - 1) & 1 + e^{-x^2}(1 - e^{x^1}) & x^3 \\ 0 & 0 & e^{-x^2} \end{pmatrix}, \\ (\Pi^{ab}) &= \begin{pmatrix} 0 & 0 & \frac{1-e^{x^2}(2-2e^{x^1}+e^{x^2})}{2} \\ 0 & 0 & \frac{1+e^{2x^2}-2e^{x^1+x^2}}{2} \\ -\frac{1-e^{x^2}(2-2e^{x^1}+e^{x^2})}{2} & -\frac{1+e^{2x^2}-2e^{x^1+x^2}}{2} & 0 \end{pmatrix}, \end{aligned} \tag{7.78}$$

we obtain the twist matrix U , and the flux is covariantly transformed

$$\mathcal{F}_A = (2, 2, 0, 0, 0, -2) = C_A^B \mathcal{F}_B^{(5|1)}. \tag{7.79}$$

Since the dual algebra is unimodular, originally we have $I^m = 0$. However, due to the dual-coordinate dependence of \bar{d} , we make the field redefinition in Eq. (6.67) and obtain

$$e^{-2\bar{d}} = e^{-2x^2}, \quad I = -\partial_3. \tag{7.80}$$

After the redefinition we obtain a solution of type IIB GSE,

$$\begin{aligned} ds^2 &= \frac{-dt^2 + dz^2}{z^2} + ds_{S^5}^2 \\ &+ \frac{t^2 \{4e^{2x^2} z^4 (e^{2x^1} dx_1^2 + dx_3^2) + [4(t^4 + z^4) + e^{4x^2} z^4] dx_2^2\}}{\Delta^2} \\ &+ \frac{4e^{x^1} t^2 z^4 [e^{x^1} (dx^1 - dx^2)^2 - e^{3x^2} dx^1 dx^2 + 2(dx^1 - dx^2) dx^2]}{\Delta^2}, \\ B_2 &= -\frac{2e^{2x^2} z^2 \{2t^4 dx^2 - z^4 (2e^{x^1} - e^{x^2} - 2) [e^{x^1} dx^1 - (e^{x^1} - 1) dx^2]\}}{\Delta^2} \wedge dx^3, \\ e^{-2\Phi} &= \frac{e^{-4x^2} \Delta^2}{4z^4}, \quad I = -\partial_3, c \quad G_3 = \frac{4e^{-2x^2} t^3 dt \wedge dx^2 \wedge dz}{z^5}, \\ G_5 &= (2e^{x^1} - e^{x^2} - 2) \left[\frac{8t^3 e^{x^1} z dt \wedge dx_1 \wedge dx^2 \wedge dx^3 \wedge dz}{\Delta^2} - 2e^{-x^2} \omega_{S^5} \right], \end{aligned} \tag{7.81}$$

which is defined on the region

$$\Delta^2 \equiv 4t^4 (e^{2x^2} + 1) z^2 + e^{2x^2} z^6 (2 - 2e^{x^1} + e^{x^2})^2 \geq 0. \tag{7.82}$$

A formal T -duality along the x^3 -direction gives a solution of type IIA supergravity,

$$\begin{aligned}
 ds^2 &= \frac{-dt^2 + dz^2}{z^2} + \frac{(t^4 + z^4)(dx^2 - dx^3)^2 + z^4 e^{2x^1}(dx^1 - dx^2 + dx^3)^2}{t^2 z^2} \\
 &\quad + z^2 e^{x^1} \frac{2(dx^2 - dx^3)(dx^1 - dx^2 + dx^3) - e^{x^2}(dx^1 - dx^2 + dx^3) dx^3}{t^2} \\
 &\quad + z^2 e^{x^2} \frac{(dx^3 - dx^2) dx^3}{t^2} + \frac{e^{-2x^2}(4t^4 + e^{4x^2} z^4) dx_3^2}{4t^2 z^2} + ds_{S^5}^2, \\
 e^{-2\Phi} &= \frac{t^2 e^{2(x^3 - x^2)}}{z^2}, \quad G_4 = -\frac{4e^{x^3 - 2x^2} t^3 dt \wedge dx^2 \wedge dx^3 \wedge dz}{z^5}.
 \end{aligned}$$

7.8. (6₀|5_{ii}): Type IIA SUGRA

Finally, we consider the redefinition

$$T'_A = C_A{}^B T_B^{(5|1)}, \quad C = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 & \frac{1}{2} \\ 0 & 1 & 0 & 1 & 0 & \frac{1}{2} \\ -1 & 0 & 1 & 0 & \frac{1}{2} & 0 \\ 1 & 0 & 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{7.83}$$

This time, we consider the parameterization¹¹

$$l = g' \tilde{g}', \quad g' = e^{x^3 T'_3} e^{x^2 T'_2} e^{(x^1 + x^2) T'_1}, \quad \tilde{g}' = e^{\tilde{x}'_1 \tilde{T}'^1} e^{(\tilde{x}'_2 - \tilde{x}'_1) \tilde{T}'^2} e^{\tilde{x}'_3 \tilde{T}'^3}, \tag{7.84}$$

which leads to

$$e^{-2\tilde{d}} = e^{-2x^1} = e^{-2(\tilde{x}'_2 - x^3)}. \tag{7.85}$$

By using

$$\begin{aligned}
 \ell &= (dx^1 + dx^2 - x^2 dx^3) T_1 + [dx^2 - (x^1 + x^2) dx^3] T_2 + dx^3 T_3, \\
 r &= (\cosh x^3 dx^1 + e^{-x^3} dx^2) T_1 + (-\sinh x^3 dx^1 + e^{-x^3} dx^2) T_2 + dx^3 T_3, \\
 v_1 &= \partial_1, \quad v_2 = \partial_2 - \partial_1, \quad v_3 = \partial_3 - x^1 \partial_1 + (x^1 + x^2) \partial_2,
 \end{aligned} \tag{7.86}$$

$$a = \begin{pmatrix} \cosh x^3 & \sinh x^3 & 0 \\ \sinh x^3 & \cosh x^3 & 0 \\ -x^2 & -x^1 - x^2 & 1 \end{pmatrix}, \quad (\Pi^{ab}) = \begin{pmatrix} 0 & x^1 & e^{-x^3} - 1 \\ -x^1 & 0 & e^{-x^3} - 1 \\ 1 - e^{-x^3} & 1 - e^{-x^3} & 0 \end{pmatrix}, \tag{7.87}$$

we can check the covariance of the flux,

$$\mathcal{F}_A = (0, 0, -2, 2, 2, 0) = C_A{}^B \mathcal{F}_B^{(5|1)}. \tag{7.88}$$

¹¹ We note that, in general, the parameterization should be carefully chosen such that the resulting twist matrix U does not break the section condition.

Since the dual algebra 5.ii is non-unimodular, the Killing vector becomes

$$I = \frac{1}{2} \tilde{f}^{ba} \tilde{v}_a = -(v_1 + v_2) = -\partial_2 \quad (\tilde{f}^{b1}_b = -2, \quad \tilde{f}^{b2}_b = -2), \quad (7.89)$$

but by absorbing the dual-coordinate dependence of \vec{d} , we obtain

$$e^{-2\vec{d}} = e^{2x^3}, \quad I^m = 0. \quad (7.90)$$

Then, after the redefinition, we obtain a solution of type IIA supergravity,

$$\begin{aligned} ds^2 &= \frac{-dt^2 + dz^2}{z^2} + t^2 \frac{e^{4x^3} dx_1^2 - 2e^{3x^3} (dx^1 + x^1 dx^3) dx^1}{z^2 [2 - 2e^{x^3} + (x_1^2 + 1)e^{2x^3}]} \\ &\quad + z^2 \frac{[(1 - e^{x^3}) dx^1 + 2 dx^2 - e^{x^3} x^1 dx^3]^2}{4 t^2 [2 - 2e^{x^3} + (x_1^2 + 1)e^{2x^3}]}, \\ &\quad + \frac{e^{2x^3} t^2 [2 dx_1^2 + 4 x^1 dx^1 dx^3 + (2 x_1^2 + 1) dx_3^2]}{z^2 [2 - 2e^{x^3} + (1 + x_1^2)e^{2x^3}]} + ds_{S^5}^2, \\ B_2 &= \frac{e^{2x^3} x^1 dx^1 \wedge dx^2 + [1 + e^{2x^3} (\sinh x^3 - \frac{1}{2})] dx^1 \wedge dx^3 + (2 - e^{x^3}) dx^2 \wedge dx^3}{2 - 2e^{x^3} + (x_1^2 + 1)e^{2x^3}}, \\ e^{-2\Phi} &= \frac{t^2 [2 - 2e^{x^3} + e^{2x^3} (x_1^2 + 1)]}{z^2}, \quad G_4 = -\frac{4 t^3 e^{2x^3} dt \wedge dx^1 \wedge dx^3 \wedge dz}{z^5}. \end{aligned} \quad (7.91)$$

8. Conclusion and outlook

8.1. Summary of results

We have discussed two approaches to the non-Abelian T -duality. One is the traditional NATD, obtained by integrating out the gauge fields associated with non-Abelian isometries, and the other is the PL T -duality/plurality, which is based on the Drinfel'd double.

In NATD, a closed-form expression for the duality rules including the R–R fields was explicitly known only for a certain isometry group, $SU(2)$, but we proposed a general formula by assuming that the isometry group freely acts on the target space. The duality rules, under the setup of Eq. (3.1), are summarized in Eqs. (5.2) and (5.3). In order to check the formula we studied many examples, particularly the NATD for non-unimodular isometry groups.

For the PL T -duality, the treatments of the R–R fields have been discussed in recent papers [138,139,142], but concrete examples have not been well studied. We first considered the case without spectator fields, and translated the known transformation rules for $\{g_{mn}, B_{mn}, \Phi\}$ into the rules for the generalized metric \mathcal{H}_{MN} and the DFT dilaton d . Then, using a result of the gauged DFT, we showed that the equations of motion are transformed covariantly under the PL T -plurality (by introducing a Killing vector I^m appropriately). We also introduced the R–R fields, and determined their transformation rule under the $O(n, n)$ PL T -plurality transformation such that the equations of motion are covariantly transformed. We further considered the case with spectator fields and, requiring some dualizability conditions, we showed that the DFT equations of motion are indeed satisfied even in the presence of spectators. Finally, we studied a concrete example of the PL T -plurality. Starting with the $AdS_5 \times S^5$ solution, we obtained the family of solutions in Fig. 2.




Fig. 2. Family of solutions for concrete PL T -plurality example.

Three of these are solutions of GSE. There are two origins of GSE: one is the Killing vector $I^i = \frac{1}{2} \tilde{f}^{ba}{}_b v_a^i$ that appears when the dual algebra is non-unimodular, and the other is the dual-coordinate dependence in \bar{d} . In the examples **(2.i|5)** and **(6₀|5.ii)**, the two contributions are canceled with each other, and they are solutions of the usual supergravity even though their dual algebras are non-unimodular. In the literature, when \bar{d} has a dual-coordinate dependence, since its interpretation is not clear in string theory or supergravity, such a Manin triple was ignored. However, in DFT we can treat the dual coordinates in the same ways as the physical coordinates, and we can lift the restriction. In this way, the PL T -plurality is a solution-generating technique of the DFT, rather than the usual supergravity.

8.2. Discussion and outlook

As we discussed, if we consider a supergravity solution that contains a four-dimensional Minkowski spacetime, $ds^2 = f^2(y) \eta_{\mu\nu} dx^\mu dx^\nu + \dots$, we can choose the coordinates such that the **(5|1)** symmetry is manifest. Then, as long as the B -field is isometric along the three Killing vectors, we will obtain a family of eight solutions similar to the case of $\text{AdS}_5 \times S^5$. Moreover, low-dimensional Drinfel'd doubles have already been classified in Refs. [52–54], and a useful list is given in Sect. 3 of Ref. [54]. If we have a DFT solution with an isometry algebra \mathfrak{g} , we may find a series of Manin triples,

$$(\mathfrak{g}|\mathbf{1}) \cong (\mathfrak{g}'|\mathfrak{g}'') \cong \dots, \tag{8.1}$$

and obtain a chain of DFT solutions. We may also start from a background with a $(\mathfrak{g}|\tilde{\mathfrak{g}})$ symmetry. For example, as discussed in Ref. [50], the Yang–Baxter deformed backgrounds are also PL T -dualizable. Indeed, a Yang–Baxter deformed background has the form

$$E^{mn} = \tilde{g}^{mn} - \beta^{mn}, \quad \beta^{mn} \equiv 2 \eta r^{ab} v_a^m v_b^n \quad (r^{ab} = -r^{ba}), \tag{8.2}$$

where η and r^{ab} are constant, and $\mathfrak{L}_{v_a} \tilde{g}_{mn} = 0$ and $[v_a, v_b] = f_{ab}{}^c v_c$ are satisfied. Then, we can show that $\mathfrak{L}_{v_a} E^{mn} = \tilde{f}^{bc}{}_a v_b^m v_c^n$ with $\tilde{f}^{bc}{}_a = 2 \eta (r^{bd} f_{da}{}^c - r^{cd} f_{da}{}^b)$, and this is a dualizable background with the $(\mathfrak{g}|\tilde{\mathfrak{g}})$ symmetry. Then, by finding a group element $g(x)$ which realizes the set of Killing vectors v_a^m as the left-invariant vector fields and β^{mn} as $\beta^{mn} = e_a^m e_b^n \Pi^{ab}$ [i.e. $(a^\top b)^{ab} = 2 \eta r^{ab}$], we can perform the PL T -plurality transformations of the Yang–Baxter-deformed background. In this

way, from a given solution we can find new solutions one after another, and the PL T -plurality is a useful solution-generating technique.

In the traditional approach to NATD, we introduced the generalized Killing vector $(V_a^M) = (v_a^m, \tilde{v}_{am})$. When the dual components \tilde{v}_{am} are present, we cannot regard the NATD as a particular case of the PL T -plurality. Also when the generalized Killing vectors depend on the spectator fields y^μ , we cannot realize them as the left-invariant vector fields. In this sense, the traditional NATD is not completely contained in the PL T -plurality discussed here. It is interesting to study whether it is possible to generalize the PL T -plurality such that the traditional NATD can be realized as a particular case. In the realm of NATD that is going beyond the PL T -plurality, it is not ensured that the dual background is a solution of DFT. By the definition of NATD, the duality rules for the metric and B -field should not be modified, but the transformation rule for the dilaton and I^m may be modified from Eq. (5.2). It will be an important task to determine the general rule for the dilaton and I^m that is consistent with the DFT equations of motion. Once the modification of the rule for the dilaton is determined, the modification of the rule for the R–R fields (by an overall factor) can also be determined, and then we can check the equation of motion for the R–R fields.

In the two approaches studied in this paper we have assumed that the isometry group acts on the target space freely, or without isotropy. If the assumption is not satisfied, we cannot take a gauge $x^i = c^i$ and we need to consider a more non-trivial gauge fixing. Treatments in such cases are discussed, for example, in Refs. [19,73,77,176] for the NATD, and in Refs. [41,45,47] for the PL T -duality. It is an interesting future direction to check whether the DFT equations of motion are covariantly rotated even in such general cases.

In the study of the PL T -plurality we have checked the covariance of the flux $\mathcal{F}_A = 2\mathcal{D}_A\bar{d}$ on a case-by-case basis. The covariance is highly non-trivial but it was indeed transformed covariantly in all of the examples, and we suspect that there is some mechanism to be clarified. To show the covariance of \mathcal{F}_A , clear understanding of the (finite) coordinate transformation $(x^i, \tilde{x}_i) \rightarrow (x'^i, \tilde{x}'_i)$ on the Drinfel'd double will be indispensable. The $2D$ diffeomorphism in DFT_{WZW} [135–137] may be useful for this purpose.

8.3. Toward non-Abelian U -duality

Another important future direction is an investigation of the non-Abelian U -duality. As an attempt toward this, let us first consider an extension of the traditional NATD. As a natural extension of Eq. (3.3), let us consider the following setup [155]:

$$\mathfrak{L}_{v_a}g_{ij} = 0, \quad \iota_{v_a}F_4 + d\hat{v}_a^{(2)} = 0, \quad \mathfrak{L}_{v_a}v_b = f_{ab}{}^c v_c, \quad \mathfrak{L}_{v_a}\hat{v}_b^{(2)} = f_{ab}{}^c \hat{v}_c^{(2)}, \quad (8.3)$$

where $F_4 \equiv dC_3$ is the four-form field strength in the eleven-dimensional supergravity. The two-form $\hat{v}_a^{(2)}$ is the generalization of the one-form \hat{v}_{am} appearing in Eq. (3.3), and the first two relations in Eq. (8.3) are understood as a form of generalized Killing equations. The remaining two equations are generalizations of the C-brackets between the generalized Killing vectors.

We define $\hat{v}_{ab}^{(1)} \equiv \iota_{v_b}\hat{v}_a^{(2)}$, and assume the following relation for simplicity:

$$\hat{v}_{(ab)}^{(1)} = 0, \quad \iota_{v_a}\hat{v}_{[bc]}^{(1)} = \iota_{v_{[a}}\hat{v}_{bc]}^{(1)}. \quad (8.4)$$

Table 1. U -duality groups $E_{n(n)}$.

n	4	5	6	7	8
U -duality group $E_{n(n)}$	SL(5)	SO(5, 5)	$E_{6(6)}$	$E_{7(7)}$	$E_{8(8)}$
Dimension D	10	16	27	56	248

We also assume the existence of the one-forms $\ell^a \equiv \ell^a_i dx^i$ that are dual to ν_a ($\iota_{\nu_a} \ell^b = \delta^b_a$), and then we find that the action¹²

$$S = \int_{\Sigma} \left[\frac{1}{2} (g_{ij} Dx^i \wedge *Dx^j + *1) + C_3 + 2y_{ab} F^a \wedge (\ell^b - A^b) \right] + \int_{\Sigma} \left[-A^a \wedge \hat{\nu}_a^{(2)} + \frac{1}{2} A^a \wedge A^b \wedge \hat{\nu}_{ab}^{(1)} - \frac{1}{3!} A^a \wedge A^b \wedge A^c \iota_{\nu_a} \hat{\nu}_{bc}^{(1)} \right] \tag{8.5}$$

is invariant under

$$\delta_{\epsilon} x^i(\sigma) = \epsilon^a(\sigma) \nu_a^i(x), \quad \delta_{\epsilon} A^a(\sigma) = d\epsilon^a(\sigma) + f_{bc}{}^a A^b(\sigma) \epsilon^c(\sigma),$$

$$\delta_{\epsilon} y_{ab} = \epsilon^c (f_{ca}{}^d y_{db} + f_{cb}{}^d y_{ad}). \tag{8.6}$$

Here, by following the approach of Ref. [177] (see also Ref. [178]), we have introduced antisymmetric Lagrange multipliers $y_{ab} = -y_{ba}$ that will ensure $F^a = 0$.

In the Abelian limit we can realize $\nu_a^i = \delta_a^i$ and $\ell^a = \delta^a_i dx^i$, and then we can always choose a gauge $x^i = 0$. By further assuming $\hat{\nu}_a^{(2)} = -\iota_{\nu_a} C_3$, the action reduces to

$$S = \int_{\Sigma} \left[\frac{1}{2} (g_{ij} A^i \wedge *A^j + *1) + \frac{1}{3!} C_{abc} A^a \wedge A^b \wedge A^c + dy_{ab} \wedge A^a \wedge A^b \right]. \tag{8.7}$$

This is precisely the action discussed in Refs. [177,178], and Eq. (8.5) can be regarded as a natural extension. However, unlike the case of the string action, it is not clear how to eliminate the gauge fields A^a , and at the present time we do not know how to obtain the dual action.

A more promising approach may be the following one based on a generalization of DFT. The U -dual version of DFT is known as the exceptional field theory (EFT) [178–185] and it is actively being studied. In DFT, the generalized coordinates are $(x^M) = (x^m, \tilde{x}_m)$ and the dual coordinates \tilde{x}_m are associated with the string winding number. On the other hand, in EFT we introduce the dual coordinates for all of the wrapped branes that are connected by U -duality transformations. For example, in M-theory on a n -torus we have the M2-brane, the M5-brane, and the Kaluza–Klein monopole, and more exotic branes in general. Correspondingly, we introduce the generalized coordinates as

$$(x^I) = (x^i, y_{i_1 i_2}, y_{i_1 \dots i_5}, y_{i_1 \dots i_7, i}, \dots) \quad (i = 1, \dots, n). \tag{8.8}$$

By understanding that the multiple indices separated by commas are totally antisymmetrized, we can easily see that the number of dimensions of the extended space x^I is the same as the dimension D of a fundamental representation of the $E_{n(n)}$ U -duality group, as shown in Table 1. The generalized

¹² In the string action of Eq. (3.4), by adding a total-derivative term the Lagrangian multiplier was introduced with derivative $d\tilde{x}_a$ (see Ref. [22] for the Abelian case), but here we only discuss the classical equations of motion without investigating such a total-derivative term.

metric \mathcal{M}_{IJ} has been constructed in such extended space in Refs. [178,182], and it contains the bosonic fields, such as the metric g_{ij} , and the three-form and six-form potentials, $C_{i_1i_2i_3}$ and $C_{i_1\dots i_6}$. It is a natural generalization of the generalized metric \mathcal{H}_{MN} in DFT.

In DFT, the section condition $\eta^{IJ} \partial_I \partial_J = 0$ reduces the doubled space to the physical subspace. The section condition in EFT (for $n \leq 6$) also has a similar form $\eta^{IJ;\hat{K}} \partial_I \partial_J = 0$, where $\eta^{IJ;\hat{K}}$ is known as the η -symbol and it has an additional index \hat{K} transforming in another representation (see Ref. [186] for the explicit form of the η -symbol). When all of the fields depend only on the coordinates x^i of Eq. (8.8), we find

$$\eta^{IJ;\hat{K}} \partial_I \partial_J = \eta^{ij;\hat{K}} \partial_i \partial_j = 0 \quad (\because \eta^{ij;\hat{K}} = 0), \tag{8.9}$$

and the section condition is satisfied. This n -dimensional solution is called the M-theory section. Another solution, called the type IIB section, was found in Ref. [187], and in order to discuss the type IIB section it is convenient to reparameterize the coordinates as¹³

$$(x^M) = (x^m, y_m^\alpha, y_{m_1 m_2 m_3}, y_{m_1 \dots m_5}, y_{m_1 \dots m_6, m}, \dots) \quad (m = 1, \dots, n - 1, \alpha = 1, 2), \tag{8.10}$$

where the dual coordinates are associated with the type IIB branes. If the fields depend only on the x^m , the section condition is again satisfied because $\eta^{mn;\hat{P}} = 0$. Since we cannot introduce any more coordinate dependence, the subspace spanned by x^m is also a maximally isotropic subspace, although it is $(n - 1)$ dimensional unlike the M-theory section. In this way, a single EFT can be understood from two viewpoints: M-theory and type IIB theory.

One of the key relations in the PL T -duality is the self-duality relation,

$$\eta_{AB} \hat{\mathcal{P}}^B = \hat{\mathcal{H}}_{AB} * \hat{\mathcal{P}}^B, \quad \hat{\mathcal{P}}(\sigma) = dl l^{-1}. \tag{8.11}$$

This is a covariant rewriting of the string equations of motion, but a similar equation for the M2- or M5-brane theory has been discussed in Refs. [189,190] for the SL(5) and SO(5, 5) case, and in Ref. [191] for higher exceptional groups. For the Mp -brane ($p = 2, 5$), it has a similar form,

$$\eta_{IJ} \wedge \mathcal{P}^J = \mathcal{M}_{IJ} * \mathcal{P}^J, \tag{8.12}$$

where η_{IJ} is some $(p - 1)$ -form that contains dx^i and the field strengths of the worldvolume gauge fields. In the case of the flat torus, the equations of motion give $d\mathcal{P}^I = 0$ and we find the on-shell expression $\mathcal{P}^I = dx^I$. On the other hand, by requiring a certain ‘‘dualizability condition’’ on \mathcal{M}_{IJ} appropriately, the equations of motion may lead to $\mathcal{P} = dl l^{-1}$, where l is an element of a certain large group \mathfrak{E} with dimension D . The corresponding algebra \mathfrak{e} will be endowed with a bilinear form, corresponding to the η -symbol. Then, the U -dual version of the PL T -plurality may be the equivalence between sigma models with n - or $(n - 1)$ -dimensional target spaces that have an isometry algebra $[T_a, T_b] = f_{ab}{}^c T_c$ satisfying $\eta^{ab;\hat{A}} = 0$. The identification of the detailed structure of the group \mathcal{E} and the systematic construction of the twist matrix U , whose flux gives the structure constant of \mathfrak{e} , are interesting future directions.

Note added

After this paper appeared on arXiv, an interesting paper, Ref. [192], appeared, which also discusses NATD from the perspective of the gauged DFT.

¹³ The explicit relation between x^I and x^M was determined in Ref. [188].

Acknowledgements

We would like to thank Falk Hassler for elucidating the approach of Refs. [138,139]. We also would like to thank Yolanda Lozano, Jeong-Hyuck Park, Jun-ichi Sakamoto, and Shozo Uehara for helpful discussions and comments. We are also grateful to the organizers and the participants of the workshop ‘‘String: T-duality, Integrability and Geometry’’ held at Tohoku University. This work is supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Numbers 18K13540 and 18H01214.

Funding

Open Access funding: SCOAP³.

Appendix A. Conventions

The symmetrization and antisymmetrization are normalized as

$$A_{(m_1 \dots m_n)} \equiv \frac{1}{n!} (A_{m_1 \dots m_n} + \dots), \quad A_{[m_1 \dots m_n]} \equiv \frac{1}{n!} (A_{m_1 \dots m_n} \pm \dots). \quad (A.1)$$

Our conventions for differential forms are as follows, both for the spacetime and the worldsheet:

$$\begin{aligned} (*\alpha_q)_{m_1 \dots m_{p+1-q}} &= \frac{1}{q!} \varepsilon^{n_1 \dots n_q} \alpha_{n_1 \dots n_q}, & d^D x &= dx^1 \wedge \dots \wedge dx^D, \\ * (dx^{m_1} \wedge \dots \wedge dx^{m_q}) &= \frac{1}{(p+1-q)!} \varepsilon^{m_1 \dots m_q} dx^{n_1} \wedge \dots \wedge dx^{n_{p+1-q}}, \\ (\iota_\nu \alpha_n) &= \frac{1}{(n-1)!} v^n \alpha_{nm_1 \dots m_{n-1}} dx^{m_1} \wedge \dots \wedge dx^{m_{n-1}}. \end{aligned} \quad (A.2)$$

The epsilon tensors on the spacetime and the worldsheet are defined as follows:

$$\varepsilon^{01} = \frac{1}{\sqrt{|\gamma|}}, \quad \varepsilon_{01} = -\sqrt{|\gamma|}, \quad \varepsilon^{1 \dots D} = -\frac{1}{\sqrt{|g|}}, \quad \varepsilon_{1 \dots D} = \sqrt{|g|}. \quad (A.3)$$

For the R–R fields, we have the R–R potential in the A-basis $A_{m_1 \dots m_p}$ and the C-basis $C_{m_1 \dots m_p}$ [160]. In terms of the polyform,

$$A \equiv \sum_p \frac{1}{p!} A_{m_1 \dots m_p} dx^{m_1} \wedge \dots \wedge dx^{m_p}, \quad C \equiv \sum_p \frac{1}{p!} C_{m_1 \dots m_p} dx^{m_1} \wedge \dots \wedge dx^{m_p}, \quad (A.4)$$

they are related as

$$A = e^{B_2 \wedge} C, \quad C = e^{-B_2 \wedge} A. \quad (A.5)$$

Their field strengths are defined as

$$F = dA, \quad G = dC + H_3 \wedge C, \quad (A.6)$$

and they are also related as

$$F = e^{B_2 \wedge} G, \quad G = e^{-B_2 \wedge} F. \quad (A.7)$$

For simplicity, in this paper we call the field strength F the Page form. In our convention, the G satisfies the self-duality relation

$$*G_p = (-1)^{\frac{p(p+1)}{2}+1} G_{10-p}, \quad G_p = (-1)^{\frac{p(p-1)}{2}} *G_{10-p}. \quad (A.8)$$

In the presence of the Killing vector I^m in the GSE, which satisfies

$$\mathfrak{L}_I g_{mn} = \mathfrak{L}_I B_2 = \mathfrak{L}_I \Phi = \mathfrak{L}_I F = \mathfrak{L}_I G = 0, \tag{A.9}$$

the relations in Eq. (A.6) are modified as

$$F = dA - \iota_I A, \quad G = dC + H_3 \wedge C - \iota_I B_2 \wedge C - \iota_I C, \tag{A.10}$$

and the Bianchi identities, which are equivalent to the equations of motion under Eq. (A.8), become

$$dF - \iota_I F = 0, \quad dG + H_3 \wedge G - \iota_I B_2 \wedge G - \iota_I G = 0. \tag{A.11}$$

The GSE for the fields in the NS–NS sector can be summarized as

$$\begin{aligned} R + 4 D^m \partial_m \Phi - 4 |\partial \Phi|^2 - \frac{1}{2} |H_3|^2 - 4 (I^m I_m + U^m U_m + 2 U^m \partial_m \Phi - D_m U^m) &= 0, \\ R_{mn} - \frac{1}{4} H_{mpq} H_n{}^{pq} + 2 D_m \partial_n \Phi + D_m U_n + D_n U_m &= T_{mn}, \\ -\frac{1}{2} D^p H_{pmn} + \partial_p \Phi H^p{}_{mn} + U^p H_{pmn} + D_m I_n - D_n I_m &= K_{mn}, \end{aligned} \tag{A.12}$$

where $U_1 \equiv U_m dx^m$ is defined as $U_1 \equiv \iota_I B_2$, and T_{mn} and K_{mn} are

$$\begin{aligned} T_{mn} &\equiv \frac{e^{2\Phi}}{4} \sum_p \left[\frac{1}{(p-1)!} G_{(m}{}^{q_1 \dots q_{p-1}} G_{n)q_1 \dots q_{p-1}} - \frac{1}{2} g_{mn} |G_p|^2 \right], \\ K_{mn} &\equiv \frac{e^{2\Phi}}{4} \sum_p \frac{1}{(p-2)!} G_{q_1 \dots q_{p-2}} G_{mn}{}^{q_1 \dots q_{p-2}}. \end{aligned} \tag{A.13}$$

In the presence of the Killing vector $(I^m) = (I^i, I^z)$, if we perform a formal T -duality along the x^z -direction then the supergravity fields are transformed as follows [149]:

$$\begin{aligned} g'_{ij} &= g_{ij} - \frac{g_{iz} g_{jz} - B_{iz} B_{jz}}{g_{zz}}, \quad g'_{iz} = \frac{B_{iz}}{g_{zz}}, \quad g'_{zz} = \frac{1}{g_{zz}}, \\ B'_{ij} &= B_{ij} - \frac{B_{iz} g_{jz} - g_{iz} B_{jz}}{g_{zz}}, \quad B'_{iz} = \frac{g_{iz}}{g_{zz}}, \\ \Phi' &= \Phi + \frac{1}{4} \ln \left| \frac{\det(g'_{mn})}{\det(g_{mn})} \right| + I^z z, \quad I'^i = I^i, \quad I'^z = 0, \\ A'_{i_1 \dots i_{p-1} z} &= e^{-I^z z} A_{i_1 \dots i_{p-1}}, \quad A'_{i_1 \dots i_p} = e^{-I^z z} A_{i_1 \dots i_p z}, \\ C'_{i_1 \dots i_{p-1} z} &= e^{-I^z z} \left[C_{i_1 \dots i_{p-1}} - (p-1) \frac{C_{[i_1 \dots i_{p-2} z] g_{i_{p-1} z]}}{g_{zz}} \right], \\ C'_{i_1 \dots i_p} &= e^{-I^z z} \left[C_{i_1 \dots i_p z} + p C_{[i_1 \dots i_{p-1}} B_{i_p] z} + p(p-1) \frac{C_{[i_1 \dots i_{p-2} z] B_{i_{p-1} z] g_{i_p z]}}{g_{zz}} \right]. \end{aligned} \tag{A.14}$$

Appendix B. Technical details of DFT

In this appendix we explain the technical details of (gauged) DFT and show the covariance of the DFT equations of motion under the PL T -plurality with spectator fields.

B.1. NS–NS sector

For convenience, let us introduce the double vielbein $(V_{\hat{A}}^M) \equiv (V_a^M, V_{\bar{a}}^M)$ as

$$\mathcal{H}_{MN} = V_M^{\hat{A}} V_N^{\hat{B}} \mathcal{H}_{\hat{A}\hat{B}}, \quad \eta_{MN} = V_M^{\hat{A}} V_N^{\hat{B}} \eta_{\hat{A}\hat{B}}, \quad V_M^{\hat{A}} V_{\hat{A}}^N = \delta_M^N, \quad (\text{B.1})$$

where $V_M^{\hat{A}}$ is an $O(D, D)$ matrix and we have defined

$$(\mathcal{H}_{\hat{A}\hat{B}}) = \begin{pmatrix} \eta_{ab} & 0 \\ 0 & \eta_{\bar{a}\bar{b}} \end{pmatrix}, \quad (\eta_{\hat{A}\hat{B}}) = \begin{pmatrix} \eta_{ab} & 0 \\ 0 & -\eta_{\bar{a}\bar{b}} \end{pmatrix}, \quad (\text{B.2})$$

and $(\eta_{ab}) \equiv (\eta_{\bar{a}\bar{b}}) \equiv \text{diag}(-1, 1, \dots, 1)$. They can be parameterized as

$$(V_a^M) = \frac{1}{\sqrt{2}} \begin{pmatrix} e_a^m \\ (g+B)_{mn} e_a^n \end{pmatrix}, \quad (V_{\bar{a}}^M) = \frac{1}{\sqrt{2}} \begin{pmatrix} e_{\bar{a}}^m \\ (-g+B)_{mn} e_{\bar{a}}^n \end{pmatrix}, \quad (\text{B.3})$$

where $e_a^m = e_{\bar{a}}^m$ is the vielbein satisfying $g_{mn} = e_m^a e_n^b \eta_{ab} = e_{\bar{m}}^{\bar{a}} e_{\bar{n}}^{\bar{b}} \eta_{\bar{a}\bar{b}}$.

The equations of motion for the DFT dilaton and the generalized metric are

$$\begin{aligned} \mathcal{R} &\equiv -2 \bar{P}^{\hat{A}\hat{B}} (2 \mathcal{D}_{\hat{A}} \mathcal{F}_{\hat{B}} - \mathcal{F}_{\hat{A}} \mathcal{F}_{\hat{B}}) - \frac{1}{3} \bar{P}^{\hat{A}\hat{B}\hat{C}\hat{D}\hat{E}\hat{F}} \mathcal{F}_{\hat{A}\hat{B}\hat{C}} \mathcal{F}_{\hat{D}\hat{E}\hat{F}} = 0, \\ \mathcal{G}^{\hat{A}\hat{B}} &\equiv -4 \bar{P}^{\hat{D}[\hat{A}} \mathcal{D}^{\hat{B}]} \mathcal{F}_{\hat{D}} + 2 (\mathcal{F}_{\hat{D}} - \mathcal{D}_{\hat{D}}) \check{\mathcal{F}}^{\hat{D}[\hat{A}\hat{B}]} - 2 \check{\mathcal{F}}^{\hat{C}\hat{D}[\hat{A}} \mathcal{F}_{\hat{C}\hat{D}}^{\hat{B}]} = 0, \end{aligned} \quad (\text{B.4})$$

where $\mathcal{D}_{\hat{A}} \equiv V_{\hat{A}}^M \partial_M$, and $\mathcal{F}_{\hat{A}}$ and $\mathcal{F}_{\hat{A}\hat{B}\hat{C}}$ are defined by

$$\mathcal{F}_{\hat{A}\hat{B}\hat{C}} \equiv 3 \Omega_{[\hat{A}\hat{B}\hat{C}]}, \quad \mathcal{F}_{\hat{A}} \equiv \Omega_{\hat{A}}^{\hat{B}} + 2 \mathcal{D}_{\hat{A}} d, \quad \Omega_{\hat{A}\hat{B}\hat{C}} \equiv -\mathcal{D}_{\hat{A}} V_{\hat{B}}^M V_{M\hat{C}}, \quad (\text{B.5})$$

and $\check{\mathcal{F}}^{\hat{A}\hat{B}\hat{C}}$ is defined similarly to Eq. (2.23). We can show that $\mathcal{R} = \mathcal{S}$ under the section condition, but the equivalence of $\mathcal{G}^{\hat{A}\hat{B}} = 0$ and $\mathcal{S}_{MN} = 0$ is non-trivial. To see the equivalence, we show that

$$V^{\bar{a}M} V^{bN} \mathcal{S}_{MN} = e^{\bar{a}m} e^{bn} s_{mn}, \quad V^{aM} V^{bN} \mathcal{S}_{MN} = 0 = V^{\bar{a}M} V^{\bar{b}N} \mathcal{S}_{MN}. \quad (\text{B.6})$$

Under the section condition, we can also find

$$\mathcal{G}^{\bar{a}b} = V^{\bar{a}M} V^{bN} \mathcal{S}_{MN}, \quad \mathcal{G}^{ab} = 0 = \mathcal{G}^{\bar{a}\bar{b}}, \quad (\text{B.7})$$

and they clearly show the equivalence of $\mathcal{G}^{\hat{A}\hat{B}} = 0$ and $\mathcal{S}_{MN} = 0$.

By using the identities [134]

$$\begin{aligned} \mathcal{Z} &\equiv \mathcal{D}^{\hat{A}} \mathcal{F}_{\hat{A}} - \frac{1}{2} \mathcal{F}_{\hat{A}} \mathcal{F}_{\hat{A}} + \frac{1}{12} \mathcal{F}_{\hat{A}\hat{B}\hat{C}} \mathcal{F}^{\hat{A}\hat{B}\hat{C}} = 0, \\ \mathcal{Z}_{\hat{A}\hat{B}} &\equiv 2 \mathcal{D}_{[\hat{A}} \mathcal{F}_{\hat{B}]} + \mathcal{F}^{\hat{C}} \mathcal{F}_{\hat{C}\hat{A}\hat{B}} - \mathcal{D}^{\hat{C}} \mathcal{F}_{\hat{C}\hat{A}\hat{B}} = 0, \end{aligned} \quad (\text{B.8})$$

which hold under the section condition, we can simplify the expressions for \mathcal{R} and $\mathcal{G}^{\hat{A}\hat{B}}$ as

$$\begin{aligned} \mathcal{R} &= \mathcal{H}^{\hat{A}\hat{B}} (2 \mathcal{D}_{\hat{A}} \mathcal{F}_{\hat{B}} - \mathcal{F}_{\hat{A}} \mathcal{F}_{\hat{B}}) \\ &\quad + \frac{1}{12} \mathcal{H}^{\hat{A}\hat{D}} (3 \eta^{\hat{B}\hat{E}} \eta^{\hat{C}\hat{F}} - \mathcal{H}^{\hat{B}\hat{E}} \mathcal{H}^{\hat{C}\hat{F}}) \mathcal{F}_{\hat{A}\hat{B}\hat{C}} \mathcal{F}_{\hat{D}\hat{E}\hat{F}}, \\ \mathcal{G}^{\hat{A}\hat{B}} &= 2 \mathcal{H}^{\hat{D}[\hat{A}} \mathcal{D}^{\hat{B}]} \mathcal{F}_{\hat{D}} - \frac{1}{2} \mathcal{H}^{\hat{D}\hat{E}} (\eta^{\hat{A}\hat{F}} \eta^{\hat{B}\hat{G}} - \mathcal{H}^{\hat{A}\hat{F}} \mathcal{H}^{\hat{B}\hat{G}}) (\mathcal{F}_{\hat{D}} - \mathcal{D}_{\hat{D}}) \mathcal{F}_{\hat{E}\hat{F}\hat{G}} \end{aligned} \quad (\text{B.9})$$

$$-\mathcal{H}_{\hat{E}}^{[\hat{A}} (\mathcal{F}_{\hat{D}} - \mathcal{D}_{\hat{D}}) \mathcal{F}^{\hat{B}]\hat{D}\hat{E}} + \frac{1}{2} (\eta^{\hat{C}\hat{E}} \eta^{\hat{D}\hat{F}} - \mathcal{H}^{\hat{C}\hat{E}} \mathcal{H}^{\hat{D}\hat{F}}) \mathcal{H}^{\hat{G}[\hat{A}} \mathcal{F}_{\hat{C}\hat{D}}^{\hat{B}]} \mathcal{F}_{\hat{E}\hat{F}\hat{G}}. \quad (\text{B.10})$$

As a side remark, we note that the equations of motion $\mathcal{G}^{\hat{A}\hat{B}} = 0$ can also be expressed as

$$\begin{aligned} \tilde{\mathcal{G}}^{\hat{A}\hat{B}} &\equiv \mathcal{H}^{\hat{A}}_{\hat{C}} \mathcal{G}^{\hat{C}\hat{B}} \\ &= \bar{P}^{(\hat{A}\hat{B})\hat{C}\hat{D}} (\bar{P}^{\hat{E}\hat{F}\hat{G}\hat{H}} \mathcal{F}_{\hat{C}\hat{E}\hat{F}} \mathcal{F}_{\hat{D}\hat{G}\hat{H}} + 2 \mathcal{D}_{(\hat{C}} \mathcal{F}_{\hat{D})}) + 2 \bar{P}^{\hat{C}\hat{D}\hat{E}(\hat{A}} (\mathcal{F}_{\hat{E}} - \mathcal{D}_{\hat{E}}) \mathcal{F}^{\hat{B})\hat{C}\hat{D}} \\ &\quad - P^{[\hat{A}\hat{B}]\hat{C}\hat{D}} [(\mathcal{F}^{\hat{E}} - \mathcal{D}^{\hat{E}}) \mathcal{F}_{\hat{E}\hat{C}\hat{D}} + 2 \mathcal{D}_{[\hat{C}} \mathcal{F}_{\hat{D}]}] = 0, \end{aligned} \quad (\text{B.11})$$

where we have defined the projectors

$$P^{\hat{A}\hat{B}\hat{C}\hat{D}} \equiv \frac{1}{2} (\eta^{\hat{A}\hat{C}} \eta^{\hat{B}\hat{D}} + \mathcal{H}^{\hat{A}\hat{C}} \mathcal{H}^{\hat{B}\hat{D}}), \quad \bar{P}^{\hat{A}\hat{B}\hat{C}\hat{D}} \equiv \frac{1}{2} (\eta^{\hat{A}\hat{C}} \eta^{\hat{B}\hat{D}} - \mathcal{H}^{\hat{A}\hat{C}} \mathcal{H}^{\hat{B}\hat{D}}). \quad (\text{B.12})$$

Now, let us decompose the double vielbein and the DFT dilaton as

$$V_M^{\hat{A}} = U_M^B(x^I) \hat{V}_B^{\hat{A}}(y^\mu), \quad d = \hat{d}(y^\mu) + \mathbf{d}(x^I), \quad (\text{B.13})$$

where the twist matrix U_M^A is an $O(D, D)$ matrix and the untwisted metric is defined by

$$\hat{\mathcal{H}}_{AB}(y) \equiv \hat{V}_A^{\hat{C}}(y) \hat{V}_B^{\hat{D}}(y) \mathcal{H}_{\hat{C}\hat{D}}. \quad (\text{B.14})$$

Then, by requiring

$$\mathcal{D}_A \hat{V}_B^{\hat{C}} = \partial_A \hat{V}_B^{\hat{C}}, \quad \mathcal{D}_A \hat{d} = \partial_A \hat{d} \quad (\mathcal{D}_A \equiv U_A^M \partial_M), \quad (\text{B.15})$$

the generalized fluxes can be decomposed as

$$\begin{aligned} \mathcal{F}_{\hat{A}} &= \hat{\mathcal{F}}_{\hat{A}}(y) + \hat{V}_{\hat{A}}^B(y) \mathcal{F}_B, \\ \mathcal{F}_{\hat{A}\hat{B}\hat{C}} &= \hat{\mathcal{F}}_{\hat{A}\hat{B}\hat{C}}(y) + \hat{V}_{\hat{A}}^D(y) \hat{V}_{\hat{B}}^E(y) \hat{V}_{\hat{C}}^F(y) \mathcal{F}_{DEF}, \end{aligned} \quad (\text{B.16})$$

where $\hat{\mathcal{F}}_{\hat{A}}(y)$ and $\hat{\mathcal{F}}_{\hat{A}\hat{B}\hat{C}}(y)$ are the generalized fluxes associated with $\{\hat{V}_A^{\hat{B}}, \hat{d}\}$,

$$\hat{\mathcal{F}}_{\hat{A}\hat{B}\hat{C}} \equiv 3 \hat{\Omega}_{[\hat{A}\hat{B}\hat{C}]}, \quad \hat{\mathcal{F}}_{\hat{A}} \equiv \hat{\Omega}^{\hat{B}}_{\hat{A}\hat{B}} + 2 \hat{D}_{\hat{A}} \hat{d}, \quad \hat{\Omega}_{\hat{A}\hat{B}\hat{C}} \equiv -\hat{D}_{\hat{A}} \hat{V}_{\hat{B}}^D \hat{V}_{\hat{C}}^E, \quad (\text{B.17})$$

and $\hat{D}_A \equiv \hat{V}_A^B \partial_B$. Then, the generalized Ricci scalar can be decomposed as

$$\begin{aligned} \mathcal{R} &= \hat{\mathcal{R}} + \frac{1}{12} \hat{\mathcal{H}}^{AD} (3 \eta^{BE} \eta^{CF} - \hat{\mathcal{H}}^{BE} \hat{\mathcal{H}}^{CF}) \mathcal{F}_{ABC} \mathcal{F}_{DEF} - \hat{\mathcal{H}}^{AB} \mathcal{F}_A \mathcal{F}_B \\ &\quad - \frac{1}{2} \mathcal{F}^A_{BC} \hat{\mathcal{H}}^{BD} \hat{\mathcal{H}}^{CE} \mathcal{D}_D \hat{\mathcal{H}}_{AE} + 2 \mathcal{F}_A \mathcal{D}_B \hat{\mathcal{H}}^{AB} - 4 \hat{\mathcal{H}}^{AB} \mathcal{F}_A \mathcal{D}_B \hat{d}. \end{aligned} \quad (\text{B.18})$$

Here, we have assumed that \mathcal{F}_A and \mathcal{F}_{ABC} are constant and have used $\mathcal{F}^A_{DE} \partial_A \hat{E}_{\hat{B}}^C(y) = 0$, which is satisfied under our setup $\mathcal{F}^\alpha_{BC} = 0$. In addition, $\hat{\mathcal{R}}$ is the generalized Ricci scalar associated with the untwisted fields $\{\hat{\mathcal{H}}_{AB}, \hat{d}\}$,

$$\hat{\mathcal{R}} \equiv \mathcal{H}^{\hat{A}\hat{B}} (2 \hat{D}_{\hat{A}} \hat{\mathcal{F}}_{\hat{B}} - \hat{\mathcal{F}}_{\hat{A}} \hat{\mathcal{F}}_{\hat{B}}) - \frac{1}{12} \mathcal{H}^{\hat{A}\hat{D}} (\mathcal{H}^{\hat{B}\hat{E}} \mathcal{H}^{\hat{C}\hat{F}} - 3 \eta^{\hat{B}\hat{E}} \eta^{\hat{C}\hat{F}}) \hat{\mathcal{F}}_{\hat{A}\hat{B}\hat{C}} \hat{\mathcal{F}}_{\hat{D}\hat{E}\hat{F}}. \quad (\text{B.19})$$

Now, let us show the covariance of the equations of motion under the $O(n, n)$ PL T -plurality transformation,

$$\hat{\mathcal{H}}_{AB} \rightarrow (C \hat{\mathcal{H}} C^T)_{AB}, \quad \hat{d} \rightarrow \hat{d}, \quad \mathcal{F}_A \rightarrow (C \mathcal{F})_A, \quad \mathcal{F}_{ABC} \rightarrow C_A^D C_B^E C_C^F \mathcal{F}_{DEF}. \quad (\text{B.20})$$

From the relation in Eq. (B.14), the first rule implies the following rule for the untwisted vielbein:

$$\hat{V}_A^{\hat{B}} \rightarrow C_A^C \hat{V}_C^{\hat{B}}, \quad \hat{V}_A^B \rightarrow \hat{V}_A^C (C^{-1})_C^B. \quad (\text{B.21})$$

Since the untwisted fields satisfy Eq. (B.15), we can show for an arbitrary untwisted field $g(y)$ that

$$\hat{\mathcal{D}}_A^{\hat{B}} g'(y) = \hat{V}_A^B (C^{-1})_B^C \partial_C g'(y) = \hat{V}_A^B \partial_B g'(y) = \hat{\mathcal{D}}_A^{\hat{B}} g'(y), \quad (\text{B.22})$$

and $\hat{\mathcal{F}}_A^{\hat{\lambda}}$ and $\hat{\mathcal{F}}_{\hat{A}\hat{B}\hat{C}}^{\hat{\lambda}}$ are invariant under the PL T -plurality transformation. Then, from Eq. (B.16), the fluxes $\mathcal{F}_{\hat{\lambda}}$ and $\mathcal{F}_{\hat{A}\hat{B}\hat{C}}^{\hat{\lambda}}$ are also invariant,

$$\mathcal{F}'_{\hat{\lambda}} = \mathcal{F}_{\hat{\lambda}}, \quad \mathcal{F}'_{\hat{A}\hat{B}\hat{C}}^{\hat{\lambda}} = \mathcal{F}_{\hat{A}\hat{B}\hat{C}}^{\hat{\lambda}}. \quad (\text{B.23})$$

Moreover, according to the constancy of \mathcal{F}_A and \mathcal{F}_{ABC} , $\mathcal{F}_{\hat{\lambda}}$ and $\mathcal{F}_{\hat{A}\hat{B}\hat{C}}^{\hat{\lambda}}$ depend only on the spectator fields, and from Eqs. (B.15), (B.22), and (B.23) we have

$$\begin{aligned} \mathcal{D}'_{\hat{\lambda}} \mathcal{F}'_{\hat{B}} &= \hat{\mathcal{D}}_{\hat{\lambda}} \mathcal{F}'_{\hat{B}} = \hat{\mathcal{D}}_{\hat{\lambda}} \mathcal{F}_{\hat{B}} = \mathcal{D}_{\hat{\lambda}} \mathcal{F}_{\hat{B}}, \\ \mathcal{D}'_{\hat{\lambda}} \mathcal{F}'_{\hat{B}\hat{C}\hat{D}} &= \hat{\mathcal{D}}_{\hat{\lambda}} \mathcal{F}'_{\hat{B}\hat{C}\hat{D}} = \hat{\mathcal{D}}_{\hat{\lambda}} \mathcal{F}_{\hat{B}\hat{C}\hat{D}} = \mathcal{D}_{\hat{\lambda}} \mathcal{F}_{\hat{B}\hat{C}\hat{D}}. \end{aligned} \quad (\text{B.24})$$

Then, as is clear from Eqs. (B.9) and (B.10), \mathcal{R} and $\mathcal{G}^{\hat{A}\hat{B}}$ are also invariant,

$$\mathcal{R}' = \mathcal{R}, \quad \mathcal{G}'^{\hat{A}\hat{B}} = \mathcal{G}^{\hat{A}\hat{B}}. \quad (\text{B.25})$$

Note that if we define the quantity

$$\mathcal{G}^{AB} \equiv \hat{V}_{\hat{C}}^A \hat{V}_{\hat{D}}^B \mathcal{G}^{\hat{C}\hat{D}}, \quad (\text{B.26})$$

we can clearly see that it transforms as $\mathcal{G}^{AB} \rightarrow (C^{-T} \mathcal{G} C^{-1})^{AB} = C^A_C \mathcal{G}^{CD} C_D^B$. This is precisely the \mathcal{G}^{AB} discussed in Eq. (2.26) when the untwisted fields are constant.

In order to show the covariance of S_{MN} , it is convenient to use the relation

$$\hat{V}^{\hat{a}C} V^{bD} U_C^M U_D^N S_{MN} = \mathcal{G}^{\hat{a}b} = \mathcal{G}'^{\hat{a}b} = \hat{V}'^{\hat{a}C} V'^{bD} U_C^M U_D^N S'_{MN}. \quad (\text{B.27})$$

From $\hat{V}'^{\hat{a}C} = \hat{V}^{\hat{a}D} (C^{-1})_D^C$, we find that

$$U_A^M U_B^N S_{MN} = (C^{-1})_A^C (C^{-1})_B^D U_C^M U_D^N S'_{MN}. \quad (\text{B.28})$$

Namely, we obtain

$$S'_{MN} = (h S h^T)_{MN}, \quad h_M^N \equiv U_M^A C_A^B U_B^N. \quad (\text{B.29})$$

Therefore, the generalized Ricci tensor transforms covariantly in the same manner as \mathcal{H}_{MN} .

B.2. R–R sector

The R–R fields in the approach of Ref. [125] are defined as

$$|F\rangle = \sum_p \frac{1}{p!} F_{m_1 \dots m_p} \Gamma^{m_1 \dots m_p} |0\rangle, \quad \Gamma^{m_1 \dots m_p} \equiv \Gamma^{[m_1 \dots m_p]}. \quad (\text{B.30})$$

Here, the gamma matrix $(\Gamma^M) \equiv (\Gamma^m, \Gamma_m)$ is real and satisfies $(\Gamma^M)^T = \Gamma_M$ and

$$\{\Gamma^M, \Gamma^N\} = \eta^{MN} \quad (\Leftrightarrow \{\Gamma^m, \Gamma_n\} = \delta_n^m, \quad \{\Gamma^m, \Gamma^n\} = 0 = \{\Gamma_m, \Gamma_n\}). \quad (\text{B.31})$$

By considering Γ^m and Γ_n as the creation and annihilation operator, we define the Clifford vacuum $|0\rangle$ as $\Gamma_m |0\rangle = 0$ that is normalized as $\langle 0|0\rangle = 1$ where $\langle 0| \equiv |0\rangle^\top$. We define the charge conjugation matrix as

$$\begin{aligned} \mathcal{C} &\equiv (\Gamma^0 \pm \Gamma_0) \cdots (\Gamma^{D-1} \pm \Gamma_{D-1}) \quad (D \text{ even/odd}), \\ \mathcal{C} \Gamma^A \mathcal{C}^{-1} &= -(\Gamma^A)^\top, \quad \mathcal{C}^{-1} = (-1)^{\frac{D(D+1)}{2}} \mathcal{C} = \mathcal{C}^\top, \end{aligned} \tag{B.32}$$

and introduce the notations

$$\langle F| \equiv (|F\rangle)^\top = \sum_p \frac{1}{p!} F_{m_1 \cdots m_p} \langle 0| (\Gamma^{m_p})^\top \cdots (\Gamma^{m_1})^\top, \quad \overline{\langle F|} \equiv \langle F| \mathcal{C}^\top. \tag{B.33}$$

In type IIA/IIB theory, the R–R field strength satisfies

$$\Gamma^{11} |F\rangle = \pm |F\rangle \quad (\text{type IIA/IIB}), \tag{B.34}$$

where the chirality operator is defined by

$$\Gamma^{11} \equiv (-1)^{N_F}, \quad N_F \equiv \Gamma^m \Gamma_m. \tag{B.35}$$

The Bianchi identity is given by

$$\partial |F\rangle = 0, \quad \partial \equiv \Gamma^M \partial_M, \tag{B.36}$$

where the nilpotency $\partial^2 = 0$ is ensured by the section condition. The R–R potential (in the A-basis) is defined through

$$|F\rangle = \partial |A\rangle, \quad |A\rangle = \sum_p \frac{1}{p!} A_{m_1 \cdots m_p} \Gamma^{m_1 \cdots m_p} |0\rangle, \tag{B.37}$$

and in terms of differential form we have¹⁴

$$F = dA, \quad d \equiv dx^m \wedge \partial_m + \iota_m \tilde{\partial}^m. \tag{B.38}$$

Under an $O(D, D)$ transformation,

$$\mathcal{H}_{MN} \rightarrow \mathcal{H}'_{MN} = (h \mathcal{H} h^\top)_{MN}, \tag{B.39}$$

the $O(D, D)$ spinors, $|F\rangle$ and $|A\rangle$, transform as

$$|F\rangle \rightarrow |F'\rangle = S_h |F\rangle, \quad |A\rangle \rightarrow |A'\rangle = S_h |A\rangle, \tag{B.40}$$

where S_h is defined through

$$S_h \Gamma_M S_h^{-1} = (h^{-1})_M{}^N \Gamma_N. \tag{B.41}$$

¹⁴ In GSE, the R–R fields have the dual-coordinate dependence as $A = e^{-l^m \tilde{x}_m} \bar{A}(x^m)$ and $F = e^{-l^m \tilde{x}_m} \bar{F}(x^m)$, and the relation $F = dA$ reproduces $\bar{F} = e^{l^m \tilde{x}_m} dA = d\bar{A} - \iota_l \bar{A}$. By considering $\{\bar{A}, \bar{F}\}$ as the dynamical fields, we obtain the relation in Eq. (A.10). See Ref. [149] for more detail.

We also define the corresponding operation \mathbb{S}_h acting on the polyform F as

$$S_h |F\rangle = |\mathbb{S}_h F\rangle. \tag{B.42}$$

The concrete expressions of S_h and \mathbb{S}_h for the $GL(D)$ -, B -, and β -transformation are as follows:

$$S_{h_M} = e^{\frac{1}{2} \rho_m^n [\Gamma^m, \Gamma_n]} = \frac{1}{\sqrt{|\det M|}} e^{\rho_m^n \Gamma^m \Gamma_n} \quad (\rho \equiv \ln M) \tag{B.43}$$

$$\Leftrightarrow h_M = \begin{pmatrix} M & 0 \\ 0 & M^{-\top} \end{pmatrix} \Leftrightarrow \mathbb{S}_{h_M} F \equiv F^{(M)} \quad (F_{m_1 \dots m_p}^{(M)} \equiv M_{m_1}^{n_1} \dots M_{m_p}^{n_p} F_{n_1 \dots n_p}),$$

$$S_{h_\omega} = e^{\frac{1}{2} \omega_{mn} \Gamma^{mn}} \Leftrightarrow h_\omega = \begin{pmatrix} \mathbf{1}_d & \omega \\ 0 & \mathbf{1}_d \end{pmatrix} \Leftrightarrow \mathbb{S}_{h_\omega} = e^{(\frac{1}{2} \omega_{mn} dx^m \wedge dx^n)^\wedge}, \tag{B.44}$$

$$S_{h_\chi} = e^{\frac{1}{2} \chi^{mn} \Gamma_{mn}} \Leftrightarrow h_\chi = \begin{pmatrix} \mathbf{1}_d & 0 \\ \chi & \mathbf{1}_d \end{pmatrix} \Leftrightarrow \mathbb{S}_{h_\chi} = e^{\frac{1}{2} \chi^{mn} \iota_{m\iota n}}. \tag{B.45}$$

The factorized T -duality along the x^z -direction is generated by

$$S_{h_z} = (\Gamma^z - \Gamma_z) \Gamma^{11} \Leftrightarrow h_z = \begin{pmatrix} \mathbf{1}_d - e_z & e_z \\ e_z & \mathbf{1}_d - e_z \end{pmatrix} \\ \Leftrightarrow \mathbb{S}_{h_z} F = F \wedge dx^z + F \vee dx^z. \tag{B.46}$$

In fact, the R–R field $|F\rangle$ is as an $O(D, D)$ spinor density with weight $1/2$.¹⁵ Correspondingly, under the $GL(D)$ transformation, the above S_{h_M} needs to be corrected as

$$\tilde{S}_{h_M} |F\rangle \equiv \sqrt{|\det M|} S_{h_M} |F\rangle = |\mathbb{S}_{h_M} F\rangle \tag{B.47}$$

when acting on the $O(D, D)$ spinor density. We can absorb the extra factor $\sqrt{|\det M|}$ into the DFT dilaton by considering a weightless $O(D, D)$ spinor, $|\mathcal{F}\rangle \equiv e^d |F\rangle$.

For later convenience, we define S_g and \mathcal{K} as

$$S_g \Gamma_M S_g^{-1} = -\mathbf{g}_M^N \Gamma_N, \quad (\mathbf{g}_{MN}) \equiv \begin{pmatrix} g_{mn} & 0 \\ 0 & g^{mn} \end{pmatrix}, \\ \mathcal{K} \Gamma_M \mathcal{K}^{-1} = -\mathcal{H}_M^N \Gamma_N, \quad \mathcal{K} = e^{\mathbf{B}} S_g e^{-\mathbf{B}}, \quad \mathbf{B} \equiv \frac{1}{2} B_{mn} \Gamma^{mn}. \tag{B.48}$$

By using the property $S_g |0\rangle = \sqrt{|g|} |0\rangle$, we can show that

$$\langle \alpha | S_g | \beta \rangle = -\sqrt{|g|} \sum_p \frac{1}{p!} g^{m_1 n_1} \dots g^{m_p n_p} \alpha_{m_1 \dots m_p} \beta_{n_1 \dots n_p} = \langle \beta | S_g | \alpha \rangle \tag{B.49}$$

for $O(D, D)$ spinors $|\alpha\rangle$ and $|\beta\rangle$. Moreover, the self-duality relation in Eq. (A.8) can be expressed as

$$|F\rangle = \mathcal{K} |F\rangle. \tag{B.50}$$

¹⁵ This can also be observed from the definition of the generalized Lie derivative,

$$\hat{\mathfrak{L}}_V |F\rangle = (V^M \partial_M + \partial_M V_N \Gamma^{MN}) |F\rangle + \frac{1}{2} \partial_M V^N |F\rangle.$$

We also define the correspondent of \mathcal{K} for the untwisted metric as

$$\mathcal{K} = S_U \hat{\mathcal{K}} S_U^{-1}, \quad \hat{\mathcal{K}} \Gamma_A \hat{\mathcal{K}}^{-1} = -\hat{\mathcal{H}}_A^B \Gamma_B, \tag{B.51}$$

where $\Gamma_A \equiv U_A^M S_U^{-1} \Gamma_M S_U = \delta_A^M \Gamma_M$.

The bosonic part of the Lagrangian in type II DFT is

$$\mathcal{L} = e^{-2d} \mathcal{S} + \frac{1}{4} \langle \overline{F} | \mathcal{K} | F \rangle, \tag{B.52}$$

and the equations of motion for $\{\mathcal{H}_{MN}, d, |A\rangle\}$ are summarized as

$$\begin{aligned} \mathcal{S}_{MN} &= \mathcal{E}_{MN}, & \mathcal{S} &= 0, & \partial \mathcal{K} | F \rangle &= 0, \\ \mathcal{E}_{MN} &\equiv -\frac{1}{4} e^{2d} \left[\langle \overline{F} | \Gamma_{(M} \mathcal{K} \Gamma_{N)} | F \rangle + \frac{1}{2} \mathcal{H}_{MN} \langle \overline{F} | \mathcal{K} | F \rangle \right]. \end{aligned} \tag{B.53}$$

Under the self-duality relation of Eq. (B.50), the equation of motion for the R–R field $\partial \mathcal{K} | F \rangle = 0$ is precisely the Bianchi identity $\partial | F \rangle = 0$.

In the gauged DFT, we consider the reduction ansatz (see Refs. [132,134,193,194])

$$| F \rangle = e^{-d(\alpha^I)} S_{U(\alpha^I)} | \hat{\mathcal{F}}(y) \rangle, \tag{B.54}$$

and assume that $| \hat{\mathcal{F}}(y) \rangle$ satisfies the condition $\mathcal{D}_A | \hat{\mathcal{F}}(y) \rangle = \partial_A | \hat{\mathcal{F}}(y) \rangle$, similar to Eq. (B.15). In the case of the twist matrix $U = R \mathbf{\Pi}$, $| \hat{\mathcal{F}} \rangle$ is explicitly given by

$$\begin{aligned} | \hat{\mathcal{F}} \rangle &= e^d S_{\mathbf{\Pi}^{-1}} S_{R^{-1}} | F \rangle \\ &= \frac{e^d}{\sqrt{|\det(e_a^m)|}} e^{\frac{1}{2} \Pi^{ab} \Gamma_{ab}} \sum_p \frac{1}{p!} e_{a_1}^{m_1} \dots e_{a_p}^{m_p} F_{m_1 \dots m_p} \Gamma^{a_1 \dots a_p} | 0 \rangle. \end{aligned} \tag{B.55}$$

In terms of the differential form, this reads as

$$\hat{\mathcal{F}} = \frac{e^d}{\sqrt{|\det(e_a^m)|}} \mathbb{S}_{U^{-1}} F. \tag{B.56}$$

In terms of the untwisted field, the self-duality relation can be expressed as

$$| \hat{\mathcal{F}} \rangle = \hat{\mathcal{K}} | \hat{\mathcal{F}} \rangle. \tag{B.57}$$

We can clearly see that this relation is preserved under the PL T -plurality transformation

$$\hat{\mathcal{K}} \rightarrow S_C \hat{\mathcal{K}} S_C^{-1}, \quad | \hat{\mathcal{F}} \rangle \rightarrow S_C | \hat{\mathcal{F}} \rangle. \tag{B.58}$$

Now, let us show the covariance of the Bianchi identity. From the reduction ansatz, we have

$$\begin{aligned} 0 = \partial | F \rangle &= e^{-d} S_U (\partial - \Gamma^A \mathcal{D}_A \mathbf{d} + S_U^{-1} \partial S_U) | \hat{\mathcal{F}} \rangle \\ \Leftrightarrow \left(\partial - \frac{1}{2} \Gamma^A \mathcal{F}_A + \frac{1}{3!} \Gamma^{ABC} \mathcal{F}_{ABC} \right) | \hat{\mathcal{F}} \rangle &= 0, \end{aligned} \tag{B.59}$$

where we have used the identity

$$S_U^{-1} \partial S_U = \frac{1}{3!} \mathcal{F}_{ABC} \Gamma^{ABC} - \frac{1}{2} \Omega_{AB}^B \Gamma^A. \tag{B.60}$$

We require “the Bianchi identity” for the untwisted field $\partial|\hat{\mathcal{F}}\rangle = 0$, and then the Bianchi identity in Eq. (B.59) becomes an algebraic relation,

$$\left(\frac{1}{3!}\Gamma^{ABC}\mathcal{F}_{ABC}-\frac{1}{2}\Gamma^A\mathcal{F}_A\right)|\hat{\mathcal{F}}\rangle=0. \quad (\text{B.61})$$

This is manifestly covariant under the PL T -plurality transformation. Since the Bianchi identity and the self-duality relation are covariantly transformed, the equation of motion for the R–R field is also satisfied in the dualized background.

Finally, we show the covariance of the energy–momentum tensor. To this end, we define

$$\hat{\mathcal{E}}_{AB}\equiv U_A{}^M U_B{}^N \mathcal{E}_{MN}=-\frac{1}{4}e^{2\hat{d}}\left[\langle\hat{\mathcal{F}}|\Gamma_{(A}\hat{\mathcal{K}}\Gamma_{B)}|\hat{\mathcal{F}}\rangle+\frac{1}{2}\hat{\mathcal{H}}_{AB}\langle\hat{\mathcal{F}}|\hat{\mathcal{K}}|\hat{\mathcal{F}}\rangle\right]. \quad (\text{B.62})$$

Under the PL T -plurality, Eq. (B.58), we can easily show that

$$U_A{}^M U_B{}^N \mathcal{E}'_{MN}=\hat{\mathcal{E}}'_{AB}=C_A{}^C C_B{}^D \hat{\mathcal{E}}_{CD}=C_A{}^C C_B{}^D U_A{}^M U_B{}^N \mathcal{E}_{MN}, \quad (\text{B.63})$$

and, similar to the generalized Ricci tensor, we obtain

$$\mathcal{E}'_{MN}=(h\mathcal{E}h^T)_{MN}, \quad h_M{}^N\equiv U_M{}^A C_A{}^B U_B{}^N. \quad (\text{B.64})$$

This completes the proof for the covariance of the equations of motion.

References

- [1] K. Kikkawa and M. Yamasaki, Phys. Lett. B **149**, 357 (1984).
- [2] N. Sakai and I. Senda, Prog. Theor. Phys. **75**, 692 (1986); **77**, 773 (1987) [erratum].
- [3] V. P. Nair, A. Shapere, A. Strominger, and F. Wilczek, Nucl. Phys. B **287**, 402 (1987).
- [4] P. Ginsparg and C. Vafa, Nucl. Phys. B **289**, 414 (1987).
- [5] A. Giveon, E. Rabinovici, and G. Veneziano, Nucl. Phys. B **322**, 167 (1989).
- [6] A. Shapere and F. Wilczek, Nucl. Phys. B **320**, 669 (1989).
- [7] T. H. Buscher, Phys. Lett. B **194**, 59 (1987).
- [8] T. H. Buscher, Phys. Lett. B **201**, 466 (1988).
- [9] S. Cecotti, S. Ferrara, and L. Girardello, Nucl. Phys. B **308**, 436 (1988).
- [10] M. J. Duff, Nucl. Phys. B **335**, 610 (1990).
- [11] M. Roček and E. Verlinde, Nucl. Phys. B **373**, 630 (1992) [arXiv:hep-th/9110053] [Search INSPIRE].
- [12] E. Alvarez, L. Alvarez-Gaumé, J. L. F. Barbón, and Y. Lozano, Nucl. Phys. B **415**, 71 (1994) [arXiv:hep-th/9309039] [Search INSPIRE].
- [13] E. Bergshoeff, C. Hull, and T. Ortín, Nucl. Phys. B **451**, 547 (1995) [arXiv:hep-th/9504081] [Search INSPIRE].
- [14] S. F. Hassan, Nucl. Phys. B **460**, 362 (1996) [arXiv:hep-th/9504148] [Search INSPIRE].
- [15] E. Bergshoeff, M. de Roo, M. B. Green, G. Papadopoulos, and P. K. Townsend, Nucl. Phys. B **470**, 113 (1996) [arXiv:hep-th/9601150] [Search INSPIRE].
- [16] S. F. Hassan, Nucl. Phys. B **568**, 145 (2000) [arXiv:hep-th/9907152] [Search INSPIRE].
- [17] A. Giveon, M. Porrati, and E. Rabinovici, Phys. Rept. **244**, 77 (1994) [arXiv:hep-th/9401139] [Search INSPIRE].
- [18] E. Alvarez, L. Alvarez-Gaume, and Y. Lozano, Nucl. Phys. B. Proc. Suppl. **41**, 1 (1995) [arXiv:hep-th/9410237] [Search INSPIRE].
- [19] X. C. de la Ossa and F. Quevedo, Nucl. Phys. B **403**, 377 (1993) [arXiv:hep-th/9210021] [Search INSPIRE].
- [20] B. E. Fridling and A. Jevicki, Phys. Lett. B **134**, 70 (1984).
- [21] E. S. Fradkin and A. A. Tseytlin, Ann. Phys. **162**, 31 (1985).
- [22] A. Giveon and M. Roček, Nucl. Phys. B **421**, 173 (1994) [arXiv:hep-th/9308154] [Search INSPIRE].
- [23] M. Gasperini, R. Ricci, and G. Veneziano, Phys. Lett. B **319**, 438 (1993) [arXiv:hep-th/9308112] [Search INSPIRE].

- [24] K. Sfetsos, Phys. Rev. D **50**, 2784 (1994) [arXiv:hep-th/9402031] [Search INSPIRE].
- [25] E. Álvarez, L. Álvarez-Gaumé, and Y. Lozano, Nucl. Phys. B **424**, 155 (1994) [arXiv:hep-th/9403155] [Search INSPIRE].
- [26] K. Sfetsos and A. A. Tseytlin, Nucl. Phys. B **427**, 245 (1994) [arXiv:hep-th/9404063] [Search INSPIRE].
- [27] E. Kiritsis and N. A. Obers, Phys. Lett. B **334**, 67 (1994) [arXiv:hep-th/9406082] [Search INSPIRE].
- [28] S. Elitzur, A. Giveon, E. Rabinovici, A. Schwimmer, and G. Veneziano, Nucl. Phys. B **435**, 147 (1995) [arXiv:hep-th/9409011] [Search INSPIRE].
- [29] E. Tyurin, Phys. Lett. B **348**, 386 (1995) [arXiv:hep-th/9411242] [Search INSPIRE].
- [30] I. Bakas and K. Sfetsos, Phys. Lett. B **349**, 448 (1995) [arXiv:hep-th/9502065] [Search INSPIRE].
- [31] Y. Lozano, Phys. Lett. B **355**, 165 (1995) [arXiv:hep-th/9503045] [Search INSPIRE].
- [32] E. Tyurin, Phys. Lett. B **364**, 157 (1995) [arXiv:hep-th/9507014] [Search INSPIRE].
- [33] S. Hewson, Class. Quant. Grav. **13**, 1739 (1996) [arXiv:hep-th/9510092] [Search INSPIRE].
- [34] K. Sfetsos, Phys. Rev. D **54**, 1682 (1996) [arXiv:hep-th/9602179] [Search INSPIRE].
- [35] E. Plauschinn, Nucl. Phys. B **893**, 257 (2015) [arXiv:1408.1715 [hep-th]] [Search INSPIRE].
- [36] B. Fraser, D. Manolopoulos, and K. Sfetsos, Nucl. Phys. B **934**, 498 (2018) [arXiv:1805.03657 [hep-th]] [Search INSPIRE].
- [37] C. Klimčík and P. Ševera, Phys. Lett. B **351**, 455 (1995) [arXiv:hep-th/9502122] [Search INSPIRE].
- [38] C. Klimčík, Nucl. Phys. B. Proc. Suppl. **46**, 116 (1996) [arXiv:hep-th/9509095] [Search INSPIRE].
- [39] C. Klimčík and P. Ševera, Phys. Lett. B **372**, 65 (1996) [arXiv:hep-th/9512040] [Search INSPIRE].
- [40] C. Klimčík and P. Ševera, Phys. Lett. B **383**, 281 (1996) [arXiv:hep-th/9605212] [Search INSPIRE].
- [41] C. Klimčík and P. Ševera, Phys. Lett. B **381**, 56 (1996) [arXiv:hep-th/9602162] [Search INSPIRE].
- [42] C. Klimčík and P. Ševera, arXiv:hep-th/9610198 [Search INSPIRE].
- [43] K. Sfetsos, Nucl. Phys. B. Proc. Suppl. **56**, 302 (1997) [arXiv:hep-th/9611199] [Search INSPIRE].
- [44] K. Sfetsos, Nucl. Phys. B **517**, 549 (1998) [arXiv:hep-th/9710163] [Search INSPIRE].
- [45] K. Sfetsos, Nucl. Phys. B **561**, 316 (1999) [arXiv:hep-th/9904188] [Search INSPIRE].
- [46] A. Bossard and N. Mohammadi, Nucl. Phys. B **619**, 128 (2001) [arXiv:hep-th/0106211] [Search INSPIRE].
- [47] R. Squellari, Nucl. Phys. B **853**, 379 (2011) [arXiv:1105.0162 [hep-th]] [Search INSPIRE].
- [48] M. A. Lledó and V. S. Varadarajan, Lett. Math. Phys. **45**, 247 (1998) [arXiv:hep-th/9803175] [Search INSPIRE].
- [49] M. A. Jafarizadeh and A. Rezaei-Aghdam, Phys. Lett. B **458**, 477 (1999) [arXiv:hep-th/9903152] [Search INSPIRE].
- [50] C. Klimčík, J. High Energy Phys. **0212**, 051 (2002) [arXiv:hep-th/0210095] [Search INSPIRE].
- [51] L. Hlavatý and I. Petr, Int. J. Mod. Phys. A **29**, 1450009 (2014) [arXiv:1308.0153 [hep-th]] [Search INSPIRE].
- [52] L. Hlavatý and L. Šnobl, Mod. Phys. Lett. A **17**, 429 (2002) [arXiv:hep-th/0110139] [Search INSPIRE].
- [53] L. Hlavatý and L. Šnobl, arXiv:math/0202209 [math.QA] [Search INSPIRE].
- [54] L. Šnobl and L. Hlavatý, Int. J. Mod. Phys. A **17**, 4043 (2002) [arXiv:math/0202210] [Search INSPIRE].
- [55] R. von Unge, J. High Energy Phys. **0207**, 014 (2002) [arXiv:hep-th/0205245] [Search INSPIRE].
- [56] L. Hlavatý and L. Šnobl, J. High Energy Phys. **0405**, 010 (2004) [arXiv:hep-th/0403164] [Search INSPIRE].
- [57] L. Hlavatý and L. Šnobl, J. High Energy Phys. **0410**, 045 (2004) [arXiv:hep-th/0408126] [Search INSPIRE].
- [58] L. Hlavatý, arXiv:hep-th/0601172 [Search INSPIRE].
- [59] L. Hlavatý, J. Hýbl, and M. Turek, Int. J. Mod. Phys. A **22**, 1039 (2007) [arXiv:hep-th/0608069] [Search INSPIRE].
- [60] L. Hlavatý and M. Turek, J. Mod. Phys. **3**, 1088 (2012) [arXiv:1201.5939 [hep-th]] [Search INSPIRE].
- [61] L. Hlavatý and L. Šnobl, Nucl. Phys. B **768**, 209 (2007) [arXiv:hep-th/0608133] [Search INSPIRE].
- [62] L. Hlavatý and I. Petr, J. High Energy Phys. **1904**, 157 (2019) [arXiv:1811.12235 [hep-th]] [Search INSPIRE].
- [63] A. Y. Alekseev, C. Klimčík, and A. A. Tseytlin, Nucl. Phys. B **458**, 430 (1996) [arXiv:hep-th/9509123] [Search INSPIRE].
- [64] E. Tyurin and R. von Unge, Phys. Lett. B **382**, 233 (1996) [arXiv:hep-th/9512025] [Search INSPIRE].

- [65] K. Sfetsos, Phys. Lett. B **432**, 365 (1998) [arXiv:hep-th/9803019] [Search INSPIRE].
- [66] C. Klimčík and G. Valent, Phys. Lett. B **565**, 237 (2003) [arXiv:hep-th/0304053] [Search INSPIRE].
- [67] G. Valent, C. Klimčík, and R. Squellari, Phys. Lett. B **678**, 143 (2009) [arXiv:0902.1459 [hep-th]] [Search INSPIRE].
- [68] K. Sfetsos and K. Siampos, J. High Energy Phys. **0906**, 082 (2009) [arXiv:0904.4248 [hep-th]] [Search INSPIRE].
- [69] S. D. Avramis, J.-P. Derendinger, and N. Prezas, Nucl. Phys. B **827**, 281 (2010) [arXiv:0910.0431 [hep-th]] [Search INSPIRE].
- [70] K. Sfetsos, K. Siampos, and D. C. Thompson, Nucl. Phys. B **827**, 545 (2010) [arXiv:0910.1345 [hep-th]] [Search INSPIRE].
- [71] L. Hlavatý, J. Navrátil, and L. Šnobl, Acta Polytech. **53**, 433 (2013) [arXiv:1212.5936 [hep-th]] [Search INSPIRE].
- [72] K. Sfetsos and D. C. Thompson, Nucl. Phys. B **846**, 21 (2011) [arXiv:1012.1320 [hep-th]] [Search INSPIRE].
- [73] Y. Lozano, E. Ó. Colgáin, K. Sfetsos, and D. C. Thompson, J. High Energy Phys. **1106**, 106 (2011) [arXiv:1104.5196 [hep-th]] [Search INSPIRE].
- [74] G. Itsios, Y. Lozano, E. Ó. Colgáin, and K. Sfetsos, J. High Energy Phys. **1208**, 132 (2012) [arXiv:1205.2274 [hep-th]] [Search INSPIRE].
- [75] Y. Lozano, E. Ó. Colgáin, D. Rodríguez-Gómez, and K. Sfetsos, Phys. Rev. Lett. **110**, 231601 (2013) [arXiv:1212.1043 [hep-th]] [Search INSPIRE].
- [76] G. Itsios, C. Núñez, K. Sfetsos, and D. C. Thompson, Phys. Lett. B **721**, 342 (2013) [arXiv:1212.4840 [hep-th]] [Search INSPIRE].
- [77] G. Itsios, C. Núñez, K. Sfetsos, and D. C. Thompson, Nucl. Phys. B **873**, 1 (2013) [arXiv:1301.6755 [hep-th]] [Search INSPIRE].
- [78] J. Jeong, Ö. Kelekci, and E. Ó. Colgáin, J. High Energy Phys. **1305**, 079 (2013) [arXiv:1302.2105 [hep-th]] [Search INSPIRE].
- [79] A. Barranco, J. Gaillard, N. T. Macpherson, C. Núñez, and D. C. Thompson, J. High Energy Phys. **1308**, 018 (2013) [arXiv:1305.7229 [hep-th]] [Search INSPIRE].
- [80] E. Gevorgyan and G. Sarkissian, J. High Energy Phys. **1403**, 035 (2014) [arXiv:1310.1264 [hep-th]] [Search INSPIRE].
- [81] N. T. Macpherson, J. High Energy Phys. **1311**, 137 (2013) [arXiv:1310.1609 [hep-th]] [Search INSPIRE].
- [82] Y. Lozano, E. Ó. Colgáin, and D. Rodríguez-Gómez, J. High Energy Phys. **1405**, 009 (2014) [arXiv:1311.4842 [hep-th]] [Search INSPIRE].
- [83] J. Gaillard, N. T. Macpherson, C. Núñez, and D. C. Thompson, Nucl. Phys. B **884**, 696 (2014) [arXiv:1312.4945 [hep-th]] [Search INSPIRE].
- [84] E. Caceres, N. T. Macpherson, and C. Núñez, J. High Energy Phys. **1408**, 107 (2014) [arXiv:1402.3294 [hep-th]] [Search INSPIRE].
- [85] Y. Lozano and N. T. Macpherson, J. High Energy Phys. **1411**, 115 (2014) [arXiv:1408.0912 [hep-th]] [Search INSPIRE].
- [86] K. Sfetsos and D. C. Thompson, J. High Energy Phys. **1411**, 006 (2014) [arXiv:1408.6545 [hep-th]] [Search INSPIRE].
- [87] Ö. Kelekci, Y. Lozano, N. T. Macpherson, and E. Ó. Colgáin, Class. Quant. Grav. **32**, 035014 (2015) [arXiv:1409.7406 [hep-th]] [Search INSPIRE].
- [88] N. T. Macpherson, C. Núñez, L. A. Pando Zayas, V. G. J. Rodgers, and C. A. Whiting, J. High Energy Phys. **1502**, 040 (2015) [arXiv:1410.2650 [hep-th]] [Search INSPIRE].
- [89] K. S. Kooner and S. Zacarías, J. High Energy Phys. **1508**, 143 (2015) [arXiv:1411.7433 [hep-th]] [Search INSPIRE].
- [90] T. R. Araujo and H. Nastase, Phys. Rev. D **91**, 126015 (2015) [arXiv:1503.00553 [hep-th]] [Search INSPIRE].
- [91] Y. Bea, J. D. Edelstein, G. Itsios, K. S. Kooner, C. Núñez, D. Schofield, and J. A. Sierra-García, J. High Energy Phys. **1505**, 062 (2015) [arXiv:1503.07527 [hep-th]] [Search INSPIRE].
- [92] Y. Lozano, N. T. Macpherson, J. Montero, and E. Ó. Colgáin, J. High Energy Phys. **1508**, 121 (2015) [arXiv:1507.02659 [hep-th]] [Search INSPIRE].
- [93] Y. Lozano, N. T. Macpherson, and J. Montero, J. High Energy Phys. **1510**, 004 (2015) [arXiv:1507.02660 [hep-th]] [Search INSPIRE].

- [94] T. R. Araujo and H. Nastase, *J. High Energy Phys.* **1511**, 203 (2015) [arXiv:1508.06568 [hep-th]] [Search INSPIRE].
- [95] N. T. Macpherson, C. Núñez, D. C. Thompson, and S. Zacarias, *J. High Energy Phys.* **1511**, 212 (2015) [arXiv:1509.04286 [hep-th]] [Search INSPIRE].
- [96] H. Dimov, S. Mladenov, R. C. Rashkov, and T. Vetsov, *Fortsch. Phys.* **64**, 657 (2016) [arXiv:1511.00269 [hep-th]] [Search INSPIRE].
- [97] L. A. Pando Zayas, V. G. J. Rodgers, and C. A. Whiting, *J. High Energy Phys.* **1602**, 061 (2016) [arXiv:1511.05991 [hep-th]] [Search INSPIRE].
- [98] Y. Lozano, N. T. Macpherson, J. Montero, and C. Núñez, *J. High Energy Phys.* **1611**, 133 (2016) [arXiv:1609.09061 [hep-th]] [Search INSPIRE].
- [99] L. A. Pando Zayas, D. Tsimpis, and C. A. Whiting, *Phys. Rev. D* **96**, 046013 (2017) [arXiv:1701.01643 [hep-th]] [Search INSPIRE].
- [100] Y. Lozano, C. Núñez, and S. Zacarias, *J. High Energy Phys.* **1709**, 008 (2017) [arXiv:1703.00417 [hep-th]] [Search INSPIRE].
- [101] R. Borsato and L. Wulff, *J. High Energy Phys.* **1808**, 027 (2018) [arXiv:1806.04083 [hep-th]] [Search INSPIRE].
- [102] B. Hoare and A. A. Tseytlin, *J. Phys. A* **49**, 494001 (2016) [arXiv:1609.02550 [hep-th]] [Search INSPIRE].
- [103] R. Borsato and L. Wulff, *Phys. Rev. Lett.* **117**, 251602 (2016) [arXiv:1609.09834 [hep-th]] [Search INSPIRE].
- [104] B. Hoare and D. C. Thompson, *J. High Energy Phys.* **1702**, 059 (2017) [arXiv:1611.08020 [hep-th]] [Search INSPIRE].
- [105] R. Borsato and L. Wulff, *J. High Energy Phys.* **1710**, 024 (2017) [arXiv:1706.10169 [hep-th]] [Search INSPIRE].
- [106] C. Klimčík, *Nucl. Phys. B* **900**, 259 (2015) [arXiv:1508.05832 [hep-th]] [Search INSPIRE].
- [107] B. Vicedo, *J. Phys. A* **48**, 355203 (2015) [arXiv:1504.06303 [hep-th]] [Search INSPIRE].
- [108] B. Hoare and A. A. Tseytlin, *Nucl. Phys. B* **897**, 448 (2015) [arXiv:1504.07213 [hep-th]] [Search INSPIRE].
- [109] K. Sfetsos, K. Siampos, and D. C. Thompson, *Nucl. Phys. B* **899**, 489 (2015) [arXiv:1506.05784 [hep-th]] [Search INSPIRE].
- [110] C. Klimčík, *Phys. Lett. B* **760**, 345 (2016) [arXiv:1606.03016 [hep-th]] [Search INSPIRE].
- [111] C. Klimčík, *Phys. Lett. B* **772**, 725 (2017) [arXiv:1706.08912 [hep-th]] [Search INSPIRE].
- [112] C. Klimčík, *Nucl. Phys. B* **939**, 191 (2019) [arXiv:1809.01614 [hep-th]] [Search INSPIRE].
- [113] C. Klimčík, arXiv:1903.00439 [hep-th] [Search INSPIRE].
- [114] W. Siegel, *Phys. Rev. D* **47**, 5453 (1993) [arXiv:hep-th/9302036] [Search INSPIRE].
- [115] W. Siegel, *Phys. Rev. D* **48**, 2826 (1993) [arXiv:hep-th/9305073] [Search INSPIRE].
- [116] W. Siegel, arXiv:hep-th/9308133 [Search INSPIRE].
- [117] C. Hull and B. Zwiebach, *J. High Energy Phys.* **0909**, 099 (2009) [arXiv:0904.4664 [hep-th]] [Search INSPIRE].
- [118] C. Hull and B. Zwiebach, *J. High Energy Phys.* **0909**, 090 (2009) [arXiv:0908.1792 [hep-th]] [Search INSPIRE].
- [119] O. Hohm, C. Hull, and B. Zwiebach, *J. High Energy Phys.* **1007**, 016 (2010) [arXiv:1003.5027 [hep-th]] [Search INSPIRE].
- [120] O. Hohm, C. Hull, and B. Zwiebach, *J. High Energy Phys.* **1008**, 008 (2010) [arXiv:1006.4823 [hep-th]] [Search INSPIRE].
- [121] I. Jeon, K. Lee, and J.-H. Park, *J. High Energy Phys.* **1104**, 014 (2011) [arXiv:1011.1324 [hep-th]] [Search INSPIRE].
- [122] O. Hohm and S. K. Kwak, *J. Phys. A* **44**, 085404 (2011) [arXiv:1011.4101 [hep-th]] [Search INSPIRE].
- [123] I. Jeon, K. Lee, and J.-H. Park, *Phys. Rev. D* **84**, 044022 (2011) [arXiv:1105.6294 [hep-th]] [Search INSPIRE].
- [124] O. Hohm, S. K. Kwak, and B. Zwiebach, *Phys. Rev. Lett.* **107**, 171603 (2011) [arXiv:1106.5452 [hep-th]] [Search INSPIRE].
- [125] O. Hohm, S. K. Kwak, and B. Zwiebach, *J. High Energy Phys.* **1109**, 013 (2011) [arXiv:1107.0008 [hep-th]] [Search INSPIRE].

- [126] O. Hohm and S. K. Kwak, *J. High Energy Phys.* **1111**, 086 (2011) [arXiv:1108.4937 [hep-th]] [Search INSPIRE].
- [127] O. Hohm and B. Zwiebach, *J. High Energy Phys.* **1205**, 126 (2012) [arXiv:1112.5296 [hep-th]] [Search INSPIRE].
- [128] I. Jeon, K. Lee, and J.-H. Park, *J. High Energy Phys.* **1209**, 079 (2012) [arXiv:1206.3478 [hep-th]] [Search INSPIRE].
- [129] O. Hohm and B. Zwiebach, *J. High Energy Phys.* **1302**, 075 (2013) [arXiv:1207.4198 [hep-th]] [Search INSPIRE].
- [130] I. Jeon, K. Lee, J.-H. Park, and Y. Suh, *Phys. Lett. B* **723**, 245 (2013) [arXiv:1210.5078 [hep-th]] [Search INSPIRE].
- [131] G. Aldazabal, W. Baron, D. Marqués, and C. Núñez, *J. High Energy Phys.* **1111**, 052 (2011); **1111**, 109 (2011) [erratum] [arXiv:1109.0290 [hep-th]] [Search INSPIRE].
- [132] D. Geissbühler, *J. High Energy Phys.* **1111**, 116 (2011) [arXiv:1109.4280 [hep-th]] [Search INSPIRE].
- [133] M. Graña and D. Marques, *J. High Energy Phys.* **1204**, 020 (2012) [arXiv:1201.2924 [hep-th]] [Search INSPIRE].
- [134] D. Geissbühler, D. Marqués, C. Núñez, and V. Penas, *J. High Energy Phys.* **1306**, 101 (2013) [arXiv:1304.1472 [hep-th]] [Search INSPIRE].
- [135] R. Blumenhagen, F. Hassler, and D. Lüst, *J. High Energy Phys.* **1502**, 001 (2015) [arXiv:1410.6374 [hep-th]] [Search INSPIRE].
- [136] R. Blumenhagen, P. du Bosque, F. Hassler, and D. Lüst, *J. High Energy Phys.* **1508**, 056 (2015) [arXiv:1502.02428 [hep-th]] [Search INSPIRE].
- [137] P. du Bosque, F. Hassler, and D. Lüst, *J. High Energy Phys.* **1602**, 039 (2016) [arXiv:1509.04176 [hep-th]] [Search INSPIRE].
- [138] F. Hassler, arXiv:1707.08624 [hep-th] [Search INSPIRE].
- [139] S. Demulder, F. Hassler, and D. C. Thompson, *J. High Energy Phys.* **1902**, 189 (2019) [arXiv:1810.11446 [hep-th]] [Search INSPIRE].
- [140] D. Lüst and D. Osten, *J. High Energy Phys.* **1805**, 165 (2018) [arXiv:1803.03971 [hep-th]] [Search INSPIRE].
- [141] B. Jurčo and J. Vysoký, *J. Geom. Phys.* **130**, 1 (2018) [arXiv:1708.04079 [hep-th]] [Search INSPIRE].
- [142] P. Ševera and F. Valach, arXiv:1810.07763 [math.DG] [Search INSPIRE].
- [143] J. J. Fernández-Melgarejo, J. Sakamoto, Y. Sakatani, and K. Yoshida, *J. High Energy Phys.* **1712**, 108 (2017) [arXiv:1710.06849 [hep-th]] [Search INSPIRE].
- [144] M. Hong, Y. Kim, and E. Ó. Colgáin, *Eur. Phys. J. C* **78**, 1025 (2018) [arXiv:1801.09567 [hep-th]] [Search INSPIRE].
- [145] G. Arutyunov, S. Frolov, B. Hoare, R. Roiban, and A. A. Tseytlin, *Nucl. Phys. B* **903**, 262 (2016) [arXiv:1511.05795 [hep-th]] [Search INSPIRE].
- [146] A. A. Tseytlin and L. Wulff, *J. High Energy Phys.* **1606**, 174 (2016) [arXiv:1605.04884 [hep-th]] [Search INSPIRE].
- [147] C. M. Hull and P. K. Townsend, *Nucl. Phys. B* **274**, 349 (1986).
- [148] J. J. Fernández-Melgarejo, J. Sakamoto, Y. Sakatani, and K. Yoshida, *Phys. Rev. Lett.* **122**, 111602 (2019) [arXiv:1811.10600 [hep-th]] [Search INSPIRE].
- [149] J. Sakamoto, Y. Sakatani, and K. Yoshida, *Prog. Theor. Exp. Phys.* **2017**, 053B07 (2017) [arXiv:1703.09213 [hep-th]] [Search INSPIRE].
- [150] B. Hoare and A. A. Tseytlin, *J. High Energy Phys.* **1510**, 060 (2015) [arXiv:1508.01150 [hep-th]] [Search INSPIRE].
- [151] Y. Sakatani, S. Uehara, and K. Yoshida, *J. High Energy Phys.* **1704**, 123 (2017) [arXiv:1611.05856 [hep-th]] [Search INSPIRE].
- [152] J.-H. Park, S.-J. Rey, W. Rim, and Y. Sakatani, *J. High Energy Phys.* **1511**, 131 (2015) [arXiv:1507.07545 [hep-th]] [Search INSPIRE].
- [153] J. Sakamoto and Y. Sakatani, *J. High Energy Phys.* **1806**, 147 (2018) [arXiv:1803.05903 [hep-th]] [Search INSPIRE].
- [154] C. M. Hull and B. Spence, *Phys. Lett. B* **232**, 204 (1989).
- [155] C. M. Hull and B. Spence, *Nucl. Phys. B* **353**, 379 (1991).
- [156] T. Kugo and B. Zwiebach, *Prog. Theor. Phys.* **87**, 801 (1992) [arXiv:hep-th/9201040] [Search INSPIRE].

- [157] M. Fukuma, T. Oota, and H. Tanaka, *Prog. Theor. Phys.* **103**, 425 (2000) [arXiv:hep-th/9907132] [Search INSPIRE].
- [158] S. F. Hassan, *Nucl. Phys. B* **583**, 431 (2000) [arXiv:hep-th/9912236] [Search INSPIRE].
- [159] S. F. Hassan, *Nucl. Phys. Proc. Suppl.* **102**, 77 (2001) [arXiv:hep-th/0103149] [Search INSPIRE].
- [160] E. Bergshoeff, R. Kallosh, T. Ortín, D. Roest, and A. Van Proeyen, *Class. Quant. Grav.* **18**, 3359 (2001) [arXiv:hep-th/0103233] [Search INSPIRE].
- [161] L. Wulff, *Phys. Lett. B* **781**, 417 (2018) [arXiv:1803.07391 [hep-th]] [Search INSPIRE].
- [162] R. Borsato and L. Wulff, *J. Phys. A: Math. Theor.* **52**, 225401 (2019) [arXiv:1812.07287 [hep-th]] [Search INSPIRE].
- [163] K. Lee and J.-H. Park, *Nucl. Phys. B* **880**, 134 (2014) [arXiv:1307.8377 [hep-th]] [Search INSPIRE].
- [164] S. M. Ko, C. Melby-Thompson, R. Meyer, and J.-H. Park, *J. High Energy Phys.* **1512**, 144 (2015) [arXiv:1508.01121 [hep-th]] [Search INSPIRE].
- [165] K. Morand and J.-H. Park, *Eur. Phys. J. C* **77**, 685 (2017); **78**, 901 (2018) [erratum] [arXiv:1707.03713 [hep-th]] [Search INSPIRE].
- [166] K. Cho, K. Morand, and J.-H. Park, *Phys. Lett. B* **793**, 65 (2019) [arXiv:1808.10605 [hep-th]] [Search INSPIRE].
- [167] D. S. Berman, C. D. A. Blair, and R. Otsuki, arXiv:1902.01867 [hep-th] [Search INSPIRE].
- [168] A. A. Tseytlin, *Phys. Lett. B* **242**, 163 (1990).
- [169] A. A. Tseytlin, *Nucl. Phys. B* **350**, 395 (1991).
- [170] C. M. Hull, *J. High Energy Phys.* **0510**, 065 (2005) [arXiv:hep-th/0406102] [Search INSPIRE].
- [171] C. M. Hull, *J. High Energy Phys.* **0707**, 080 (2007) [arXiv:hep-th/0605149] [Search INSPIRE].
- [172] N. B. Copland, *J. High Energy Phys.* **1204**, 044 (2012) [arXiv:1111.1828 [hep-th]] [Search INSPIRE].
- [173] C. Klimčík and P. Ševera, *Nucl. Phys. B* **488**, 653 (1997) [arXiv:hep-th/9609112] [Search INSPIRE].
- [174] C. M. Hull and R. A. Reid-Edwards, *J. High Energy Phys.* **0909**, 014 (2009) [arXiv:0902.4032 [hep-th]] [Search INSPIRE].
- [175] R. A. Reid-Edwards, arXiv:1001.2479 [hep-th] [Search INSPIRE].
- [176] N. Mohammedi, *Phys. Lett. B* **414**, 104 (1997) [arXiv:hep-th/9709071] [Search INSPIRE].
- [177] M. J. Duff and J. X. Lu, *Nucl. Phys. B* **347**, 394 (1990).
- [178] D. S. Berman and M. J. Perry, *J. High Energy Phys.* **1106**, 074 (2011) [arXiv:1008.1763 [hep-th]] [Search INSPIRE].
- [179] P. West, *Class. Quant. Grav.* **18**, 4443 (2001) [arXiv:hep-th/0104081] [Search INSPIRE].
- [180] P. West, *Phys. Lett. B* **575**, 333 (2003) [arXiv:hep-th/0307098] [Search INSPIRE].
- [181] C. Hillmann, arXiv:0902.1509 [hep-th] [Search INSPIRE].
- [182] D. S. Berman, H. Godazgar, M. J. Perry, and P. West, *J. High Energy Phys.* **1202**, 108 (2012) [arXiv:1111.0459 [hep-th]] [Search INSPIRE].
- [183] D. S. Berman, M. Cederwall, A. Kleinschmidt, and D. C. Thompson, *J. High Energy Phys.* **1301**, 064 (2013) [arXiv:1208.5884 [hep-th]] [Search INSPIRE].
- [184] P. West, *J. High Energy Phys.* **1212**, 068 (2012) [arXiv:1206.7045 [hep-th]] [Search INSPIRE].
- [185] O. Hohm and H. Samtleben, *Phys. Rev. Lett.* **111**, 231601 (2013) [arXiv:1308.1673 [hep-th]] [Search INSPIRE].
- [186] Y. Sakatani and S. Uehara, *Prog. Theor. Exp. Phys.* **2017**, 113B01 (2017) [arXiv:1708.06342 [hep-th]] [Search INSPIRE].
- [187] C. D. A. Blair, E. Malek, and J.-H. Park, *J. High Energy Phys.* **1401**, 172 (2014) [arXiv:1311.5109 [hep-th]] [Search INSPIRE].
- [188] Y. Sakatani and S. Uehara, *Prog. Theor. Exp. Phys.* **2017**, 043B05 (2017) [arXiv:1701.07819 [hep-th]] [Search INSPIRE].
- [189] M. Hatsuda and K. Kamimura, *J. High Energy Phys.* **1211**, 001 (2012) [arXiv:1208.1232 [hep-th]] [Search INSPIRE].
- [190] M. Hatsuda and K. Kamimura, *J. High Energy Phys.* **1306**, 095 (2013) [arXiv:1305.2258 [hep-th]] [Search INSPIRE].
- [191] Y. Sakatani and S. Uehara, *Prog. Theor. Exp. Phys.* **2018**, 033B05 (2018) [arXiv:1712.10316 [hep-th]] [Search INSPIRE].
- [192] A. Catal-Ozer, arXiv:1904.00362 [hep-th] [Search INSPIRE].
- [193] A. Çatal-Özer, *J. High Energy Phys.* **1709**, 044 (2017) [arXiv:1705.08181 [hep-th]] [Search INSPIRE].
- [194] A. Çatal-Özer, *J. High Energy Phys.* **1802**, 179 (2018) [arXiv:1706.08883 [hep-th]] [Search INSPIRE].