

## Landmark role of $Z = 101/100$ nuclides in the transuranic region

K. Vijay Sai\*, R. Gowrishankar and P. C. Sood

<sup>1</sup>Department of Physics, Sri Satya Sai Institute of Higher Learning, Prasanthi Nilayam, A.P. – 515 134, INDIA

\* email: [kvijaysai@sssihl.edu.in](mailto:kvijaysai@sssihl.edu.in)

None of the transuranic ( $Z > 92$ ) species occur in nature; they are all produced in the laboratory through nuclear reactions. However, while the  $93 \leq Z \leq 101$  nuclides can be reached in  $\alpha$  (or lighter) particle induced reactions, exploration of the still higher  $Z$  domain is a highly challenging, albeit competitive, task.

Considering this fact, International Union of Pure and Applied Chemistry (IUPAC) and that of Physics (IUPAP) had constituted [1] a joint Transfermium Working Group (TWG) for authentication of discovery of  $Z > 100$  elements. A factual summary of experiments on the first identification, and the ones confirming the same, for each nuclide in the  $Z > 100$  isotopic sequences has since been presented by Thoenissen [2]. Exhaustive information across all nuclear regions is available in various nuclear data bases [3-5]. Spurred by findings in some of our recent reports [6,7], herein we examine the occurrence of a LANDMARK around  $Z = 100/101$  in the vast nuclear landscape. To start this exercise, we first look at some basic properties and processes.


A survey [3-5] of the half-lives ( $t_{1/2}$ ) of  $Z > 92$  nuclides reveals that such data for the

Longest-Lived Isotope (LLI) in each sequence fall in 3 zones: for  $Z < 100$  species, LLI is in years, for  $Z = 100/101$ , it is 100/51 days and for  $Z > 101$ , it progressively appears in h/s/ms. This observation clearly indicates a landmark at  $Z = 100/101$  along this trajectory.


Another evidence for such an occurrence is provided by consideration of their production procedures. It is seen that while the  $Z \leq 101$  species are produced in ‘stepping’ process wherein  $n/d/\alpha$  incident on a newly synthesized high  $Z$  material produces the next high  $Z$  species, the  $Z > 101$  species need Heavy-Ion accelerators for their production.

Next we turn our attention to the higher end of the periodic table. In our 2011 report [6], we had traced the trajectory starting from each of the 4 Naturally Occurring Radioactive Series (NORS) across the  $Z = 100/101$  domain right upto  $Z = 113$  isotopes synthesized in Cold Fusion (CF).

Our recent (2017) study of  $^{254}\text{Md}$  level structures [7] provides evidence for a landmark role of  $Z = 101/100$  [ $^{254}\text{Md} (\varepsilon) ^{254}\text{Fm}$ ] locale in the decay path of (4n+2) SHE, as shown in Fig. 1.




**Fig. 1:** Sequential  $\alpha$ -decays from SHE to NORS Head with marked discontinuities at Md ( $\varepsilon$ ) Fm for (4n) & (4n+2) series.



**Fig. 2:** Sequential  $\alpha$ -decays from SHE to NORS Head with marked discontinuities at  $\text{Md} (\epsilon) \text{ Fm}$  for  $4n+1$  and  $4n+3$  series


In the analysis therein [7], we expressly took note of the fact that decay of each CF-produced SHE proceeds via a genetically correlated sequential  $\alpha$ -chain. This process is depicted in the  $^{278}\text{Nh}$  ( $6\alpha$ )  $^{254}\text{Md}$  segment in our plot of Fig. 1. This plot, further on, shows an  $\alpha$ -discontinuity at the  $^{254}\text{Md} (\epsilon) ^{254}\text{Fm}$  transition point, and then proceeds via  $^{254}\text{Fm}$  ( $4\alpha$ )  $^{238}\text{U}$  to the sequential  $^{238}\text{U} \rightarrow ^{208}\text{Pb}$  ( $4n+2$ ) NORS. The Landmark feature at  $Z=101/100$  across the transuranic region is thus unmistakably established through the following process:



Another new feature, highlighting the Landmark at  $Z=100/101$ , brought out in our earlier study [7] was that all the  $Z \geq 101$  nuclides in this chain are odd-odd whereas all the  $Z < 101$  nuclides are even-even.

The third distinctive feature from SHE decay is that  $\alpha$  and SF are dominant decay modes for all  $Z > 101$  nuclei, whereas SF is usually minor/insignificant decay channel for  $Z \leq 101$  members thereof. It is of interest to add that, a collateral decay path of another  $4n+2$  SHE can be traced as  $^{266}\text{Mt}$  ( $4\alpha$ )  $^{250}\text{Md} (\epsilon) ^{250}\text{Fm}$ , again with a discontinuity at  $Z=101/100$ .

Results from similar analysis for each of the  $4n$ ,  $4n+1$  and  $4n+3$  SHE decay chain, are plotted in Fig.s 1 & 2. The decay chain for  $4n$  SHE, as shown in Fig. 1 corresponding to



replicates all the above discussed three features of  $4n+2$  chain, and thus confirms the Landmark role for the  $Z=101/100$  in the  $4n$  series as well.

The odd- $A$  ( $4n+1$ ) and ( $4n+3$ ) SHE, namely  $^{277}\text{Cn}$  and  $^{271}\text{Ds}$  respectively, decay chains shown in Fig. 2 again clearly depict the discontinuity at  $Z=100$ . This feature results from the fact that CF produces only  $n$ -deficient SHE and the odd- $A$  nuclides are predominantly even  $Z$  / odd  $N$  assemblies, and hence the sequential  $6/5$   $\alpha$ -decays terminate at an  $\epsilon$ -decaying  $Z=100$  isotope.

In summary, our analysis identifies five observed factors namely  $t_{1/2}$ , production process, SHE  $\alpha$ -chains terminations, odd-odd/ even-even transitions, and % SF branching. These factors, individually and collectively, establish the Landmark role for the respective  $Z=101/100$  isotope in the transuranic domain.

## References

- [1] Pure & Appl. Chem. **63** (1991) 879; **65** (1993) 1757
- [2] M. Thoennessen, ADNDT, **99** (2013) 312
- [3] KPL Chart of Nuclides, 17<sup>th</sup> ed. (2009)
- [4] ENSDF and XUNDL, August 2018 Version
- [5] G. Audi *et al.*, Chin. Phys. C **41** (2017) 030001
- [6] K. Vijay Sai *et al.*, Proc. DAE Symp. Nucl. Phys. **56** (2011) 186
- [7] P. C. Sood and R. Gowrishankar, Phys. Rev. C **95** (2017) 024317