

## A viscous blast-wave model for heavy-ion collisions

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 779 012065

(<http://iopscience.iop.org/1742-6596/779/1/012065>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 131.169.5.251

This content was downloaded on 13/02/2017 at 21:16

Please note that [terms and conditions apply](#).

You may also be interested in:

[Jet Measurements in Heavy-ion Collisions with CMS](#)

Raghav Kunnawalkam Elayavalli and CMS collaboration

[Resonances as Probes of Heavy-Ion Collisions at ALICE](#)

A G Knospe and ALICE Collaboration

[Analysis of Collective Flow in High Energy Heavy-Ion Collisions in the Relativistic VUU Approach](#)

Hong-Bo Zhou, Zhu-Xia Li and Yi-Zhong Zhuo

[Hard physics with heavy ions in CMS](#)

A David

[Locally anisotropic momentum distributions of hadrons at freeze-out in relativistic heavy-ion collisions](#)

Maciej Rybczynski and Wojciech Florkowski

[Recent development on heavy quark theory in heavy-ion collisions](#)

Shanshan Cao

[Jet correlation measurement in heavy-ion collisions: from RHIC to LHC](#)

Constantinos A Loizides

[Femtoscopy of the system shape fluctuations](#)

Sergei A Voloshin

[Indication of transverse radial flow in high-multiplicity proton--proton collisions at the Large Hadron Collider](#)

Premomoy Ghosh, Sanjib Muhuri, Jajati K Nayak et al.

# A viscous blast-wave model for heavy-ion collisions

Amaresh Jaiswal<sup>1</sup> and Volker Koch<sup>2</sup>

<sup>1</sup>GSI, Helmholtzzentrum für Schwerionenforschung, Planckstrasse 1, D-64291 Darmstadt, Germany

<sup>2</sup>Lawrence Berkeley National Laboratory, Nuclear Science Division, MS 70R0319, Berkeley, California 94720, USA

E-mail: jaiswal.amaresh@gmail.com, vkoch@lbl.gov

**Abstract.** We present a generalization of the blast-wave model by incorporating viscous effects in the fluid velocity profile as well as in the Cooper-Frye freeze-out. We apply this model to study the identified particles spectra and anisotropic flow at the Large Hadron Collider (LHC). We show that this improved viscous blast-wave model leads to good description of the transverse momentum distribution of particle multiplicities and elliptic as well as triangular flow. Within this model, we estimate the shear viscosity to entropy density ratio  $\eta/s \simeq 0.24$  at the LHC.

## 1. Introduction

The quark-gluon plasma (QGP) formed in relativistic heavy-ion collisions exhibit strong collective behaviour and hence can be studied within the framework of relativistic hydrodynamics. The hydrodynamical modelling of heavy-ion collisions suggests that the QGP behaves like a nearly perfect fluid having an extremely small shear viscosity to entropy density ratio  $\eta/s$  [1]. Apart from hydrodynamics, the collective behaviour of QGP can also be studied within the so-called blast-wave model [2, 3, 4, 5, 6]. Here we generalize the blast-wave model to include viscous effects by using a viscosity-based survival scale for initial geometrical anisotropies, formed in relativistic heavy-ion collisions, to parametrize the radial flow velocity. We employ this viscous blast-wave model to obtain the transverse momentum dependence of particle yields and flow harmonics for the Large Hadron Collider (LHC). We fix the model parameters by fitting the transverse momentum distribution of identified particle spectra. Subsequently, we demonstrate that this leads to reasonably good agreement with transverse momentum dependence of elliptic and triangular flow for various centralities. Within the present model, we estimate the shear viscosity to entropy density ratio  $\eta/s \simeq 0.24$  at the LHC.

## 2. The model

We work in the Milne co-ordinate system where,  $\tau = \sqrt{t^2 - z^2}$ ,  $\eta_s = \tanh^{-1}(z/t)$ ,  $r = \sqrt{x^2 + y^2}$ ,  $\varphi = \text{atan2}(y, x)$ . The most important feature of the blast-wave model is the Hubble like parametrization of transverse velocity,  $u^r \sim r$ , which is found to be in agreement with hydro results [7]. In addition, we include angular anisotropy in the radial fluid velocity profile in the transverse plane. The hydrodynamic fields are parametrized as [8]

$$T = T_f, \quad u^r = u_0 \frac{r}{R} \left[ 1 + 2 \sum_{n=1}^{\infty} u_n \cos[n(\varphi - \psi_n)] \right], \quad u^\varphi = u^{\eta_s} = 0, \quad u^\tau = \sqrt{1 + (u^r)^2}, \quad (1)$$



where  $R$  is the transverse radius of the fireball at freeze-out. The condition  $u^\mu u_\mu = 1$  leads to the expression for  $u^\tau$ . In order to determine  $u_n$ , we use the fact that the initial geometrical anisotropies eventually converts to anisotropies in the radial fluid velocity. The participant anisotropies,  $\varepsilon_n$ , is defined in terms of the Fourier expansion for a single-particle distribution as

$$f(\varphi) = \frac{1}{2\pi} \left[ 1 + 2 \sum_{n=1}^{\infty} \varepsilon_n \cos[n(\varphi - \psi_n)] \right], \quad (2)$$

where  $\psi_n$  is the angle between the  $x$  axis and the major axis of the participant distribution. Next, we determine the conversion efficiency of the initial geometrical eccentricity to final anisotropy in the radial fluid velocity, i.e.,  $u_n/\varepsilon_n$ .

We start from the dispersion relation for sound in a viscous medium,  $\omega = c_s k + ik^2 \frac{1}{T} \left( \frac{2}{3} \frac{\eta}{s} \right)$ , where  $c_s$  is the speed of sound in the medium,  $\eta$  is the coefficient of shear viscosity and  $s = (\epsilon + P)/T$  is the entropy density. Using a Fourier ansatz, one finds that the amplitudes of the stress tensor harmonics, with momentum  $k$ , are attenuated by a factor [9]

$$\delta T^{\mu\nu}(t, k) = \exp \left[ - \left( \frac{2}{3} \frac{\eta}{s} \right) \frac{k^2 t}{T} \right] \delta T^{\mu\nu}(0, k), \quad (3)$$

where the oscillatory pre-factor has been ignored. Each harmonics is a damped oscillator and form standing waves on the fireball circumference throughout the evolution, i.e.,  $k = n/R$ . Note that the presence of momentum squared in the exponent results in enhanced effect of viscosity for the higher harmonics. Therefore, at freeze-out time  $t_f$ , the wave amplitude reaction is [10]

$$\frac{\delta T^{\mu\nu}|_{t=t_f}}{\delta T^{\mu\nu}|_{t=0}} = \exp \left[ -n^2 \left( \frac{2}{3} \frac{\eta}{s} \right) \frac{t_f}{R^2 T_f} \right], \quad (4)$$

where  $T_f$  is the freeze-out temperature. The above equation indicates a viscosity-based survival scale for anisotropic structures formed by point like perturbations. The final radial fluid velocity is due to initial geometrical perturbations and, in the absence of viscosity, the conversion efficiency remains the same for all harmonics. Therefore, for a viscous medium, the conversion efficiency must be proportional to the wave amplitude reaction,

$$\frac{u_n}{\varepsilon_n} = \alpha_0 \exp \left[ -n^2 \left( \frac{2}{3} \frac{\eta}{s} \right) \frac{t_f}{R^2 T_f} \right], \quad (5)$$

where  $\alpha_0$  is the constant of proportionality. The above equation, along with Eq. (1), leads to the fluid velocity profile which is used in the freeze-out prescription to obtain particle spectra and anisotropic flow coefficients.

### 3. Particle spectra and anisotropic flow

Hadron spectra can be obtained using the Cooper-Frye prescription for particle production [11]

$$\frac{dN}{d^2 p_T dy} = \frac{1}{(2\pi)^3} \int p_\mu d\Sigma^\mu f(x, p), \quad (6)$$

where  $d\Sigma_\mu$  is the freeze-out hyper-surface and  $f(x, p)$  is the distribution function of the particles. For a system close to equilibrium,  $f = f_0 + \delta f$  where  $\delta f \ll f_0$ . For the equilibrium distribution function,  $f_0$ , we use Fermi-Dirac statistics ( $a = +1$ ) for baryons and Bose-Einstein statistics ( $a = -1$ ) for mesons. For the non-equilibrium part, we employ the Grad's 14-moment approximation with viscous corrections up to first-order [12]

$$\delta f_1 = \frac{f_0 \tilde{f}_0}{T^3} \left( \frac{\eta}{s} \right) p^\alpha p^\beta \nabla_{\langle \alpha} u_{\beta \rangle}. \quad (7)$$

where  $\tilde{f}_0 = 1 - af_0$  and the angular brackets denote traceless symmetric projection orthogonal to  $u^\mu$ . In the case of blast-wave model, the form of  $p^\alpha p^\beta \nabla_{\langle\alpha} u_{\beta\rangle}$  is explicitly calculated in Refs. [5, 8].

The anisotropic flow is defined as

$$v_n(p_T) \equiv \frac{\int_{-\pi}^{\pi} d\phi \cos[n(\phi - \Psi_n)] \frac{dN}{dy p_t dp_T d\phi}}{\int_{-\pi}^{\pi} d\phi \frac{dN}{dy p_t dp_T d\phi}}, \quad (8)$$

where  $\Psi_n$  is the event-plane angle for the  $n$ -th harmonic. Up to first order in viscosity [5],

$$v_n(p_T) = v_n^{(0)}(p_T) \left( 1 - \frac{\int d\phi \frac{dN^{(1)}}{dy p_t dp_T d\phi}}{\int d\phi \frac{dN^{(0)}}{dy p_t dp_T d\phi}} \right) + \frac{\int d\phi \cos[n(\phi - \Psi_n)] \frac{dN^{(1)}}{dy p_t dp_T d\phi}}{\int d\phi \frac{dN^{(0)}}{dy p_t dp_T d\phi}}, \quad (9)$$

where the superscript ‘(0)’ and ‘(1)’ denote quantities calculated using the ideal distribution function and first-order viscous correction, Eq. (7), respectively.

#### 4. Initial conditions and numerical results

We consider two identical relativistic nuclei with mass number  $A$  colliding with impact parameter  $b$ . The transverse expansion of the fireball is obtained by using the unperturbed radial velocity,


$$u^r \equiv \frac{dr}{d\tau} = u_0 \frac{r}{R} \quad \Rightarrow \quad \int_{r_0}^R \frac{dr}{r} = \int_0^{\tau_f} \frac{u_0}{R} d\tau \quad \Rightarrow \quad R = r_0 \exp\left(\frac{u_0 \tau_f}{R}\right), \quad (10)$$

where  $r_0 = \frac{1}{2} \left( b^2 - 2bR_0 \sqrt{2 + b/R_0} + 4R_0^2 \right)^{1/2}$  and  $R_0 = 1.25A^{1/3}$  fm is the radius of each colliding nuclei [8]. In order to determine the freeze-out times for non-central collisions, we employ the analytical result of Bjorken expansion  $\epsilon \propto \tau^{-4/3}$ . Assuming the initial thermalization time and the freeze-out energy density to be same for all collisions, we get


$$\tau_f = \tau_{f0} \left( \frac{\epsilon_i}{\epsilon_{i0}} \right)^{3/4}, \quad (11)$$

where  $\tau_{f0}$  is the freeze-out time for most central collisions which is fitted to match the transverse momentum distribution of particle multiplicities. The ratio  $\epsilon_i/\epsilon_{i0}$  is the initial central energy density for all centrality, scaled by its corresponding value in most central collisions. While the Bjorken estimate for the absolute values of the freeze-out times is rather crude, we have used the above equation to fix the freeze-out time for non-central collisions relative to that of the most central ones. In the present case, it seems to be a reliable approximation as is evident from Fig. 1. Therefore, in order to fit the transverse momentum spectra, the parameters that needs to be fixed within the viscous-blast wave model are the freeze-out temperature  $T_f$ , the freeze-out time for central collision  $\tau_{f0}$  and the maximum radial flow velocity  $u_0$ . An interplay between  $\alpha_0$  in Eq. (5) and  $\eta/s$  is important to reproduce the transverse momentum dependence of flow harmonics.

Figure 1 shows the transverse momentum distribution of pions, kaons, and protons spectra for Pb+Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV in (a): 0 – 5% and (b): 20 – 30% centrality. We observe that, for a freeze-out temperature of 120 MeV, the spectra for  $\pi^+$  and  $K^+$  from the viscous blast-wave model are in good overall agreement with the ALICE data [13]. On the other hand, the freeze-out temperature for protons is considered to be 135 MeV in order to



**Figure 1.** Transverse momentum distribution of  $\pi^+$ ,  $K^+$  and  $p$  multiplicities in Pb+Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV in two centrality ranges, (a): 0–5% and (b): 20–30%. The symbols represent ALICE data [13] at mid-rapidity and the lines correspond to viscous blast-wave results.



**Figure 2.** Transverse momentum dependence of the anisotropic flow coefficients  $v_n(p_T)$  of charged hadrons, for  $n = 2$  and  $3$ , calculated at various centralities in Pb+Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV in the viscous blast-wave model (lines) with  $\eta/s = 0.24$  as compared to the ATLAS data [14] (symbols).

fit the experimental data. Figure 2 shows our results for the  $v_n(p_T)$ , for various centralities, in comparison with the ATLAS data [14]. For  $\alpha_0 = 0.4$  and  $\eta/s = 0.24$ , we find fairly good agreement with the experimental data for elliptic flow,  $v_2(p_T)$ , and triangular flow,  $v_3(p_T)$ , at the LHC. We have used the root-mean square values of initial eccentricities,  $\varepsilon_n$ , obtained from the Monte-Carlo Glauber model [15].

*Acknowledgements:-* A.J. was supported by the Frankfurt Institute for Advanced Studies (FIAS), Germany. The work of V.K. was supported by the ExtreMe Matter Institute (EMMI) and by the Office of Nuclear Physics in the US Department of Energy’s Office of Science under Contract No. DE-AC02-05CH11231.

## References

- [1] U. Heinz and R. Snellings, *Ann. Rev. Nucl. Part. Sci.* **63**, 123 (2013).
- [2] E. Schnedermann, J. Sollfrank and U. W. Heinz, *Phys. Rev. C* **48**, 2462 (1993).
- [3] P. Huovinen, P. F. Kolb, U. W. Heinz, P. V. Ruuskanen and S. A. Voloshin, *Phys. Lett. B* **503**, 58 (2001).
- [4] C. Adler *et al.* [STAR Collaboration], *Phys. Rev. Lett.* **87**, 182301 (2001).
- [5] D. Teaney, *Phys. Rev. C* **68**, 034913 (2003).
- [6] X. Sun, H. Masui, A. M. Poskanzer and A. Schmah, *Phys. Rev. C* **91**, 024903 (2015).
- [7] D. Teaney, J. Lauret and E. V. Shuryak, *nucl-th/0110037*.
- [8] A. Jaiswal and V. Koch, *arXiv:1508.05878 [nucl-th]*; *EPJ Web Conf.* **120**, 06001 (2016).
- [9] P. Staig and E. Shuryak, *Phys. Rev. C* **84**, 034908 (2011).
- [10] E. Shuryak and I. Zahed, *Phys. Rev. C* **88**, 044915 (2013).
- [11] F. Cooper and G. Frye, *Phys. Rev. D* **10**, 186 (1974).
- [12] H. Grad, *Comm. Pure Appl. Math.* **2**, 331 (1949).
- [13] B. Abelev *et al.* [ALICE Collaboration], *Phys. Rev. C* **88**, 044910 (2013).
- [14] G. Aad *et al.* [ATLAS Collaboration], *Phys. Rev. C* **86**, 014907 (2012).
- [15] E. Retinskaya, M. Luzum and J. Y. Ollitrault, *Phys. Rev. C* **89**, 014902 (2014).