

The MAJORANA DEMONSTRATOR: A Search for Neutrinoless Double-beta Decay of Germanium-76

J F Wilkerson^{15,12,8}, E Aguayo¹⁰, F T Avignone III^{16,8}, H O Back^{7,12}, A S Barabash³, J R Beene⁸, M Bergevin⁶, F E Bertrand⁸, M Boswell⁵, V Brudanin⁴, M Busch^{2,12}, Y-D Chan⁶, C D Christofferson¹¹, J I Collar¹⁴, D C Combs^{7,12}, R J Cooper⁸, J A Detwiler⁶, P J Doe¹⁹, Yu Efremenko¹⁸, V Egorov⁴, H Ejiri⁹, S R Elliott⁵, J Esterline^{2,12}, J E Fast¹⁰, N Fields¹⁴, P Finnerty^{15,12}, F M Fraenkle^{15,12}, V M Gehman⁵, G K Giovanetti^{1,2}, M P Green^{15,12}, V E Guiseppe¹⁷, K Gusey⁴, A L Hallin¹³, R Hazama⁹, R Henning^{15,12}, E W Hoppe¹⁰, M Horton¹¹, S Howard¹¹, M A Howe^{15,12}, R A Johnson¹⁹, K J Keeter¹, C Keller¹⁷, M F Kidd⁵, A Knecht¹⁹, O Kochetov⁴, S I Konovalov³, R T Kouzes¹⁰, B D LaFerriere¹⁰, B H LaRoque⁵, J Leon¹⁹, L E Leviner^{7,12}, J C Loach⁶, S MacMullin^{15,12}, M G Marino¹⁹, R D Martin⁶, D-M Mei¹⁷, J H Merriman¹⁰, M L Miller¹⁹, L Mizouni^{16,10}, M Nomachi⁹, J L Orrell¹⁰, N R Overman¹⁰, D G Phillips II^{15,12}, A W P Poon⁶, G Perumpilly¹⁷, G Prior⁶, D C Radford⁸, K Rielage⁵, R G H Robertson¹⁹, M C Ronquest⁵, A G Schubert¹⁹, T Shima⁹, M Shirchenko⁴, K J Snavely^{15,12}, D Steele⁵, J Strain^{15,12}, K Thomas¹⁷, V Timkin⁴, W Tornow^{2,12}, I Vanyushin³, R L Varner⁸, K Vetter⁶, K Vorren^{5,12}, E Yakushev⁴, A R Young^{7,12}, C-H Yu⁸, V Yumatov³ and C Zhang¹⁷

¹ Department of Physics, Black Hills State University, Spearfish, SD, USA

² Department of Physics, Duke University, Durham, NC, USA

³ Institute for Theoretical and Experimental Physics, Moscow, Russia

⁴ Joint Institute for Nuclear Research, Dubna, Russia

⁵ Los Alamos National Laboratory, Los Alamos, NM, USA

⁶ Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

⁷ Department of Physics, North Carolina State University, Raleigh, NC, USA

⁸ Oak Ridge National Laboratory, Oak Ridge, TN, USA

⁹ Research Center for Nuclear Physics and Department of Physics, Osaka University, Ibaraki, Osaka, Japan

¹⁰ Pacific Northwest National Laboratory, Richland, WA, USA

¹¹ South Dakota School of Mines and Technology, Rapid City, SD, USA

¹² Triangle Universities Nuclear Laboratory, Durham, NC, USA

¹³ Centre for Particle Physics, University of Alberta, Edmonton, AB, Canada

¹⁴ Department of Physics, University of Chicago, Chicago, IL, USA

¹⁵ Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC, USA

¹⁶ Department of Physics and Astronomy, University of South Carolina, Columbia, SC, USA

¹⁷ Department of Physics, University of South Dakota, Vermillion, SD, USA

¹⁸ Department of Physics and Astronomy, University of Tennessee, Knoxville, TN, USA

¹⁹ Center for Experimental Nuclear Physics and Astrophysics, and Department of Physics, University of Washington, Seattle, WA, USA

E-mail: jfw@unc.edu

Abstract. The observation of neutrinoless double-beta decay would show that neutrinos are Majorana particles and provide information on neutrino mass. Attaining sensitivities for neutrino masses in the inverted hierarchy region requires large, tonne scale detectors with extremely low backgrounds, at the level of 10^{-3} counts $\text{keV}^{-1} \text{t}^{-1} \text{y}^{-1}$ or lower in the region of the signal. The MAJORANA collaboration is constructing the DEMONSTRATOR, an array consisting of 40 kg of p-type point contact germanium detectors, at least half of which will be enriched to 86% in ^{76}Ge . The primary aim is to show the feasibility for a future tonne scale measurement. With a sub-keV energy threshold, the array should also be able to search for light WIMP dark matter. This paper presents a brief update on the status of constructing the DEMONSTRATOR including an electroforming facility that is now operating underground at the Sanford Underground Research Facility.

1. The MAJORANA DEMONSTRATOR

Neutrinoless double-beta decay ($0\nu\beta\beta$) is the only viable technique to determine the Dirac-Majorana nature of the neutrino[1, 2]. The observation of $0\nu\beta\beta$ in two or more different isotopes would provide compelling evidence that neutrinos are Majorana particles and that lepton number is violated. Reaching the neutrino mass scale associated with the inverted mass hierarchy will require a half-life sensitivity on the order of 10^{27} y for ^{76}Ge , corresponding to a signal of a few counts or less per tonne-year in the $0\nu\beta\beta$ peak located at 2038-keV. To observe such a small signal, one will need to limit backgrounds in the region of interest to ~ 1 count per tonne-year (2.5×10^{-4} counts $\text{keV}^{-1} \text{t}^{-1} \text{y}^{-1}$). The MAJORANA collaboration is constructing the MAJORANA DEMONSTRATOR with the aim of achieving such low backgrounds while also testing the Klapdor-Kleingrothaus claim[3], of neutrinoless double-beta decay in ^{76}Ge .

The DEMONSTRATOR array will consist of an array of *natural* ^{76}Ge and 86% enriched ^{76}Ge high purity germanium (HPGe) detectors contained in an ultra-low background structure that maximizes the concentration of crystals while minimizing the amounts of structural materials[4, 5, 6]. Two ultra-pure electroformed copper cryostats are being fabricated, each of which will contain up to 35 close-packed P-type point-contact (P-PC) germanium detectors[7]. The cryostats are within a graded shield surrounded by structural and additional gamma and neutron shielding materials, which are themselves surrounded by an active veto system (Figure 1). The array will be located at the 4850 ft level of the Sanford Underground Research Facility (SURF) in Lead, SD.

2. Underground Electroforming

In typical materials uranium (U) and thorium (Th) decay-chain contaminants are found at levels of $\mu\text{g/g}$ to ng/g , which will produce unacceptable backgrounds in the DEMONSTRATOR. It has been shown that electroforming copper in a carefully controlled and clean environment allows one to produce ultra clean copper, with U and Th below the level of 10^{-12} g/g[8]. Copper has mechanical, thermal, and electrical properties that are suitable for the DEMONSTRATOR's cryostats, detector mounts, and inner shield.

We have completed and put into operation an advanced electroforming facility that is located underground near the Ross shaft at the 4850 ft level at SURF. At this depth, cosmogenic backgrounds are negligible compared to those that would be produced by cosmic ray bombardment of copper on the surface of the earth. The electroforming facility is divided into a class 1000 cleanroom and an anteroom, which contains data acquisition (DAQ) computers and a gowning area. Particle counts within the clean room are well below 1000 per cubic foot, with typical values well below 100 during normal operations. A slow controls and




Figure 1. Cross-sectional view of the MAJORANA DEMONSTRATOR, with one cryostat inserted and another being positioned for insertion into the shield. Cryostats are mounted on moveable transporters allowing assembly and testing before installation into the shield. The cryostat vacuum system is visible for the cryostat on the right. The graded shield consists of electroformed copper, commercial copper, lead, a radon exclusion volume, an active muon scintillator veto, and a polyethylene neutron moderator. For scale, the inner Cu volume where the cryostats are inserted is 20" high, 20" wide, and 30" in length.

environmental monitoring system continuously monitors a variety of parameters including temperature, humidity, conductivities of baths, voltages and currents of electroforming power supplies, leakage into electroforming baths secondary containment, nitrogen flow rate, particle count, radon level, and oxygen levels. This system allows remote access, monitoring and control. A similar electroforming facility has also been established in a shallow underground laboratory, 37 m.w.e., at Pacific Northwest National Laboratory (PNNL).

Ten large electroforming baths have been setup and are now operating at SURF, Figure 2), and another 6 baths are in operation at the PNNL facility. Each bath consists of a rectangular tank, 24"L x 24"W x 36"H, a cylindrical baffle, and cover, all contained in a secondary container. The baths will accommodate cylindrical stainless steel mandrels up to 13" diameter and 23" long. All components are constructed of high density polyethylene. Each bath and mandrel undergoes a rigorous cleaning and etching procedure prior to use. The baths contain an electrolyte solution composed of a simple acid sulfate system of ultra pure copper sulfate, sulfuric acid, and water. The electroformed copper that is being produced will be used to fabricate cryostats and internal components, detector mounting and string hardware, as well as the inner shield plates (typically 20" x 20" x 0.5") for the DEMONSTRATOR. Deposition rates are conservatively estimated at 0.002" per day.

3. Construction Status

Over 20 kg of *natural* Ge P-PC HPGe detectors have been acquired and characterized. For the enriched detectors, an initial order of 30 kg of GeO₂ enriched to 88% ⁷⁶Ge has been produced at the electrochemical plant in Zelonogorsk, Russia. The material was shipped to Oak Ridge, TN using ground-transportation in a steel-shielded container. It is now being stored in an underground location with 120 ft of rock overburden. (Additional material will be acquired in 2012.) The GeO₂ will be reduced to Ge and zone-refined into electronic grade Ge in partnership with Electrochemical Systems Incorporated, where a facility has been established to reduce, refine, and recycle the enriched material. A test run using natural GeO₂ has been successfully completed. A prototype cryostat is currently being fabricated using the same techniques planned for the ultra-clean cryostats. This cryostat will also be used to test the thermosyphon based



Figure 2. On the left is a view of the electroforming facility with 7 of the 10 baths visible in the photograph. On the right is a mandrel that has been lifted out of its bath for an initial inspection of the copper growth.

cooling system. It will be populated with 2-3 strings of natural P-PC detectors and mounted on a monolith transporter. A multi-stage glove box will be used to build the strings and insert them into the cryostats while within a Rn suppressed N_2 atmosphere. A dedicated underground laboratory for the DEMONSTRATOR at the Davis Campus at SURF has been excavated and outfitting is underway, with expected beneficial occupancy in the spring of 2012. The first underground cryostat, containing a mixed array of natural and enriched germanium, is expected to begin taking data in mid-2013.

Acknowledgments

We acknowledge support from the Office of Nuclear Physics in the DOE Office of Science under grant numbers DE-AC02-05CH11231, DE-FG02-97ER41041, DE-FG02-97ER41033, DE-FG02-97ER4104, DE-FG02-97ER41042, DE-SC0005054, DE-FG02-10ER41715, and DE-FG02-97ER41020. We acknowledge support from the Particle and Nuclear Astrophysics Program of the National Science Foundation through grant numbers PHY-0919270, PHY-1003940, 0855314, and 1003399. We gratefully acknowledge support from the Russian Federal Agency for Atomic Energy. We gratefully acknowledge the support of the U.S. Department of Energy through the LANL/LDRD Program. N. Fields is supported by the DOE/NNSA SSGF program. G. K. Giovanetti is supported by the Department of Energy Office of Science Graduate Fellowship Program (DOE SCGF), made possible in part by the American Recovery and Reinvestment Act of 2009, administered by ORISE-ORAU under contract no. DE-AC05-06OR23100.

References

- [1] Leslie Camilleri, Eligio Lisi, and John F. Wilkerson, *Annu. Rev. Nucl. Part. Sci.* **58** (2008) 343-69.
- [2] Frank T. Avignone, III, Steven R. Elliott, and Jonathan Engel, *Rev. Mod. Phys.* **80** (2008) 481-516.
- [3] H Klapdor-Kleingrothaus and I Krivosheina, *Mod. Phys. Lett.* **A21** (2006) 1547-66.
- [4] A.G. Schubert et al. 2011, To be published in AIP Conference Proceedings, 19th Particles & Nuclei International Conference (PANIC 2011), Massachusetts Institute of Technology, Cambridge, MA, USA, July 24-29, 2011; arXiv:1109.1567

- [5] D G Phillips II et al. 2011, To be published in Rutherford Centennial Conference on Nuclear Physics Proceedings. University of Manchester, Manchester UK August 8-12, 2011; arXiv:1111.5578
- [6] E. Aguayo et al. 2011, To be published in the Proceedings of the DPF-2011 Conference, Providence, RI, USA, August 8-13, 2011; arXiv:1109.6913
- [7] G.K Giovanetti, these proceedings.
- [8] E Hoppe et al. 2009 *J. Radioanal. Nucl. Chem.* **282** 315-320