
Article

Particles of Negative and Zero Energy in Black Holes and Cosmological Models

Andrey A. Grib and Yuri V. Pavlov

Special Issue

The Friedmann Cosmology: A Century Later

Edited by

Prof. Dr. Galina L. Klimchitskaya, Prof. Dr. Vladimir M. Mostepanenko and
Prof. Dr. Sergey V. Sushkov



Article

Particles of Negative and Zero Energy in Black Holes and Cosmological Models

Andrey A. Grib ^{1,2}  and Yuri V. Pavlov ^{3,*} 

¹ Theoretical Physics and Astronomy Department, The Herzen University, 48 Moika, St. Petersburg 191186, Russia; andrei_grib@mail.ru

² A. Friedmann Laboratory for Theoretical Physics, St. Petersburg 191186, Russia

³ Institute of Problems in Mechanical Engineering of Russian Academy of Sciences, 61 Bolshoy, V.O., St. Petersburg 199178, Russia

* Correspondence: yuri.pavlov@mail.ru

Abstract: Particles with negative energies are considered for three different cases: inside the horizon of a Schwarzschild black hole, Milne's coordinates in flat Minkowski space–time (Milne's universe using nonsynchronous coordinates) and in the cosmological Gödel model of the rotating universe. It is shown that, differently from the Gödel model with a nondiagonal term, where it occurs that negative energies are impossible, they are present in all other cases considered in the paper. Particles with zero energy are also possible in the first two cases.

Keywords: negative energy; zero energy; black hole; Milne universe; Gödel universe

1. Introduction



Citation: Grib, A.A.; Pavlov, Y.V. Particles of Negative and Zero Energy in Black Holes and Cosmological Models. *Universe* **2023**, *9*, 217. <https://doi.org/10.3390/universe9050217>

Academic Editors: Galina L. Klimchitskaya, Vladimir M. Mostepanenko and Sergey V. Sushkov

Received: 18 March 2023

Revised: 28 April 2023

Accepted: 28 April 2023

Published: 1 May 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

It appears that in order to have negative energy of the relativistic particle with nonzero mass, one must have a very strong external field leading to large potential energy, as is the case for rotating black holes. However, it is well known that the value of the energy depends on the choice of the reference frame and the time coordinate or Killing vector in case of conserved energy. It leads to the situation whereby states with negative energies in the relativistic case occur in the case of a rotating coordinate system outside the static limit [3], where an effect analogical to the Penrose effect is found to be observable [4], and in a nonsynchronous coordinate system in cosmology [5,6].

It seems from these examples that negative energies arise in the case of the existence of nondiagonal terms in a metrical tensor (gravymagnetism), but, in this paper, we show that in the Gödel Universe, in spite of the presence of such terms, negative energies are absent. The negative energies are present in cases of the movement inside of the horizon of the Schwarzschild black hole and in Milne's universe where nondiagonal terms are present in a nonsynchronous coordinate frame.

2. Negative Energy in Nonrotating Black Hole

A nonrotating black hole of mass M in Schwarzschild coordinates is described by the metric

$$ds^2 = \left(1 - \frac{r_g}{r}\right)c^2 dt^2 - \frac{dr^2}{1 - \frac{r_g}{r}} - r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (1)$$

where $r_g = 2GM/c^2$ is the gravitational radius of the black hole, G is the gravitational constant, and c is the light velocity. The geodesic complete space–time of the nonrotating black hole can be described in Kruskal–Szekeres coordinates, $\{u, v\} \in (-\infty, +\infty)$, which, in region $u > |v| \geq 0$, are connected with the Schwarzschild coordinate in $r > r_g$ in the following way:

$$u = \sqrt{\frac{r}{r_g} - 1} \exp\left(\frac{r}{2r_g}\right) \cosh \frac{ct}{2r_g}, \quad v = \sqrt{\frac{r}{r_g} - 1} \exp\left(\frac{r}{2r_g}\right) \sinh \frac{ct}{2r_g}. \quad (2)$$

For $r < r_g$ and $v > |u| \geq 0$, the transformation from the Schwarzschild coordinate into the Kruskal–Szekeres coordinates has the form

$$u = \sqrt{1 - \frac{r}{r_g}} \exp\left(\frac{r}{2r_g}\right) \sinh \frac{ct}{2r_g}, \quad v = \sqrt{1 - \frac{r}{r_g}} \exp\left(\frac{r}{2r_g}\right) \cosh \frac{ct}{2r_g}. \quad (3)$$

Schwarzschild coordinates are singular at $r = r_g$. Regarding their connection with Kruskal–Szekeres coordinates for other u, v , see [7], Section 31.5.

Any possible movement of physical bodies and particles must satisfy the condition $ds^2 > 0$ leading to

$$r > r_g \Rightarrow \left| \frac{dr}{dt} \right| \leq c \left(1 - \frac{r_g}{r}\right), \quad (4)$$

$$r < r_g \Rightarrow \left| \frac{dr}{dt} \right| \geq c \left(\frac{r_g}{r} - 1\right). \quad (5)$$

For $r < r_g$, the coordinate ct is space and r/c is the time coordinate.

Geodesic equations in Schwarzschild coordinates in the plane $\theta = 0$ are [8]

$$\frac{dt}{d\lambda} = \frac{r}{r - r_g} \frac{E}{c^2}, \quad \frac{d\varphi}{d\lambda} = \frac{J}{r^2}, \quad \left(\frac{dr}{d\lambda} \right)^2 = \frac{E^2}{c^2} + \frac{r_g - r}{r^3} J^2 + \frac{r_g - r}{r} m^2 c^2, \quad (6)$$

where E is the energy of a moving particle (measured by the static observer in r, θ, ϕ coordinates), J is the conserved projection of the particle's angular momentum on the axis orthogonal to the plane of motion, m is the particle mass, and λ is an affine parameter on the geodesic. For massive particles, $\lambda = \tau/m$, where τ is the proper time.

In an external region of the black hole ($r > r_g$) for any physical object, the time coordinate t is always increasing and so the energy E of the particle is positive (see (6)). Inside the horizon of the black ($r < r_g$), where t is space-like ($g_{tt} < 0$), one has movement as in increasing as in decreasing t . As is seen from the first formula in (6) for a particle moving inside the horizon in the direction of a decrease in the coordinate t , the energy E of the particle will be positive, while, for increasing coordinate t , the energy E is negative. For constant t inside the black hole, $E = 0$ due to formula (6).

Surely, t inside of the black hole is space-like and E is proportional to the t -component of the momentum. Inside a black hole, one can use other reference frames and corresponding energies [9]. However, for the observer outside the black hole, the conserved E along all trajectories of the free fall is equal (see formula (88.9) in [10]) to

$$E = mc^2 \sqrt{\left(1 - \frac{r_g}{r}\right) / \left(1 - \frac{\mathbf{v}^2}{c^2}\right)}, \quad (7)$$

where \mathbf{v} is the velocity measured by the observer at rest in the Schwarzschild coordinates. Thus, we can call E inside the black hole, following [7], the “energy at infinity”. For a discussion of other ways to determine the energy within the horizon and the movement of particles there, see, for example, the articles [9,11–13].

In Figure 1, the trajectories for radial movement with positive, zero and negative energies in Kruskal–Szekeres coordinates are represented by red, green and blue lines.




Figure 1. Trajectories of particles with positive ($B^+H^+F^+$), zero ($M^\pm ON^\pm$) and negative ($B^-H^-F^-$) energy, $t_1^\pm = \mp 0.5r_g/c$. On the left, one can see it falling from the rest at $|E| = 0.5mc^2$; on the right, $|E| = mc^2$ with the corresponding initial velocity from the point $r = 1.15r_g$. On lines ($M^\pm ON^\pm$), the coordinate $t = \pm r_g/c$.

As one can see from (2) and (3), the coordinate lines of constant t in Kruskal–Szekeres coordinates are straight lines through the origin of the coordinates. In region II, coordinate t decreases when moving from H^+ to F^+ (positive E) and increases when moving from H^- to F^- (negative value of E). Direct lines ($M^\pm ON^\pm$) correspond to constant $t = \pm r_g/c$ and therefore $E = 0$.

Let us consider the problem of the back influence of falling particles on the metric of the black hole space–time. For macroscopic bodies with 4-velocity (u^i) , with the energy density ε and pressure p in space–time with metric g_{ik} , the energy–momentum density tensor is [10]

$$T_{ik} = (\varepsilon + p)u_i u_k - p g_{ik}, \quad (8)$$

$i, k = 0, 1, 2, 3$. The trace of the energy–momentum tensor

$$T_i^i = \varepsilon - 3p \quad (9)$$

is invariant and it will be negative for $\varepsilon - 3p < 0$ —in particular, for dust-like matter ($p = 0$) with negative energy $\varepsilon < 0$. The back influence of falling particles with negative energy will be determined by the energy–momentum tensor in the right-hand side of the Einstein equations. The notion of the existence of particles with negative energies as it is known was used by S. Hawking to predict the Hawking effect for black holes [14].

For a discussion of other ways to determine the energy within the horizon and the movement of particles there, see, for example, works

3. Negative and Zero Energies in Flat Space–Time

The geodesic line equations can be obtained for space–time with metric g_{ik} from the Lagrangian

$$L = \frac{g_{ik}}{2} \frac{dx^i}{d\lambda} \frac{dx^k}{d\lambda}, \quad (10)$$

where λ is the affine parameter on the geodesic [8]. The energy of the particle E is equal to the zero covariant component of the momentum (p_i) multiplied on the light velocity

$$p_i = \frac{\partial L}{\partial \left(\frac{dx^i}{d\lambda} \right)} = g_{ik} \frac{dx^k}{d\lambda}, \quad (11)$$

$$E = cp_0 = c g_{0k} \frac{dx^k}{d\lambda}. \quad (12)$$

Defining the affine parameter for the massive particle as $\lambda = \tau/m$, where τ is the proper time of the moving particle, one obtains

$$p_i p^i = m^2 c^2 \quad (13)$$

and the energy of the particle is

$$E = mc g_{0k} \frac{dx^k}{d\tau}. \quad (14)$$

Using notation $(\zeta^i) = (1, 0, 0, 0)$ for the translation in the time coordinate generator, one can write (12) for the energy of the particle as

$$E = c(p, \zeta). \quad (15)$$

If the metric components do not depend on the time coordinate x^0 , then ζ is the time-like Killing vector and the energy E is conserved on the geodesic. For time-like vector ζ and a massive particle, one has [15]

$$\sqrt{(\zeta, \zeta)} \leq \frac{E}{mc^2} < +\infty \quad (16)$$

and the energy (15) is positive. For space-like vector ζ , as it takes place in the ergosphere of a rotating black hole, the arbitrary positive and negative values are possible (see [15], p. 325).

Note that in spite of the invariance of the scalar product (15), the value (14) of the energy depends on the choice of the reference frame. This occurs due to the fact that by changing the reference frame in which the physical measurements are made, the observer is changing vector ζ . The analysis of the situation in a rotating coordinate system in flat space-time is provided in [3].

In Minkowski space-time in the Galilean coordinate system or any other coordinate system with $g_{00} = 1, g_{0\alpha} = 0, (\alpha = 1, 2, 3)$, the energy (12) is

$$E = c^2 \frac{dt}{d\lambda} \quad (17)$$

and it is always positive in movement “forward” in time because in the future light cone, one has $dt/d\lambda > 0$.

Consider the coordinate system in which the metric of flat space-time has the form of the metric of the expanding homogeneous isotropic Universe—the Milne universe [16]:

$$ds^2 = c^2 dt^2 - c^2 t^2 (d\chi^2 + \sinh^2 \chi d\Omega^2), \quad (18)$$

where $d\Omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2$, and coordinate χ is changing from 0 to $+\infty$. In new coordinates

$$T = t \cosh \chi, \quad r = ct \sinh \chi, \quad cT > r > 0 \quad (19)$$

the interval (18) takes the form of a Minkowski interval

$$ds^2 = c^2 dT^2 - dr^2 - r^2 d\Omega^2. \quad (20)$$

This space–time with coordinate $t \geq 0, \chi \geq 0$ corresponds to the future cone in coordinates cT, r .

The radial distance between points $\chi = 0$ and χ in metric (18) is $D = ct\chi$. Taking D as the radial coordinate [17], one obtains the interval as

$$ds^2 = \left(1 - \frac{D^2}{c^2 t^2}\right) c^2 dt^2 + 2 \frac{D}{t} dt dD - dD^2 - c^2 t^2 \sinh^2\left(\frac{D}{ct}\right) d\Omega^2. \quad (21)$$

From the condition $ds^2 \geq 0$, one obtains that if D is larger than $D_s = ct$, no physical object can be at rest in coordinates t, D, θ, ϕ . The value D_s corresponds to $\chi = 1$ and it plays the role of the static limit for the rotating black hole in Boyer–Lindquist [18] coordinates.

The energy of the particle with mass m in coordinates t, D, θ, ϕ is

$$E = c g_{0k} \frac{dx^k}{d\lambda} = mc^2 \frac{dt}{d\tau} \left(1 - \frac{D^2}{c^2 t^2} + \frac{D}{c^2 t} \frac{dD}{dt}\right) = mc^2 \frac{dt}{d\tau} \left(1 + \chi t \frac{d\chi}{dt}\right). \quad (22)$$

From (18), one obtains, for any physical object, the inequality

$$t \left| \frac{d\chi}{dt} \right| \leq 1. \quad (23)$$

Thus, a particle can have negative energy only for $\chi > 1$, i.e., out of the static limit, if

$$\frac{d\chi}{dt} < -\frac{1}{\chi t}. \quad (24)$$

Note that the components of metric (21) depend on time and the energy (22) in general is not conserved on the geodesics. If the energy is zero, then the particle is moving noninertial according to the law

$$\frac{d\chi}{dt} = -\frac{1}{\chi t} \Leftrightarrow \chi = \sqrt{\chi_0^2 - 2 \log(t/t_0)}, \quad t \in [t_0, t_0 \exp((\chi_0^2 - 1)/2)]. \quad (25)$$

The trajectory of such movement for case $\chi_0 = 2, t_0 = 0.11$ is represented by the curve in Figure 2 in coordinates T, r (see (19)).




Figure 2. Possible region of movement of particle with negative and zero energies in the reference frame t, D in flat coordinate T, r .

In the event that the inertial movement trajectory in these coordinates is a straight line, the possible region of movement of particles with negative and zero energies in the reference frame t, D is defined in the coordinate T, r by conditions $1 \leq cT/r \leq \coth 1 \approx 1.313$.

Velocities of movement in coordinates T, r and t, χ satisfy condition [19]

$$t \frac{d\chi}{dt} = \frac{\frac{dr}{dT} - c \tanh \chi}{c - \frac{dr}{dT} \tanh \chi}. \quad (26)$$

Thus, for

$$\chi \tanh \chi \geq 1 \quad (27)$$

particles at rest in inertial frame T, r will have negative energies in the frame t, D . This region can be seen in Figure 2 as the region above the blue line in the red district. Zero energy of the particle at rest in T, r coordinates is possible only if the blue line is defined by the root of equation $\chi \tanh \chi = 1$, i.e., $\chi \approx 1.1997$.

Thus, one can see that for a specific choice of coordinates, one can obtain negative and zero energies for particles at rest in an inertial frame.

Note that for small distances $(D/(ct)) = \chi \ll 1$, the metric (21) becomes the metric of a comoving spherical coordinate system of Minkowski space-time

$$ds^2 = c^2 dt^2 - dD^2 - D^2 d\Omega^2, \quad (28)$$

and the energy (22) will be equal to the usual energy in the inertial system of flat space-time

$$E_u = mc^2 \frac{dt}{d\tau} \approx mc^2 \frac{dT}{d\tau}, \quad (29)$$

because, for $\chi \ll 1$, one has $t \approx T$.

The decay of the body on two bodies, one with negative energy and the other with the positive energy being larger than the energy of the initial decaying one, corresponds to the Penrose process. This process occurs outside the static limit on distance $D > ct$. However, later, these two products of the decay move inside the static limit, and, during flight in the direction of the origin, where the metric is that of Minkowski space, they change their energies in such a manner that the result will be the same as in the inertial frame. In fact, due to (22)

$$E = E_u + mc^2 \chi t \frac{d\chi}{d\tau}. \quad (30)$$

Here, E_u is the energy in the reference frame t, χ , such that $g_{0i} = 0, i \neq 0$, and g_{00} does not depend on time. Thus, E_u is conserved. At the point of decay, both energies E and E_u are conserved. When body 2 with the positive energy arrives at the coordinate origin $\chi = 0$, its energy E (30) will be equal to E_u and no growth in the energy will be observed.

Body 1, with the negative energy E due to (22) after decay, will have a negative value of velocity $d\chi/dt$ larger (in absolute value) than that of body 2. This means that it will arrive at the origin before the arrival of body 2. Its energy in the origin of the coordinate frame will be also positive and the full energy of 1 and 2 will be equal to that of the decaying body. Thus, at the origin, one will not observe any effect that makes this situation similar to the situation for Kerr's black hole.

In fact, for rotating black hole [1], as in the case of rotating coordinate system [4], the energy is conserved. In this case, when body 2 travels out of the ergosphere, far from it, body 1, with negative energy, moves further inside the horizon of the black hole or goes to infinity in case of rotating coordinates in Minkowski space. Thus, the energy of the body with positive energy, due to the conservation of the energy, will be always larger than that of the initial decaying body.

Note that the existence of states with negative energies for Milne's metric leads to an effect similar to the Hawking effect [14] for the Schwarzschild metric. Particle creation in quantum theory will occur and the detector of particles checks them (see Section 5.3 in [20]).

This will be the creation of virtual particles (see Section 9.8 in [21]) so no change in the metric due to them can be observed.

4. Negative Energy in Gödel Universe

The metric of the Gödel cosmological model of the rotating Universe proposed in 1949 (see [22] or [23]) can be written as

$$ds^2 = c^2 dt^2 - dx_1^2 + \frac{\exp(2\sqrt{2}\omega x_1/c)}{2} dx_2^2 + 2 \exp(\sqrt{2}\omega x_1/c) c dt dx_2 - dx_3^2, \quad (31)$$

where ω is constant. Such a metric is the exact solution of Einstein's equation with background matter as an ideal liquid without pressure and negative cosmological constant Λ

$$R_{ik} - \frac{1}{2} R g_{ik} + \Lambda g_{ik} = -8\pi \frac{G}{c^4} T_{ik}, \quad (32)$$

where

$$-\Lambda = \left(\frac{\omega}{c}\right)^2 = 4\pi \frac{G}{c^2} \rho, \quad T_{ik} = \rho c^2 u_i u_k, \quad (33)$$

$u^i = \delta_0^i$. Here, ω has the sense of the angular velocity of rotation of the vector of fluid of the background matter u^i .

Taking, instead of t, x_1, x_2 , new coordinates t', r, ϕ ,

$$\exp(\sqrt{2}\omega x_1/c) = \cosh 2r + \cos \phi \sinh 2r, \quad (34)$$

$$\omega x_2 \exp(\sqrt{2}\omega x_1/c) = \sin \phi \sinh 2r, \quad (35)$$

$$\tan \frac{1}{2}(\phi + \omega t - \sqrt{2}t') = \exp(-2r) \tan \frac{\phi}{2}, \quad (36)$$

one writes the interval (31) in the form [22,24]

$$ds^2 = 2\left(\frac{c}{\omega}\right)^2 \left(dt'^2 - dr^2 + (\sinh^4 r - \sinh^2 r) d\phi^2 + 2\sqrt{2} \sinh^2 r d\phi dt' \right) - dx_3^2, \quad (37)$$

where $-\infty < t' < \infty, 0 \leq r < \infty, 0 \leq \phi < 2\pi$ and identifying $\phi = 0$ and $\phi = 2\pi$.

Now, consider the general case of space-time t', r, ϕ, z with the interval

$$ds^2 = a^2 \left[(dt' + \Phi(r) d\phi)^2 - dr^2 - dz^2 - R^2(r) d\phi^2 \right], \quad (38)$$

where a is constant, $-\infty < t < \infty, 0 \leq r < \infty, -\infty < z < \infty, 0 \leq \phi \leq 2\pi$, and identifying $\phi = 0$ and $\phi = 2\pi$. Let us say $\Phi(r) > 0$ and $R(r) > 0$ for $r > 0$. For Gödel Universe $a = \sqrt{2}c/\omega, z = x_3/a$ and

$$\Phi(r) = \sqrt{2} \sinh^2 r, \quad R(r) = \sinh r \cosh r. \quad (39)$$

The metrical tensor is

$$(g_{ik}) = a^2 \begin{pmatrix} 1 & \Phi & 0 & 0 \\ \Phi & \Phi^2 - R^2 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad (40)$$

$$(g^{ik}) = \frac{1}{a^2 R^2(r)} \begin{pmatrix} R^2 - \Phi^2 & \Phi & 0 & 0 \\ \Phi & -1 & 0 & 0 \\ 0 & 0 & -R^2 & 0 \\ 0 & 0 & 0 & -R^2 \end{pmatrix}, \quad (41)$$

where indexes $i, k = 0, 1, 2, 3$ correspond to t', ϕ, r, z . Note that for any $r > 0$, the metrical tensor is not degenerate $\det(g_{ik}) = -a^8 R^2(r) < 0$. The degeneration for $r = 0$ in the Gödel Universe is coordinate degeneracy. The eigenvalues of the g_{ik} tensor are

$$\begin{aligned} \lambda_{1,2} &= \frac{a^2}{2} \left(\Phi^2 - R^2 + 1 \pm \sqrt{(\Phi^2 - R^2 + 1)^2 + 4R^2} \right), \\ \lambda_{3,4} &= -a^2. \end{aligned} \quad (42)$$

For $r > 0$, one has

$$\lambda_1 \geq a^2, \quad 0 > \lambda_2 \geq -a^2 R^2. \quad (43)$$

Note that although $g_{\phi\phi}$ is positive for $\Phi(r) > R(r)$, the signature of g_{ik} for all $r > 0$ is the standard $(+, -, -, -)$.

Possible movement of particles is defined by $ds^2 \geq 0$, so, for the interval (38), one has

$$dt'^2 + (\Phi^2(r) - R^2(r)) d\phi^2 + 2\Phi(r) d\phi dt' - dr^2 - dz^2 \geq 0. \quad (44)$$

It is important that for any coordinate system with interval (38), the physical body for any values of r, ϕ, z can be at rest, i.e., there is no static limit. However, in (38), there is nondiagonal term $dt' d\phi$ as in Kerr's metric. However, differently from the case of rotating coordinate system [3], there is the possibility of a change in the sign before $d\phi^2$.

From (44), one obtains

$$\left(\frac{dt'}{d\phi} \right)^2 + 2\Phi(r) \frac{dt'}{d\phi} + \Phi^2(r) - R^2(r) \geq 0. \quad (45)$$

The solution of this inequality is the union of two intervals

$$\frac{dt'}{d\phi} \in (-\infty, -(\Phi(r) + R(r))] \cup [R(r) - \Phi(r), +\infty). \quad (46)$$

Considering cases of different signs of $d\phi$, one obtains the following sets of solutions of (45):

$$d\phi \geq 0 \Rightarrow dt' \geq (R - \Phi)d\phi \vee dt' \leq -(R + \Phi)d\phi, \quad (47)$$

$$d\phi \leq 0 \Rightarrow dt' \geq -(R + \Phi)d\phi \vee dt' \leq (R - \Phi)d\phi. \quad (48)$$

These sets define light "cones" of the future and past for the metric (38). The form of these cones in cases $\Phi \ll R$, $\Phi = R$ and $\Phi > R$ is shown in Figure 3 for the Gödel Universe with

$$\Phi(r) > R(r) \Leftrightarrow r > r_0 = \log(1 + \sqrt{2}). \quad (49)$$

Let us find limitations on possible values of the energy of particles moving in such a Universe. The coordinate t' is dimensionless, so the "physical energy" of the particle is expressed through the time component of the momentum as

$$E = p_0 \frac{c}{a} = g_{0k} \frac{c}{a} \frac{dx^k}{d\lambda}. \quad (50)$$

For the frame with coordinates (38), the covariant t', ϕ, z components are conserved, because the components of the metric depend only on r . Thus, the conserved energy on the geodesic for the interval (38) is

$$E = ca \left(\frac{dt'}{d\lambda} + \Phi(r) \frac{d\phi}{d\lambda} \right). \quad (51)$$

From (47) and (48), for the case of movement “forward” in time, i.e., in the future light cone, one obtains

$$dt' + \Phi d\phi \geq R |d\phi|, \quad (52)$$

so

$$E \geq caR \frac{|d\phi|}{d\lambda}. \quad (53)$$

It means that for a particle moving in the future cone in the Gödel Universe, the energy is not negative.

For movement “back in time”, the energy is limited from above by

$$E \leq -caR \frac{|d\phi|}{d\lambda} \quad (54)$$

and so it can be less than or equal to zero. However, such movement physically is inconsistent. The “time machine” effect in the Gödel Universe corresponds to continuous movement in the future cone. Thus, for $r > r_0$, where $\Phi(r) > R(r)$ closed loops (they are not geodesic lines) $r = \text{const}, z = \text{const}$, called Gödel cycles, from $\phi = 0$ to $\phi = 2\pi$, are closed time-like curves [24]. Particles moving along such a cycle are moving “forward” in time but, due to the identification of values ϕ different on 2π , it occurs in the past after the whole cycle. Its energy is positive due to (53). Such a “time machine” is different from that moving in the past by the sign of the particle energy.




Figure 3. Light “cones” of future (blue color) and past (yellow color) for Gödel Universe in coordinates t', ϕ for cases $r = 10^{-3}$ (left), $r = r_0$ (center) and $r = 2r_0$ (right).

5. Conclusions

Three different cases are investigated concerning the possibility of the existence of particles with negative and zero energies.

1. Schwarzschild black hole of the mass M . Trajectories of particles with negative and zero energies exist inside the horizon of the black hole, which can be shown in Kruskal–Szekeres coordinates.
2. Flat space–time in Milne’s coordinates. Here, one also has the possibility of the existence of particles with negative and zero energies if a nonsynchronous system of coordinates is used.
3. Gödel cosmological model with rotation. Here, we proved that, in this model, in Gödel’s coordinates, particles with negative and zero energies do not exist.

As for observations of the discussed effects, one can say that even the well-established Penrose and Hawking effects are still not observed. However, we hope that in the future

development of observational astrophysics, one will see the consequences of the existence of negative and zero energies.

Author Contributions: A.A.G. and Y.V.P. have contributed equally to all parts of this work. All authors have read and agreed to the published version of the manuscript.

Funding: This research is supported by the Russian Science Foundation (Grant No. 22-22-00112).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Penrose, R. Gravitational collapse: The role of general relativity. *Riv. Nuovo C.* **1969**, *1*, 252–276.
2. Penrose, R.; Floyd, R.M. Extraction of rotational energy from a black hole. *Nat. Phys. Sci.* **1971**, *229*, 177–179. [\[CrossRef\]](#)
3. Grib, A.A.; Pavlov, Y.V. Comparison of particle properties in Kerr metric and in rotating coordinates. *Gen. Relativ. Gravit.* **2017**, *49*, 78. [\[CrossRef\]](#)
4. Grib, A.A.; Pavlov, Y.V. Static limit and Penrose effect in rotating reference frames. *Theor. Math. Phys.* **2019**, *200*, 1117–1125. [\[CrossRef\]](#)
5. Grib, A.A.; Pavlov, Y.V. Particle properties outside of the static limit in cosmology. *Int. J. Mod. Phys. A* **2020**, *35*, 2040044. [\[CrossRef\]](#)
6. Grib, A.A.; Pavlov, Y.V. Particles with negative energies in nonrelativistic and relativistic cases. *Symmetry* **2020**, *12*, 528. [\[CrossRef\]](#)
7. Misner, C.W.; Thorne, K.S.; Wheeler, J.A. *Gravitation*; Freeman: San Francisco, CA, USA, 1973.
8. Chandrasekhar, S. *The Mathematical Theory of Black Holes*; Oxford University Press: Oxford, UK, 1983.
9. Grib, A.A.; Pavlov, Y.V. Particles with negative energies in black holes. *Int. J. Mod. Phys. D* **2011**, *20*, 675–684. [\[CrossRef\]](#)
10. Landau, L.D.; Lifshitz, E.M. *The Classical Theory of Fields*; Pergamon Press: Oxford, UK, 1983.
11. Toporensky, A.V.; Zaslavskii, O.B. Zero-momentum trajectories inside a black hole and high energy particle collisions. *JCAP* **2019**, *12*, 063. [\[CrossRef\]](#)
12. Augousti, A.T.; Gusin, P.; Kuśmierz, B.; Masajada, J.; Radosz, A. On the speed of a test particle inside the Schwarzschild event horizon and other kinds of black holes. *Gen. Relativ. Gravit.* **2018**, *50*, 131. [\[CrossRef\]](#)
13. Radosz, A.; Gusin, P.; Augousti, A.T.; Formalik, F. Inside spherically symmetric black holes or how a uniformly accelerated particle may slow down. *Eur. Phys. J. C* **2019**, *79*, 876. [\[CrossRef\]](#)
14. Hawking, S.W. Particle creation by black holes. *Commun. Math. Phys.* **1975**, *43*, 199–220. [\[CrossRef\]](#)
15. Lightman, A.P.; Press, W.H.; Price, R.H.; Teukolsky, S.A. *Problem Book in Relativity and Gravitation*; Princeton University Press: Princeton, NJ, USA, 1979.
16. Milne, E.A. *Relativity, Gravitation and World-Structure*; Clarendon Press: Oxford, UK, 1935.
17. Ellis, G.F.R.; Rothman, T. Lost horizons. *Am. J. Phys.* **1993**, *61*, 883–893. [\[CrossRef\]](#)
18. Boyer, R.H.; Lindquist, R.W. Maximal analytic extension of the Kerr metric. *J. Math. Phys.* **1967**, *8*, 265–281. [\[CrossRef\]](#)
19. Grib, A.A.; Pavlov, Y.V. Some properties of energies in nonsynchronous reference frames in cosmology. *J. Phys. Conf. Ser.* **2021**, *2081*, 012004. [\[CrossRef\]](#)
20. Birrell, N.D.; Davies, P.C.W. *Quantum Fields in Curved Space*; Cambridge University Press: Cambridge, UK, 1982.
21. Grib, A.A.; Mamayev, S.G.; Mostepanenko, V.M. *Vacuum Quantum Effects in Strong Fields*; Friedmann Laboratory Publishing: St. Petersburg, Russia, 1994.
22. Gödel, K. An example of a new type of cosmological solutions of Einstein's field equations of gravitation. *Rev. Mod. Phys.* **1949**, *21*, 447–450. [\[CrossRef\]](#)
23. Gödel, K. *Collected Works. Volume II. Publications 1938–1974: Editor-in-chief Solomon Feferman*; Oxford University Press: Oxford, UK, 1990; Volume 2, p. 190.
24. Hawking, S.W.; Ellis, G.F.R. *The Large Scale Structure of Space-Time*; Cambridge University Press: Cambridge, UK, 1973.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.