



mathematics



Article

Scalar Field Cosmology from a Modified Poisson Algebra

Genly Leon, Alfredo D. Millano and Andronikos Paliathanasis

Special Issue

Noncommutativity, Gravity and Cosmology

Edited by




Dr. Seyed Meraj Mousavi Rasouli and Prof. Dr. Paulo Vargas Moniz



<https://doi.org/10.3390/math11010120>

Article

Scalar Field Cosmology from a Modified Poisson Algebra

Genly Leon ^{1,2,*} , Alfredo D. Millano ¹  and Andronikos Paliathanasis ^{1,2} 

¹ Departamento de Matemáticas, Universidad Católica del Norte, Avda. Angamos 0610, Casilla 1280, Antofagasta 1240000, Chile

² Institute of Systems Science, Durban University of Technology, Durban 4000, South Africa

* Correspondence: genly.leon@ucn.cl

Abstract: We investigate the phase space of a scalar field theory obtained by minisuperspace deformation. We consider quintessence or phantom scalar fields in the action that arises from minisuperspace deformation on the Einstein–Hilbert action. We use a modified Poisson algebra where Poisson brackets are the α -deformed ones and are related to the Moyal–Weyl star product. We discuss early- and late-time attractors and reconstruct the cosmological evolution. We show that the model can have the Λ CDM model as a future attractor if we initially consider a massless scalar field without a cosmological constant term.

Keywords: cosmology; scalar field; modified Poisson algebra; dynamical analysis

PACS: 98.80.-k; 95.35.+d; 95.36.+x

MSC: 83D05; 17B63; 37J25; 37J39



Citation: Leon, G.; Millano, A.D.; Paliathanasis, A. Scalar Field Cosmology from a Modified Poisson Algebra. *Mathematics* **2023**, *11*, 120. <https://doi.org/10.3390/math11010120>

Academic Editors: Seyed Rasouli and Paulo Vargas Moniz

Received: 6 December 2022

Revised: 22 December 2022

Accepted: 24 December 2022

Published: 27 December 2022



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Cosmological observations indicate that the universe has gone through two acceleration phases [1–4], an early acceleration phase known as inflation and the present acceleration phase. The source of the cosmic acceleration is unknown. In the context of General Relativity, cosmic acceleration occurs when the cosmic fluid is dominated by a vacuum-like source known as dark energy (DE) with the property of having a negative value of the equation of state (EoS) parameter.

The cosmological constant Λ leads to the Λ -cosmology being indeed the simplest candidate for DE; however, it suffers from two problems, the fine-tuning and the coincidence problems [5,6]. Furthermore, the detailed analysis of the recent cosmological observations shows that the Λ -cosmology cannot solve tensions arising from the statistical analysis of the data, such as the H_0 -tension [7]. There are various DE alternatives to the cosmological constant, which have been proposed to overpass the above-mentioned problems; see, for instance [8–17] and the references therein.

Scalar fields play a significant role in the description of cosmic acceleration. Indeed, introducing a scalar field in the field equations provides new degrees of freedom in the gravitational dynamics that provide acceleration effects. The most straightforward mechanism for describing the early acceleration phase of the universe, that is of the inflationary epoch, is that of the inflaton field [18–25]. During inflation [26,27], the scalar field dominates the cosmological dynamics and provides the antigravity effects. Similarly, for the description of the late-time acceleration [28], a tracker scalar field can be introduced [12], which roles down the potential energy $V(\phi)$ to have DE effects [29–33]. Another novelty of the scalar fields is that they can reproduce various DE alternatives such as the Chaplygin gas and others [34,35].

In quintessence scalar field cosmology [8], the EoS parameter of the scalar field is constrained to the range $|w_\phi| \leq 1$, where $w_\phi = 1$ corresponds to a stiff fluid, where only

the kinetic part of the scalar field dominates, while the limit $w_\phi = -1$ corresponds to the case where only the scalar field potential dominates, leading to the Λ -cosmology. Recall that acceleration occurs when $-1 \leq w_{eff} < -1/3$. There is a family of scalar field models, known as phantom scalar fields, where w_ϕ can cross the limit -1 and take smaller values, which is possible, for example, when there exists a negative kinetic energy [36–39].

During the very early stages of the universe, we expect that quantum effects play an important role in cosmic evolution. Until now, there is not a unique theory of quantum gravity; that is why various approaches have been considered in the literature by various groups [40–48]. String theory, double-special relativity, and the generalized uncertainty principle require the existence of a minimum length scale of the order of the Planck length l_{pl} [49–56]. As a result of the modification of the Heisenberg uncertainty in the latter approaches, a deformation parameter is introduced, which leads to the deformation of the coordinate representation of the operators of the momentum position, that is to a deformation of the Poisson algebra [57].

Noncommutative theories, quantum cosmology, quantum deformation, deformed phase space, Brans–Dicke theory, and noncommutative minisuperspace, as alternatives of the cosmological constant, have been treated, for example, in [58–64] and the references therein.

In [65], the phase space for the cosmological dynamics in quintessence cosmology was modified by a deformed Poisson algebra among the coordinates and the canonical momenta. The main result was that the deformation parameter is related to the accelerating scale factor provided by the deformed Poisson algebra in the absence of a cosmological constant. A similar result was determined recently in [66] and the case of a phantom scalar field.

The Moyal–Weyl star product provides a simple prescription for constructing noncommutative field theories on the noncommutative manifold [65] with $[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}$. One replaces all the pointwise products in ordinary field theory with one of the star products. For example, the noncommutative action for a real massless scalar field Φ in four dimensions is

$$S_\phi = \frac{1}{2} \int d^4x D^\mu \Phi \star_\alpha D_\mu \Phi, \quad (1)$$

where the ordinary derivative ∂_μ appearing in the commutative scalar field action is replaced by the noncommutative covariant derivative D_μ and the action is invariant under the noncommutative gauge transformation.

In this paper, we are interested in studying the effects of the deformed Poisson algebra in the cosmological evolution. Specifically, we perform a detailed phase space analysis to investigate the existence of equilibrium points and reconstruct the cosmological parameters' evolution. Such an analysis provides important information about the theory's viability and can give us important results for the nature of the deformation parameter. For this analysis, one can introduce auxiliary variables, which transform the cosmological equations into an autonomous dynamical system [67–81]. Hence, we obtain a system of the form $\mathbf{X}' = \mathbf{f}(\mathbf{X})$, where \mathbf{X} is the column vector of the auxiliary variables and $\mathbf{f}(\mathbf{X})$ is an autonomous vector field. The derivative is with respect to a logarithmic time scale. The stability analysis comprises several steps. First, the critical points \mathbf{X}_c are extracted under the requirement of $\mathbf{X}' = \mathbf{0}$. Then, one considers linear perturbations around \mathbf{X}_c as $\mathbf{X} = \mathbf{X}_c + \mathbf{U}$, with \mathbf{U} the column vector of the auxiliary variable's perturbations. Therefore, up to first order, we obtain $\mathbf{U}' = \mathbf{\Xi} \cdot \mathbf{U}$, where the matrix $\mathbf{\Xi}$ contains the coefficients of the perturbed equations. Finally, the type stability of each hyperbolic critical point is determined by the eigenvalues of $\mathbf{\Xi}$. That is, the point is stable (unstable) if the real parts of the eigenvalues are negative (positive) or a saddle point if the eigenvalues have real parts with different signs.

The structure of the paper is as follows. In Section 2, we introduce the modified Poisson algebra. In Section 3, we derive the modified field equations in the case of scalar field cosmology in an isotropic and homogeneous spatially flat universe. Sections 4 and 5 include the main results of this study, where we present the detailed analysis of the phase

space for the modified field equations. Finally, in Section 6, we summarize our results and conclude.

2. Modified Poisson Algebra

We consider the modified Poisson algebra [65]:

$$\{x_i, x_j\}_\alpha = \theta_{ij}, \tag{2}$$

$$\{p_i, p_j\}_\alpha = \beta_{ij}, \tag{3}$$

$$\{x_i, p_j\}_\alpha = \delta_{ij} + \sigma_{ij}, \tag{4}$$

where the Moyal–Weyl brackets are defined through the relation:

$$\{f, g\}_\alpha = f \star_\alpha g - g \star_\alpha f \tag{5}$$

in which the product between f and g is substituted by the Moyal–Weyl star product:

$$(f \star_\alpha g) = \exp\left[\frac{1}{2}\alpha^{ab}\partial_a^{(1)}\partial_b^{(2)}\right]f(x_1)g(x_2)\Big|_{x_1=x_2=x'} \tag{6}$$

such that

$$\alpha = \begin{pmatrix} \theta_{ij} & \delta_{ij} + \sigma_{ij} \\ -\delta_{ij} - \sigma_{ij} & \beta_{ij} \end{pmatrix}, \tag{7}$$

where θ_{ij} and β_{ij} are 2×2 antisymmetric matrices indicating the noncommutativity in the coordinates and momenta, respectively. Particular deformations:

$$\theta_{ij} = -\theta\epsilon_{ij}, \beta_{ij} = \beta\epsilon_{ij}, \tag{8}$$

where ϵ_{ij} is the two-index Levi-Civita symbol, are considered.

By removing the sub-index in \star_α , the \star -Friedman equations can be derived for the \star -FLRW metric as follows [82].

$$R_{\mu\nu} - \frac{1}{2}R \star g_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu} \tag{9}$$

with the energy–momentum tensor:

$$T_{\mu\nu} = p \star g_{\mu\nu} + (\rho + p) \star U_\mu \star U_\nu \tag{10}$$

where $U_\mu = \delta_\mu^0$ is the co-moving observer and p and ρ are the total pressure and fluid energy three-density, respectively.

To avoid the complexities of \star -algebras, one may consider the field equations arising from the point-like action for a scalar field with action [83]:

$$S = \int dt N \left(-3\frac{a\dot{a}^2}{N^2}\right) + \frac{1}{2} \int dt N a^3 \left(\epsilon\frac{\dot{\phi}^2}{N^2} - 2V(\phi)\right). \tag{11}$$

We define the point-like Lagrangian [83]:

$$\mathcal{L}(N, a, \phi, \dot{a}, \dot{\phi}) := \frac{1}{N} \left(-3a\dot{a}^2 + \frac{1}{2}a^3\epsilon\dot{\phi}^2\right) - a^3NV(\phi), \tag{12}$$

while, for simplicity, we consider a constant potential $V(\phi) = \tilde{\Lambda}$. The sign $\epsilon = 1$ corresponds to quintessence, and the sign $\epsilon = -1$ corresponds to the phantom field.

With the variation with respect to $\{N, a, \phi\}$ and the replacement $N = 1$ after variation, we obtain the Euler–Lagrange equations:

$$\frac{1}{2} \left(6a\dot{a}^2 - a^3 (2\tilde{\Lambda} + \epsilon\dot{\phi}^2) \right) = 0, \tag{13}$$

$$\frac{3}{2} \left(4a\ddot{a} + 2\dot{a}^2 + a^2 (\epsilon\dot{\phi}^2 - 2\tilde{\Lambda}) \right) = 0, \tag{14}$$

$$-\epsilon a^2 (3\dot{a}\dot{\phi} + a\ddot{\phi}) = 0. \tag{15}$$

Introducing the Hubble parameter $H = \dot{a}/a$, the previous equations can be written as [83]:

$$3H^2 = \tilde{\Lambda} + \frac{1}{2}\epsilon\dot{\phi}^2, \tag{16}$$

$$2\dot{H} = -3H^2 + \tilde{\Lambda} - \frac{1}{2}\epsilon\dot{\phi}^2, \tag{17}$$

$$\ddot{\phi} + 3H\dot{\phi} = 0. \tag{18}$$

For the Lagrangian function (12), we define the generalized momenta by $p_i = \frac{\partial \mathcal{L}}{\partial \dot{q}^i}$, where $q^i \in \{a, \phi\}$, $p_i \in \{p_a, p_\phi\}$, namely

$$p_a \equiv -\frac{6a\dot{a}}{N}, \quad p_\phi \equiv \frac{\epsilon a^3 \dot{\phi}}{N}. \tag{19}$$

Hence, we can introduce the Hamiltonian function $\mathcal{H} = p_a \dot{a} + p_\phi \dot{\phi} - \mathcal{L}$, which is written as

$$\mathcal{H} = N \left(-\frac{p_a^2}{12a} + \frac{\epsilon p_\phi^2}{2a^3} + \tilde{\Lambda} a^3 \right) \tag{20}$$

We define the canonical coordinates [83]:

$$x = \lambda^{-1} (\epsilon a)^{3/2} \sinh(\sqrt{\epsilon} \lambda \phi), \quad y = \lambda^{-1} a^{3/2} \cosh(\sqrt{\epsilon} \lambda \phi), \tag{21}$$

with the inverse:

$$a = \left(\lambda y \sqrt{1 - \frac{\epsilon x^2}{y^2}} \right)^{2/3}, \quad \phi = \frac{\tanh^{-1}(\sqrt{\epsilon} x/y)}{\lambda \sqrt{\epsilon}}, \tag{22}$$

where $\lambda^{-1} = \sqrt{8/3}$, and we consider the simpler case where the matter content is an ordinary ($\epsilon = +1$) or a phantom ($\epsilon = -1$) scalar field in the action. Then, (12) becomes

$$\mathcal{L}(N, a, \phi, \dot{a}, \dot{\phi}) := \frac{1}{N} \left(-3a\dot{a}^2 + \frac{1}{2}a^3\epsilon\dot{\phi}^2 \right) - a^3 N V(\phi). \tag{23}$$

Generalized momenta are given by

$$P_x = \frac{\epsilon \dot{x}}{N}, \quad P_y = -\frac{\dot{y}}{N}. \tag{24}$$

Hence, the problem can be formulated from the canonical Hamiltonian:

$$\mathcal{H}_c = \epsilon N \left(\frac{1}{2} P_x^2 + \frac{\omega^2}{2} x^2 \right) - N \left(\frac{1}{2} P_y^2 + \frac{\omega^2}{2} y^2 \right), \tag{25}$$

where $\omega^2 = -3\tilde{\Lambda}/4$, and we use the comoving frame $N = 1$. For the choice $\epsilon = +1$, see the related work [65].

We have the evolution equations for (\dot{x}, \dot{y}) as given by (24):

$$\dot{x} = \epsilon P_x, \quad \dot{y} = -P_y \tag{26}$$

Hamilton’s equations $\dot{p}_i = -\frac{\partial \mathcal{H}}{\partial q^i}$, where $\mathcal{H} = \mathcal{H}_c$ and $q^i \in \{x, y\}$, $p_i \in \{P_x, P_y\}$, lead to

$$\dot{P}_x = -\epsilon\omega^2x, \dot{P}_y = \omega^2y. \tag{27}$$

which lead to the following equations for $\epsilon = \pm 1$:

$$\ddot{x} + \omega^2x = 0, \ddot{y} + \omega^2y = 0, \tag{28}$$

with conserved quantity:

$$\tilde{y}^2 - \epsilon\tilde{x}^2 = 1, (\tilde{x}, \tilde{y}) = \lambda a^{-3/2} \cdot (x, y). \tag{29}$$

By the definition $\omega^2 = -3/4\tilde{\Lambda}$, the solutions are

$$x(t) = c_1e^{\frac{1}{2}\sqrt{3}\sqrt{\tilde{\Lambda}}t} + c_2e^{-\frac{1}{2}\sqrt{3}\sqrt{\tilde{\Lambda}}t}, \tag{30}$$

$$y(t) = c_3e^{\frac{1}{2}\sqrt{3}\sqrt{\tilde{\Lambda}}t} + c_4e^{-\frac{1}{2}\sqrt{3}\sqrt{\tilde{\Lambda}}t}. \tag{31}$$

Then,

$$\phi(t) = \frac{\coth^{-1}\left(\frac{e^{3/2}(c_3e^{\sqrt{3}\sqrt{\tilde{\Lambda}}t} + c_4)}{c_1e^{\sqrt{3}\sqrt{\tilde{\Lambda}}t} + c_2}\right)}{\lambda\sqrt{\epsilon}}, \tag{32}$$

$$a(t) = e^{-\sqrt{\frac{\tilde{\Lambda}}{3}}t} \left(\lambda(c_3e^{\sqrt{3}\sqrt{\tilde{\Lambda}}t} + c_4)\right)^{2/3} \left(1 - \frac{\epsilon(c_1e^{\sqrt{3}\sqrt{\tilde{\Lambda}}t} + c_2)^2}{(c_3e^{\sqrt{3}\sqrt{\tilde{\Lambda}}t} + c_4)^2}\right)^{1/3}, \tag{33}$$

such that

$$\phi \sim \frac{\ln\left(1 - \frac{2c_1}{c_1 - c_3e^{3/2}}\right)}{2\lambda\sqrt{\epsilon}}, a \sim c_3^{2/3}\lambda^{2/3}3\sqrt[3]{1 - \frac{c_1^2\epsilon}{c_3^2}e^{\frac{\sqrt{\tilde{\Lambda}}t}{\sqrt{3}}}} \tag{34}$$

as $t \rightarrow \infty$. That is, a de Sitter solution is obtained.

The elements of the new configuration space, (x, y) , and their conjugate momenta fulfil the following commutation relations based on the Poisson bracket:

$$\{x_k, x_j\} = 0, \{P_{x_k}, P_{x_j}\} = 0, \{x_k, P_{x_j}\} = \delta_{kj} \tag{35}$$

where k and j can take 1 and 2, that is $(x_1, x_2) = (x, y)$ and δ_{kj} is the usual Kronecker delta.

To obtain a modified scenario, we take classical phase space variables (x, y, P_x, P_y) and perform the transformation (see the related work [65]):

$$\begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix} + \frac{\theta}{2} \begin{pmatrix} P_y \\ -P_x \end{pmatrix} \tag{36}$$

and

$$\begin{pmatrix} \hat{P}_x \\ \hat{P}_y \end{pmatrix} = \frac{\beta}{2} \begin{pmatrix} -y \\ x \end{pmatrix} + \begin{pmatrix} P_x \\ P_y \end{pmatrix} \tag{37}$$

The modified Poisson Algebra is given by

$$\{\hat{y}, \hat{x}\} = \theta, \{\hat{P}_y, \hat{P}_x\} = \beta \tag{38}$$

and

$$\{\hat{x}, \hat{P}_x\} = 1 + \sigma, \{\hat{y}, \hat{P}_y\} = 1 + \sigma, \tag{39}$$

where $\sigma = \theta\beta/4$. Now, we change the notation $\{\hat{x}, \hat{y}, \hat{P}_x, \hat{P}_y\}$ to $\{x, y, p_x, p_y\}$.

The modified Hamiltonian will be

$$\mathcal{H}_{mod.} = \frac{1}{2}\epsilon p_x^2 - \frac{1}{2}p_y^2 - \frac{\omega_1^2}{2}(xp_y + \epsilon yp_x) + \frac{\omega_2^2}{2}(\epsilon x^2 - y^2), \tag{40}$$

where

$$p_x = \dot{x} + \frac{1}{2}\omega_1 y, \quad p_y = \frac{-x\omega_1^2\epsilon - 2\dot{y}}{2\epsilon}, \tag{41}$$

and we define the parameters

$$\omega_1^2 = \frac{4\beta - 4\epsilon\omega^2\theta}{4 - \epsilon\omega^2\theta^2}, \quad \omega_2^2 = \frac{4\omega^2 - \epsilon\beta^2}{4 - \epsilon\omega^2\theta^2}, \tag{42}$$

$$\Lambda = -\frac{4((\beta - 1)\beta\epsilon + (\theta - 4)\omega^2)}{3(\theta^2\omega^2\epsilon - 4)}. \tag{43}$$

If $\omega = 0$, the latter definitions are

$$\omega_1^2 = \beta, \quad \omega_2^2 = -\frac{\epsilon\beta^2}{4}, \quad \Lambda = \frac{(\beta - 1)\beta\epsilon}{3}. \tag{44}$$

We can infer from these that the cosmological constant term is introduced from the modification of the Poisson algebra if our initial model does not include a cosmological constant term. The equations of motion derived from $\mathcal{H}_{mod.}$ are

$$\ddot{x} + \omega_1^2 \dot{y} - \frac{3}{4}\Lambda x = 0, \tag{45}$$

and

$$\ddot{y} + \epsilon\omega_1^2 \dot{x} - \frac{3}{4}\Lambda y = 0, \tag{46}$$

These equations have the solutions:

$$x(t) = c_1 \cosh\left(\frac{t\omega_1^2}{2}\right) \cosh\left(\frac{1}{2}t\sqrt{3\Lambda + \omega_1^4}\right) - c_3 \sinh\left(\frac{t\omega_1^2}{2}\right) \cosh\left(\frac{1}{2}t\sqrt{3\Lambda + \omega_1^4}\right) + \frac{\sinh\left(\frac{1}{2}t\sqrt{3\Lambda + \omega_1^4}\right) \left((c_3\omega_1^2 + 2c_2) \cosh\left(\frac{t\omega_1^2}{2}\right) - (c_1\omega_1^2 + 2c_4) \sinh\left(\frac{t\omega_1^2}{2}\right) \right)}{\sqrt{3\Lambda + \omega_1^4}}, \tag{47}$$

$$y(t) = c_3 \cosh\left(\frac{t\omega_1^2}{2}\right) \cosh\left(\frac{1}{2}t\sqrt{3\Lambda + \omega_1^4}\right) - c_1 \sinh\left(\frac{t\omega_1^2}{2}\right) \cosh\left(\frac{1}{2}t\sqrt{3\Lambda + \omega_1^4}\right) + \frac{\sinh\left(\frac{1}{2}t\sqrt{3\Lambda + \omega_1^4}\right) \left((c_1\omega_1^2 + 2c_4) \cosh\left(\frac{t\omega_1^2}{2}\right) - (c_3\omega_1^2 + 2c_2) \sinh\left(\frac{t\omega_1^2}{2}\right) \right)}{\sqrt{3\Lambda + \omega_1^4}}. \tag{48}$$

Some solutions of this form have been found before in the literature, e.g., [84,85].

3. Modified Friedmann Equations

Equations (45) and (46) are equivalent to

$$\ddot{a} = -\frac{\dot{a}^2}{2a} - \frac{1}{6}a \left(4\lambda\epsilon\dot{\phi} \left(\lambda\dot{\phi} + \omega_1^2 \right) + 4\omega_2^2 + \omega_1^2\epsilon \right), \tag{49}$$

$$\ddot{\phi} = -\frac{3\dot{a}(2\lambda\dot{\phi} + \omega_1^2)}{2\lambda a} \tag{50}$$

with the first integral

$$-3H^2 + \frac{1}{2}\epsilon\dot{\phi}^2 - \frac{4\omega_2^2}{3} - \frac{\omega_1^2\epsilon}{3} = 0. \tag{51}$$

where $H = \dot{a}/a$ is the Hubble parameter.

3.1. Vacuum Case

Using the reparameterization (43), the modified Friedman equation reads

$$3H^2 = \frac{1}{2}\epsilon\dot{\phi}^2 + \Lambda. \tag{52}$$

The modified Klein–Gordon equation is

$$\ddot{\phi} = -H(\sqrt{6}\omega_1^2 + 3\dot{\phi}) \tag{53}$$

The Raychaudhuri equation is

$$\dot{H} = -\frac{3H^2}{2} - \frac{\epsilon\omega_1^2\dot{\phi}}{\sqrt{6}} - \frac{1}{4}\epsilon\dot{\phi}^2 + \frac{\Lambda}{2}, \tag{54}$$

Alternatively, by removing H^2 and using (52), we obtain

$$\dot{H} = -\frac{\epsilon\omega_1^2\dot{\phi}}{\sqrt{6}} - \frac{1}{2}\epsilon\dot{\phi}^2. \tag{55}$$

With the definitions:

$$\rho_\phi = \frac{1}{2}\epsilon\dot{\phi}^2 + \Lambda, \quad P_\phi = \frac{1}{2}\epsilon\dot{\phi}^2 + \sqrt{\frac{2}{3}}\omega_1^2\epsilon\dot{\phi} - \Lambda, \tag{56}$$

the Klein–Gordon equation can be written as the conservation equation:

$$\dot{\rho}_\phi + 3H(\rho_\phi + P_\phi) = 0. \tag{57}$$

Moreover, we define the effective EoS parameter of ϕ as

$$w_\phi := \frac{P_\phi}{\rho_\phi} = \frac{\epsilon\dot{\phi}^2 + 2\sqrt{\frac{2}{3}}\omega_1^2\epsilon\dot{\phi} - 2\Lambda}{\epsilon\dot{\phi}^2 + 2\Lambda}. \tag{58}$$

3.2. Including Matter

The Friedman equation reads

$$3H^2 = \frac{1}{2}\epsilon\dot{\phi}^2 + \rho_m + \Lambda. \tag{59}$$

The modified Klein–Gordon equation is

$$\ddot{\phi} = -H(\sqrt{6}\omega_1^2 + 3\dot{\phi}) \tag{60}$$

which can be written using (56) as

$$\dot{\rho}_\phi + 3H(\rho_\phi + P_\phi) = 0. \tag{61}$$

We have the matter conservation equation:

$$\dot{\rho}_m + 3H(1 + w_m)\rho_m = 0. \tag{62}$$

The Raychaudhuri equation is

$$\dot{H} = -\frac{\omega_1^2 \epsilon \dot{\phi}}{\sqrt{6}} - \frac{1}{2} \epsilon \dot{\phi}^2 - \frac{1}{2} (w_m + 1) \rho_m. \tag{63}$$

4. Dynamical Systems' Analysis in the Vacuum Case

In this section, we proceed with the analysis of the phase space for the modified cosmological field equations. In order to perform such an analysis, we define dimensionless variables in the Hubble normalization approach, that is

$$\Sigma_\phi = \frac{\dot{\phi}}{\sqrt{6}H}, \quad \Sigma = \frac{\omega_1^2}{H}, \tag{64}$$

which satisfies the constraint equation:

$$\Sigma_\phi^2 \epsilon + \mu \Sigma^2 = 1, \tag{65}$$

where we have introduced the constant $\mu = \Lambda / (3\omega_1^4)$, or, alternatively,

$$\Sigma_\phi^2 \epsilon + \Omega_\Lambda = 1. \tag{66}$$

where, for convenience, we define the fractional energy density of Λ as

$$\Omega_\Lambda := \frac{\Lambda}{3H^2} = \mu \Sigma^2, \quad \mu > 0. \tag{67}$$

Thus, the dynamical system (52), (53), and (55) can be written as a dynamical system:

$$\Sigma'_\phi = (3\Sigma_\phi + \Sigma) (\Sigma_\phi^2 \epsilon - 1), \tag{68}$$

$$\Sigma' = \epsilon \Sigma_\phi \Sigma (3\Sigma_\phi + \Sigma), \tag{69}$$

where we have introduced the new time derivative $f' = H^{-1} \dot{f}$.

4.1. Analysis of the 2D Flow

In this section, we analyze the 2D flow associated with the dynamical system (68) and (69). We obtain the (lines of) equilibrium points of the system (68) and (69), which are summarized in Table 1 for $\epsilon = \pm 1$ along with their coordinates, eigenvalues, and stability.

Table 1. Equilibrium points of System (68)–(69) for $\epsilon = +1$ with their eigenvalues and stability. P_5 is a sink, but does not satisfy the condition (65).

Label	Existence	Coordinates (Σ_ϕ, Σ)	Eigenvalues	Stability
$P_{1,2}$	$\epsilon = +1$	$(\pm 1, 0)$	$\{3, 6\}$	Unstable
L_1	always	$(\Sigma_\phi, -3\Sigma_\phi)$	$\{-3, 0\}$	Stable
$P_{3,6}$	$\epsilon = \pm 1$	$\left(\frac{\epsilon}{\sqrt{9\mu+\epsilon}}, -\frac{3\epsilon}{\sqrt{9\mu+\epsilon}}\right)$	$\{-3, 0\}$	Stable
$P_{4,7}$	$\epsilon = \pm 1$	$\left(-\frac{\epsilon}{\sqrt{9\mu+\epsilon}}, \frac{3\epsilon}{\sqrt{9\mu+\epsilon}}\right)$	$\{-3, 0\}$	Stable
P_5	\nexists	$(0, 0)$	$\{-3, 0\}$	Stable

4.1.1. Case $\epsilon = +1$

In the case $\epsilon = +1$, there exist kinetic-dominated solutions, given by the points P_1 and P_2 . They represent stiff solutions ($w_\phi = 1$).

There is a line of equilibrium points L_1 , which corresponds to

$$\dot{\phi} + \frac{\sqrt{6}}{3}\omega_1^2 = 0 \implies \phi = -\frac{\sqrt{6}}{3}\omega_1^2 t + c_1. \tag{70}$$

Then, from (55), we have at the lines of the equilibrium points

$$\dot{H} = 0 \implies H = H_0, \implies a = e^{H_0(t-t_0)}. \tag{71}$$

where H_0 satisfies $3H_0^2 = \Lambda + \frac{\omega_1^4 \epsilon}{3}$. That is a de Sitter solution.

Moreover, imposing the condition (65), the lines are reduced to the points P_3 and P_4 that belong to the lines of the equilibrium points $3\Sigma_\phi + \Sigma = 0$.

For these equilibrium points, we have $\dot{\phi} = \sqrt{6}\Sigma_{\phi_{1,2}}H$, where $\Sigma_{\phi_{1,2}} = \pm 1/\sqrt{9\mu + 1}$. Hence,

$$\dot{H} + \Sigma_{\phi_{1,2}}H \left(3\Sigma_{\phi_{1,2}}H + \sqrt{\Lambda/(3\mu)} \right) = 0. \tag{72}$$

That is,

$$H = \frac{\sqrt{3}H_0\sqrt{\frac{\Lambda}{\mu}}}{\left(9H_0\Sigma_{\phi_{1,2}} + \sqrt{3}\sqrt{\frac{\Lambda}{\mu}} \right) e^{(t-t_0)\Sigma_{\phi_{1,2}}\sqrt{\frac{\Lambda}{3\mu}}} - 9H_0\Sigma_{\phi_{1,2}}}. \tag{73}$$

The line L_1 also contains the point P_5 , which is a sink, but does not satisfy the condition (65).

Figure 1 presents a phase plot for System (68)–(69) for $\epsilon = +1$ and different values of μ . The dashed black line corresponds to L_1 .




Figure 1. Phase plot for System (68)–(69) for $\epsilon = +1$ and different values of μ . The dashed black line corresponds to L_1 .

4.1.2. Case $\epsilon = -1$

For the case $\epsilon = -1$, the equilibrium points of the system (68) and (69) are, as before, the line L_1 , which contains P_5 (which does not satisfy (65)). Moreover, imposing the condition (65), the lines are reduced to the points P_6 and P_7 that belong to the lines of equilibrium points $3\Sigma_\phi + \Sigma = 0$, where $\Sigma_{\phi_{1,2}} = \mp 1/\sqrt{9\mu - 1}$.

For these equilibrium points, we have $\dot{\phi} = \sqrt{6}\Sigma_{\phi_{1,2}}H$. Hence,

$$\dot{H} - \Sigma_{\phi_{1,2}}H \left(3\Sigma_{\phi_{1,2}}H + \sqrt{\Lambda/(3\mu)} \right) = 0. \tag{74}$$

That is,

$$H = \frac{\sqrt{3}H_0\sqrt{\frac{\Lambda}{\mu}}}{\left(9H_0\Sigma_{\phi_{1,2}} + \sqrt{3}\sqrt{\frac{\Lambda}{\mu}} \right) e^{-(t-t_0)\Sigma_{\phi_{1,2}}\sqrt{\frac{\Lambda}{3\mu}}} - 9H_0\Sigma_{\phi_{1,2}}}. \tag{75}$$

Figure 2 presents a phase plot for System (68)–(69) for $\epsilon = -1$ and different values of μ . The dashed black line corresponds to L_1 .




Figure 2. Phase plot for System (68)–(69) for $\epsilon = -1$ and different values of μ . The dashed black line corresponds to L_1 .

4.2. The 1D Reduced System

Using (65) to reduce the dimensionality and for $\Sigma \geq 0$, we have

$$\Sigma = \sqrt{(1 - \Sigma_\phi^2 \epsilon) / \mu}. \tag{76}$$

Then, we have the reduced dynamical system:

$$\Sigma'_\phi = -\left(3\Sigma_\phi + \sqrt{(1 - \Sigma_\phi^2 \epsilon) / \mu}\right) \left(1 - \Sigma_\phi^2 \epsilon\right), \tag{77}$$

This patch covers only the equilibrium points (68)–(69) with $\Sigma > 0$ and $\Sigma_\phi = 0$, say $P_1 : \Sigma_\phi = 1$, $P_2 : \Sigma_\phi = -1$ and $P_4 : \Sigma_\phi = -1 / \sqrt{9\mu + 1}$. Moreover, the equilibrium point of the 1D system (77) is $P_6 : \Sigma_\phi = -1 / \sqrt{9\mu - 1}$.

Table 2 presents the equilibrium points of System (77) for $\epsilon = \pm 1$ with their eigenvalues and stability.

Table 2. Equilibrium points of System (77) for $\epsilon = \pm 1$ with their eigenvalues and stability.

Label	Existence	Coordinates Σ_ϕ	Eigenvalue	Stability
$P_{1,2}$	$\epsilon = +1$	± 1	6	Unstable
P_3	$\epsilon = +1, \mu > -\frac{1}{9}$	$\frac{1}{\sqrt{9\mu+1}}$	-3	Stable
P_6	$\epsilon = -1, \mu > \frac{1}{9}$	$-\frac{1}{\sqrt{9\mu-1}}$	-3	Stable

Figure 3 presents a phase plot for the reduced 1D equation (77) for $\epsilon = \pm 1$ and $\mu = 2$.




Figure 3. Phase plot for the reduced 1D equation (77) for $\epsilon = \pm 1$ and $\mu = 2$.

On the other hand, using

$$\Sigma = -\sqrt{(1 - \Sigma_\phi^2 \epsilon) / \mu}, \tag{78}$$

we have the reduced dynamical system:

$$\Sigma'_\phi = -\left(3\Sigma_\phi - \sqrt{(1 - \Sigma_\phi^2\epsilon)/\mu}\right)\left(1 - \Sigma_\phi^2\epsilon\right), \tag{79}$$

This patch covers only the equilibrium points (68)–(69) with $\Sigma < 0$ and $\Sigma_\phi = 0$, say $P_1 : \Sigma_\phi = 1, P_2 : \Sigma_\phi = -1$ and $P_3 : \Sigma_\phi := 1/\sqrt{9\mu + 1}$. Moreover, the equilibrium point of the 1D system (79) is $P_7 : \Sigma_\phi = 1/\sqrt{9\mu - 1}$, which belongs to the lines of the equilibrium points $3\Sigma_\phi + \Sigma = 0$.

Table 3 presents the equilibrium points of System (79) for $\epsilon = \pm 1$ with their eigenvalues and stability.

Table 3. Equilibrium points of System (79) for $\epsilon = \pm 1$ with their eigenvalues and stability.

Label	Existence	Coordinates Σ_ϕ	Eigenvalue	Stability
$P_{1,2}$	± 1	$\epsilon = +1$	6	Unstable
P_4	$\epsilon = +1, \mu > -\frac{1}{9}$	$-\frac{1}{\sqrt{9\mu+1}}$	-3	Stable
P_7	$\epsilon = -1, \mu > \frac{1}{9}$	$\frac{1}{\sqrt{9\mu-1}}$	-3	Stable

Figure 4 presents a phase plot for the reduced 1D equation (79) for $\epsilon = \pm 1$ and $\mu = 2$.




Figure 4. Phase plot for the reduced 1D equation (79) for $\epsilon = \pm 1$ and $\mu = 2$.

5. Dynamical Systems' Analysis by Including Matter

We define

$$\Sigma_\phi = \frac{\dot{\phi}}{\sqrt{6H}}, \Sigma = \frac{\omega_1^2}{H}, \Omega_m = \frac{\rho_m}{3H^2}, \tag{80}$$

which satisfy

$$\Sigma_\phi^2\epsilon + \mu\Sigma^2 + \Omega_m = 1. \tag{81}$$

5.1. The 3D System

The system (52), (53), and (52) can be written as the dynamical system given by

$$\Sigma_\phi' = \frac{3}{2}(w_m + 1)\Sigma_\phi\Omega_m + (3\Sigma_\phi + \Sigma)\left(\Sigma_\phi^2\epsilon - 1\right), \tag{82}$$

$$\Sigma' = \frac{3}{2}(w_m + 1)\Sigma\Omega_m + \Sigma_\phi\Sigma\epsilon(3\Sigma_\phi + \Sigma), \tag{83}$$

$$\Omega'_m = \Omega_m(3w_m(\Omega_m - 1) + 2\Sigma_\phi\epsilon(3\Sigma_\phi + \Sigma) + 3(\Omega_m - 1)). \tag{84}$$

Table 4 presents the equilibrium points of System (82), (83), and (84) for $\epsilon = \pm 1$ with their eigenvalues and stability.

Table 4. Equilibrium points of System (82), (83), and (84) for $\epsilon = \pm 1$ with their eigenvalues and stability.

Label	Existence	Coordinates $(\Sigma_\phi, \Sigma, \Omega_m)$	Eigenvalues	Stability
$P_{1,2}$	$\epsilon = +1$	$(\pm 1, 0, 0)$	$\{3, 6, 3(1 - \omega_m)\}$	Unstable
L_1	always	$(\Sigma_\phi, -3\Sigma_\phi, 0)$	$\{-3, 0, -3(\omega_m + 1)\}$	Stable
$P_{3,6}$	$\epsilon = \pm 1$	$\left(\frac{\epsilon}{\sqrt{9\mu+\epsilon}}, -\frac{3\epsilon}{\sqrt{9\mu+\epsilon}}, 0\right)$	$\{-3, 0, -3(\omega_m + 1)\}$	Stable
$P_{4,7}$	$\epsilon = \pm 1$	$\left(-\frac{\epsilon}{\sqrt{9\mu+\epsilon}}, \frac{3\epsilon}{\sqrt{9\mu+\epsilon}}, 0\right)$	$\{-3, 0, -3(\omega_m + 1)\}$	Stable
P_5	\nexists	$(0, 0, 0)$	$\{-3, 0, -3(\omega_m + 1)\}$	Stable
M	$\epsilon = \pm 1$	$(0, 0, 1)$	$\left\{\frac{3(\omega_m-1)}{2}, \frac{3(\omega_m+1)}{2}, 3(\omega_m + 1)\right\}$	Stable for $\omega_m = -1$ Unstable for $\omega_m = 1$ Saddle otherwise

5.1.1. Case $\epsilon = +1$

For the case $\epsilon = +1$, the equilibrium points of the system (82), (83), and (84) are P_1, P_2 , which are kinetic-dominated solutions. The line of equilibrium points L_1 represents the de Sitter solutions. This line contains the points P_3, P_4 , and P_5 . Additionally, we have the matter-dominated solution $M : (\Sigma_\phi, \Sigma, \Omega_m) = (0, 0, 1)$.

Figure 5 presents a 3D phase plot for System (82), (83), and (84) for $\epsilon = 1, \mu = 2$, and different values of ω_m .




Figure 5. The 3D phase plot for System (82), (83), and (84) for $\epsilon = +1, \mu = 2$, and different values of ω_m .

5.1.2. Case $\epsilon = -1$

For the case $\epsilon = -1$, the equilibrium points of the system (82), (83), and (84) are the line of equilibrium points L_1 , which represents the de Sitter solutions. This line contains the points P_5 , P_6 , and P_7 . Additionally, we have the matter-dominated solution $M : (\Sigma_\phi, \Sigma, \Omega_m) = (0, 0, 1)$.

Figure 6 presents a 3D phase plot for System (82), (83), and (84) for $\epsilon = -1, \mu = 2$, and different values of ω_m .




Figure 6. The 3D phase plot for System (82), (83), and (84) for $\epsilon = -1, \mu = 2$, and different values of ω_m .

5.2. Reduced 2D System

Eliminate Ω_m from (81) to obtain the reduced system:

$$\Sigma_\phi' = -\frac{1}{2}(\Sigma_\phi^2 \epsilon - 1)(3(\omega_m - 1)\Sigma_\phi - 2\Sigma) - \frac{3}{2}\mu(\omega_m + 1)\Sigma_\phi \Sigma^2, \tag{85}$$

$$\Sigma' = \frac{3}{2}(\omega_m + 1)\Sigma(1 - \Sigma_\phi^2 \epsilon - \mu \Sigma^2) + \Sigma_\phi \Sigma \epsilon (3\Sigma_\phi + \Sigma). \tag{86}$$

5.2.1. Case $\epsilon = +1$

The equilibrium points of the system (85) and (86) are $P_{1,2}, L_1, P_3, P_4$, and M , as summarized in Table 5.

Table 5. Equilibrium points of System (85) and (86) for $\epsilon = +1$ with their eigenvalues and stability.

Label	Coordinates $(\Sigma\phi, \Sigma)$	Eigenvalues	Stability
$P_{1,2}$	$(\pm 1, 0)$	$\{3, 6\}$	Unstable
L_1	$(\Sigma\phi, -3\Sigma\phi)$	$\{-3, -3(\omega + 1)\}$	Stable
P_3	$\left(\frac{1}{\sqrt{9\mu+1}}, -\frac{3}{\sqrt{9\mu+1}}\right)$	$\{-3, -3(\omega + 1)\}$	Stable
P_4	$\left(-\frac{1}{\sqrt{9\mu+1}}, \frac{3}{\sqrt{9\mu+1}}\right)$	$\{-3, -3(\omega + 1)\}$	Stable
M	$(0, 0)$	$\left\{\frac{3(\omega_m-1)}{2}, 3(\omega_m + 1)\right\}$	Stable for $\omega = -1$ Unstable for $\omega = 1$ Saddle otherwise

Figure 7 presents the 2D projections of the system (85)–(86) for $\epsilon = +1, \mu = 2$, and different values of ω_m .




Figure 7. The 2D projections of the system (85)–(86) for $\epsilon = +1, \mu = 2$, and different values of ω_m .

5.2.2. Case $\epsilon = -1$

The equilibrium points of the system (85) and (86) for $\epsilon = -1$ are summarized in Table 6.

Table 6. Equilibrium points of System (85) and (86) for $\epsilon = -1$ with their eigenvalues and stability.

Label	Coordinates (Σ_ϕ, Σ)	Eigenvalues	Stability
L_1	$(\Sigma_\phi, -3\Sigma_\phi)$	$\{-3, -3(\omega_m + 1)\}$	Stable
P_6	$\left(\frac{1}{\sqrt{9\mu-1}}, -\frac{3}{\sqrt{9\mu-1}}\right)$	$\{-3, -3(\omega + 1)\}$	Stable
P_7	$\left(-\frac{1}{\sqrt{9\mu-1}}, \frac{3}{\sqrt{9\mu-1}}\right)$	$\{-3, -3(\omega + 1)\}$	Stable
M	$(0, 0)$	$\left\{\frac{3(\omega_m-1)}{2}, 3(\omega_m + 1)\right\}$	Stable for $\omega = -1$ Unstable for $\omega = 1$ Saddle otherwise

Figure 8 presents the 2D projection of (85)–(86) for $\epsilon = -1, \mu = 2$, and different values of ω_m .




Figure 8. The 2D projection of (85)–(86) for $\epsilon = -1, \mu = 2$, and different values of ω_m .

6. Conclusions

In this study, we investigated the effects of the modification of the Poisson algebra on the dynamics of scalar field cosmology. Specifically, we performed a detailed phase space analysis by studying the equilibrium points and their stability, reconstructing the cosmological history.

The modified Poisson algebra modifies the field equations, introducing a cosmological constant term. The pressure component of the scalar field's energy-momentum tensor is different from that of the canonical scalar field. Moreover, a mass term for the scalar field is introduced, which is described by the cosmological constant.

As a result, the equilibrium points provided by the modified field equations are different from those of the usual scalar field model. From the analysis, we can conclude that the modified equations can provide more than one accelerating universe, described by the de Sitter solution. Hence, cosmic inflation and late-time acceleration are provided by the specific theory.

In the matter-less case, we divided the study into two subcases, one for $\epsilon = 1$ and one for $\epsilon = -1$. We have six families of physically acceptable equilibrium points that can describe stiff fluid solutions and de Sitter spacetime in the asymptotic regime.

In the case with the matter, we also considered the subcases $\epsilon = \pm 1$, and in total, we obtained eight families of equilibrium points, the same ones as in the case without matter and one additional equilibrium point that describes matter.

In future work, we plan to further investigate the modified field equations with the introduction of a nonzero scalar field potential. In contrast, an interacting term between the scalar field and the matter source will be considered.

The steps in this paper allow exploring the cosmological models' feasibility in concordance with the observational data set from measurements of Supernovae Ia, Cosmic Chronometers, baryon acoustic oscillation and cosmic microwave background. However, the observational test is out of the scope of the present research.

Author Contributions: G.L. managed the project, was involved in the asymptotic solutions, and wrote the final version of the paper. A.D.M. was involved in the asymptotic solutions, dynamical system analysis, and numerical integrations. A.P. organized the project, made the physical interpretation of the results and wrote the first draft. All authors have read and agreed to the published version of the manuscript.

Funding: Genly Leon was funded by Vicerrectoría de Investigación y Desarrollo Tecnológico (Vridt) at Universidad Católica del Norte (UCN) through Concurso De Pasantías De Investigación Año 2022, Resolución Vridt No. 040/2022, and through Resolución Vridt No. 054/2022. He also thanks the support of Núcleo de Investigación Geometría Diferencial y Aplicaciones, Resolución Vridt No. 096/2022. Andronikos Paliathanasis acknowledges Vridt-UCN through Concurso de Estadías de Investigación, Resolución VRIDT No. 098/2022. Alfredo David Millano was supported by Agencia Nacional de Investigación y Desarrollo (ANID) Subdirección de Capital Humano/Doctorado Nacional/año 2020 folio 21200837, Gastos operacionales proyecto de tesis/2022 folio 242220121, and Vridt-UCN.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tegmark, M.; Blanton, M.R.; Strauss, M.A.; Hoyle, F.; Schlegel, D.; Scoccimarro, R.; Vogeley, M.S.; Weinberg, D.H.; Zehavi, I.; Berlind, A.; et al. The 3-D power spectrum of galaxies from the SDSS. *Astrophys. J.* **2004**, *606*, 702. [[CrossRef](#)]
2. Kowalsk, M.; Rubin, D.; Aldering, G.; Agostinho, R.J.; Amadon, A.; Amanullah, R.; Bolland, C.; Barbary, K.; Blanc, G.; Challis, P.J.; et al. Improved Cosmological Constraints from New, Old, and Combined Supernova Data Sets. *Astrophys. J.* **2008**, *686*, 749. [[CrossRef](#)]
3. Komatsu, E.; Dunkley, J.; Nolta, M.R.; Bennett, C.L.; Gold, B.; Hinshaw, G.; Jarosik, N.; Larson, D.; Limon, M.; Page, L.; et al. Five-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation. *Astrophys. J. Suppl. Ser.* **2009**, *180*, 330. [[CrossRef](#)]
4. Ade, P.A.R.; Aghanim, N.; Armitage-Caplan, C.; Arnaud, M.; Ashdown, M.; Atrio-Barandela, F.; Aumont, J.; Baccigalupi, C.; Banday, A.J.; Barreiro, R.B.; et al. Planck 2013 results. XV. CMB power spectra and likelihood. *Astron. Astrophys.* **2014**, *571*, A15.
5. Padmanabhan, T. Cosmological constant—The weight of the vacuum. *Phys. Rep.* **2003**, *380*, 235. [[CrossRef](#)]
6. Weinberg, S. The Cosmological Constant Problem. *Rev. Mod. Phys.* **1989**, *61*, 1. [[CrossRef](#)]

7. Valentino, E.D.; Mena, O.; Pan, S.; Visinelli, L.; Yang, W.; Melchiorri, A.; Mota, D.F.; Reiss, A.G.; Silk, J. In the realm of the Hubble tension—A review of solutions. *Class. Quantum Grav.* **2021**, *38*, 153001.
8. Ratra, B.; Peebles, P.J.E. Cosmological Consequences of a Rolling Homogeneous Scalar Field. *Phys. Rev D* **1988**, *37*, 3406. [[CrossRef](#)]
9. Chen, W.; Wu, Y.-S. Implications of a cosmological constant varying as $R^{**(-2)}$. *Phys. Rev. D* **1990**, *41*, 695–698; Erratum in *Phys. Rev. D* **1992**, *45*, 4728. [[CrossRef](#)]
10. Basilakos, S.; Plionis, M.; Solà, S. Hubble expansion & Structure Formation in Time Varying Vacuum Models. *Phys. Rev. D* **2009**, *80*, 3511.
11. Wetterich, C. The Cosmon model for an asymptotically vanishing time dependent cosmological ‘constant’. *Astron. Astrophys.* **1995**, *301*, 321.
12. Caldwell, R.R.; Dave, R.; Steinhardt, P.J. Cosmological imprint of an energy component with general equation of state. *Phys. Rev. Lett.* **1998**, *80*, 1582. [[CrossRef](#)]
13. Brax, P.; Martin, J. Quintessence and supergravity. *Phys. Lett.* **1999**, *B468*, 40. [[CrossRef](#)]
14. Caldwell, R.R. A Phantom Menace? Cosmological consequences of a dark energy component with super-negative equation of state. *Phys. Rev. Lett. B* **2002**, *545*, 23. [[CrossRef](#)]
15. Lima, J.A.S.; Silva, F.E.; Santos, R.C. Accelerating Cold Dark Matter Cosmology ($\Omega_\Lambda \equiv 0$). *Class. Quant. Grav.* **2008**, *25*, 205006. [[CrossRef](#)]
16. Brookfield, A.W.; van de Bruck, C.; Mota, D.F.; Tocchini-Valentini, D. Cosmology with massive neutrinos coupled to dark energy. *Phys. Rev. Lett.* **2006**, *96*, 061301. [[CrossRef](#)]
17. Amendola, L.; Tsujikawa, S. *Dark Energy Theory and Observations*; Cambridge University Press: Cambridge, UK, 2010.
18. Linde, A.D. Chaotic Inflation. *Phys. Lett. B* **1983**, *129*, 177. [[CrossRef](#)]
19. Liddle, A.R. Power Law Inflation With Exponential Potentials. *Phys. Lett. B* **1989**, *220*, 502. [[CrossRef](#)]
20. Charters, T.; Mimoso, J.P.; Nunes, A. Slow roll inflation without fine tuning. *Phys. Lett. B* **2000**, *472*, 21. [[CrossRef](#)]
21. Barrow, J.D.; Saich, P. Scalar field cosmologies. *Class. Quantum Grav.* **1993**, *10*, 279. [[CrossRef](#)]
22. Chervon, S.V.; Zhuravlev, V.M.; Shchigolev, V.K. New exact solutions in standard inflationary models. *Phys. Lett. B* **1997**, *398*, 269. [[CrossRef](#)]
23. Kallosh, R.; Linde, A. Superconformal generalization of the chaotic inflation model $\frac{\lambda}{4}\phi^4 - \frac{\xi}{2}\phi^2 R$. *JCAP* **2013**, *13*, 027. [[CrossRef](#)]
24. Paliathanasis, A. New inflationary exact solution from Lie symmetries. *Mod. Phys. Lett. A* **2022**, *37*, 2250119. [[CrossRef](#)]
25. de Haro, J.; Amorós, J.; Pan, S. Simple inflationary quintessential model. *Phys. Rev. D* **2016**, *93*, 084018. [[CrossRef](#)]
26. Starobinsky, A.A. A New Type of Isotropic Cosmological Models Without Singularity. *Phys. Lett. B* **1980**, *91*, 99. [[CrossRef](#)]
27. Guth, A. The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems. *Phys. Rev. D* **1981**, *23*, 347. [[CrossRef](#)]
28. Dolgov, A.D. An attempt to get rid of the Cosmological Constant. In *The Very Early Universe*; Gibbons, G., Hawking, S.W., Tiklos, S.T., Eds.; Cambridge University Press: Cambridge UK, 1982.
29. Jassal, H.K.; Bagla, J.S.; Padmanabhan, T. Observational constraints on low redshift evolution of dark energy: How consistent are different observations? *Phys. Rev.* **2005**, *D72* 103503. [[CrossRef](#)]
30. Jassal, H.K.; Bagla, J.S.; Padmanabhan, T. Understanding the origin of CMB constraints on dark energy. *Mon. Not. R. Astron. Soc. Lett.* **2005**, *356*, L11–L16. [[CrossRef](#)]
31. Samushia, L.; Ratra, B. Cosmological Constraints from Hubble Parameter versus Redshift Data. *Astrophys. J.* **2006**, *650*, L5. [[CrossRef](#)]
32. Samushia, L.; Ratra, B. Constraints on Dark Energy from Galaxy Cluster Gas Mass Fraction versus Redshift Data. *Astrophys. J.* **2008**, *680*, L1. [[CrossRef](#)]
33. Simon, J.; Verde, L.; Jiménez, R. Constraints on the redshift dependence of the dark energy potential. *Phys. Rev.* **2005**, *D71*, 123001. [[CrossRef](#)]
34. Barrow, J.D.; Paliathanasis, A. Observational Constraints on New Exact Inflationary Scalar-field Solutions. *Phys. Rev. D* **2016**, *94*, 083518. [[CrossRef](#)]
35. Pan, S.; Yang, W.; Paliathanasis, A. Imprints of an extended Chevallier–Polarski–Linder parametrization on the large scale of our universe. *EPJC* **2020**, *80*, 274. [[CrossRef](#)]
36. Faraoni, V. Superquintessence. *Int. J. Mod. Phys. D* **2002**, *11*, 471. [[CrossRef](#)]
37. Lima, J.A.S.; Alcaniz, J.S. Thermodynamics and spectral distribution of dark energy. *Phys. Lett. B* **2004**, *600*, 191. [[CrossRef](#)]
38. Pereira, S.H.; Lima, J.A.S. On Phantom Thermodynamics. *Phys. Lett. B* **2008**, *669*, 266. [[CrossRef](#)]
39. Paliathanasis, A.; Tsamparlis, M.; Basilakos, S. Dynamical symmetries and observational constraints in scalar field cosmology. *Phys. Rev. D* **2014**, *90*, 103524. [[CrossRef](#)]
40. Martin, J.; Brandenberger, R.H. The TransPlanckian problem of inflationary cosmology. *Phys. Rev. D* **2001**, *63*, 123501. [[CrossRef](#)]
41. Niemeyer, J.C. Inflation with a Planck scale frequency cutoff. *Phys. Rev. D* **2001**, *63*, 123502. [[CrossRef](#)]
42. Kempf, A. Mode generating mechanism in inflation with cutoff. *Phys. Rev. D* **2001**, *63*, 083514. [[CrossRef](#)]
43. Kempf, A.; Niemeyer, J. Perturbation spectrum in inflation with cutoff. *Phys. Rev. D* **2001**, *64*, 103501. [[CrossRef](#)]
44. Ashoorioon, A.; Kempf, A.; Mann, R.B. Minimum length cutoff in inflation and uniqueness of the action. *Phys. Rev. D* **2005**, *71*, 023503. [[CrossRef](#)]

45. Ashoorioon, A.; Hovdebo, J.L.; Mann, R.B. Running of the spectral index and violation of the consistency relation between tensor and scalar spectra from trans-Planckian physics. *Nucl. Phys. B* **2005**, *727*, 63–76. [[CrossRef](#)]
46. Zampeli, A.; Paliathanasis, A. Quantization of inhomogeneous spacetimes with cosmological constant term. *Class. Quantum Grav.* **2021**, *38*, 165012. [[CrossRef](#)]
47. Paliathanasis, A. Quantum potentiality in Inhomogeneous Cosmology. *Universe* **2021**, *7*, 52. [[CrossRef](#)]
48. Zampeli, A.; Pailas, T.; Terzis, P.A.; Christodoulakis, T. Conditional symmetries in axisymmetric quantum cosmologies with scalar fields and the fate of the classical singularities. *JCAP* **2016**, *05*, 066. [[CrossRef](#)]
49. Mukhi, S. String theory: A perspective over the last 25 years. *Class. Quant. Grav.* **2011**, *28*, 153001. [[CrossRef](#)]
50. Kowalski-Glikman, J. Introduction to doubly special relativity. *Lect. Notes Phys.* **2005**, *669*, 131–159.
51. Amelino-Camelia, G. Doubly-Special Relativity: Facts, Myths and Some Key Open Issues. *Symmetry* **2010**, *2*, 230–271. [[CrossRef](#)]
52. Bekenstein, J.D. Black holes and entropy. *Phys. Rev. D* **1973**, *7*, 2333. [[CrossRef](#)]
53. Bekenstein, J.D. Black Holes and the Second Law. *Lett. Nuovo C* **1972**, *4*, 737–740. [[CrossRef](#)]
54. Maggiore, M. A Generalized uncertainty principle in quantum gravity. *Phys. Lett. B* **1993**, *304*, 65. [[CrossRef](#)]
55. Giacomini, A.; Leon, G.; Paliathanasis, A.; Pan, S. Dynamics of Quintessence in Generalized Uncertainty Principle. *Eur. Phys. J. C* **2020**, *80*, 931. [[CrossRef](#)]
56. Paliathanasis, A.; Leon, G.; Khylllep, W.; Dutta, J.; Pan, S. Interacting quintessence in light of generalized uncertainty principle: Cosmological perturbations and dynamics. *Eur. Phys. J. C* **2021**, *81*, 607. [[CrossRef](#)]
57. Masood, S.; Faizal, M.; Zal, Z.; Ali, A.F.; Raza, J.; Shah, M.B. The most general form of deformation of the Heisenberg algebra from the generalized uncertainty principle. *Phys. Lett. B* **2016**, *763*, 218. [[CrossRef](#)]
58. Rasouli, S.M.M.; Ziaie, A.H.; Marto, J.; Moniz, P.V. Gravitational Collapse of a Homogeneous Scalar Field in Deformed Phase Space. *Phys. Rev. D* **2014**, *89*, 044028. [[CrossRef](#)]
59. Jalalzadeh, S.; Rasouli, S.M.M.; Moniz, P.V. Quantum cosmology, minimal length and holography. *Phys. Rev. D* **2014**, *90*, 023541. [[CrossRef](#)]
60. Rasouli, S.M.M.; Farhoudi, M.; Vargas Moniz, P. Modified Brans–Dicke theory in arbitrary dimensions. *Class. Quant. Grav.* **2014**, *31*, 115002. [[CrossRef](#)]
61. Rasouli, S.M.M.; Vargas Moniz, P. Noncommutative minisuperspace, gravity-driven acceleration, and kinetic inflation. *Phys. Rev. D* **2014**, *90*, 083533. [[CrossRef](#)]
62. Rasouli, S.M.M.; Ziaie, A.H.; Jalalzadeh, S.; Moniz, P.V. Non-singular Brans–Dicke collapse in deformed phase space. *Ann. Phys.* **2016**, *375*, 154–178. [[CrossRef](#)]
63. Rasouli, S.M.M.; Vargas Moniz, P. Gravity-Driven Acceleration and Kinetic Inflation in Noncommutative Brans–Dicke Setting. *Odessa Astron. Pub.* **2016**, *29*, 19. [[CrossRef](#)]
64. Jalalzadeh, S.; Capistrano, A.J.S.; Moniz, P.V. Quantum deformation of quantum cosmology: A framework to discuss the cosmological constant problem. *Phys. Dark Univ.* **2017**, *18*, 55–66. [[CrossRef](#)]
65. Pérez-Payán, S.; Sabido, M.; Yee-Romero, C. Effects of deformed phase space on scalar field cosmology. *Phys. Rev. D* **2013**, *88*, 027503. [[CrossRef](#)]
66. Tajahmad, B. Late-time-accelerated expansion esteemed from minisuperspace deformation. *Eur. Phys. J. C* **2022**, *82*, 965. [[CrossRef](#)]
67. Tavakol, R. *Introduction to Dynamical Systems*; Cambridge University Press: Cambridge, UK, 1997; pp. 84–104.
68. Copeland, E.J.; Liddle, A.R.; Wands, D. Exponential potentials and cosmological scaling solutions. *Phys. Rev. D* **1998**, *57*, 4686–4690. [[CrossRef](#)]
69. Coley, A.A. *Dynamical Systems and Cosmology*; Kluwer: Dordrecht, The Netherlands, 2003.
70. Leon, G.; Fadrugas, C.R. *Cosmological Dynamical Systems*; LAP LAMBERT Academic Publishing: Saarbrücken, Germany, 2012; ISBN 978-3-8473-0233-9.
71. Gong, Y.; Wang, A.; Zhang, Y.-Z. Exact scaling solutions and fixed points for general scalar field. *Phys. Lett. B* **2006**, *636*, 286–292. [[CrossRef](#)]
72. Setare, M.R.; Saridakis, E.N. Quintom dark energy models with nearly flat potentials. *Phys. Rev. D* **2009**, *79*, 043005. [[CrossRef](#)]
73. Chen, X.M.; Gong, Y.-G.; Saridakis, E.N. Phase-space analysis of interacting phantom cosmology. *JCAP* **2009**, *04*, 001.
74. Gupta, G.; Saridakis, E.N.; Sen, A.A. Non-minimal quintessence and phantom with nearly flat potentials. *Phys. Rev. D* **2009**, *79*, 123013. [[CrossRef](#)]
75. Farajollahi, H.; Salehi, A.; Tayebi, F.; Ravanpak, A. Stability Analysis in Tachyonic Potential Chameleon cosmology. *JCAP* **2011**, *05*, 017. [[CrossRef](#)]
76. Arturo Urena-Lopez, L. Unified description of the dynamics of quintessential scalar fields. *JCAP* **2012**, *03*, 035. [[CrossRef](#)]
77. Escobar, D.; Fadrugas, C.R.; Leon, G.; Leyva, Y. Phase space analysis of quintessence fields trapped in a Randall-Sundrum Braneworld: A refined study. *Class. Quant. Grav.* **2012**, *29*, 175005. [[CrossRef](#)]
78. Escobar, D.; Fadrugas, C.R.; Leon, G.; Leyva, Y. Phase space analysis of quintessence fields trapped in a Randall-Sundrum Braneworld: Anisotropic Bianchi I brane with a Positive Dark Radiation term. *Class. Quant. Grav.* **2012**, *29*, 175006. [[CrossRef](#)]
79. Xu, C.; Saridakis, E.N.; Leon, G. Phase-Space analysis of Teleparallel Dark Energy. *JCAP* **2012**, *07*, 005. [[CrossRef](#)]
80. Leon, G.; Saavedra, J.; Saridakis, E.N. Cosmological behavior in extended nonlinear massive gravity. *Class. Quant. Grav.* **2013**, *30*, 135001. [[CrossRef](#)]

81. Burd, A.B.; Barrow, J.D. Inflationary Models with Exponential Potentials. *Nucl. Phys. B* **1988**, *308*, 929–945; Erratum in *Nucl. Phys. B* **1989**, *324*, 276. [[CrossRef](#)]
82. de Vegvar, P.G.N. Commutatively deformed general relativity: Foundations, cosmology, and experimental tests. *Eur. Phys. J. C* **2021**, *81*, 786. [[CrossRef](#)]
83. Basilakos, S.; Tsamparlis, M.; Paliathanasis, A. Using the Noether symmetry approach to probe the nature of dark energy. *Phys. Rev. D* **2011**, *83*, 103512. [[CrossRef](#)]
84. Paliathanasis, A.; Tsamparlis, M. Two scalar field cosmology: Conservation laws and exact solutions. *Phys. Rev. D* **2014**, *90*, 043529. [[CrossRef](#)]
85. Paliathanasis, A.; Leon, G.; Pan, S. Exact Solutions in Chiral Cosmology. *Gen. Rel. Grav.* **2019**, *51*, 106. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.