

December 8 1987

NIKHEF-H187-23



## CHARMONIUM PRODUCTION IN PHOTON-PHOTON COLLISIONS

TPC/Two-Gamma Collaboration

H. Aihara, M. Alston-Garnjost, R.E. Avery, A. Barbaro-Galtieri, A.R. Barker, B.A. Barnett, D.A. Bauer, H-U. Bengtsson, G.J. Bobbink, C.D. Buchanan, A. Buijs, D.O. Caldwell, H-Y. Chao, S-B. Chun, A.R. Clark, G.D. Cowan, D.A. Crane, O.I. Dahl, M. Daoudi, K.A. Derby, J.J. Eastman, P.H. Eberhard, T.K. Edberg, A.M. Eisner, R. Enomoto, F.C. Ern , K.H. Fairfield, J.M. Hauptman, W. Hofmann, J. Hylen, T. Kamae, H.S. Kaye, R.W. Kenney, S. Khacheryan, R.R. Kofler, W.G.J. Langeveld, J.G. Layter, W.T. Lin, F.L. Linde, S.C. Loken, A. Lu, G.R. Lynch, R.J. Madaras, B.D. Magnuson, G.E. Masek, L.G. Mathis, J.A.J. Matthews, S.J. Maxfield, E.S. Miller, W. Moses, D.R. Nygren, P.J. Oddone, H.P. Paar, S.K. Park, D.E. Pellett, M. Pripstein, M.T. Ronan, R.R. Ross, F.R. Rouse, K.A. Schwitkis, J.C. Sens, G. Shapiro, B.C. Shen, W.E. Slater, J.R. Smith, J.S. Steinman, R.W. Stephens, M.L. Stevenson, D.H. Stork, M.G. Strauss, M.K. Sullivan, T. Takahashi, S. Toutounchi, R. van Tyen, G.J. VanDalen, W. Vernon, W. Wagner, E.M. Wang, Y-X. Wang, W.A. Wenzel, Z.R. Wolf, H. Yamamoto, S.J. Yellin, C. Zeitlin

Lawrence Berkeley Laboratory, Berkeley, California 94720 and University of California, Davis, California 95616 and University of California Intercampus Institute for Research at Particle Accelerators, Stanford, California 94305 and University of California, Los Angeles, California 90024 and University of California, Riverside, California 92521 and University of California, San Diego, California 92093 and University of California, Santa Barbara, California 93106 and Ames Laboratory, Iowa State University, Ames, Iowa 50011 and Johns Hopkins University, Baltimore, Maryland 21218 and University of Massachusetts, Amherst, Massachusetts 01003 and National Institute for Nuclear and High Energy Physics, Amsterdam, The Netherlands and University of Tokyo, Tokyo, Japan

### ABSTRACT

We have searched for the two-photon production of the  $\eta_c$ ,  $\chi_0$  and  $\chi_2$  charmonium states at the  $e^+e^-$  collider PEP in the channels  $\gamma\gamma \rightarrow K^\pm K_S^0 \pi^\mp$ ,  $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^-$ ,  $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^+ \pi^-$  and  $\gamma\gamma \rightarrow K^+ K^- K^+ K^-$ . We identify four  $\eta_c$  candidates in the  $K^+ K^- K^+ K^-$  channel on a negligible background; the one  $\phi\phi$  event among them implies a 95% C.L. lower limit for  $\Gamma_{\gamma\gamma}(\eta_c)$  of 1.7 keV. In the other channels we find no evidence for any of the three states. We establish 95% C.L. upper limits  $\Gamma_{\gamma\gamma}(\eta_c) < 15.5$  keV,  $\Gamma_{\gamma\gamma}(\chi_0) < 17.0$  keV and  $\Gamma_{\gamma\gamma}(\chi_2) < 4.2$  keV. From all channels combined, we obtain the value  $\Gamma_{\gamma\gamma}(\eta_c) = 6.4_{-3.4}^{+5.0}$  keV.

PACS numbers: 14.40.Gx, 13.25.+m, 13.60.Le

# CHARMONIUM PRODUCTION IN PHOTON-PHOTON COLLISIONS

TPC/Two-Gamma Collaboration

H. Aihara, M. Alston-Garnjost, R.E. Avery, A. Barbaro-Galtieri, A.R. Barker, B.A. Barnett,  
D.A. Bauer, H-U. Bengtsson, G.J. Bobbink, C.D. Buchanan, A. Buijs, D.O. Caldwell,  
H-Y. Chao, S-B. Chun, A.R. Clark, G.D. Cowan, D.A. Crane, O.I. Dahl, M. Daoudi,  
K.A. Derby, J.J. Eastman, P.H. Eberhard, T.K. Edberg, A.M. Eisner, R. Enomoto,  
F.C. Erne, K.H. Fairfield, J.M. Hauptman, W. Hofmann, J. Hylen, T. Kamae, H.S. Kaye,  
R.W. Kenney, S. Khacheryan, R.R. Kofler, W.G.J. Langeveld, J.G. Layter, W.T. Lin,  
F.L. Linde, S.C. Loken, A. Lu, G.R. Lynch, R.J. Madaras, B.D. Magnuson,  
G.E. Masek, L.G. Mathis, J.A.J. Matthews, S.J. Maxfield, E.S. Miller,  
W. Moses, D.R. Nygren, P.J. Oddone, H.P. Paar, S.K. Park, D.E. Pellett,  
M. Pripstein, M.T. Ronan, R.R. Ross, F.R. Rouse, K.A. Schwitkis, J.C. Sens,  
G. Shapiro, B.C. Shen, W.E. Slater, J.R. Smith, J.S. Steinman,  
R.W. Stephens, M.L. Stevenson, D.H. Stork, M.G. Strauss, M.K. Sullivan,  
T. Takahashi, S. Toutouchi, R. van Tyen, G.J. VanDalen, W. Vernon,  
W. Wagner, E.M. Wang, Y-X. Wang, W.A. Wenzel,  
Z.R. Wolf, H. Yamamoto, S.J. Yellin, C. Zeitlin

*Lawrence Berkeley Laboratory, Berkeley, California 94720 and University of California, Davis,  
California 95616 and University of California Intercampus Institute for Research at Particle  
Accelerators, Stanford, California 94305 and University of California, Los Angeles, California  
90024 and University of California, Riverside, California 92521 and University of California,  
San Diego, California 92093 and University of California, Santa Barbara, California 93106  
and Ames Laboratory, Iowa State University, Ames, Iowa 50011 and Johns Hopkins University,  
Baltimore, Maryland 21218 and University of Massachusetts, Amherst, Massachusetts 01003  
and National Institute for Nuclear and High Energy Physics, Amsterdam, The Netherlands  
and University of Tokyo, Tokyo, Japan*

## ABSTRACT

We have searched for the two-photon production of the  $\eta_c$ ,  $\chi_0$  and  $\chi_2$  charmonium states at the  $e^+e^-$  collider PEP in the channels  $\gamma\gamma \rightarrow K^\pm K_S^0 \pi^\mp$ ,  $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^-$ ,  $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^+ \pi^-$  and  $\gamma\gamma \rightarrow K^+ K^- K^+ K^-$ . We identify four  $\eta_c$  candidates in the  $K^+ K^- K^+ K^-$  channel on a negligible background; the one  $\phi\phi$  event among them implies a 95% C.L. lower limit for  $\Gamma_{\gamma\gamma}(\eta_c)$  of 1.7 keV. In the other channels we find no evidence for any of the three states. We establish 95% C.L. upper limits  $\Gamma_{\gamma\gamma}(\eta_c) < 15.5$  keV,  $\Gamma_{\gamma\gamma}(\chi_0) < 17.0$  keV and  $\Gamma_{\gamma\gamma}(\chi_2) < 4.2$  keV. From all channels combined, we obtain the value  $\Gamma_{\gamma\gamma}(\eta_c) = 6.4_{-3.4}^{+5.0}$  keV.

PACS numbers: 14.40.Gx, 13.25.+m, 13.60.Le

The two-photon widths  $\Gamma_{\gamma\gamma}$  of the  $\eta_c$ ,  $\chi_0$  and  $\chi_2$  are of interest for understanding the dynamics of charmonium states. In a non-relativistic potential model,<sup>[1]</sup> one considers the  $\eta_c$  as a  $c\bar{c}$   $^1S_0$  bound state with the same radial wave function as the  $J/\psi$ ; the electromagnetic width of the  $J/\psi$  then implies  $\Gamma_{\gamma\gamma}(\eta_c) \approx 6.7$  keV. Such a model also gives relations between electromagnetic and hadronic widths, leading to predictions of  $\Gamma_{\gamma\gamma}$  for the  $^3P_J$  states  $\chi_0$  and  $\chi_2$ , though with larger uncertainties. Calculations of  $\Gamma_{\gamma\gamma}$  have also been made in relativistic potential models<sup>[2]</sup> and in dispersion-relation models based upon QCD,<sup>[3,4]</sup> yielding values in the 3 to 6 keV range for  $\eta_c$ . Predictions for  $\chi_0$  and  $\chi_2$  have ranged from 4 to 8 keV and 1 to 2 keV, respectively.

The experimental observation of charmonium states in photon-photon collisions is difficult because the two-photon luminosity at these relatively high masses is low and because the charmonium states have no dominant decay modes. The decays are thought to proceed via two intermediate gluons; consequently, branching fractions in the most common hadronic final states are only a few percent.<sup>[5]</sup> Among photon-photon collision experiments, several<sup>[6,7,8]</sup> have obtained only upper limits on  $\Gamma_{\gamma\gamma}(\eta_c)$ , while the Pluto experiment<sup>[9]</sup> has reported a signal in the  $K^\pm K_S^0 \pi^\mp$  ( $K_S^0 \rightarrow \pi^+ \pi^-$ ) channel. A value has also been reported from the reaction  $p\bar{p} \rightarrow \gamma\gamma$  at the ISR.<sup>[10]</sup> Two-photon widths of  $\chi$ -states have been reported from radiative  $\psi'$  decay<sup>[11]</sup> and from  $p\bar{p}$  annihilation.<sup>[10]</sup>

We report here a measurement of the two-photon reaction  $e^+e^- \rightarrow e^+e^-R$  (where  $R$  stands for  $\eta_c$ ,  $\chi_0$ , or  $\chi_2$ ) with the TPC/Two-Gamma facility<sup>[12]</sup> at the SLAC  $e^+e^-$  storage ring PEP, with incident  $e^+$  and  $e^-$  energies of 14.5 GeV. Using the particle-identification capabilities of the Time Projection Chamber

(TPC), we have studied four final states:  $K^\pm K_S^0 \pi^\mp$  ( $K_S^0 \rightarrow \pi^+ \pi^-$ ),  $K^+ K^- \pi^+ \pi^-$ ,  $\pi^+ \pi^- \pi^+ \pi^-$ , and  $K^+ K^- K^+ K^-$ . The sums of known branching fractions<sup>[13,14]</sup> to these final states are 4.5%, 6.7%, and 4.1% for  $\eta_c$ ,  $\chi_0$  and  $\chi_2$ , respectively.

The data described here correspond to an integrated luminosity of  $69 \text{ pb}^{-1}$ . The TPC tracked charged particles in a 13.25 kG solenoidal magnetic field with a momentum resolution at large angles of  $(\sigma_p/p)^2 \approx (1.5\%)^2 + (0.65\% \times p)^2$ , with  $p$  in GeV. The TPC also sampled energy loss ( $dE/dx$ ) along particle trajectories, with a typical resolution of 3.5%. Two proportional-mode pole-tip calorimeters (PTC) and a hexagonal Geiger-mode calorimeter (HEX) covered polar angles above 260 mrad. For some events, the detection of a scattered  $e^\pm$  tagged one of the virtual photons (with four-momentum  $q_1$  or  $q_2$ ). These ‘tags’ were detected in two arrays of NaI crystals in the polar angle range 25 – 90 mrad and lead-scintillator shower counters from 100 – 180 mrad, and in two arrays of fifteen drift chamber planes in front of these detectors.

The trigger required at least two charged tracks in the TPC, each with polar angle  $\theta > 26^\circ$  and projecting back to the interaction point within 20 cm along the beam axis. Tracks with  $\theta > 45^\circ$  were required to be in coincidence with hits in a drift chamber outside the coil of the solenoid.

In this four-prong analysis, we selected events with two positively and two negatively charged tracks coming from the vertex. Two tracks were required to have  $\theta > 30^\circ$ . Minimum momenta of 100 and 300 MeV were required for pions and kaons, respectively, with momentum uncertainty less than 30%. For particle identification, the measured  $dE/dx$  (defined as the average of the lowest 65% of the samples) and momentum were compared to empirically-determined curves

for the various particle types ( $e, \mu, \pi, K, p$ ). A confidence level was calculated for each of the four final states of interest, and in addition for the background states  $p\bar{p}\pi^+\pi^-$ ,  $e^+e^-e^+e^-$ , and  $\pi^+\pi^-\gamma$ ,  $K^+K^-\gamma$  where the photon converts to an  $e^+e^-$  pair. We required the confidence level for a final state to be above 10%. Additional requirements were imposed on the differences between that confidence level and those for other possible final states. For the  $K^\pm K_S^0 \pi^\mp$  channel, two oppositely charged pions had to have an effective mass within 40 MeV of the  $K_S^0$  mass. Events with photon candidates having energy depositions larger than 150 MeV in the HEX or larger than 300 MeV in the PTC were rejected. We required the summed transverse momentum for the four charged particles and the tag (if present) to be less than 0.2 GeV.

To determine the acceptance, Monte Carlo events were generated according to the luminosity function for transversely polarized photons.<sup>[15]</sup> We used a  $\gamma\gamma$  cross-section based on the matrix element  $\sqrt{(q_1 q_2)^2 - q_1^2 q_2^2} F(q_1^2, q_2^2)$  for  $\eta_c$  production<sup>[16]</sup> and  $F(q_1^2, q_2^2)$  for  $\chi_0, \chi_2$  production<sup>[17]</sup>, where  $F^2(0,0) \propto \Gamma_{\gamma\gamma}$ . For the  $q_i^2$  dependence of  $F$ ,  $1/[(1 - q_1^2/m_{J/\psi}^2)(1 - q_2^2/m_{J/\psi}^2)]$  was assumed; this is numerically similar to predictions for the  $\eta_c$  from charmonium sum rules.<sup>[3]</sup> Generated events were processed through a detector simulation program (which included resolution effects, energy loss, multiple scattering, nuclear interactions in the detector materials and a trigger simulation) and then passed through the same cuts as the data. Overall hadronic final state acceptances ranged from 3 to 7%.

The mass spectrum of the selected  $K^+K^-K^+K^-$  events shown in Fig. 1 contains six entries. The mass resolution determined from errors on measured

quantities is 22 MeV, which agrees with Monte Carlo expectations. Four of these events have an effective mass within 1.5 standard deviations from the nominal<sup>[18]</sup>  $\eta_c$  mass of 2981 MeV; we ascribe the two events at 2903 and 3330 MeV to background. The average background in the mass range  $2981 \pm 33$  MeV is estimated to contribute about 0.1 event to the signal. The  $K^+K^-$  mass spectrum of the six  $K^+K^-K^+K^-$  events is shown in Fig. 2. Given a  $K^+K^-$  mass resolution of about 10 MeV, one of the  $\eta_c$  events can be classified as  $\phi\phi$ , two are  $\phi K^+K^-$ , and one is  $K^+K^-K^+K^-$  with no  $\phi$  content. . The two background events are both consistent with  $\phi K^+K^-$ . Two of the six events, including one  $\eta_c$  candidate, have a tag.

The observation of the  $\eta_c$  in the  $K^+K^-K^+K^-$  channel at present hardly constrains the value of  $\Gamma_{\gamma\gamma}(\eta_c)$ , since only the part of the  $\eta_c$  branching fraction to  $K^+K^-K^+K^-$  through the  $\phi\phi$  intermediate state is known. We have taken a weighted average of the measurements in Ref. 14, including systematic uncertainties, to compute a branching fraction  $B(\eta_c \rightarrow \phi\phi) = (0.39 \pm 0.14)\%$ . Based upon our single  $\eta_c \rightarrow \phi\phi$  event, we arrive at a 95% C.L. lower limit  $\Gamma_{\gamma\gamma}(\eta_c) > 1.7$  keV. The one event represents a value  $\Gamma_{\gamma\gamma}(\eta_c) = 39_{-28}^{+50}$  keV.<sup>[18]</sup>

We found no evidence for charmonium in the mass spectra of  $K^+K^-\pi^+\pi^-$ ,  $\pi^+\pi^-\pi^+\pi^-$  and  $K^\pm K_S^0 \pi^\mp$ . We also investigated the  $\rho^0\rho^0$  final state (with subsequent decay into  $\pi^+\pi^-\pi^+\pi^-$ ), for which an  $\eta_c$  branching fraction has been reported.<sup>[19]</sup> Upper limits on  $\Gamma_{\gamma\gamma}(R) \times B(R \rightarrow \text{finalstate})$  from the investigated final states were obtained by a Monte Carlo method as follows. We fitted each spectrum with a smooth continuum to describe the background. The number of events expected from the sum of the background and a signal calculated for

a given value of  $\Gamma_{\gamma\gamma} \times B$  in a 125 MeV mass bin centered at a state  $R$  was generated according to a Poisson distribution. Errors on acceptance (15%) and luminosity (10%) were included in the procedure by throwing each time a value for each of these according to a gaussian distribution, cut off on the low side at two standard deviations. The number of Monte Carlo events was then compared with the number of events in the corresponding mass bin in the data and a probability was obtained for the number of Monte Carlo events to exceed the number of data events for the assumed value of  $\Gamma_{\gamma\gamma} \times B$ . The procedure was repeated for different values of  $\Gamma_{\gamma\gamma} \times B$  so that a 95% C.L. upper or lower limit could be found. The values derived from the separate channels are shown in Table I.<sup>[20]</sup>

The sum of the mass spectra for the final states  $\phi\phi$ ,  $\pi^+\pi^-\pi^+\pi^-$ ,  $K^+K^-\pi^+\pi^-$  and  $K^\pm K_S^0 \pi^\mp$  is shown in Fig. 3. To extract a best value and upper limit for  $\Gamma_{\gamma\gamma}(\eta_c)$ , a likelihood technique was used to simultaneously treat these four final states. In this case the Monte Carlo procedure also took into account the uncertainties in the branching fractions (typically 30%). Care was taken to treat separately those uncertainties which are correlated between channels, including that in  $B(J/\psi \rightarrow \gamma\eta_c)$ <sup>[13]</sup>, which is a common ingredient in all values of  $\eta_c$  decay branching fractions. For  $K^\pm K_S^0 \pi^\mp$ , isospin conservation and the  $K_S^0 \rightarrow \pi^+\pi^-$  decay reduce the  $K\bar{K}\pi$  branching fraction by a factor of 0.23, while the  $\phi \rightarrow K^+K^-$  decay reduces the  $\phi\phi$  branching fraction by a factor of 0.25. We find a 95% C.L. upper limit  $\Gamma_{\gamma\gamma}(\eta_c) < 15.5 \text{ keV}$  and a central value  $\Gamma_{\gamma\gamma}(\eta_c) = 6.4_{-3.4}^{+5.0} \text{ keV}$ <sup>[18,22]</sup>. This is lower than Pluto's measured value of  $28 \pm 15 \text{ keV}$ <sup>[9,21]</sup> while it is in agreement with the value of  $4.3_{-3.7}^{+3.4} \pm 2.4 \text{ keV}$ <sup>[10]</sup> from  $p\bar{p}$  annihilation.

From the combined spectra of  $\pi^+\pi^-\pi^+\pi^-$  and  $K^+K^-\pi^+\pi^-$ , and the known branching fractions,<sup>[18]</sup> we find the 95% C.L. upper limits  $\Gamma_{\gamma\gamma}(\chi_0) < 17.0$  keV and  $\Gamma_{\gamma\gamma}(\chi_2) < 4.2$  keV. The upper limit on  $\Gamma_{\gamma\gamma}(\chi_0)$  is compatible with the value of  $4.0 \pm 2.8$  keV quoted in Ref. 11. The upper limit on  $\Gamma_{\gamma\gamma}(\chi_2)$  is consistent with the values of  $2.8 \pm 2.0$  and  $2.9^{+1.3}_{-1.0} \pm 1.7$  keV quoted in Refs. 10 and 11 respectively. Our findings on the three charmonium states are in accord with the predictions of potential models<sup>[1,2]</sup> and the dispersion relation approach.<sup>[8,4]</sup>

We thank the PEP staff for their dedication and productive running of the machine, and our engineers and technicians for their efforts in the construction and maintenance of the detector. This work was supported in part by the United States Department of Energy, the National Science Foundation, the Joint Japan-United States Collaboration in High Energy Physics, and the Foundation for Fundamental Research on Matter in the Netherlands.

## REFERENCES

- 1) J.D. Jackson, Proceedings of the Summer Institute on Particle Physics, SLAC Rep. **198**, Nov. 1976, p.147.
- 2) L. Bergstrom, H. Snellman and G. Tengstrand, Phys. Lett. **82B**, 419 (1979).
- 3) R. Kirschner and A. Schiller, Z. Phys. **C16**, 141 (1982).
- 4) M.A. Shifman, Z. Phys. **C4**, 345 (1980) and (erratum) Z. Phys.**C6**, 282 (1980); L.J. Reinders, H.R. Rubinstein and S. Yazaki, Phys. Lett. **113B**, 411 (1982); V.A. Novikov et al., Phys. Lett. **67B**, 409 (1977) and Phys. Rep. **41C**, 1 (1978).
- 5) C. Quigg and J.L. Rosner, Phys. Rev. **D16**, 1497 (1977).
- 6) J.E. Olsson, Proceedings of the 5th Int. Workshop on  $\gamma\gamma$  Interactions, edited by Ch. Berger, Lecture Notes in Physics, Vol. 191, (Springer Verlag, Aachen, 1983), p.45; M. Althoff et al., Z. Phys. **C29**, 189 (1985).
- 7) H. Aihara et al., "Production of Four-Prong Final States in Photon-Photon Collisions", to be published in Physical Review D. These results are consistent with the present ones. They are based on data with an integrated luminosity comparable to that of the present set, but with a smaller acceptance and significantly worse mass resolution, mainly due to the lower magnetic field (3.9 kG).
- 8) A.E. Blinov et al., contribution to the 23rd International Conference on High Energy Physics, Berkeley (July 1986); G. Gidal et al., Proceedings of the 23rd International Conference on High Energy Physics, edited by S.C. Loken (World Scientific, Berkeley, 1987), p. 1220.

- 9) Ch. Berger et al., Phys. Lett. **167B**, 120 (1986). We have used a more recent determination of the  $B(\eta_c \rightarrow K^\pm \pi^\mp K_S^0 (K_S^0 \rightarrow \pi^+ \pi^-))$  to update their value for  $\Gamma_{\gamma\gamma}(\eta_c)$ .
- 10) C. Baglin et al., Phys. Lett. **187B**, 191 (1987).
- 11) R.A. Lee, "Radiative Decays of the Psi Prime to All-Photon Final States", Ph.D. Thesis, **SLAC-282** (1985).
- 12) H. Aihara et al., I.E.E.E. Trans. Nucl. Sci. **NS 30**, 63, 67, 76, 117, 153, and 162 (1983); M.P. Cain et al., Phys. Lett. **147B**, 232 (1984).
- 13) Review of Particle Properties, M. Aguilar-Benitez et al., Phys. Lett. **170B** (1986).
- 14) R.M. Baltrusaitis et al., Phys. Rev. Lett. **52**, 2126 (1984); D. Bisello et al., Phys. Lett. **179B**, 289 (1986). Only the branching fraction  $B(J/\psi \rightarrow \gamma \eta_c \rightarrow \gamma \phi \phi)$  is taken from these references.
- 15) V.M. Budnev et al., Phys. Rep. **15C**, 181 (1975).
- 16) G. Köpp et al., Nucl. Phys. **B70**, 461 (1974).
- 17) M. Poppe, Int. J. Mod. Phys. **1**, 545 (1986); we have assumed helicity two production for  $\chi_2$ .
- 18) The errors represent 84% C.L. upper and lower limits, chosen to resemble one standard deviation Gaussian errors.
- 19) B. Jean-Marie, Proceedings of the 23rd International Conference on High Energy Physics, op.cit., p. 689. We have used the  $\rho^0 \rho^0$  selection algorithm from this reference.

- 20) The event selection procedure was insensitive to the possible presence of low energy photons. No evidence for such photons was found in the  $\phi\phi$  event. Low energy calorimeter signals in the  $\phi K^+ K^-$  events are compatible with the effects from secondary interactions by charged particles in the calorimeters. However, because a calorimeter signal in the  $K^+ K^- K^+ K^-$  event cannot easily be explained in this manner, we quote only an upper limit on  $\Gamma_{\gamma\gamma}(\eta_c)$  for this channel.
- 21) From Table I, our 95% C.L. upper limit on  $\Gamma_{\gamma\gamma}(\eta_c) \times B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$  is 0.33 keV, compared to Pluto's measurement of  $0.5_{-0.15}^{+0.2} \pm 0.1$  keV.
- 22) The combined analysis implies a 95% C.L. lower limit of  $\approx 1.5$  keV.


**Table I**

Values and 95% C.L. upper limits on  
 $\Gamma_{\gamma\gamma}(R) \times B(R \rightarrow \text{final state})$  in keV.  
(See Ref. 18 for meaning of errors.)

final state	$\eta_c(2981)$	$\chi_0(3415)$	$\chi_2(3555)$
$\phi\phi$	$0.15^{+0.20}_{-0.11}$	$< 0.62$	$< 0.14$
$\phi K^+ K^-$	$0.36^{+0.25}_{-0.19}$	$< 0.52$	$< 0.12$
$K^+ K^- K^+ K^-$	$< 0.29$	$< 0.27$	$< 0.06$
$\pi^+ \pi^- \pi^+ \pi^-$	$< 0.29$	$< 0.71$	$< 0.11$
$\rho^0 \rho^0$	$< 0.20$	$< 0.45$	$< 0.11$
$K^\pm K_S^0 \pi^\mp$	$< 0.33$	$< 0.33$	$< 0.15$
$K^+ K^- \pi^+ \pi^-$	$< 0.30$	$< 0.60$	$< 0.11$

## FIGURE CAPTIONS

1. Effective mass of the  $K^+K^-K^+K^-$  events. The bin size (25 MeV) is approximately equal to the mass resolution.
2. Effective  $K^+K^-$  mass of the six  $K^+K^-K^+K^-$  events. The histogram has four and the scatter plot has two entries per event. The closed circles (shaded histogram) denote combinations from the four  $\eta_c$  events while the open circles (unshaded histogram) are from the two background events.
3. Effective mass of the  $\phi\phi \rightarrow K^+K^-K^+K^-$ ,  $K^\pm K_S^0 \pi^\mp (K_S^0 \rightarrow \pi^+\pi^-)$ ,  $K^+K^-\pi^+\pi^-$  and  $\pi^+\pi^-\pi^+\pi^-$  events combined. The curve is the sum of the Monte Carlo generated continuum and the expected signal for  $\Gamma_{\gamma\gamma}(\eta_c) = 15.5 \text{ keV}$ .



10-87

$M_{K^+K^-K^+K^-}$  (GeV)

5887A1

**FIGURE 1**





FIGURE 2



10-87

5887A3

FIGURE 3