

READINGS OF THE LICHNEROWICZ–YORK EQUATION

NIALL Ó MURCHADHA

Physics Department, University College, Cork, Ireland
email: `niall@ucc.ie`

(Received November 16, 2004)

*I would like to dedicate this article to Professor Staruszkiewicz
on the occasion of his 65th birthday*

James York, in a major extension of André Lichnerowicz's work, showed how to construct solutions to the constraint equations of general relativity. The York method consists of choosing a 3-metric on a given manifold; a divergence-free, tracefree (TT) symmetric 2-tensor wrt this metric; and a single number, the trace of the extrinsic curvature. One then obtains a quasi-linear elliptic equation for a scalar function, the Lichnerowicz–York (L–Y) equation. The solution of this equation is used as a conformal factor to transform the data into a set that satisfies the constraints. If the manifold is compact and without boundary, one quantity that emerges is the volume of the physical space. This article reinterprets the L–Y equation as an eigenvalue equation so as to get a set of data with a preset physical volume. One chooses the conformal metric, the TT tensor, and the physical volume, while regarding the trace of the extrinsic curvature as a free parameter. The resulting equation has extremely nice uniqueness and existence properties. A even more radical approach would be to fix the base (conformal) metric, the physical volume, and the trace. One also selects a TT tensor, but one is free to multiply it by a constant (unspecified). One then solves the L–Y equation as an eigenvalue equation for this constant. A third choice would be to fix the TT tensor and multiply the base metric by a constant. Each of these three formulations has good uniqueness and existence properties.

PACS numbers: 04.20.Cv, 04.20.Fy

1. Introduction

When general relativity is considered as a dynamical system, the Einstein equations, just like the Maxwell equations, split into constraint and evolution equations. In this article I focus entirely on the constraints. The initial data

consists of a pair (g_{ij}, K^{ij}) , defined on a given 3-manifold, where g_{ij} is a Riemannian 3-metric and K^{ij} is a symmetric tensor. K^{ij} is the extrinsic curvature of the 3-manifold when regarded as an embedded hypersurface of a pseudo-Riemannian 4-manifold that satisfies the Einstein equations [1].

The initial data cannot be specified freely: they must satisfy both a scalar and vector equation, known respectively as the Hamiltonian and momentum constraints, *i.e.*,

$$R - K_{ij}K^{ij} + K^2 = 0, \quad (1)$$

$$\nabla_i(K^{ij} - g^{ij}K) = 0, \quad (2)$$

where R is the scalar curvature and $K = g_{ij}K^{ij}$ is the trace of the extrinsic curvature. This trace K will play a key role in the rest of this article.

I (implicitly) define the extrinsic curvature by writing the relation between the extrinsic curvature and the time derivative of the metric as

$$\partial_t g_{ij} = 2NK_{ij} + \nabla_i N_j + \nabla_j N_i, \quad (3)$$

where N is the lapse and N^i is the shift. This means that my sign convention for K^{ij} agrees with Wald [3] and is the opposite of Misner, Thorne, Wheeler [2]. In particular

$$\frac{1}{\sqrt{g}}\partial_t \sqrt{g} = NK + \nabla_i N^i. \quad (4)$$

The first significant step in understanding how to solve the constraints was achieved by André Lichnerowicz in 1944 [4]. He realised that if the extrinsic curvature was tracefree, then the momentum constraint, Eq. (2), required that the extrinsic curvature be both divergence-free and tracefree (described as transverse tracefree, hence TT). TT tensors are conformally covariant. Given a TT tensor wrt a given metric, on making the following transformations

$$\overline{g_{ij}} = \phi^4 g_{ij}, \quad \overline{K_{TT}^{ij}} = \phi^{-10} K_{TT}^{ij}, \quad (5)$$

then $\overline{K_{TT}^{ij}}$ is TT wrt the transformed metric $\overline{g_{ij}}$. Thus it continues to satisfy the tracefree momentum constraint. Lichnerowicz had the idea of choosing a base metric and a TT tensor on it, and making a conformal transformation so that the transformed metric would satisfy the Hamiltonian constraint, *i.e.*,

$$\overline{R} - \overline{K_{TT}^{ij}} \overline{K_{ij}^{TT}} = 0. \quad (6)$$

This reduces to the well-known Lichnerowicz equation for the conformal factor ϕ

$$8\nabla^2 \phi - R\phi + K_{TT}^{ij} K_{ij}^{TT} \phi^{-7} = 0. \quad (7)$$

Of course, since the solution ϕ is to be used as a conformal factor, one is only interested in solutions which are everywhere positive.

It turns out that for either the asymptotically flat case or for the compact, without boundary, case, the Lichnerowicz equation can only be solved for a restricted class of data. The choice of TT data is essentially irrelevant, but the choice of the base metric is important. The ‘tracefree’ Hamiltonian constraint Eq. (6) requires the solution scalar curvature to be everywhere nonnegative. There exists a conformal invariant (the Yamabe constant) [5]

$$Y = \inf \frac{\int R\theta^2 + 8(\nabla\theta)^2 dv}{[\int \theta^6 dv]^{\frac{1}{3}}}, \quad (8)$$

where the infimum is taken over smooth functions in the compact case and over smooth functions of compact support in the asymptotically flat case. The minimizing function solves the Yamabe equation, and, when used as a conformal factor, transforms the manifold into a compact one with constant scalar curvature. On such a manifold, the minimizing function can be chosen equal to 1, and then we get

$$Y = \frac{R_0 V_0}{V_0^{\frac{1}{3}}} = R_0 V_0^{\frac{2}{3}}. \quad (9)$$

A given metric can be conformally transformed into a metric with positive scalar curvature if and only if the Yamabe constant is positive. As a result, the Lichnerowicz equation can be solved if and only if the metric belongs to the positive Yamabe class.

In 1970, James York [6] extended the Lichnerowicz approach significantly when he realised that one could add a constant trace term, K , to the extrinsic curvature without disturbing the conformal covariance of the momentum constraint, *i.e.*,

$$\overline{g_{ij}} = \phi^4 g_{ij}, \quad \overline{K^{ij}} = \phi^{-10} K_{TT}^{ij} + \frac{1}{3} K \overline{g^{ij}}. \quad (10)$$

Note that the constant trace term is a conformal invariant! The equation for the conformal factor becomes the Lichnerowicz–York equation

$$8\nabla^2\phi - R\phi + K_{TT}^{ij}K_{ij}^{TT}\phi^{-7} - \frac{2}{3}K^2\phi^5 = 0. \quad (11)$$

The Hamiltonian constraint no longer places any restriction on the sign of the scalar curvature when the trace of the extrinsic curvature is nonzero. Therefore the Lichnerowicz–York equation can almost invariably be solved with any choice of base metric, any choice of TT tensor, and any choice of K .

It works both in the compact and the asymptotically flat cases. The initial data that is constructed has constant trace of the extrinsic curvature and is thus called a constant mean curvature (CMC) solution to the constraints.

The standard method is quite straightforward: one chooses the base metric, a TT tensor, and a value of K ; one then solves for the conformal factor and constructs the physical metric; given the physical metric, one can find the physical volume. The aim of this article, however, is to show that the ‘standard’ interpretation of the Lichnerowicz–York equation, at least in the compact case, is not the only reasonable one. It turns out that one can trade off knowledge of K for a specification of the physical volume ($V_p = \int \phi^6 dv$). This means reading Eq. (11) as an eigenvalue equation for K by putting a normalization condition on the conformal factor so that the physical volume has a prespecified value.

There are even more radical interpretations. If one is given a tensor which is TT wrt some metric, then one can multiply it by any constant and it remains TT . Now solve the Lichnerowicz–York equation, specifying the base metric, the TT tensor defined up to a multiplicative constant, the value of K , and the physical volume. The freedom to choose the factor multiplying the TT tensor means that this is also a wellposed system.

Finally, one could fix the TT tensor, but allow the base metric be defined up to a multiplicative constant. Again, both K and V can be fixed and the constant regarded as an eigenvalue. These three options will be discussed further.

2. Trading off K for V_p

As part of his analysis of the constraints, James York [6] suggested that conformal superspace — the space of Riemannian 3-metrics, modulo diffeomorphisms, and conformal transformations [8] — should be regarded as the natural configuration space of GR. Recently, a number of us [9] have found an action on conformal superspace which generates GR in the CMC gauge. A major difference here is that we need to specify the physical volume as part of our initial data, while the value of K emerges as part of the solution. An obvious question is whether this switch loses us the extremely nice existence and uniqueness properties of the original form of the Lichnerowicz–York equation. The answer, as revealed here, is that we lose essentially nothing.

As can be seen from Eq. (4), and as previously pointed out by York [6], K and the volume can be regarded as canonically conjugate variables; therefore switching from K to V_p should be viewed as a Legendre transformation, and so should not cause any major disruption of the system. In [9] we write down an action, so we are working in the Lagrangian rather than the

Hamiltonian framework. Therefore it seems more natural to use the volume (a metric quantity) rather than specifying K , which is part of the conjugate momentum. Nevertheless, these sorts of arguments can only be regarded as poor substitutes for showing that the equation itself is well behaved after the switch from K to V_p .

What we need to do is to fix the (conformal) metric and the TT tensor, and to allow K^2 to span the entire range from 0 to ∞ , solving the Lichnerowicz–York equation for each value of K^2 . We should then investigate what happens to the physical volume as a function of K^2 . Note that the sign of K does not matter in the Lichnerowicz–York equation: only K^2 appears. The sign plays a role only in the evolution, when we determine whether the universe either expands or contracts to the future. Let us start with a perturbation calculation.

We begin with a solution to the constraints, *i.e.*, we are given a metric, a TT tensor, and a constant K , which satisfy

$$R - K_{ij}^{TT} K_{TT}^{ij} + \frac{2}{3} K^2 = 0. \quad (12)$$

We perturb this data by changing only K^2 , by an amount δK^2 , and solve the perturbed Lichnerowicz–York equation (remember we are perturbing around $\phi \equiv 1$)

$$8\nabla^2 \delta\phi - R\delta\phi - 7K_{TT}^{ij} K_{ij}^{TT} \delta\phi - \frac{10}{3} K^2 \delta\phi = \frac{2}{3} \delta K^2. \quad (13)$$

This is an inhomogeneous linear elliptic equation for $\delta\phi$. We can simplify it by eliminating R using Eq. (12) to get

$$8\nabla^2 \delta\phi - \left(8K_{TT}^{ij} K_{ij}^{TT} + \frac{8}{3} K^2 \right) \delta\phi = \frac{2}{3} \delta K^2. \quad (14)$$

This is a particularly nice elliptic equation, because the coefficient of the linear term is everywhere negative, which, in turn, guarantees that a unique solution to the inhomogeneous equation exists. This should not come as a surprise, because we know that the nonlinear equation always has a regular solution. What is interesting, however, is that we can use the maximum principle to control the sign of $\delta\phi$. Let us assume that δK^2 is positive. If $\delta\phi$ were positive in any region, we would have a point where $\delta\phi$ has a positive maximum. However, at a positive maximum, $\nabla^2 \delta\phi$ is nonpositive, and the linear term is negative, which is incompatible with a positive right hand side. Therefore, we know that $\delta\phi < 0$. This means that as K^2 increases ϕ decreases at each point, and the physical volume monotonically decreases. The mapping between V_p and K^2 is one to one.

We have a fixed base metric, a fixed TT tensor, and a K^2 which can become either unboundedly large or go to zero. The next question to ask

is what happens when K^2 becomes large. It can be shown that ϕ shrinks monotonically to 0 everywhere, and the physical volume vanishes. The easiest way to see this is to consider the alternative. For each choice of K^2 we solve for ϕ , which we know decreases. Let us assume that this ϕ does not go to zero everywhere in the limit. In the region where ϕ is bounded away from zero, the physical scalar curvature and the physical TT tensor remain bounded, while K^2 blows up. This is not compatible with solving the Hamiltonian constraint. Therefore ϕ shrinks everywhere to zero, and the physical volume does likewise.

We can do better. For ease of computation let us first make a conformal transformation so that the base metric has constant scalar curvature. The Lichnerowicz–York equation then becomes

$$8\nabla^2\phi - R_0\phi + K_{TT}^{ij}K_{ij}^{TT}\phi^{-7} - \frac{2}{3}K^2\phi^5 = 0. \quad (15)$$

At the maximum of ϕ we have $\nabla^2\phi \leq 0$. At this point we have

$$K_{TT}^{ij}K_{ij}^{TT} \geq R_0\phi_{\max}^8 + \frac{2}{3}K^2\phi_{\max}^{12}. \quad (16)$$

The $K_{TT}^{ij}K_{ij}^{TT}$ term is evaluated at the point where ϕ reaches its maximum. We can replace this in inequality (16) by the maximum value over the whole manifold to give

$$K_{TT}^{ij}K_{ij}^{TT}(\max) \geq R_0\phi_{\max}^8 + \frac{2}{3}K^2\phi_{\max}^{12}. \quad (17)$$

This means that as K^2 increases $\phi_{\max}$ must decrease, and therefore the physical volume shrinks to zero.

In the nonnegative Yamabe classes, where $R_0 \geq 0$, one can just discard the R_0 term in inequality (17) to get

$$K_{TT}^{ij}K_{ij}^{TT}(\max) \geq \frac{2}{3}K^2\phi_{\max}^{12}. \quad (18)$$

From this one gets

$$V_p = \int \phi^6 dv \leq \phi_{\max}^6 V_0 = \sqrt{\frac{3K_{TT}^{ij}K_{ij}^{TT}(\max)}{2K^2}} V_0. \quad (19)$$

This estimate holds always in the positive and zero Yamabe classes; in particular, it shows how the volume goes to zero as K^2 becomes large. In the negative Yamabe class such an estimate will continue to hold, but only

in the large K^2 limit. This can be seen by looking at the polynomial on the right hand side of (17) assuming $R_0 = -C^2$.

The final question is what happens when K^2 shrinks to zero. While we know that the conformal factor increases and drags the volume with it, we have no guarantee that the volume becomes infinitely large. The Lichnerowicz–York equation, when K^2 equals zero, reduces to the original Lichnerowicz equation. Therefore we know that the Yamabe constant of the base metric must be the determining factor. If the base metric is in the positive Yamabe class, then the Lichnerowicz equation has a regular solution with finite physical volume, and a maximal slice emerges. This will form the limiting solution to the sequence of Lichnerowicz–York equations. The volume of the maximal solution is the largest volume we can specify. It, of course, depends both on the choice of base metric and on the specified TT tensor.

If the Yamabe class of the base metric is nonpositive, then no regular solution to the Lichnerowicz equation exists. The conformal factor blows up as K^2 goes towards zero, and the physical volume becomes unboundedly large.

Let us consider the case where the Yamabe constant is negative. This means that one can make a conformal transformation to a metric where the scalar curvature is constant and negative, *i.e.*, $R_0 = -C = \text{constant}$. One could pick the constant to equal -1 , but this is not important. This base metric will have total volume V_0 . As usual, one solves the Lichnerowicz–York equation for the conformal factor ϕ , and calculates the physical volume V_p . I will show that

$$V_p > \left(\frac{3C}{2K^2} \right)^{\frac{3}{2}} V_0 = \left(\frac{3|Y|}{2K^2} \right)^{\frac{3}{2}}. \quad (20)$$

The equality in Eq. (20) follows from the definition of the Yamabe constant, Eq. (9). This guarantees that the volume diverges as $K^2 \rightarrow 0$.

Start with a set of data, a metric with constant negative scalar curvature, a TT tensor, and K^2 . First make a conformal transformation with a constant conformal factor $\phi_0 = (3C/2K^2)^{1/4}$. This transforms the base manifold to one where the scalar curvature $\bar{R} = -2K^2/3$ and the volume $\bar{V} = (3C/2K^2)^{3/2}V_0$. The Lichnerowicz–York equation in this frame becomes

$$8\bar{\nabla}^2\bar{\phi} + \frac{2}{3}K^2\bar{\phi} + \overline{K^{TT}K_{TT}}\bar{\phi}^{-7} - \frac{2}{3}K^2\bar{\phi}^5 = 0, \quad (21)$$

where $\overline{K^{TT}K_{TT}} = K^{TT}K_{TT}(2K^2/3C)^3$. From the min-max principle we know that at the minimum of $\bar{\phi}$ we have $\bar{\nabla}^2\bar{\phi} \geq 0$. Therefore we have

$$\frac{2}{3}K^2(\bar{\phi}_{\min}^5 - \bar{\phi}_{\min}) - \overline{K^{TT}K_{TT}}\bar{\phi}_{\min}^{-7} \geq 0. \quad (22)$$

This guarantees that $\bar{\phi}_{\min} \geq 1$ and thus $\bar{\phi} \geq 1$. In turn, this means that the physical volume is greater than $\bar{V} = (3C/2K^2)^{3/2}V_0$.

We can do somewhat better than that by looking at the maximum of $\bar{\phi}$. In this case we have

$$\frac{2}{3}K^2(\bar{\phi}_{\max}^5 - \bar{\phi}_{\max}) - \overline{K^{TT}K_{TT}}\bar{\phi}_{\max}^{-7} \leq 0. \quad (23)$$

This can be rearranged to give

$$\bar{\phi}_{\max}^7(\bar{\phi}_{\max}^5 - \bar{\phi}_{\max}) \leq \frac{3}{2}\overline{K^{TT}K_{TT}}/K^2 = K^{TT}K_{TT}(2K^2/3)^2 \frac{1}{C^3}. \quad (24)$$

This tells us that $\bar{\phi}_{\max}$ cannot be very large. As $K^2 \rightarrow 0$, the right hand side of Eq. (24) shrinks to zero and $\bar{\phi}_{\max}$ is pushed down to 1. Thus the volume of the physical space converges to the volume of $\bar{V} = (3C/2K^2)^{3/2}V_0 = (3Y/2K^2)^{3/2}$. Another way of saying this is that in the limit, as K becomes small, the conformal factor becomes large, which renders the TT tensor unimportant. The physical geometry then more and more approximates a large-volume 3-manifold with constant negative scalar curvature $\bar{R} = -2K^2/3$. An alternative way of expressing this is that $V_p K^3 \approx (3Y/2)^{3/2} = \text{constant}$.

In summary: If the base metric is in the positive Yamabe class, and if we pick any physical volume which is less than or equal to the volume of the maximal solution, there exists a unique choice of K^2 , which, on solving the Lichnerowicz–York equation, generates a solution to the constraints with the chosen volume. If the base metric has nonpositive Yamabe constant, one can choose any volume between 0 and ∞ ; then there exists a unique choice of K^2 which delivers this volume, on solving the Lichnerowicz–York equation.

3. Other choices

In addition to trading off between the physical volume and the value of K , there are several other specifications of data for the Lichnerowicz–York equation that work. Given a base metric, a TT tensor, and a constant K , one can find a complete family of TT tensors by simply multiplying the given tensor by a constant. Then, for each choice of the constant, one can solve the Lichnerowicz–York equation and find the physical volume. One can specify the volume and solve for the constant multiplying the TT tensor.

3.1. Scaling the TT tensor

We start off with a modified Lichnerowicz–York equation

$$8\nabla^2\phi - R\phi + \alpha^2 K_{TT}^{ij}K_{ij}^{TT}\phi^{-7} - \frac{2}{3}K^2\phi^5 = 0, \quad (25)$$

and we let α^2 range from 0 to ∞ . We now look for the relationship between the physical volume and the value of α^2 . Let us again perform a perturbation calculation, *i.e.*, we start with a solution to the Hamiltonian constraint

$$R - K_{ij}^{TT} K_{TT}^{ij} + \frac{2}{3} K^2 = 0, \quad (26)$$

and perturb Eq. (25) around it, holding the metric, the TT tensor, and K^2 fixed. We only perturb α^2 and consider the change of ϕ away from $\phi \equiv 1$. We get

$$8\nabla^2 \delta\phi - R\delta\phi - 7K_{TT}^{ij} K_{ij}^{TT} \delta\phi - \frac{10}{3} K^2 \delta\phi = -\delta\alpha^2 K_{TT}^{ij} K_{ij}^{TT}. \quad (27)$$

By using Eq. (26) to eliminate R , we can simplify this equation to get

$$8\nabla^2 \delta\phi - \left(8K_{TT}^{ij} K_{ij}^{TT} + \frac{8}{3} K^2 \right) \delta\phi = -\delta\alpha^2 K_{TT}^{ij} K_{ij}^{TT}. \quad (28)$$

We can again use the min-max principle on this equation and if $\delta\alpha^2 > 0$ we have $\delta\phi > 0$. In this case, as distinct from the ‘changing K^2 ’ case, as we increase the K^{TT} term the volume monotonically increases. This guarantees uniqueness.

In order to investigate whether there is any restriction on the allowed range of the volume, we need to look at what happens at the two extremes, *i.e.*, when $\alpha^2 \rightarrow \infty$ and $\alpha^2 \rightarrow 0$. Let us start with the case where $\alpha^2 \rightarrow \infty$. The Hamiltonian constraint, after solving the equation, is

$$\overline{R} - \alpha^2 \phi^{-12} K_{ij}^{TT} K_{TT}^{ij} + \frac{2}{3} K^2 = 0. \quad (29)$$

If ϕ were to remain bounded on the support of K^{TT} as $\alpha^2 \rightarrow \infty$, the central term in Eq. (29) blows up while the other two terms remain finite. This cannot happen; therefore ϕ must blow up on the support of K^{TT} , and the volume of the physical space becomes unboundedly large. This happens with any choice of base metric.

The other extreme is when $\alpha^2 \rightarrow 0$. In this case the TT term drops out of the Hamiltonian constraint, and we get

$$\overline{R} + \frac{2}{3} K^2 = 0. \quad (30)$$

This is a manifold with constant negative scalar curvature, and this is only achievable if the Yamabe constant of the base metric is negative.

Therefore if the base metric is in the negative Yamabe class, the conformal factor tends to a finite limit and generates a manifold of constant

negative scalar curvature. This manifold of constant negative scalar curvature is the manifold of least volume in the sequence, and any physical volume we pick must be at least as large as this. We can find an expression for this minimum volume easily in terms of the Yamabe constant. On a manifold of constant curvature the Yamabe constant is achieved with a constant function. Therefore we have

$$\frac{\overline{R}V_p}{V_p^{\frac{1}{3}}} = -\frac{2K^2}{3}V_p^{\frac{2}{3}} = Y. \quad (31)$$

From this we get

$$V_p = \left(\frac{3|Y|}{2K^2} \right)^{\frac{3}{2}}. \quad (32)$$

This is the minimum volume we can specify if the Yamabe constant is negative.

It is clear, from Eq. (32), that this minimum volume shrinks to zero as $|Y| \rightarrow 0$. We can, however, do somewhat better. Let us assume that we are given a base metric in the positive Yamabe class. Let us make a conformal transformation so as to map the base metric to one with constant positive scalar curvature. Let us further arrange that this constant $R_0 = 2K^2/3$. This is not really necessary, but it makes the algebra somewhat easier. The L-Y equation now becomes

$$8\nabla^2\phi - \frac{2}{3}K^2\phi + \alpha^2 K^{TT} K_{TT}\phi^{-7} - \frac{2}{3}K^2\phi^5 = 0. \quad (33)$$

At the maximum of ϕ we have that $\nabla^2\phi \leq 0$. This gives

$$\alpha^2 K^{TT} K_{TT} \geq \frac{2}{3}K^2(\phi_{\max}^8 + \phi_{\max}^{12}), \quad (34)$$

where, of course, the left hand side of Eq. (24) is evaluated at the point where ϕ achieves its maximum. First, we see that ϕ achieves its maximum on the support of K^{TT} . Further, we can get an upper bound for $\phi_{\max}$ in terms of the maximum of $K^{TT} K_{TT}$ from

$$\frac{3\alpha^2}{2K^2}(K^{TT} K_{TT})_{\max} \geq \phi_{\max}^8. \quad (35)$$

This is not in any way sharp. A ‘better’ estimate can be obtained by turning inequality (34) into a cubic equation. This, however, offers no real improvement, because we are really interested in the case when $\alpha \rightarrow 0$; since then $\phi_{\max} \rightarrow 0$ and the ϕ^{12} term can be neglected wrt the ϕ^8 term.

From (35) we get the following estimate for the physical volume

$$V_p = \int \phi^6 dv \leq \phi_{\max}^6 \int dv = \phi_{\max}^6 V_0 \leq \left[\frac{3\alpha^2}{2K^2} (K^{TT} K_{TT})_{\max} \right]^{\frac{3}{4}} V_0, \quad (36)$$

where V_0 is the volume of the base, constant curvature, metric. This shows that V_p goes to zero as α goes to zero like $\alpha^{3/2}$. One could replace V_0 in Eq. (36) by the Yamabe constant to give

$$V_p \leq \alpha^{\frac{3}{2}} \left[\frac{3}{2K^2} \right]^{\frac{9}{4}} [(K^{TT} K_{TT})_{\max}]^{\frac{3}{4}} Y^{\frac{3}{2}}. \quad (37)$$

3.2. Scaling the base metric

A third way of interpreting the Lichnerowicz–York equation is to choose a base metric g_b , a TT tensor, a physical volume V_p , and the trace of the extrinsic curvature K , but to assume the base metric is defined only up to a multiplicative constant $g'_b = \beta^4 g_b$. The given TT tensor will still be TT wrt the adjusted metric. The value of β is a freely chosen parameter.

The behaviour of this system exactly mirrors the second system analysed, where the TT tensor is scaled. This is because the following global scaling

$$(g_b, K_{ij}^{TT}, K) \rightarrow (\beta^4 g_b, \beta^{-2} K_{ij}^{TT}, K) \quad (38)$$

leaves the physical metric, and also the physical volume, unchanged. Therefore there is a tradeoff between scaling on g_b and a scaling of K^{TT} . We have the following chain

$$(g_b, K_{ij}^{TT}, K) \rightarrow (\beta^4 g_b, K_{ij}^{TT}, K) \rightarrow (g_b, \beta^2 K_{ij}^{TT}, K). \quad (39)$$

The first link is the nontrivial change we wish to analyse. The second link is an identity, which leaves V_p unchanged. Therefore scaling the metric by β^4 is exactly equivalent to scaling the TT tensor by β^2 .

This means that as $\beta \rightarrow \infty$, the physical volume monotonically increases and becomes unboundedly large. As $\beta \rightarrow 0$, the behaviour depends on the Yamabe number of the base metric. If the Yamabe number is negative the volume shrinks to a finite volume, represented by a manifold of constant negative scalar curvature, while if the Yamabe number is nonnegative, the manifold shrinks to zero volume. The estimates given above in Subsection 3.1 continue to hold, except that α has to be replaced by β^2 .

For many years I have been indebted to the relativity group of the Institute of Theoretical Physics of the Jagellonian University. Since the first of my many visits, in 1987, I have found the Institute and its members to be welcoming and friendly as well as providing a stimulating place in which to work. Much of the credit for maintaining this environment is surely due to its director and senior academic, Professor Andrzej Staruszkiewicz. I would like to dedicate this article to Professor Staruszkiewicz on the occasion of his 65th birthday.

REFERENCES

- [1] R. Arnowitt, S. Deser, C. Misner in *Gravitation: an Introduction to Current Research*, ed. L. Witten, Wiley, New York, 1962 [[arXiv:gr-qc/0405109](#)].
- [2] C. Misner, K. Thorne, J.A. Wheeler, *Gravitation*, Freeman, San Francisco 1973.
- [3] R. Wald, *General Relativity*, U. Chicago Press, Chicago 1984.
- [4] A. Lichnerowicz, *J. Math. Pures Appl.* **23**, 37 (1944).
- [5] H. Yamabe, *Osaka Math. J.* **12**, 21 (1960); R. Schoen, *J. Diff. Geom.* **20**, 479 (1984); J.M. Lee, T.H. Parker, *Bull. Am. Math. Soc.* **17**, 37 (1987).
- [6] J.W. York, *Phys. Rev. Lett.* **26**, 1656 (1971); J.W. York, *Phys. Rev. Lett.* **28**, 1082 (1972); J.W. York, *J. Math. Phys.* **14**, 456 (1973).
- [7] N. Ó Murchadha, J.W. York, *J. Math. Phys.* **14**, 1551 (1972).
- [8] J.W. York, *Ann. Inst. Henri Poincaré* **21**, 319 (1974); A.E. Fischer, V. Moncrief, *Gen. Rel. Grav.* **28**, 221 (1996).
- [9] E. Anderson, J. Barbour, B.Z. Foster, B. Kelleher, N. Ó Murchadha, [arXiv:gr-qc/0407104](#).