

physics



Article

New Insights into the Lamb Shift: The Spectral Density of the Shift

G. Jordan Maclay

Special Issue

Vacuum Fluctuations

Edited by

Prof. Dr. G. Jordan Maclay and Prof. Dr. Roberto Passante



<https://doi.org/10.3390/physics4040081>

Article

New Insights into the Lamb Shift: The Spectral Density of the Shift

G. Jordan Maclay 

Quantum Fields LLC, 147 Hunt Club Drive, St. Charles, IL 60174, USA; jordanmaclay@quantumfields.com

Abstract: In an atom, the interaction of a bound electron with the vacuum fluctuations of the electromagnetic field leads to complex shifts in the energy levels of the electron, with the real part of the shift corresponding to a shift in the energy level and the imaginary part to the width of the energy level. The most celebrated radiative shift is the Lamb shift between the $2s_{1/2}$ and the $2p_{1/2}$ levels of the hydrogen atom. The measurement of this shift in 1947 by Willis Lamb Jr. proved that the prediction by Dirac theory that the energy levels were degenerate was incorrect. Hans Bethe's non-relativistic calculation of the shift using second-order perturbation theory demonstrated the renormalization process required to deal with the divergences plaguing the existing theories and led to the understanding that it was essential for theory to include interactions with the zero-point quantum vacuum field. This was the birth of modern quantum electrodynamics (QED). Numerous calculations of the Lamb shift followed including relativistic and covariant calculations, all of which contain a nonrelativistic contribution equal to that computed by Bethe. The semi-quantitative models for the radiative shift of Welton and Power, which were developed in an effort to demonstrate physical mechanisms by which vacuum fluctuations lead to the shift, are also considered here. This paper describes a calculation of the shift using a group theoretical approach which gives the shift as an integral over frequency of a function, which is called the "spectral density of the shift." The energy shift computed by group theory is equivalent to that derived by Bethe yet, unlike in other calculations of the non-relativistic radiative shift, no sum over a complete set of states is required. The spectral density, which is obtained by a relatively simple computation, reveals how different frequencies of vacuum fluctuations contribute to the total energy shift. The analysis shows, for example, that half the radiative shift for the ground state $1S$ level in H comes from virtual photon energies below 9700 eV, and that the expressions of Power and Welton have the correct high-frequency behavior, but not the correct low-frequency behavior, although they do give approximately the correct value for the total shift.

Keywords: Bethe; radiative shift; shift spectral density; spectral density; vacuum fluctuations; vacuum field; mass renormalization; Lamb shift; QED; radiative reaction; zero point fluctuations; hydrogen atom



Citation: Maclay, G.J. New Insights into the Lamb Shift: The Spectral Density of the Shift. *Physics* **2022**, *4*, 1253–1277. <https://doi.org/10.3390/physics4040081>

Received: 22 April 2022

Accepted: 31 August 2022

Published: 19 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In astronomy, in quantum theory, and in quantum electrodynamics (QED), there have been periods of great progress in which solutions to challenging problems have been obtained, and the fields have moved forward. However, in some cases getting the right answers can still leave fundamental questions unanswered. The Big Bang explained the origin of cosmic background radiation but left the problem of why the universe appears to be made of matter and not equal amounts of matter and antimatter [1]. In quantum theory, physicists can compute the behavior of atoms yet cannot describe a measurement in a self-consistent way [2], or make sense of the collapse of a photon wavefunction from a near infinite volume to a point [3]. In quantum electrodynamics, we can compute the Lamb shift of the H atom to 15 decimal places [4], yet we are left with the paradox of using perturbation theory to remove infinite terms or to understand a quantum vacuum with infinite energy. In this paper, several different approaches to the computation of the

non-relativistic Lamb shift are examined. For these approaches, the Lamb shift can be expressed in different ways as an integral over the frequency of a spectral density. This paper analyzes the differences in the non-relativistic spectral densities for the different approaches as a function of frequency and compares the spectral densities to those obtained using a group theoretical analysis. The integral of the spectral density over all frequencies gives the corresponding value of the non-relativistic Lamb shift.

Feynman called the three-page long 1947 non-relativistic Lamb shift calculation by Hans Bethe the most important calculation in quantum electrodynamics because it tamed the infinities plaguing earlier attempts [5,6]. When the sum over all states is evaluated numerically, it gives a finite prediction that agreed with experiment [7,8]. Dirac said it “fundamentally changed the nature of theoretical physics” [6]. Quoting the classic text of *Quantum Electrodynamics*, Volume 4 in the Landau and Lifshitz series on Theoretical Physics: “This work provided the initial stimulus for the whole subsequent development of quantum electrodynamics” [5]. Yet when this calculation is explored more deeply, questions arise about it and about other fundamentally different calculations of the Lamb shift, for example, those by Welton [9] and Power [10], that employ different conceptual approaches that have similar high-frequency but different low-frequency behavior from Bethe’s result yet give approximately the same value for the level shift [6]. These three approaches to the Lamb shift and the corresponding vacuum energy densities have also been considered in [11].

There is an intimate relationship between radiative shifts and vacuum fluctuations. The shift can be interpreted as arising from virtual transitions induced by the quantum fluctuations of the electromagnetic field. Since the vacuum field contains all frequencies, virtual transitions to all states, bound and scattering, are possible. These short-lived virtual transitions result in a slight shift in the average energy of the atom, a shift which is the Lamb shift [12]. The Lamb shift can also be described as an interaction of the electron with its own radiation field, yielding the same results as the vacuum field [6].

Shortly after Bethe’s calculation, Dyson published, as a problem assigned by Bethe, a calculation of the Lamb shift for a spinless electron [13]. Formal and rigorous relativistic perturbation theory calculations to first-order in the radiation field and to fourth order in the ratio of the velocity of the atomic electron to the velocity of light, based on the Dirac equation, including spin and relativistic effects, were carried out independently in 1949 by J. French and V. Weisskopf [14] and N. Kroll and W. Lamb [15]. They used a high-frequency cutoff for the non-relativistic contribution which also acted as a low-frequency cutoff for the relativistic portion, and when both contributions were added together, the sum was independent of the cutoff, and equaled a term corresponding to the Bethe log plus constants corresponding to relativistic corrections. Although these calculations were difficult and cumbersome, they have stood the test of time [16]. These relativistic calculations gave a value of 1052 MHz for the Lamb shift, compared to Bethe’s result of 1040 MHz, about a 1% difference. The determination of the second-order corrections for the Lamb shift is quite complicated. The most complete tabulation and systematic derivation was carried out by Erickson and Yennie [17]. Today the Lamb shift is computed precisely to over 10 decimal places to be about 1057 MHz, the largest error being due to the uncertainty in the radius of the proton [4].

At about the same time as the first relativistic calculations were published, Schwinger published a general covariant approach to quantum electrodynamics which he applied to the Lamb shift computation [18]. Within a year, three different approaches to quantum electrodynamics were independently developed that were relativistic and could deal with divergences with some success. Schwinger, Tomonaga, and Feynman each had proposed a manifestly covariant method, and shown its capability to address a broader range of QED problems than just the energy levels of the H atom [19]. Freeman Dyson showed that these three methods had essential similarities and were mutually consistent [20].

The classic text, *Quantum Electrodynamics*, Volume 4 of the Landau and Lifshitz series on Theoretical Physics, second edition published in 1979, does the Lamb shift calculation by dividing the vacuum energy spectrum into a low and high-frequency region with a

cutoff [5]. For the high-energy region, they use scattering theory based on Schwinger's approach. For the low-frequency region they use non-relativistic kinematics and start with the expression for dipole radiation, which they transform with some mathematical operations on delta function; then they integrate over frequency and obtain essentially the same result as Bethe plus constants corresponding to vacuum polarization and other corrections that are not considered here. When the high and low-frequency results are added, the cutoff cancels. The final result contains the Bethe log. A slightly different approach is taken but with similar results in [21]. Another important text, published in 1980 by Itzykson and Zuber, starts with a modified Dirac equation with a fictitious photon mass, and that includes vacuum polarization and vertex corrections for an electron in a slowly varying external field [22]. This equation yields a level shift with two terms that correspond to the high-energy region and low-energy region. After evaluation, the sum of the terms is taken, the fictitious mass cancels, and the result is the Bethe log and a constant.

Modern QED texts use similar approaches. Weinberg uses a fictitious photon mass in the photon propagator which leads to a high and low-energy term [16]. When added, the fictitious mass cancels, yielding a constant and the Bethe log term. The evolution of Lamb shift calculations is outlined in [12] and the detailed status is in [23].

For the purposes of the present paper, which is a non-relativistic calculation of the radiative Lamb shift, not including vacuum polarization, it is important to note that Bethe's non-relativistic result and the numerically evaluated Bethe log have been obtained in virtually all Lamb shift calculations, including all relativistic calculations [5,6,16,22]. The Bethe log represents the non-relativistic radiative contribution in all these diverse calculations. The Lamb shift spectral density obtained here can be integrated to obtain the non-relativistic Lamb shift, a quantity given by the Bethe logarithm. Furthermore, the integral can be taken to the desired energy to be consistent with a particular relativistic calculation. Thus, the spectral density computed is a standalone fundamental quantity that relates to all Lamb shift calculations described in the literature. Consequently, there is no need to consider the numerous relativistic calculations further in this paper.

Bethe's non-relativistic calculation to order α in the vacuum field (one virtual photon) was based on second-order perturbation theory applied to the minimal coupling of the atom with the vacuum field, $(e/mc)\mathbf{A} \cdot \mathbf{p}$, and a dipole approximation; here m , e , and \mathbf{p} are the mass, charge and momentum of the electron, respectively, c is the speed of light in a vacuum, and \mathbf{A} is the vector potential. This interaction leads to the emission and absorption of virtual photons corresponding to virtual transitions. The shift is expressed as a sum over the infinite number of intermediate states, bound and scattering, reached by virtual transitions. The predicted shift is divergent, but Bethe subtracted the term that corresponded to the linearly divergent vacuum energy shift for a free bare electron, essentially doing a mass renormalization to remove this higher-order divergence in the spectral density for the shift. For S states, the resulting spectral density has a $1/\text{frequency}$ behavior for frequencies above about $1000 \text{ eV}/\hbar$, (with \hbar the reduced Planck's constant) giving a logarithmic divergence in the shift. Bethe used a high-frequency cutoff of $\omega = mc^2/\hbar$.

The models of Welton and Power embody different perspectives on the role of vacuum fluctuations than the Bethe calculation. Itzykson and Zuber note: "Charged particles interact with the fluctuations of the quantized electromagnetic field ... we may give, following Welton, a qualitative description of the main effect: the Lamb shift" [22]. Welton's semi-quantitative model for computing the Lamb shift was based on the perturbation of the motion of a bound electron in the H atom due to the quantum vacuum fluctuations altering the location of the electron, which resulted in a slight shift of the bound state energy [6,9,12]. This simplified intuitive model predicts a spectral density proportional to $1/\text{frequency}$ for all frequencies and a shift only for S states. The approach of Feynman [24], interpreted by Power [10], considers a large box containing H atoms and is based on the shift in the energy in the quantum vacuum field due to the change in the index of refraction arising from the presence of H atoms. This approach predicts that the shift in the energy in the vacuum field around the H atoms exactly equals the radiative shift [6,11]. It gives a spectral density with

the same high-frequency dependence as Bethe, but a different low-frequency dependence. A similar calculation to Power's models the Lamb shift as a Stark shift [6].

The Lamb shift has been previously computed using $O(4)$ symmetry [25] and by using $SO(4,2)$ symmetry [26] using a different approach from that presented here. In both these cases, the Bethe log is obtained for low-lying states as a converging series. The results presented here describe a calculation of the Lamb shift for all states that is based on a $SO(4,2)$ group theoretical analysis of the H atom that allows us to determine the dependence of the shift on the frequency with no sum over states [27]. The spectral shift due to all virtual transitions is computed analytically.

The degeneracy group of the non-relativistic H atom is $O(4)$, with generators angular momentum operator L and Runge-Lenz vector A . A representation of $O(4)$ of dimension n^2 exists for each value of the principal quantum number n , where the angular momentum L has values from 0 to $n - 1$, and there are $2L + 1$ possible values of $L_z = m$, the azimuthal quantum number. If this group is extended by adding a 4-vector of generators the non-invariance group $SO(4,1)$ results, which has representations that include all states of different n and L and operators that connect states with different principal quantum numbers. Adding a 5-vector of additional generators produces the group $SO(4,2)$ and allows us to express Schrödinger's equation in terms of the new generators, and to make effective group theoretical calculations [27]. The basis states used permit both bound and scattering states to be included seamlessly [28] and no sum over states appears in the final expression for the spectral density. One advantage of this approach is that for each energy level it is possible to readily compute a spectral density for the shift whose integral over frequency from 0 to $mc^2/2\pi\hbar$ is the radiative shift that includes transitions to all possible states. This reveals how different frequencies of the vacuum field contribute to the radiative shift.

In this paper, the different results of Bethe, Welton and Power are compared to the group theoretical spectral density for the non-relativistic Lamb shift for the 1S ground state, and the 2S and 2P levels. With this new picture of the Lamb shift, differences between the various approaches are seen. Knowing the spectral density of the shift provides new insights into understanding the Lamb shift.

2. Background of Radiative Shift Calculations

The first calculation of the Lamb shift of a hydrogen atom was carried out by Bethe in 1947 [8], who assumed the shift was due to the interaction of the atom with the vacuum field. He calculated the shift using second-order perturbation theory, assuming that there was minimal coupling in the Hamiltonian

$$H_{\text{int}} = -\frac{e}{mc} \mathbf{A} \cdot \mathbf{p} + \frac{e^2}{2mc^2} \mathbf{A}^2, \quad (1)$$

where \mathbf{A} is the vector potential in the dipole approximation for the vacuum field in a large quantization volume V :

$$\mathbf{A} = \sum_{\mathbf{k}, \lambda} \left(\frac{2\pi\hbar}{\omega_k V} \right)^{1/2} (a_{\mathbf{k}, \lambda} + a_{\mathbf{k}, \lambda}^\dagger) \mathbf{e}_{\mathbf{k}, \lambda}, \quad (2)$$

where the sum is over the virtual photon frequency ω_k , and the polarization λ ; $a_{\mathbf{k}, \lambda}$ and $a_{\mathbf{k}, \lambda}^\dagger$ are the annihilation and creation operators, and $\mathbf{e}_{\mathbf{k}, \lambda}$ is a unit vector in the direction of polarization of the electric field. The shift from the \mathbf{A}^2 term is independent of the state of the atom and is therefore neglected. The total shift $\Delta E_{n\text{Tot}}$ for energy level n of the atom in state $|n\rangle$ is given by second-order perturbation theory as

$$\Delta E_{n\text{Tot}} = -\frac{2}{3\pi^2} \frac{\alpha}{m^2 c^2} \sum_m |\mathbf{p}_{mn}|^2 \int \frac{E dE}{E_m - E_n + E}, \quad (3)$$

where the quantum vacuum field energy is $E = \hbar\omega$ and the momentum matrix elements are $|\mathbf{p}_{mn}| = |\langle m|\mathbf{p}|n\rangle|$. The sum is over all intermediate states $|m\rangle$, scattering and bound, where $m \neq n$ and the integral is over the energy E of the vacuum field. The fine structure constant is $\alpha = e^2/\hbar c$.

The integrand in Equation (3) has a linear divergence. Bethe observed that this divergence corresponded to the integral that occurs when the binding energy vanishes so $(E_m - E_n) \rightarrow 0$ and the electrons are free:

$$\Delta E_{\text{free}} = -\frac{2}{3\pi^2} \frac{\alpha}{m^2 c^2} \sum_m |\mathbf{p}_{mn}|^2 \int dE. \quad (4)$$

He subtracted this divergent term ΔE_{free} from the total shift $\Delta E_{n\text{Tot}}$,

$$\Delta E_{nL} = \Delta E_{n\text{Tot}} - \Delta E_{\text{free}}, \quad (5)$$

to obtain a finite observable shift ΔE_{nL} for the state $|nL\rangle$:

$$\Delta E_{nL} = \frac{2\alpha}{3\pi(mc)^2} \sum_m |\mathbf{p}_{nm}|^2 \int_0^{\hbar\omega_c} dE \frac{(E_m - E_n)}{E_m - E_n + E - i\epsilon}, \quad (6)$$

where ω_c is a cutoff frequency for the integration that Bethe took as $\hbar\omega_c = mc^2$. Using an idea from Kramers [6,29], Bethe did this renormalization, taking the difference between the terms with a potential present and without a potential present, essentially performing the free electron mass renormalization. He reasoned that relativistic retardation could be neglected and the radiative shift could be reasonably approximated using a non-relativistic approach and he cut the integration off at an energy corresponding to the mass of the electron. He obtained a finite result that required a numerical calculation over all states, bound and scattering, that gave good agreement with measurements [7,8,30].

The spectral density in the Bethe formalism, which is analysed in this paper, is the quantity in Equation (6) being integrated over E . It includes the sum over states m and the constants. The term for m represents the contribution to the Lamb shift for the virtual transition from state n to state m . Note that since the ground state is the lowest state, all intermediate states have higher energies so the ground state shift has to be positive.

For the purposes of comparison to the other calculations of the Lamb shift, it is helpful to show the next steps Bethe took to evaluate the shift ΔE_n for S states, which have the largest shifts. Note that the spectral density in Equation (6) that will be analyzed is not affected by the subsequent approximations Bethe made to evaluate the integral. First, the E -integration is carried out:

$$\Delta E_n^{\text{Bethe}} = \frac{2\alpha}{3\pi} \left(\frac{1}{mc}\right)^2 \sum_m |\mathbf{p}_{nm}|^2 (E_m - E_n) \ln \frac{(mc^2 + E_m - E_n)}{|E_m - E_n|}. \quad (7)$$

To simplify the evaluation Bethe assumed $|E_m - E_n| \ll mc^2$ in the logarithm and that the logarithm would vary slowly with the index m so it could be replaced by an average value,

$$\widehat{\Delta E}_n^{\text{Bethe}} = \frac{2\alpha}{3\pi} \left(\frac{1}{mc}\right)^2 \ln \frac{mc^2}{|E_m - E_n|_{\text{ave}}} \sum_m |\mathbf{p}_{nm}|^2 (E_m - E_n), \quad (8)$$

where the hat over the ΔE indicates this is an approximation to Equation (7). Now that the E -integration is done, the spectral density is no longer manifest. The summation can be evaluated using the dipole sum rule:

$$2 \sum_m |\mathbf{p}_{nm}|^2 (E_m - E_n) = \hbar^2 \langle n | \nabla^2 V | n \rangle. \quad (9)$$

where $\langle \dots \rangle$ indicates an expectation value. The Laplacian with a Coulomb potential $V = -Ze^2/r$ is $\nabla^2 V(r) = 4\pi Ze^2 \delta(\mathbf{r})$ which gives

$$\langle n | \nabla^2 V | n \rangle = 4\pi Ze^2 |\psi_n(0)|^2, \quad (10)$$

where Z is the positive charge of the nucleus, $\psi(r)$ is the wave function for the Coulomb potential and $|\psi_n(0)|^2$ is zero except for S states,

$$|\psi_n(0)|^2 = \frac{1}{\pi} \left(\frac{Z\alpha mc}{n\hbar} \right)^3. \quad (11)$$

For S states, this gives an energy shift equal to [6]

$$\widehat{\Delta E}_n^{\text{Bethe}} = \frac{4mc^2}{3\pi} \alpha (Z\alpha)^4 \frac{1}{n^3} \ln \frac{mc^2}{|E_m - E_n|_{\text{ave}}}. \quad (12)$$

The so-called “Bethe log” for an S state with principal quantum number n is

$$\ln \frac{mc^2}{|E_m - E_n|_{\text{ave}}} = \frac{\sum_m |\mathbf{p}_{nm}|^2 (E_m - E_n) \ln \frac{mc^2}{|E_m - E_n|}}{\sum_m |\mathbf{p}_{nm}|^2 (E_m - E_n)}, \quad (13)$$

where the sum is over all states, bound and scattering. Bethe also has extended the formalism for shifts for states that are not S states [30].

Regarding the approximations Bethe made to obtain Equation (8) from Equation (7) and the use of the Bethe log Equation (13) he commented: “The important values of $|E_m - E_n|$ will be of the order of the ground state binding energy for a hydrogenic atom. This energy is very small compared to mc^2 so the log [in Equation (7) here] is very large and not sensitive to the exact value of $(E_m - E_n)$. In the numerator we neglect $(E_m - E_n)$ altogether and replace it by an average energy [30]”. This study shows that Bethe was correct that the relative contribution from energies of the order of the ground state is very important, but the contribution from higher energy scattering states is quite significant, and therefore the approximation $|E_m - E_n| \ll mc^2$ is not valid for higher energy scattering states for which E_m increases to the value mc^2 . I am not aware of any quantitative estimates of the error in the approximation. The difference, 0.3%, between the value obtained here for the total $1S$ shift and that of Bethe may be due to this approximation, although this has not been verified. On the other hand, Bethe’s approximation may have made his non-relativistic approach viable.

To provide a more intuitive and qualitative physical picture of the shift, Welton considered [9] the effect of a zero-point vacuum field on the motion of an electron bound in a Coulomb potential $V(\mathbf{r})$ at a location \mathbf{r} . The perturbation $\xi = (\xi_x, \xi_y, \xi_z)$ in the position of the bound electron due to the random zero-point vacuum field \mathbf{E}_0 causes a variation in the potential energy,

$$V(\mathbf{r} + \xi) = V(\mathbf{r}) + \xi \cdot \nabla V(\mathbf{r}) + \frac{1}{2} (\xi \cdot \nabla)^2 V(\mathbf{r}) + \dots. \quad (14)$$

Because of the harmonic time dependence of the vacuum field $\langle \xi \rangle$ vanishes and the radiative shift is given approximately by the vacuum expectation value of the last term:

$$\Delta E_n^{\text{Welton}} = \frac{\langle \xi^2 \rangle}{6} \langle \nabla^2 V(\mathbf{r}) \rangle_n. \quad (15)$$

Since the potential has spherical symmetry $\langle \xi_x^2 \rangle = \langle \xi_y^2 \rangle = \langle \xi_z^2 \rangle = \langle \xi^2 \rangle / 3$. Equation (15) gives $\Delta E_n^{\text{Welton}}$ as the product of two factors, the first depending on the nature of the fluctuations in the position of the bound electron due to the vacuum field and the second

depending on the structure of the system. ξ is determined by $m\ddot{\xi} = e\mathbf{E}_0$, where top double-dot denotes the second order time derivative.

With a Fourier decomposition of E_0 and ξ , and integrating over the frequency distribution of the vacuum field, the vacuum expectation value can be computed [6,12]:

$$\langle (\vec{\xi})^2 \rangle = \frac{2\alpha}{\pi} \left(\frac{\hbar}{mc} \right)^2 \int_0^{mc^2} \frac{dE}{E}. \quad (16)$$

With the results in Equations (10) and (11) the Laplacian in Equation (15) can be evaluated giving a shift for S states equal to [6]:

$$\Delta E_n^{\text{Welton}} = \frac{4mc^2}{3\pi} \alpha (Z\alpha)^4 \frac{1}{n^3} \int_0^{mc^2} \frac{dE}{E}. \quad (17)$$

Equation (17) shows that the spectral density for the Welton approach is proportional to $1/E$. For the upper limit of integration, we assume the maximum wavelength equals the Compton wavelength corresponding to an energy mc^2 , which Bethe used. The lower limit of 0 gives a divergent shift, clearly showing that the model of the bound state is deficient at low energies. To give an approximate lower limit, note that the large wavelength modes are sensitive to low-lying electronic states, which suggests a wavelength cutoff about equal to the radius a of the ground state so $E = \hbar c/a = \alpha mc^2 = 4.01$ keV. For the $n = 2$ level of hydrogen, this gives a shift of 660 MHz, about 60% of the observed shift [22], which confirms the model is reasonable. On the other hand, if one happens to compare Equation (17) to Equation (12) it is clear that with the lower limit $|E_m - E_n|_{\text{ave}}$, as defined in the Bethe log Equation (13), Welton's model gives exactly the same total S state shift as in the approximate Bethe formalism Equation (12). With these limits, the RMS (root mean square) amplitude of oscillation of the electron bound in the Coulomb potential, $\sqrt{\langle (\vec{\xi})^2 \rangle}$, is about 60 fm, which is about 0.037% of the mean radius of the 2S electron orbit.

Feynman proposed another approach for computing the Lamb shift based on a fundamental observation about the interaction of matter and the vacuum field [24]. He considered a large box containing a low density of atoms in the quantum vacuum. The atoms cause a change in the index of refraction, which leads to changes in the frequencies of the vacuum field. The wavelengths remain the same. Feynman maintained that the change in the energy of the zero point vacuum field in the box due to the frequency changes resulting from a weak perturbing background of atoms acting as a refracting medium would correspond to the self energy of the atoms, which is precisely the Lamb shift.

Power, based on the suggestion by Feynman, considered the change in vacuum energy when N hydrogen atoms are placed in a volume V using the Kramers-Heisenberg expression for the index of refraction $n(\omega_k)$ [6,10]. The H atoms cause a change in the index of refraction and therefore a change in the frequencies of the vacuum fluctuations present. The corresponding change in vacuum energy ΔE is

$$\Delta E = \sum_k \frac{1}{n(\omega_k)} \frac{1}{2} \hbar \omega_k - \frac{1}{2} \hbar \omega_k, \quad (18)$$

where the sum is over all frequencies ω_k present. For a dilute gas of atoms in a level n , the index of refraction is [6]

$$n(\omega_k) = 1 + \frac{4\pi N}{3\hbar} \sum_m \frac{\omega_{mn} |\mathbf{d}_{mn}|^2}{\omega_{mn}^2 - \omega_k^2}, \quad (19)$$

where $\omega_{mn} = (E_m - E_n)/\hbar$ and $\mathbf{d}_{mn} = e\mathbf{x}_{mn}$, the transition dipole moment. Substituting $n(\omega_k)$ into Equation (18) gives a divergent result for the energy shift. Following Bethe's approach, Power subtracted from ΔE the energy shift for the N free electrons, which equals the shift when $\omega_{mn} \rightarrow 0$, with no binding energy. After making this subtraction and

converting the sum over ω_k to an integral over ω , and letting $NV \rightarrow 1$ the observable shift in energy is obtained [6]:

$$\Delta E_n^{\text{Power}} = -\frac{2}{3\pi c^3} \sum_m \omega_{mn}^3 |\mathbf{d}_{mn}|^2 \int_0^{mc^2/\hbar} \frac{\omega d\omega}{\omega_{mn}^2 - \omega^2}. \quad (20)$$

Noting that

$$\langle m | \frac{\mathbf{p}}{m} | n \rangle = \frac{i}{\hbar} \langle m | [H, \mathbf{x}] | n \rangle = \frac{i}{\hbar} (E_m - E_n) \langle m | \mathbf{x} | n \rangle, \quad (21)$$

it follows

$$|\mathbf{p}_{mn}|^2 = m^2 \omega_{mn}^2 |\mathbf{x}_{mn}|^2 = \frac{m^2 \omega_{mn}^2}{e^2} |\mathbf{d}_{mn}|^2. \quad (22)$$

This allows us to write Power's result Equation (20) as

$$\Delta E_n^{\text{Power}} = -\frac{2e^2}{3\pi m^2 c^3} \sum_m \omega_{mn} |\mathbf{p}_{mn}|^2 \int_0^{mc^2/\hbar} \frac{\omega d\omega}{\omega_{mn}^2 - \omega^2}. \quad (23)$$

Writing this equation in terms of $E = \hbar\omega$ instead of ω yields:

$$\Delta E_n^{\text{Power}} = -\frac{2\alpha}{3\pi} \left(\frac{1}{mc} \right)^2 \sum_m |\mathbf{p}_{mn}|^2 (E_m - E_n) \int_0^{mc^2} \frac{E dE}{(E_m - E_n)^2 - E^2}. \quad (24)$$

Below, in Section 5, this equation will be used to analyze the spectral density for Power's method, showing the spectral density is different from Bethe's at low frequencies but the same at high frequencies. When Equation (24) is integrated with respect to E , taking the principal value, one obtains

$$\Delta E_n^{\text{Power}} = \frac{2\alpha}{3\pi} \left(\frac{1}{mc} \right)^2 \sum_m |\mathbf{p}_{mn}|^2 (E_m - E_n) \ln \left[\frac{mc^2 + (E_m - E_n)}{E_m - E_n} \cdot \frac{mc^2 - (E_m - E_n)}{E_m - E_n} \right]^{1/2}. \quad (25)$$

Except for the argument in the \ln function, which corresponds to the upper limit of integration, this is the same as Bethe's expression (7) for the shift. Assuming $mc^2 \gg E_m - E_n$, as Bethe did, then both expressions for the total shift are identical. It is clear, however, that this approximation is not valid at high energies since the second factor in the \ln function in Equation (25) may even become less than one making the \ln term negative. Feynman's approach highlights the changes in the vacuum field energy due to the interactions with the H atoms.

One assumption in the computation by Power is that the index of refraction in the box containing the atoms is spatially uniform. In Section 6, this assumption will be revisited and a model suggested that predicts for a single atom the changes in the vacuum field energy as a function of position for each spectral component of the radiative shift.

3. Spectral Density of the Lamb Shift

Our goal is to develop an expression for the energy shift of a level, in terms of the generators of the group $SO(4,2)$, that is an integral over frequency. Then the integrand will be the spectral density of the shift, and group theoretical techniques can be used to evaluate it [27]. This approach yields a generating function for the shifts for all levels. The initial focus is on the ground state 1S level as an illustration of the results. At ordinary temperatures and pressures, most atoms are in the ground state. The radiative shift for the 1S level is [27]

$$\Delta E_1 = \frac{4mc^2 \alpha (Z\alpha)^4}{3\pi} \int_0^{\phi_c} d\phi e^\phi \sinh \phi \int_0^\infty ds e^{se^{-\phi}} \frac{d}{ds} \frac{1}{(\coth \frac{s}{2} + \cosh \phi)^2}, \quad (26)$$

where the dimensionless normalized frequency variable ϕ is defined as

$$\phi = \frac{1}{2} \ln \left[1 + \frac{\hbar\omega}{|E_1|} \right], \quad (27)$$

and E_1 is the ground state energy of the H atom -13.6 eV. The cutoff ϕ_c corresponds to $E = \hbar\omega_c = mc^2 = 511$ keV corresponding to the electron mass.

The group theoretical expression for the Lamb shift (26) is directly derived from the Klein-Gordon equations of motion using a non-relativistic dipole approximation, assuming infinite proton mass, and minimal coupling with the vacuum field. Basis states of $(1/Z\alpha)$ are used since they have no scattering states and have the same quantum numbers as the usual bound energy eigenstates [27]. The level shift is obtained as the difference between the mass renormalization for a spinless meson bound in the desired state and the mass renormalization for a free meson. Second-order perturbation theory is not used. Near the end of the derivation, an equation that is equivalent to Bethe's result (6) for the radiative shift can be derived by inserting a complete set of Schrödinger energy eigenstates. Thus, we expect the fundamental results from Bethe's spectral density (with no approximations) and the group's theoretical spectral density to be in agreement [12,27]. For convenience, in Appendix A an explanation of the basis states used to derive Equation (26) is given, and in Appendix B, the derivation of Equation (26) is given, since the derivation in [27] is spread in steps throughout the paper as the group theory methods are developed.

Equation (26) can be written as an integral over $E = \hbar\omega$, which is the energy of the vacuum field in eV. The definite integral over s can be evaluated analytically for different values of ϕ or $E = \hbar\omega$. The ground state Lamb shift ΔE_1 is measured in eV so the spectral density of the shift $d\Delta E_1/dE$ is measured in eV/eV which is dimensionless:

$$\Delta E_1 = \int_0^{mc^2} \frac{d\Delta E_1}{dE} dE, \quad (28)$$

where the ground state spectral density from Equation (26) is

$$\frac{d\Delta E_1}{dE} = \frac{4\alpha^3}{3\pi} e^{-2\phi} \sinh \phi \int_0^\infty ds e^{se^{-\phi}} \frac{1}{\sinh^2(\frac{s}{2})} \frac{1}{(\coth \frac{s}{2} + \cosh \phi)^3}. \quad (29)$$

Figure 1 shows a logarithmic plot (ordinate is a log, abscissa is linear) of the spectral density $\frac{d\Delta E_1}{dE}$ of the ground state Lamb shift with $Z = 1$ over the entire range of energy E computed from Equation (29) using Mathematica. The spectral density is largest at the lowest energies, and decreases monotonically by about 4 orders of magnitude as the energy increases to 511 keV. The ground state shift is the integral of the spectral density from energy 0 to 511 keV.

Figure 2 is a log-log plot of the same information. The use of the log-log plot expands the energy range for each decade, revealing that for energy above about 1000 eV the slope is approximately -1 , indicating that the spectral density is nearly proportional to $1/E$. For energy below about 10 eV, the spectral density in Figure 2 is almost flat, corresponding to a linear decrease with E , with a maximum at the lowest energy computed, as shown in Figure 3. Figure 2 shows that there are essentially two different behaviors of the spectral density, one for values of the energy E of the vacuum field that are less than 10 eV, which is less than $E_m - E_n$, the energy difference for all bound state transitions, and another region where E is much larger than the bound state energies, and the spectral density goes as $1/E$. The origin of this behavior is clear mathematically from the factor $B = 1/(E_m - E_n + E)$. In the expression for the spectral shift (6) for $E < E_m - E_n$, $B \propto 1 - E/(E_m - E_n)$ and for $E > E_m - E_n$, $B \propto 1/E$.

Figure 3 shows linear plots of the spectral density of the shift for the ground state computed from Equation (29) for several lower energy regions. Figure 3a shows a linear increase in the spectral density as the energy decreases over the small energy interval plotted. Figure 3b shows a linear increase of about 15% as the energy decreases from 3 eV to

0 eV. Figure 3c shows that the spectral density increases by a factor of about 4 as the energy decreases from 100 eV to 0 eV. In the low-frequency limit, the spectral density increases linearly to a constant value as the energy is reduced.




Figure 1. Plot of the log of the spectral density of the ground state Lamb shift from the group theoretical expression (29) versus the energy from 0 to 510 keV.




Figure 2. Log-log plot of the spectral density of the ground state shift from the group theoretical expression (29) versus the energy. For energies above about 1000 eV, the behavior is dominated by a $1/\text{energy}$ dependence. From about 10 eV to 0 eV, there is a slow linear increase in the spectral density.

From explicit evaluations, Section 4 shows that for shifts in S states with principal quantum number n , the asymptotic spectral density for large E is proportional to $\alpha(Z\alpha)^4(1/n^3)/E$, and Section 5 shows that as the energy E goes to zero, the spectral density increases linearly, reaching a maximum value that is proportional to $\alpha(Z\alpha)^2(1/n^2)$. An approximate fit to the ground state data in Figure 1 is

$$\frac{d\Delta E_1^{\text{fit}}}{dE} = D \frac{(1 + e^{-HE})}{(E + C)}, \quad (30)$$

where $D = 4.4008 \times 10^{-6}$, $H = 0.08445/\text{eV}$, $C = 106.79 \text{ eV}$. The fit is quite good at the asymptotes and within 10% over the entire energy range.

The spectral density shown in Figures 1 or 2 can be used to determine the contribution to the total ground state shift from different energy regions. Integrating the spectral density from 0 eV to energy E gives the value of the partial shift $\Delta_1(E)$ that these energies (0 eV to E eV) contribute to the total shift ΔE_1 for the ground state. In Figure 4, $\Delta_1(E)/\Delta E_1$, which is the fraction of the total shift ΔE_1 due to the contributions from energies below E , is plotted as a function of E . Figure 4a shows that almost 80% of the shift comes from energies below about 100,000 eV. Figure 4b shows that about half the total shift is from energies below 9050 eV. Figure 4c shows that energies below 100 eV contribute about 10% of the total shift. Energies below 13.6 eV contribute about 2.5% while energies below 1 eV contribute about 1/4% of the total. As Figure 4c shows, the fraction of the total shift increases linearly for $E < 10 \text{ eV}$, corresponding to the nearly horizontal portion of the shift

density for $E < 10$ eV, as shown in Figure 2. The contribution to the total 1S shift for the visible spectral interval 400–700 nm (1.770 eV to 3.10 eV) is about 1.00342×10^{-7} eV or about 3/10 % of the total shift.




Figure 3. Linear plots of the ground state spectral density calculated from group theory as a function of energy for low and mid energies. From about 10 eV to 0 eV, the spectral density increases linearly to its maximum value. The value of the abscissa at the origin is 0 eV for all graphs. (a): Linear change in ground state spectral density at very low energies. (b): Near linear change in ground state spectral density for visible and near infra-red energies. The contribution to the total shift for energies below 3 eV is about 0.7%. (c): Ground state spectral density calculated for energies below 80 eV, which contribute about 8.6% to the total shift.




Figure 4. The ordinate is the fraction of the ground state shift ΔE_1 due to vacuum field energies between 0 and E , plotted as a function of E . This plot is obtained by integration of the spectral density Equation (29), shown in Figure 1. The plot is linear in the ordinate and abscissa. The origin corresponds to (0,0) for all plots. (a): Fraction of the 1S shift due to energies from 0 to E for $0 < E < 510$ keV. (b): Fraction of the 1S shift due to energies below E , for $0 < E < 9000$ eV. (c): Fraction of the 1S shift due to energies from 0 to E , for $0 < E < 100$ eV. Energies below 30 eV account for about 0.05 of the total shift. The variation is linear for $E < 10$ eV.

The relative contribution to the total shift per eV is much greater for lower energies. For example, half the 1S shift corresponds to energies 0 to 9000 eV, but only about 0.2% corresponds to 500,000 to 509,000 eV. The largest contribution to the shift per eV is at the lowest energies, which have the steepest slope of the spectral density curve in Figure 1, about 1000 times greater than the slope for the largest values of the energy. However, the total range for the large energies, from 9050 to 510,000 eV is so large that the absolute contribution to the total shift for large energies is significant.

For the ground state, Figure 5 shows how the dominant terms for different m in the Bethe sum over states in Equation (6) contribute to the full spectral density obtained from group theory Equation (29). Each such term in the Bethe sum could be interpreted as corresponding to the shift resulting from virtual transitions from state n to state m occurring due to the vacuum field. Each term shown has a behavior similar to that of the full spectral density, but the magnitudes decrease as the transition probabilities decrease.




Figure 5. Log-log plot of the 1S spectral density from group theory Equation (29) in black, and the contributions to this shift in the Bethe formalism for the transition $1S \rightarrow 2P$ (blue), $1S \rightarrow 4P$ (red), $1S \rightarrow 8P$ (green). The dashed blue line shows the high-frequency $1/E$ asymptote. The black line is the complete spectral density which is the summation of the contributions from all transitions.

Figure 6 shows the spectral densities for 1S (black) and 2S (orange) shifts. The shapes are similar but the spectral density for the 1S shift is about eight times as large at high frequencies and about four times as large at low frequencies, factors that are derived explicitly in Sections 4 and 5 by considering the asymptotic forms of the spectra density for S states with different principal quantum numbers. Both S states have a $1/E$ high-frequency behavior. The s -integration in the group theoretical calculation for the 2S state diverges for energies below 10.2 eV due to a non-relativistic approximation, but the spectral density of the shift can be obtained from a low-energy approximation (48) to the group theory result, which is derived in Section 5.




Figure 6. A log-log plot of the group theoretical spectral density for the 1S (black) and 2S (orange) shifts versus energy. The dashed orange curve below 1 eV is a 2S low-energy approximation (47) from group theory or the Bethe formula. The blue is the largest single contribution in the Bethe formalism to the 2S shift for the transition $2S \rightarrow 3P$.

The spectral density $d\Delta E_n/dE$ for a state n can be defined in a convenient form suggested by Equation (29),

$$\frac{d\Delta E_n}{dE} = \frac{4\alpha^3}{3\pi} \int_0^\infty ds W_n(s, \phi_n), \quad \phi_n = \ln \left[1 + \frac{E}{|E_n|} \right], \quad (31)$$

where the energy for state n is $E_n = -mc^2(Z\alpha)^2/2n^2$. The group theoretical results give a spectral density for the 2S–2P Lamb shift [27]:

$$W_{2S-2P}(s, \phi_2) = \frac{4e^{(2se^{-\phi_2} + \phi_2)} \sinh^3(\phi_2) \operatorname{csch}^2\left(\frac{s}{2}\right)}{[\cosh(\phi_2) + \coth\left(\frac{s}{2}\right)]^5}, \quad (32)$$

and for the 2P shift [27]:

$$W_{2P}(s, \phi_2) = - \frac{e^{(2se^{-\phi_2} + \phi_2)} \sinh(\phi_2) \operatorname{csch}^4\left(\frac{s}{2}\right) [\cosh(\phi_2) \sinh(s) + \cosh(s) - 3]}{2 [\cosh(\phi_2) + \coth\left(\frac{s}{2}\right)]^5}. \quad (33)$$

The spectral density of the 2P shift has a very different behavior from the spectral density of the 2S shift (Figure 7). It is negative and it falls off as $1/E^2$. The shift is negative because the dominate contribution to the shift is from virtual transitions from the 2P state to the lower 1S state, with an energy difference of about 10.2 eV. For energies below 20 eV, the absolute value of the spectral density of the 2P shift increases rapidly in magnitude as the energy is reduced and is much bigger than the spectral density for the 2S shift. The 2S shift cannot have a negative contribution from the lower 1S state since the transition $2S \rightarrow 1S$ is forbidden by the conservation of angular momentum. The classic Lamb shift arises from the difference between the two spectral densities, so the negative 2P spectral density actually increases the 2S–2P Lamb shift as the energy decreases (Figure 8). In effect, the 2S–2P shift is dominated by vacuum energies below about 100 eV. The total 2P shift is about 0.3% percent of the 2S shift. Bethe also computed a negative contribution for the shift from the 2P state [30].




Figure 7. Log-log plot of the absolute value of the spectral density versus the energy for the 2S shift (orange), which goes as $1/E$ for large E , and for the 2P shift (green), which goes as $1/E^2$ for large E . At 511 keV, the 2P spectral density is about 5 orders of magnitude smaller than the 2S spectral density. Below 20 eV, the absolute value of the 2P spectral density is greater than the 2S spectral density. Note that the 2P spectral density is actually negative and the 2S spectral density is positive.




Figure 8. Log-log plot of the spectral density for the 2S shift (orange) and the 2S–2P Lamb shift (blue) versus energy. The solid black line is the $1/E$ asymptote.

Comparing the Ground State Group Theoretical Lamb Shift Calculations to Those of Bethe, Welton, and Feynman

Integrating the group theoretical spectral density Equation(29) from near zero energy (5.4×10^{-7} eV) to 511 keV, about the rest mass energy of the electron, gives the 1S shift of

3.4027×10^{-5} eV, in agreement with the numerical result of Bethe and Salpeter summing over states and using the Bethe log approximation, 3.392×10^{-5} eV, to about 0.3% [8].

Bethe and Salpeter [30] reported that the ground state Bethe log, as defined in Equation 13, which is a logarithmically weighted average value of the excitation of the energy levels contributing to the radiative 1S shift, was equivalent to $|E_m - E_n|_{ave}$ of 19.77 Ry or 269 eV [30]. Because of the weighting, it is not clear how to interpret this value, other than it indicates that high-energy photons and scattering states contribute significantly to the shift. The group theoretical method does not provide an equivalent weighted average value for direct comparison.

Although the methods of Bethe, Welton, and Power as defined all give approximately the same value for the 1S shift, which equals the integral of the spectral density, they differ significantly in their frequency dependence, which is examined in Section 4.

4. The Spectral Density of The Lamb Shift at High Frequency

The form for $d\Delta E_n/dE$, which is the Lamb shift spectral density for level n , can be obtained at high energies from (i) the classic calculation by Bethe using second-order perturbation theory; (ii) the calculation by Welton of the Lamb shift; (iii) the calculation by Power of the Lamb shift based on Feynman's approach; and (iv) the group theoretical calculation.

The spectral density for level n can be written from Bethe's expression (6):

$$\frac{\Delta E_n^{\text{Bethe}}}{\Delta E} = \frac{2\alpha}{3\pi} \left(\frac{1}{mc} \right)^2 \sum_m |\mathbf{p}_{mn}|^2 (E_n - E_m) \frac{1}{E_n - E_m - E}. \quad (34)$$

For the ground state spectral density $n = 1$, $Z = 1$, $E_1 = -13.613$ eV, and for the bound states with principal quantum number m , $E_m = -13.613 \text{ eV}/m^2$. For scattering states, E_m is positive. Hence the denominator is negative for all terms in the sum over m and never vanishes, and the spectral density is positive so the ground state shift is positive as it must be. For large values of E , we can make the approximation,

$$\frac{\Delta E_n^{\text{Bethe}}}{\Delta E} \Big|_{E \rightarrow \infty} = \frac{2\alpha}{3\pi} \left(\frac{1}{mc} \right)^2 \sum_m |\mathbf{p}_{mn}|^2 (E_m - E_n) \frac{1}{E}. \quad (35)$$

The summation can be evaluated using the dipole sum rule (9)–(11) for the Coulomb S state wavefunction, obtaining the final result for the high-frequency spectral density for S states with the principal quantum number, n :

$$\frac{d\Delta E_n^{\text{Bethe}}}{dE} \Big|_{E \rightarrow \infty} = \frac{4mc^2}{3\pi} \alpha (Z\alpha)^4 \frac{1}{n^3} \frac{1}{E}. \quad (36)$$

The result highlights the $1/E$ divergence at high frequencies and shows the presence of a coefficient proportional to $1/n^3$. To put a scale on the coefficient, the high-frequency spectral density can be written as $(8/3\pi)(\alpha(Z\alpha)^2/n)(E_n/E)$.

The spectral density for all frequencies from Welton's qualitative model, Equation (17), is identical to this high-frequency limit of Bethe's calculation. Thus, at low frequencies, the spectral density for Welton's semi-quantitative calculation diverges as $1/E$. Because of the expectation value of the Laplacian, Welton's approach predicts a shift only for S states. Its appeal is that it gives a physical picture of the primary role of vacuum fluctuations in the Lamb shift and shows the presence of the $1/E$ characteristic behavior. Its treatment of bound states at low energies is incomplete and inaccurate. To obtain a level shift, it requires providing a low-energy limit for the integration. As noted previously, if the lower limit is Bethe's log average excitation energy, 269 eV for $n = 1$, and the upper limit mc^2 then Welton's total 1S shift agrees with Bethe's. A choice of this type works since: (i) it does not include any contributions from energies below 269 eV and (ii) it gives a compensating contribution for energies from 269 eV to about 1000 eV that is larger than the actual spectral density, as shown in Figures 4 and (iii) above about 1000 eV, Welton's model gives the same $1/E$ spectral density as Bethe.

The spectral density for Power's model can be obtained from Equation (24):

$$\frac{\Delta E_n^{\text{Power}}}{dE} = -\frac{2\alpha}{3\pi} \frac{1}{(mc)^2} \sum_m |\mathbf{p}_{mn}|^2 (E_m - E_n) \frac{E}{(E_m - E_n)^2 - E^2}. \quad (37)$$

Letting E become large gives a result identical to the high-frequency limit (35) for the Bethe formalism and the Welton model, namely

$$\frac{\Delta E_n^{\text{Power}}}{dE} \Big|_{E \rightarrow \infty} = \frac{4mc^2}{3\pi} \frac{\alpha(Z\alpha)^4}{n^3} \frac{1}{E}. \quad (38)$$

Thus, the models of Bethe, Welton, and Power predict S states with the same $1/E$ dependence of the high-frequency spectral density, corresponding to the logarithmic divergence at high-frequency. The asymptotic theoretical result can be written in a form allowing convenient comparison to the calculated group theoretical spectral density

$$\frac{d\Delta E_n^{\text{Bethe}}}{dE} \Big|_{E \rightarrow \infty} = \frac{4mc^2}{3\pi} \frac{\alpha(Z\alpha)^4}{n^3} \frac{1}{E}. \quad (39)$$

The spectral density goes as $1/n^3$ for S states. For the ground state $n = 1$, $Z = 1$, this gives:

$$\frac{d\Delta E_1^{\text{Bethe}}}{dE} \Big|_{E \rightarrow \infty} = 4.488 \times 10^{-6} \frac{1}{E}. \quad (40)$$

A fit to the last two data points near 510 keV in the group theoretical calculations (GT-calc) gives:

$$\frac{d\Delta E_1^{\text{GTcalc}}}{dE} \Big|_{E \rightarrow \infty} = 4.4008 \times 10^{-6} \frac{1}{E}. \quad (41)$$

The coefficients differ by about 2%. Figure 9 is a plot of the ground state group theoretical calculated spectral density (red) from Equation (29) and the theoretical asymptotic behavior from Bethe, Power and Welton, Equation (40) (black), and the difference times of a factor of 10. The asymptotic theoretical result agrees with the full group theoretical calculation from Equation (29) to within about 2% at 511 keV, and to about 6% at 50 KeV. It is notable that the high-frequency form is a reasonable approximation down to 50 keV. Indeed, the Welton qualitative approach is based on this observation; it has the same $1/E$ energy dependence at all energies.




Figure 9. Top red curve is the 1S group theoretical calculated spectral density (29), slightly lower black curve is the $1/E$ asymptotic model Equation (39), and the bottom green curve is the difference times 10, plotted for the interval 50–510 keV. Both axes are linear.

5. Spectral Density of the Lamb Shift at Low Frequency

To obtain a low-frequency limit of the spectral density of the Lamb shift from the Bethe spectral density (34), expand the spectral density to first-order in E , giving

$$\frac{\Delta E_n^{\text{Bethe}}}{dE}|_{E \rightarrow 0} = \frac{2\alpha}{3\pi} \left(\frac{1}{mc} \right)^2 \sum_m |\mathbf{p}_{mn}|^2 \left(1 - \frac{E}{E_m - E_n} \right). \quad (42)$$

Since the sum is over a complete set of states m , including scattering states, the first term in parenthesis can be evaluated using the sum rule,

$$\sum_m |\mathbf{p}_{mn}|^2 = -2mE_n = (mc)^2 \frac{(Z\alpha)^2}{n^2}. \quad (43)$$

To evaluate the second term, Equation (22) and the Thomas–Reiche–Kuhn sum rule [31] can be used, giving

$$\sum_m \omega_{mn} |\mathbf{d}_{mn}|^2 = \frac{3e^2 \hbar}{2m}. \quad (44)$$

The final result for $E \rightarrow 0$ is

$$\frac{\Delta E_n^{\text{Bethe}}}{dE}|_{E \rightarrow 0} = \frac{2\alpha}{3\pi} \frac{(Z\alpha)^2}{n^2} - \frac{\alpha}{\pi mc^2} E. \quad (45)$$

The corresponding spectral density for $n = 1$, $Z = 1$, is

$$\frac{d\Delta E_1^{\text{Bethe}}}{dE}|_{E \rightarrow 0} = \frac{4\alpha \times 13.6}{3\pi mc^2} \left(1 - \frac{3E}{4 \times 13.6} \right) = 8.253 \times 10^{-8} (1 - 0.0551E). \quad (46)$$

As E decreases to zero, the spectral density increases linearly to a constant value,

$$4\alpha |E_n| / 3\pi mc^2 = 2\alpha^3 Z^2 / 3\pi n^2 = 8.253 \times 10^{-8} / n^2.$$

The intercept goes as $1/n^2$, but the slope is $\alpha / \pi mc^2$, which has a remarkably simple form and is independent of n .

Taking the low-frequency limit of the group theoretical result analytically gives exactly the same result as Equation (45) from the Bethe formulation:

$$\frac{d\Delta E_1^{\text{Gcalc}}}{dE}|_{E \rightarrow 0} = \frac{d\Delta E_1^{\text{Bethe}}}{dE}|_{E \rightarrow 0} = \frac{2\alpha}{3\pi} \frac{(Z\alpha)^2}{n^2} - \frac{\alpha}{\pi mc^2} E. \quad (47)$$

Figure 3 shows the results of group theoretical calculations of the spectral density of the ground state Lamb shift for different energy regions, showing the near linear increase in the spectral density as the frequency decreases from 80 eV to 10^{-5} eV. For low values of E , the slopes and intercept agree with Equation (47) within about two tenths of a percent.

To explore Power's approach at low frequencies, let E become small in the spectral density Equation (37), giving

$$\frac{\Delta E_n^{\text{Power}}}{dE}|_{E \rightarrow 0} = -\frac{2NV}{3\pi c^3} \sum_m |\mathbf{p}_{mn}|^2 \frac{E}{E_m - E_n}, \quad (48)$$

which is identical to the second term in the low E approximation to the Bethe result (45) so

$$\frac{\Delta E_n^{\text{Power}}}{dE}|_{E \rightarrow 0} = -\frac{1}{\pi} \frac{\alpha}{mc^2} E. \quad (49)$$

The result Equation (49) is identical to the frequency-dependent term in Equation (47) which is the low-frequency spectral density from the Bethe approach and from the group theoretical expression. However, in the low-frequency limit based on Power's expression for the spectral density, the constant term that is present in the other approaches does not appear. This is a consequence of the form used for the index of refraction, which assumes that real photons are present that can excite the atom with resonant transitions. More sophisticated implementations of Feynman's proposal may avoid this issue.

6. Comparison of the Spectral Energy Density of the Vacuum Field and the Spectral Density of the Radiative Shift

The theory of Feynman proposes that the vacuum energy density in a large box containing H atoms, which are assumed to be in the 1S ground state, increases uniformly with the addition of the atoms. Feynman maintains that the total vacuum energy in the box increases by the Lamb shift times the number of atoms present [6,10]. If there were only one atom in a very large box, one would not expect the change in energy density to be spatially uniform but more concentrated near the atom. To develop a model of the spatial dependence of the change in energy density for one atom, the close relationship between the vacuum field and the radiative shift can be used. The spectral densities of the ground state shift and of the quantum vacuum with no H atoms present are both known. In the box, the vacuum field density must increase so that the integral gives the 1S Lamb shift. The spectral energy density of the vacuum field with no H atom present is equal to [6]

$$\rho_0(\omega) = \frac{\hbar\omega^3}{2\pi^2c^3}, \quad (50)$$

where c is in cm/sec and ω is in sec^{-1} . If frequency is measured in eV, so $\hbar\omega = E$, then the vacuum spectral energy density in $1/cc$ is

$$\rho_0(E) = \frac{E^3}{2\pi^2\hbar^3c^3}, \quad (51)$$

and $\int_{E_1}^{E_2} \rho_0(E)dE$ would be the energy density eV/cc in the energy interval E_1 to E_2 . The question being addressed is: what volume of vacuum energy of density $\rho_0(E)$ is required to supply the amount of energy needed for the radiative shift? The total radiative shift ΔE_1 be expressed as the integral of the vacuum energy density $\rho_0(E)$ over an effective volume $V_1(E)$,

$$\Delta E_1 = \int_0^{mc^2} dE \rho_0(E) V_1(E), \quad (52)$$

where the same upper limit for E is used as in previous calculations. Recall the definition (28) of the spectral shift:

$$\Delta E_1 = \int_0^{mc^2} dE \frac{\Delta E_1}{dE}. \quad (53)$$

Comparing Equations (52) and (53) and insuring the energy balance at each energy E , gives the effective spectral volume $V_1(E)$

$$V_1(E) = \frac{d\Delta E_1}{dE} \frac{1}{\rho_0(E)}. \quad (54)$$

The spectral volume $V_1(E)$ has the dimensions of cc and contains the amount of vacuum energy at energy value E that corresponds to the ground state spectral density at the same energy E . In Figure 10, for the 1S ground state radiative shift, the log of the spectral volume $V_1(E)$, in cubic Angstroms (\AA^3), is plotted versus the log of the energy, E .

For energies above about 100 eV, the spectral volume is less than 1\AA^3 , approximately the volume of the ground state wavefunction. For an energy of 1 eV, the spectral volume is 11850\AA^3 , corresponding to a sphere of radius about 14 \AA . This calculation predicts that there is a sphere of positive vacuum energy of radius 14 \AA around the atom corresponding to the 1 eV shift spectral density. Figure 11 shows the radius of the spherical spectral volume $V_1(E)$ for energies from 0.05 eV, with spectral radius of 278 \AA , to 23 eV, with radius 0.49 \AA .




Figure 10. Log-log plot of the spectral volume $V_1(E)$ as a function of E . The spectral volume $V_1(E)$ contains the free field vacuum energy at energy value E that corresponds to the ground state shift spectral density at the same energy E .




Figure 11. Log of the radius of the spherical spectral volume $V_1(E)$, as a function of the vacuum field energy E , from 0.05 eV to 23 eV.

7. Conclusions

The non-relativistic Lamb shift can be interpreted as due to the interaction of an atom with the fluctuating electromagnetic field of the quantum vacuum. We introduce the concept of a spectral shift density which is a function of frequency ω or energy $E = \hbar\omega$ of the vacuum field. The integral of the spectral density from $E = 0$ to the rest mass energy of an electron, 511 keV, gives the radiative shift. We report on calculations of the spectral density of the level shifts for 1S, 2S and 2P states based on a group theoretical analysis and compare the results to the spectral densities implicit in previous non-relativistic calculations of the Lamb shift by Bethe, Welton, and Power. The group theoretical calculation provides an explicit form for the spectral density over the entire spectral range with no summation over intermediate states. Bethe's approach requires a summation over an infinite number of states, including all bound and all scattering states, to obtain a comparable spectral density. The different approaches for asymptotic cases, for very large and very small energies E , are compared.

The calculations of the shift spectral density provide a new perspective on radiative shifts. The group theory approach as well as the approaches of Bethe, Power, and Welton all show the same $1/E$ high-frequency behavior for S states above about $E = \hbar\omega = 1000$ eV to 511 keV, namely an asymptotic spectral density for S states equal to $(4/3\pi)(\alpha(Z\alpha)^4 mc^2/n^3)(1/E)$ for principal quantum number n . The group theory calculation shows that about 76% of the ground state 1S shift is contributed by E above 1000 eV, which is essentially why all the approaches give approximately the same result for the 1S Lamb shift.

Only the Bethe and group theory calculations have the correct low-frequency behavior. For S states the spectral density increases linearly as E decreases to zero. Its maximum value is at $E = 0$ and for S states equals $(2\alpha/3\pi)(Z\alpha)^2/(n^2)$. This maximum value is about

$1/(Z\alpha)^2$ or about 2×10^4 larger than the high-frequency spectral density at $E = 511$ keV. Thus, low energies contribute much more to the shift for a given spectral interval than high energies. Energies below 13.6 eV contribute about 2.5 %. Because of the huge spectral range contributing to the shift, contributions to the shift from high energies are very important. Half the contribution to the 1S shift is from energies above 9050 eV.

The 2P shift has a very different spectral density from an S state: it is negative and has an asymptotic behavior that goes as $1/E^2$ rather than as $1/E$. Below about 20 eV, the absolute value of the 2P spectral density is much larger than the 2S spectral density and it dominates the 2S–2P shift spectral density, yet the total 2P shift is only about 0.3% of the total 2S shift.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Acknowledgments: I thank Peter Milonni for many insightful and enjoyable discussions, particularly about the resonant behavior of the index of refraction and the volume of vacuum energy corresponding to the spectral density, and I thank Lowell S. Brown for his observations, especially about the $1/\text{frequency}$ asymptotic behavior.

Conflicts of Interest: The author declares no conflicts of interest.

Appendix A. Eigenstates $|nlm;a\rangle$ of $1/Z\alpha$

To obtain an equation for these basis states $|nlm;a\rangle$, write Schrödinger's equation for a charged non-relativistic particle with momentum p and energy $E = -\frac{a^2}{2m}$ [27,28] in a Coulomb potential:

$$\left[p^2 + a^2 - \frac{2m\hbar c Z\alpha}{r} \right] |a\rangle = 0. \quad (\text{A1})$$

Solutions for $|a\rangle$ exist for certain critical values of the energy $E_n = -\frac{a_n^2}{2m}$ or equivalently for critical values of $a = a_n$ where $\frac{a_n}{mcZ\alpha} = \frac{1}{n}$. These are the usual energy eigenstates which are labeled as $|nlm;a_n\rangle$. Conversely a be fixed in value and $Z\alpha$ may have different values. If it has certain eigenvalues $Z\alpha_n$ then for any value of a there is another set of eigenvectors corresponding to eigenvalues $\frac{a}{mcZ\alpha_n} = \frac{1}{n}$. To demonstrate this we start by inserting factors of $1 = \sqrt{ar} \frac{1}{\sqrt{ar}}$ in Schrödinger's Equation (A1) obtaining

$$\left(\sqrt{ar}(p^2 + a^2)\sqrt{ar} - 2amZ\alpha \right) \frac{1}{\sqrt{ar}} |a\rangle = 0. \quad (\text{A2})$$

We can rewrite this equation, multiplying successively from the left by $\frac{1}{\sqrt{ar}}$, $\frac{1}{p^2 + a^2}$, and $\frac{1}{\sqrt{ar}}$, and then multiplying by a^2 , and dividing by $mcZ\alpha$, multiplying by $\sqrt{n\hbar}$ obtaining

$$\left[\frac{a}{mcZ\alpha} - K_1(a) \right] \sqrt{\frac{n\hbar}{ar}} |a\rangle = 0, \quad (\text{A3})$$

where

$$K_1(a) = \frac{1}{\sqrt{ar}} \frac{2a^2\hbar}{p^2 + a^2} \frac{1}{\sqrt{ar}}. \quad (\text{A4})$$

Solutions exist to this equation for eigenvalues of $1/Z\alpha$ such that $\frac{a}{mcZ\alpha_n} = \frac{1}{n}$:

$$\left(\frac{1}{n} - K_1(a) \right) |nlm;a\rangle = 0, \quad (\text{A5})$$

where

$$\sqrt{\frac{n\hbar}{ar}} |nlm;a\rangle = |nlm;a\rangle.$$

The $n\hbar$ in the square root insures the new states are also normalized to 1. The kernel $K_1(a)$ is bounded and Hermitian with respect to the eigenstates $|nlm; a\rangle$ of $1/Z\alpha$, therefore these eigenstates of $1/Z\alpha$ form a complete orthonormal basis for the hydrogen atom. Because the kernel is bounded, there are no continuum states in this representation. To show they have the same quantum numbers as the usual states, note that when $a = a_n$ the eigenstates of $K_1(a_n)$ becomes $|nlm; a_n\rangle$ and these corresponds to the usual energy eigenstates $|nlm; a_n\rangle$. The value of a in Equation (A5) can be changed to obtain these eigenstates using the dilation operator $D(\lambda) = e^{iS\lambda}$, where the dimensionless operator S , which is also a generator of transformations of $SO(4,2)$, is

$$S = \frac{1}{2\hbar}(\mathbf{p} \cdot \mathbf{r} + \mathbf{r} \cdot \mathbf{p}). \quad (\text{A6})$$

S transforms the canonical variables:

$$D(\lambda)\mathbf{p}D^{-1}(\lambda) = e^{-\lambda}\mathbf{r},$$

$$D(\lambda)\mathbf{r}D^{-1}(\lambda) = e^{\lambda}\mathbf{r}.$$

Operating on $K_1(a)$ with $D(\lambda)$ gives:

$$D(\lambda)K_1(a)D^{-1}(\lambda) = K_1(ae^{\lambda}).$$

If λ is chosen as

$$\lambda_n = \ln(a_n/a),$$

then $ae^{\lambda_n} = a_n$. Thus, operating with $D(\lambda_n)$ on Equation (A5) yields:

$$\left(\frac{1}{n} - K_1(a_n)\right)D(\lambda_n)|nlm; a\rangle = 0. \quad (\text{A7})$$

This is the equation for the usual Schrödinger energy eigenstates, so

$$D(\lambda_n)|nlm; a\rangle = |nlm; a_n\rangle = \sqrt{\frac{n\hbar}{a_n r}}|nlm; a_n\rangle. \quad (\text{A8})$$

The usual Schrödinger energy eigenstates $|nlm; a_n\rangle$ can be expressed in terms of the eigenstates of $1/Z\alpha$ as

$$|nlm; a_n\rangle = \sqrt{\frac{a_n r}{n\hbar}}D(\lambda_n)|nlm; a\rangle. \quad (\text{A9})$$

This relationship shows that the complete basis functions $|nlm; a\rangle$ of $1/Z\alpha$ are proportional to the ordinary bound state energy wavefunctions and therefore have the same quantum numbers as the ordinary bound states [27,28]. A comparable set of $1/Z\alpha$ eigenstates useful for momentum space calculations is derived in [27].

Appendix B. Derivation of Group Theoretical Formula for the Shift Spectral Density

The group theoretical approach is based solely on the Schrödinger and Klein-Gordon equations of motion in the non-relativistic dipole approximation. We obtain a result [27]:

$$\Delta E_{NL} = \frac{2\alpha}{3\pi(mc)^2} \int_0^{\hbar\omega_c} dE \langle NL | p_i \frac{H - E_N}{H - (E_N - E) - i\epsilon} p_i | NL \rangle, \quad (\text{A10})$$

where $E = \hbar\omega$, $H = \frac{p^2}{2m} - \frac{Z\alpha\hbar c}{r}$ and the states $|NL\rangle$ are the usual H atom energy eigenstates. ω_c is a cutoff frequency for the integration that is taken as $\hbar\omega_c = mc^2$. Inserting a complete set of states in this expression yields Bethe's result (6) a step avoided with the group

theoretical approach. Adding and subtracting E from the numerator in Equation (A10), we find the real part of the shift is

$$\Delta E_{NL} = \frac{2\alpha}{3\pi(mc)^2} \text{Re} \int_0^{\hbar\omega_c} dE [\langle NL|p^2|NL\rangle - E\Omega_{NL}], \quad (\text{A11})$$

where

$$\Omega_{NL} = \langle NL|p_i \frac{1}{H - E_N + \hbar\omega - i\epsilon} p_i|NL\rangle. \quad (\text{A12})$$

The goal is to convert the matrix element Ω_{NL} to a matrix element of a function of $\text{SO}(4,2)$ generators taken between a new set of basis states $|nlm;a\rangle$, which are complete with no scattering states, where $a = \sqrt{2m|E|}$, and n, l, m have their usual meaning and values. The new basis states $|nlm;a\rangle$ are eigenstates of $(Z\alpha)^{-1}$ [27,28]. Sometimes we write them as $|nlm\rangle$ with the a implicit.

A generator of $\text{SO}(4,1)$ is $\Gamma_0 = 1/K_1(a)$, defined in Equation (A4), so

$$(\Gamma_0 - n)|nlm;a\rangle = 0. \quad (\text{A13})$$

This is Schrödinger's equation in the language of $\text{SO}(4,2)$.

Several more generators need to be defined. Since the algebra of $\text{SO}(4,2)$ generators closes, commutators of generators must also be generators. To find Γ_4 calculate $\Gamma_4 = -i[S, \Gamma_0]$, where the generator S is defined in Equation (A6), obtaining

$$\Gamma_4 = \frac{1}{2\hbar} \left(\frac{\sqrt{r}p^2\sqrt{r}}{a} - ar \right), \quad \Gamma_0 = \frac{1}{2\hbar} \left(\frac{\sqrt{r}p^2\sqrt{r}}{a} + ar \right). \quad (\text{A14})$$

The generators $(\Gamma_4, S, \Gamma_0) = (j_1, j_2, j_3)$ form a $\text{O}(2,1)$ subgroup of $\text{SO}(4,2)$ and $S = i[\Gamma_4, \Gamma_0]$, $\Gamma_0 = -i[S, \Gamma_4]$ and for our representations, $\Gamma_0^2 - \Gamma_4^2 - S^2 = L^2 = l(l+1)$. The scale change S transforms $\Gamma_0 \equiv \Gamma_0(a)$ according to the equation,

$$e^{i\lambda S} \Gamma_0(a) e^{-i\lambda S} = \Gamma_0(e^\lambda a) = \Gamma_0 \cosh \lambda - \Gamma_4 \sinh \lambda, \quad (\text{A15})$$

and similarly,

$$e^{i\lambda S} \Gamma_4(a) e^{-i\lambda S} = \Gamma_4(e^\lambda a) = \Gamma_4 \cosh \lambda - \Gamma_0 \sinh \lambda. \quad (\text{A16})$$

Finally define a three vector of generators proportional to the momentum

$$\Gamma_i = \frac{1}{\hbar} \sqrt{r} p_i \sqrt{r}. \quad (\text{A17})$$

The quantity $\Gamma = (\Gamma_0, \Gamma_1, \Gamma_2, \Gamma_3, \Gamma_4)$ is a 5-vector of generators under transformations generated by $\text{SO}(4,2)$. For the representation of $\text{SO}(4,2)$ based on the states $|nlm\rangle$, all generators are Hermitian, and $\Gamma^2 = \Gamma_A \Gamma^A = -\Gamma_0^2 + \Gamma_1^2 + \Gamma_2^2 + \Gamma_3^2 + \Gamma_4^2 = 1$ for our representation, and $g_{AB} = (-1, 1, 1, 1, 1)$ for $A, B = 0, 1, 2, 3, 4$. The commutators of the components of the five vector are also generators of $\text{SO}(4,2)$ transformations.

Inserting factors of $1 = \sqrt{ar} \frac{1}{\sqrt{ar}}$ and using the definitions of the generators we can transform Equation (A12) to

$$\Omega_{NL} = \frac{mv}{N^2} (NL | \Gamma_i \frac{1}{\Gamma n(\xi) - \nu} \Gamma_i | NL), \quad (\text{A18})$$

where

$$n^0(\xi) = \frac{2 + \xi}{2\sqrt{1 + \xi}} = \cosh \phi, \quad n^i = 0, \quad n^4(\xi) = -\frac{\xi}{2\sqrt{1 + \xi}} = -\sinh \phi, \quad (\text{A19})$$

and

$$\xi = \frac{\hbar\omega}{|E_N|} \quad \nu = \frac{N}{\sqrt{1 + \xi}} = Ne^{-\phi}. \quad (\text{A20})$$

From the definitions, $\phi = \frac{1}{2} \ln(1 + \xi) > 0$ and $n_A(\xi)n^A(\xi) = -1$. The contraction over i in Ω_{NL} may be evaluated using the group theoretical formula [27]:

$$\sum_B \Gamma_B f(n\Gamma) \Gamma^B = \frac{1}{2} (n\Gamma + 1)^2 f(n\Gamma + 1) + \frac{1}{2} (n\Gamma - 1)^2 f(n\Gamma - 1) - (n\Gamma)^2 f(n\Gamma). \quad (\text{A21})$$

Applying the contraction formula to the the integral representation,

$$f(n\Gamma) = \frac{1}{\Gamma n - \nu} = \int_0^\infty ds e^{\nu s} e^{-n\Gamma s}, \quad (\text{A22})$$

gives the result

$$\Gamma_A \frac{1}{\Gamma n - \nu} \Gamma^A = -2\nu \int_0^\infty ds e^{\nu s} \frac{d}{ds} \left(\sinh^2 \frac{s}{2} e^{-n\Gamma s} \right). \quad (\text{A23})$$

Applying this to Equation (A18) gives:

$$\begin{aligned} \Omega_{NL} = & -2 \frac{mv^2}{N^2} \int_0^\infty ds e^{\nu s} \frac{d}{ds} \left(\sinh^2 \frac{s}{2} M_{NL}(s) \right) \\ & - m \frac{\nu}{N^2} (NL | \Gamma_4 \frac{1}{\Gamma n(\xi) - \nu} \Gamma_4 | NL) + m \frac{\nu}{N^2} (NL | \Gamma_0 \frac{1}{\Gamma n(\xi) - \nu} \Gamma_0 | NL) \end{aligned} \quad (\text{A24})$$

where

$$M_{NL}(s) = (NL | e^{-\Gamma n(\xi)s} | NL). \quad (\text{A25})$$

In order to evaluate the last two terms in Equation (A24) use $\Gamma_0 | NL) = N | NL)$ and express the action of Γ_4 on our states as $\Gamma_4 = N - (1/\sinh \phi)(\Gamma n(\xi) - \nu)$. This expression for Γ_4 is derived from Equations (A18) and (A19): $\Gamma n(\xi) - \nu = \Gamma_0 \cosh \phi - \Gamma_4 \sinh \phi - \nu$, and then substituting Equation (A20), $\nu = Ne^{-\phi}$. Using the virial theorem $(NLM | p^2 | NLM) = a_N^2$, we find that the term in p^2 in Equation (A11) exactly cancels the last two terms in Ω_{NL} , yielding the result for the level shift:

$$\Delta E_{NL} = \frac{4mc^2 \alpha (Z\alpha)^4}{3\pi N^4} \int_0^{\phi_c} d\phi \sinh \phi e^\phi \int_0^\infty ds e^{\nu s} \frac{d}{ds} \left(\sinh^2 \frac{s}{2} M_{NL}(s) \right) \quad (\text{A26})$$

where

$$\phi_c = \frac{1}{2} \ln \left(1 + \frac{\hbar \omega_c}{|E_N|} \right) = \frac{1}{2} \ln \left(1 + \frac{2N^2}{(Z\alpha)^2} \right). \quad (\text{A27})$$

To derive a generating function for the shifts for any eigenstate characterized by N and L multiply Equation (A26) by $N^4 e^{-\beta N}$ and sum over all N , $N \geq L + 1$. To simplify the right side of the resulting equation, use the definition (A25) and the fact that Γ_4 , S , and Γ_0 form an $O(2,1)$ algebra, obtaining:

$$\sum_{N=L+1}^\infty e^{-\beta N} M_{NL} = \sum_{N=L+1}^\infty (NL | e^{-j \cdot \psi} | NL), \quad (\text{A28})$$

where

$$e^{-j \cdot \psi} \equiv e^{-\beta \Gamma_0} e^{-s \Gamma n(\xi)}. \quad (\text{A29})$$

Perform a j transformation generated by $e^{i\phi S}$, such that $e^{-j \cdot \psi} \rightarrow e^{-j_3 \psi} = e^{-\Gamma_0 \psi}$. The trace is invariant with respect to this transformation, therefore:

$$\sum_{N=L+1}^\infty e^{-\beta N} M_{NL} = \sum_{N=L+1}^\infty (NL | e^{-j_3 \psi} | NL) = \sum_{N=L+1}^\infty e^{-N\psi} = \frac{e^{-\psi(L+1)}}{1 - e^{-\psi}}, \quad (\text{A30})$$

where $(NL | \Gamma_0 | NL) = N$ is used.

In order to find a particular M_{NL} , expand the right hand side of Equation (A30) in powers of $e^{-\beta}$ and equate the coefficients to those on the left hand side. Using the

isomorphism between j and the Pauli σ matrices $(\Gamma_4, S, \Gamma_0) \rightarrow (j_1, j_2, j_3) \rightarrow (\frac{1}{2}\sigma_1, \frac{1}{2}\sigma_2, \frac{1}{2}\sigma_3)$ gives the result:

$$\cosh \frac{\psi}{2} = \cosh \frac{\beta}{2} \cosh \frac{s}{2} + \sinh \frac{\beta}{2} \sinh \frac{s}{2} \cosh \phi. \quad (\text{A31})$$

Rewriting this equation gives:

$$e^{+\frac{1}{2}\psi} = d e^{\frac{1}{2}\beta} + b e^{-\frac{1}{2}\beta} - e^{-\frac{1}{2}\psi}, \quad (\text{A32})$$

where

$$\begin{aligned} d &= \cosh \frac{s}{2} + \sinh \frac{s}{2} \cosh \phi, \\ b &= \cosh \frac{s}{2} - \sinh \frac{s}{2} \cosh \phi. \end{aligned} \quad (\text{A33})$$

Let β become large, which implies large ψ , and iterate the equation for $e^{-\frac{1}{2}\psi}$ to obtain

$$e^{-\psi} = A e^{-\beta} \left[1 + A_1 e^{-\beta} + A_2 e^{-2\beta} + \dots \right], \quad (\text{A34})$$

where $A = 1/d^2$ and $A_1 = -(2/d)(b - d^{-1})$. To obtain M_{NL} , expand the right side of Equation (A30) in powers of ψ :

$$\frac{e^{-\psi(L+1)}}{1 - e^{-\psi}} = \sum_{m=1}^{\infty} e^{-\psi(m+L)}. \quad (\text{A35})$$

For large β , it follows from Equations (A30), (A34) and (A35) that

$$\sum_{N=L+1}^{\infty} e^{-\beta N} M_{NL} = \sum_{m=1}^{\infty} \left[e^{-\beta} A (1 + A_1 e^{-\beta} + A_2 e^{-2\beta} + \dots) \right]^{m+L}. \quad (\text{A36})$$

Using the multinomial theorem [32] the right side of Equation (A36) becomes

$$\sum_{m=1}^{\infty} A^{m+L} \sum_{r,s,t,\dots} \frac{(m+L)!}{r!s!t! \dots} A_1^s A_2^t \dots e^{-\beta(m+L+s+2t+\dots)}, \quad (\text{A37})$$

where $r + s + t + \dots = m + L$. To obtain the expression for M_{NL} , we note N is the coefficient of β so $N = m + L + s + 2t + \dots = r + 2s + 3t + \dots$. Accordingly we find

$$M_{NL} = \sum_{r,s,t,\dots} A^{(r+s+t+\dots)} \frac{(r+s+t+\dots)!}{r!s!t! \dots} A_1^s A_2^t \dots, \quad (\text{A38})$$

where $r + s + t + \dots = N$ and $r + s + t + \dots > L$. As Equation (A26) indicates, the 1S shift corresponds to the matrix element M_{10} , which multiplies $e^{-\beta}$, so $M_{10} = A$. For the 2S shift $M_{20} = A^2 + A A_1$, and for the 2P shift $M_{21} = A^2$. Therefore the radiative shift for the 1S ground state is

$$\text{Re} \Delta E_{10} = \frac{4mc^2 \alpha (Z\alpha)^4}{3\pi} \int_0^{\phi_c} d\phi e^{\phi} \sinh \phi \int_0^{\infty} ds e^{se^{-\phi}} \frac{d}{ds} \frac{1}{(\coth \frac{s}{2} + \cosh \phi)^2}. \quad (\text{A39})$$

The shift for the 2S–2P level is

$$\text{Re}(\Delta E_{20} - \Delta E_{21}) = \frac{m\alpha(Z\alpha)^4}{6\pi} \int_0^{\phi_c} d\phi e^{\phi} \sinh^3 \phi \int_0^{\infty} ds e^{2se^{-\phi}} \frac{d}{ds} \frac{1}{(\coth \frac{s}{2} + \cosh \phi)^4}. \quad (\text{A40})$$

References

1. Choudhuri, A.R. *Astrophysics for Physicists*; Cambridge University Press: Cambridge, UK, 2010. [CrossRef]
2. Carroll, S.M. Addressing the quantum measurement problem. *Phys. Today* **2022**, *75*, 62–64. [CrossRef]
3. D’Espagnat, B. *Veiled Reality: An Analysis of Present-Day Quantum Mechanical Concepts*; CRC Press/Taylor & Francis Group: Boca Raton, FL, USA, 2013. [CrossRef]

4. Beyer, A.; Maisenbacher, L.; Matveev, A.; Pohl, R.; Khabarova, K.; Grinin, A.; Lamour, T.; Yost, D.C.; Hänsch, T.W.; Kolachevsky, N.; et al. The Rydberg constant and proton size from atomic hydrogen. *Science* **2017**, *358*, 79–85. [[CrossRef](#)] [[PubMed](#)]
5. Berestetskii, V.B.; Lifshitz, E.M.; Pitaevskii, L.P. *Quantum Electrodynamics. Course of Theoretical Physics: Volume 4, Second Edition*; Pergamon Press: Oxford, UK, 1982. [[CrossRef](#)]
6. Milonni, P.W. *The Quantum Vacuum. An Introduction to Quantum Electrodynamics*; Academic Press, Inc.: San Diego, CA, USA, 1994. [[CrossRef](#)]
7. Lamb, W.E.; Retherford, R.C. Fine structure of the hydrogen atom by a microwave method. *Phys. Rev.* **1947**, *72*, 241–243. [[CrossRef](#)]
8. Bethe, H.A. The electromagnetic shift of energy levels. *Phys. Rev.* **1947**, *72*, 339–341. [[CrossRef](#)]
9. Welton, T.A. Some observable effects of the quantum-mechanical fluctuations of the electromagnetic field. *Phys. Rev.* **1948**, *74*, 1157–1168. [[CrossRef](#)]
10. Power, E.A. Zero-point energy and the Lamb shift. *Am. J. Phys.* **1966**, *34*, 516–518. [[CrossRef](#)]
11. Compagno, G.; Passante, R.; Persico, F., *Atom-Field Interactions and Dressed Atoms*; Cambridge University Press: Cambridge, UK, 1995. [[CrossRef](#)]
12. Maclay, G.J. History and some aspects of the Lamb shift. *Physics* **2020**, *2*, 105–149. [[CrossRef](#)]
13. Dyson, F.J. The electromagnetic shift of energy levels. *Phys. Rev.* **1948**, *73*, 617–626. [[CrossRef](#)]
14. French, J.B.; Weisskopf, V.F. The electromagnetic shift of energy levels. *Phys. Rev.* **1949**, *75*, 1240–1248. [[CrossRef](#)]
15. Kroll, N.M.; Lamb, W.E., Jr. On the self-energy of a bound electron. *Phys. Rev.* **1949**, *75*, 388–398. [[CrossRef](#)]
16. Weinberg, S. *The Quantum Theory of Fields. Volume 1: Foundations*; Cambridge University Press: Cambridge, UK, 1995. [[CrossRef](#)]
17. Erickson, G.W.; Yennie, D.R. Radiative level shifts, I. Formulation and lowest order Lamb shift. *Ann. Phys.* **1965**, *35*, 271–313. [[CrossRef](#)]
18. Schwinger, J. Quantum electrodynamics. I. A covariant formulation. *Phys. Rev.* **1948**, *74*, 1439–1461. [[CrossRef](#)]
19. Feynman, R.P. The development of the space-time view of quantum electrodynamics. *Phys. Today* **1966**, *19*, 31–44. [[CrossRef](#)]
20. Dyson, F.J. The radiation theories of Tomonaga, Schwinger, and Feynman. *Phys. Rev.* **1949**, *75*, 486–502. [[CrossRef](#)]
21. Akhiezer, A.I.; Berestetskii, V.B. *Quantum Electrodynamics*; Interscience Publishers/John Wiley and Sons, Inc.: New York, NY, USA, 1965. Available online: <https://archive.org/details/akhiezer-berestetskii-quantum-electrodynamics/page/n7/mode/2up> (accessed on 30 August 2022).
22. Itzykson, C.; Zuber, J.-B. *Quantum Field Theory*; McGraw-Hill Inc.: New York, NY, USA, 1980.
23. Grotch, H. Status of the theory of the Hydrogen Lamb shift. *Found. Phys.* **1994**, *24*, 249–272. [[CrossRef](#)]
24. Feynman, R.P. The present status of quantum electrodynamics. In Proceedings of the 12th La Théorie Quantique des Champs. Douzième Conseil de Physique, Brussels, Belgium, 9–14 October 1961; Stoops, R., Ed.; Interscience Publishers/John Wiley and Sons, Inc.: New York, NY, USA, 1962; pp. 61–99. Available online: http://www.solvayinstitutes.be/html/solvayconf_physics.html (accessed on 30 August 2022).
25. Lieber, M. $O(4)$ symmetry of the hydrogen atom and the Lamb shift. *Phys. Rev.* **1968**, *174*, 2037–2054. [[CrossRef](#)]
26. Huff, R.W. Simplified calculation of Lamb shift using algebraic techniques. *Phys. Rev.* **1969**, *186*, 1367–1379. [[CrossRef](#)]
27. Maclay, G.J., Dynamical symmetries of the H atom, one of the most important tools of modern physics: $SO(4)$ to $SO(4,2)$, background, theory, and use in calculating radiative shifts. *Symmetry* **2020**, *12*, 1323. [[CrossRef](#)]
28. Brown, L.S. Bounds on screening corrections in beta decay. *Phys. Rev.* **1964**, *135*, B314–B319. [[CrossRef](#)]
29. Schweber, S.S., *QED and the men who made it: Dyson, Feynman, Schwinger, and Tomonaga*; Princeton University Press: Princeton, NJ, USA, 1994.
30. Bethe, H.A.; Salpeter, E.E., *The Quantum Mechanics of One- and Two-Electron Atoms*; Springer-Verlag: Berlin/Heidelberg, Germany, 1957. [[CrossRef](#)]
31. Sakurai, J.J. *Modern Quantum Mechanics*; Addison-Wesley Publishing Company, Inc.: New York, NY, USA, 1994. Available online: <https://www.fisica.net/mecanica-quantica/Sakurai%20-%20Modern%20Quantum%20Mechanics.pdf> (accessed online on 30 August 2022).
32. Morse, P.M.; Feshbach, H. *Methods of Theoretical Physics*; McGraw-Hill Book Company, Inc.: New York, NY, USA, 1953; Volume 2.