



First Measurements of Differential Cross Sections in Kinematic Imbalance Variables with the MicroBooNE Detector



Maitreyee Mahasweta Moudgalya on behalf of the MicroBooNE Collaboration

1. Introduction


- The Micro Booster Neutrino Experiment (MicroBooNE) is a Liquid Argon Time Projection Chamber (LArTPC) based at Fermilab.
- MicroBooNE ran from 2015 to 2020, forming the largest dataset of neutrino interactions with argon in the world to date.
- It has been designed for precision neutrino physics measurements.



- The scintillation light produced at the time of the interaction is collected by 32 PMTs.
- The ionisation trails are carried to the anode wire planes by the electric field.
- These charged particle trajectories are reconstructed using the known positions of the wires and the recorded drift time of the ionisation.


2. Neutrino Interaction Modeling Challenge

- Broad range of neutrino fluxes
- Many known unknowns that must be accurately simulated:
 - Ground states, Fermi motion
 - Neutrino interaction mechanisms
 - Final state interactions (FSI)
 - ...




Any mismodeling can limit experimental sensitivity → Issue for future flagship experiments like DUNE.

3. Transverse Kinematic Imbalance (TKI)



4. CC1 μ 1p0 π Differential Cross Section Results



- Can use the MicroBooNE detector to set constraints:
 - First double-differential single-proton cross section measurement on argon.
 - Uses ~ 50% of available dataset and the Booster Neutrino Beam (BNB) at Fermilab.
 - Identified phase-space regions that are separately sensitive to Fermi motion and FSI.


BNB Run 1 Data
Run 5127 Subrun 138 Event 6904

CC1p0 π

Candidate μ

Candidate p



7 cm



Similar ongoing analysis with ν_e (the appearance signal)!

Interesting regions of QE-dominated phase-space identified using TKI truth variables:

- Cross section measurement in such a “non-FSI region” is likely to yield small uncertainties.
- Can we leverage this to perform an oscillation analysis and to reduce the overall systematic uncertainty?



QE-like Signal Definition:

- 1 electron
- 1 proton with $P_p > 0.3$ GeV/c
- No π^\pm with $P_\pi > 70$ GeV/c
- No π^0 or heavier mesons
- Any number of neutrons

More resonant events passed the reco selection cuts than expected.
→ Still under investigation.

Purity ~ 28.12%
Efficiency ~ 24.37%

Need sufficient resolution (and statistics) to perform an oscillation measurement.
→ especially for the electron showers (harder to reconstruct).