

On the perturbed second Painlevé equation*

Joshua Holroyd¹  and Nalini Joshi^{1,***} 

School of Mathematics and Statistics F07, University of Sydney, Sydney, NSW 2006, Australia

E-mail: nalini.joshi@sydney.edu.au

Received 3 September 2022; revised 3 January 2023

Accepted for publication 6 January 2023

Published 23 January 2023



Abstract

We consider a perturbed version of the second Painlevé equation (P_{II}), which arises in applications, and show that it possesses solutions analogous to the celebrated Hastings–McLeod and tritronquée solutions of P_{II} . The Hastings–McLeod-type solution of the perturbed equation is holomorphic, real-valued and positive on the whole real-line, while the tritronquée-type solution is holomorphic in a large sector of the complex plane. These properties also characterise the corresponding solutions of P_{II} and are surprising because the perturbed equation does not possess additional distinctive properties that characterise P_{II} , particularly the Painlevé property.

Keywords: asymptotics, exponential asymptotics, Painlevé equations, Hastings–McLeod solution, tritronquée solution

1. Introduction

The second Painlevé equation (P_{II}) has a rich history as a mathematical model, arising in a very wide range of applications (see, for example, [9, 15, 31, 32]). In many of these applications, the particular case of P_{II} that arises is the case $\alpha = 0$ of

$$P_{II} : \quad y''(x) = 2y(x)^3 + xy(x) + \alpha,$$

* Dedicated to Sir Michael V. Berry on the occasion of his 80th birthday.

¹ NJ's research was supported by an Australian Research Council Discovery Project #DP210100129.

** Author to whom any correspondence should be addressed.



Original Content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

where primes denote differentiation in x and α is a given parameter. In this paper, we consider a perturbed version of P_{II} with $\alpha = 0$:

$$P_{II}(\mu) : \quad y''(x) = 2y(x)^3 + x^\mu y(x), \quad y : \mathbb{C} \rightarrow \mathbb{C}, \quad \mu \in \mathbb{Z}^+.$$

The main results of this paper show that there exists (i) a holomorphic, real-valued, positive solution of $P_{II}(\mu)$ on the real-line $x \in \mathbb{R}$; and (ii) a solution that is asymptotic to a power-series expansion in x , as $|x| \rightarrow \infty$, in a sector of angular extent $4\pi/(\mu + 2)$ in \mathbb{C} . These properties mirror two well-known solutions of P_{II} , called the Hastings–McLeod solution [19] and Boutroux’s tritronquée solution [7] respectively, which are distinguished by their global properties.

P_{II} belongs to a class of six ordinary differential equations (ODEs), called the Painlevé equations, whose solutions are known to have only poles as movable singularities [11, 21]. Every non-rational solution of P_{II} is known to possess an infinite number of movable poles [20], which are typically distributed across the interior of each sector S_n of the complex plane given by [7]

$$S_n = \{x \in \mathbb{C} \mid (n-1)\pi/3 \leq \arg x \leq n\pi/3\}, \quad n \in \mathbb{Z}, \quad \text{mod } 6. \quad (1.1)$$

(Here, the word ‘movable’ refers to the location of each pole depending on initial conditions.) Therefore, solutions that are free of movable poles across one or more sectors S_n are uncommon, while those free of poles in a domain containing the real line are of particular interest for physical applications. The tritronquée solution is free of poles in a wide sector of the complex plane, given by four contiguous sectors S_n , while the Hastings–McLeod solution is free of poles and analytic on the whole real-line.

The Hastings–McLeod solution now plays a critical role in modern random matrix theory, due to its appearance in the Tracy–Widom distribution, describing the distribution of the largest eigenvalue of Gaussian unitary ensembles of random matrices, as the size of each matrix approaches infinity. The Hastings–McLeod solution also arises in the analogous distribution functions for the Gaussian orthogonal and symplectic ensembles [9, 13, 31].

There are many more physical models in which P_{II} arises. We mention, in particular, fluid dynamics [1, 8, 9, 14, 18, 26], mathematical physics [15, 28–30, 32], and electrodynamics [4, 5]. In particular, the second Painlevé equation arises in a model of steady one-dimensional two-ion electro-diffusion [4]. A review may be found in [11]. In connection with plasma physics, de Boer and Ludford [12] posed the question of the existence of a solution of a generalised form of P_{II} with specific boundary conditions. Hastings and McLeod [19] not only showed the existence of a solution to this problem, but they also proved it to be unique; this is the now-famous Hastings–McLeod solution of P_{II} . It is natural in many of the physical contexts to consider small perturbations of the model of interest. We describe such a perturbation of Bass’ electro-diffusion model in the [appendix](#) and show how $P_{II}(\mu)$ arises naturally.

A perturbed version of the first Painlevé equation has been previously considered [22, 24]. Tronquée and tritronquée solutions of perturbed P_I were shown to exist (and be unique in the tritronquée case) in [22]. Meanwhile, [24] studies the full exponentially-improved asymptotic expansion corresponding to a formal series solution of perturbed P_I . These tronquée solutions are uniquely characterised by the Stokes multiplier, which is an arbitrary constant multiplying the exponentially small terms. Asymptotic approximations for the locations of singularities along boundaries of validity are deduced in terms of the Stokes multipliers [24].

In the case of $P_{II}(\mu)$, tronquée solutions are associated with a formal power-series solution $y_f \sim (-x^\mu/2)^{1/2}$ as $|x| \rightarrow \infty$ (given in full in section 3), and we mention that the (singly) exponentially small perturbation is given by

$$y \sim y_f + kx^{-\mu/4} \exp\left(\frac{2}{\mu+2} (-2x^{\mu+2})^{1/2} + o(1)\right),$$

where the square root branch depends on the sector of the complex plane, and k is the free constant multiplying exponentially small behaviour. The tritronquée solution, which we study in section 3, corresponds to the unique $k=0$ case of this expansion.

1.1. Main results

We identify a tritronquée solution of $P_{II}(\mu)$, through asymptotic expansions of solutions as $|x| \rightarrow \infty$; showing that a true solution asymptotic to such a series in the sector $|\arg(x)| \leq 2\pi/(\mu+2)$ exists and is unique. These results are given in propositions 3.4 and 3.7. We also consider a boundary-value problem on the real-line; as $x \rightarrow -\infty$, the solution is asymptotic to a power-series expansion, and as $x \rightarrow +\infty$, the solution decays to zero exponentially fast. We show that this problem has a unique solution in the case of $\mu = 2k+1$, $k \in \mathbb{N}$, which generalises the Hastings–McLeod solution. We give this result in theorem 4.2.

1.2. Background

The Painlevé equations were found in a search for ODEs that defined new higher transcendental functions a century ago. Here the word ‘new’ refers to these functions not being able to be expressed in terms of any classical operations applied to previously known such functions (we refer to definitions of these terms in [33]). Both the Hastings–McLeod and tritronquée solutions are examples of such transcendental functions.

For special values of the parameter α , P_{II} admits hierarchies of rational solutions [2, 3, 10, 16] and special solutions related to the classical Airy functions [17]. Beyond special solutions, transcendental solutions that vanish as $x \rightarrow +\infty$ satisfy the Airy equation to leading order, in the case when $\alpha=0$. The Hastings–McLeod solution first attracted attention in the asymptotic analysis of such solutions.

The asymptotic analysis of the Painlevé equations was initiated by Boutroux [7], who also briefly considered the generalised equation $P_{II}(\mu)$. Boutroux showed that the transformation of variables

$$y(x) = x^{1/2}u(z) \quad \text{and} \quad z = \frac{2}{3}x^{3/2},$$

leads to a transformed version of P_{II} which is more amenable to asymptotic analysis, that is

$$u''(z) = 2u^3(z) + u(z) + \frac{2\alpha}{3z} - \frac{u'(z)}{z} + \frac{u(z)}{9z^2}.$$

The limit $|x| \rightarrow \infty$ now becomes $|z| \rightarrow \infty$, and direct observation shows that a (Jacobi) elliptic function describes the leading-order behaviour of $u(z)$, solving the ODE

$$w''(z) = 2w^3(z) + w(z).$$

Boutroux went on to describe how lines of movable poles curve around to approach the boundaries of the sectors S_n described by equation (1.1).

1.3. Outline

In section 2, we describe decaying solutions of $P_{II}(\mu)$ as $|x| \rightarrow \infty$ on the real-line. We prove the existence and uniqueness of the analogue of Boutroux’s tritronquée solution for $P_{II}(\mu)$ and its asymptotic validity in a broad sector of the complex plane in section 3. In section 4, we prove the existence and uniqueness of a natural generalisation of the Hastings–McLeod

solution. The paper's results are summarised in section 5. Finally, we show how $P_{II}(\mu)$ arises in a physical setting in the [appendix](#).

2. Asymptotics of decaying solutions on the real-line

Since $\mu > 0$, the decaying solutions of $P_{II}(\mu)$ satisfy the leading order equation

$$y''(x) \sim x^\mu y(x),$$

as $x \rightarrow \pm\infty$ on the real-line. We analyse the asymptotic behaviours of $y(x)$, which satisfy this leading-order equation.

The linear ODE $\phi''(x) = x^\mu \phi(x)$ is a generalised Airy equation and is satisfied by two linearly independent functions

$$f(x) = \sqrt{\frac{2x}{\pi\alpha}} K_{1/(2\alpha)}\left(\frac{x^\alpha}{\alpha}\right) \quad \text{and} \quad g(x) = \sqrt{\frac{2\pi x}{\alpha}} I_{1/(2\alpha)}\left(\frac{x^\alpha}{\alpha}\right), \quad (2.1)$$

where $\alpha = (\mu + 2)/2$, and I_ν and K_ν are the modified Bessel functions of the first and second kind respectively. We choose this specific scaling of basis solutions as it leads to a simplification of their asymptotic expansions in the limit $x \rightarrow +\infty$; these expansions are readily available using known results for I_ν and K_ν , see [27, §10.40(i)]. The functions $f(x)$ and $g(x)$ are analogues of the well known Airy functions $\text{Ai}(x)$ and $\text{Bi}(x)$, reducing to scaled versions of these when $\alpha = 3/2$ (or equivalently $\mu = 1$).

Recalling that μ is positive and real, we find that

$$y(x) \sim kf(x) = kx^{-\mu/4} \exp\left(-\frac{2}{\mu+2}x^{(\mu+2)/2} + o(1)\right), \quad \text{as } x \rightarrow +\infty, \quad (2.2)$$

for some arbitrary constant k . As is well known for Airy functions, a solution $y(x)$ of $P_{II}(\mu)$ that decays to zero as $|x| \rightarrow \infty$ may not be exponentially small in other directions. In particular, for every odd, positive integer μ , the negative real semi-axis is an anti-Stokes line [6], meaning that the exponential contribution is purely oscillatory as $|x| \rightarrow \infty$. Given an odd positive integer $\mu > 1$, standard asymptotic techniques lead to

$$y(x) \sim c_1|x|^{-\mu/4} \sin\left(\frac{2}{\mu+2}|x|^{(\mu+2)/2} - c_2 + o(1)\right) \quad \text{as } x \rightarrow -\infty, \quad (2.3)$$

for arbitrary constants c_1 and c_2 . In the case $\mu = 1$, more detailed results were described by Hastings and McLeod [19].

3. Tritronquée solutions

The main object of study in this section is an asymptotic series, which solves $P_{II}(\mu)$ given in equation (3.2). We prove the existence and uniqueness of a true solution that is asymptotic to this series for $|\arg(x)| \leq 2\pi/(\mu + 2)$ as $|x| \rightarrow \infty$. This solution corresponds to the tritonquée solution in the case $\mu = 1$.

We refer the reader to Boutroux [7] for detailed descriptions of the solutions he called *intégrales tronquées*, which are pole-free (for sufficiently large x) within two adjacent sectors of the form S_n defined in equation (1.1). The region $S_n \cup S_{n+1}$ ($n \in \mathbb{Z} \bmod 6$) described by such a pair of sectors has a bisector given by a ray $\arg x = n\pi/3$. The term tritonquée arises from the fact that any line of poles that originally lies on such a ray must be truncated as $|x|$ increases (due to the pole-free nature of the solution).

There are three such rays in four contiguous sectors $S_n \cup S_{n+1} \cup S_{n+2} \cup S_{n+3}$ ($n \in \mathbb{Z} \bmod 6$). Boutroux showed that, for each choice of such a region, there exists a unique solution he called *tritronquée* (tritronquée), which is asymptotically pole-free in the region. For $P_{II}(\mu)$, these special rays are given by $\arg(x) = n\pi/(\mu + 2)$ for $n \in \mathbb{Z}$.

We deduce an asymptotic series expansion of a tritronquée solution of $P_{II}(\mu)$ in proposition 3.2. In propositions 3.4 and 3.7 we prove the existence and uniqueness of such a solution for arbitrary $\mu \in \mathbb{Z}^+$. The proof of existence relies on Wasow's theorem [34, theorem 12.1]. The further step required for uniqueness is analogous to the proof given by Joshi and Kitaev in [23] for the first Painlevé equation. We start by giving a generalisation of Boutroux's transformation of variables.

Definition 3.1 (Boutroux variables). Let $y(x)$ be a solution of $P_{II}(\mu)$. We define new variables $u(z)$ and z by

$$y(x) = x^{\mu/2} u(z) \quad \text{and} \quad z = \frac{2}{\mu+2} x^{(\mu+2)/2}.$$

Substitution into $P_{II}(\mu)$ shows that the function $u(z)$ satisfies

$$u''(z) = 2u(z)^3 + u(z) - \frac{3\mu}{\mu+2} \frac{u'(z)}{z} - \frac{\mu(\mu-2)}{(\mu+2)^2} \frac{u(z)}{z^2}. \quad (3.1)$$

In the following proposition, we consider the asymptotic expansion of $u(z)$ as $|z| \rightarrow \infty$.

Proposition 3.2. *The formal series*

$$u_f(z) = \sum_{n=0}^{\infty} \frac{a_n}{z^{2n}},$$

where $a_0 = i/\sqrt{2}$, $a_1 = -a_0 \frac{\mu(\mu-2)}{2(\mu+2)^2}$, and the coefficients a_n , for all $n > 1$, are given by the recurrence relation

$$a_n = \sigma(n) a_{n-1} + a_0 \sum_{k=1}^{n-1} a_k a_{n-k} + \sum_{j=1}^{n-1} \sum_{k=0}^{n-j} a_j a_k a_{n-j-k},$$

where $\sigma(n) = -(2n-1)(n-1) + \frac{3\mu}{\mu+2}(n-1) - \frac{\mu(\mu-2)}{2(\mu+2)^2}$,

is a formal solution of equation (3.1).

Proof. Substitution of the given series into equation (3.1). □

This gives rise to a formal series solution of $P_{II}(\mu)$ in the original variables:

$$y_f(x) = \sqrt{x^\mu} \sum_{n=0}^{\infty} \frac{\left(\frac{\mu+2}{2}\right)^{2n} a_n}{x^{(\mu+2)n}}, \quad (3.2)$$

with coefficients a_n described in proposition 3.2.

Remark 3.3. Recalling that a_0 is purely imaginary, we note that for x on the negative real semi-axis, $y_f(x)$ may only be real-valued for the case when μ is odd.

3.1. Existence

We now prove the existence of true solutions asymptotic to the given power-series in a given sector of angular width up to $2\pi/(\mu + 2)$ in the following proposition.

Proposition 3.4 (tronquée solutions). *Given $\mu \in \mathbb{Z}^+$ and a sector \mathcal{S} of angle less than $2\pi/(\mu+2)$, there exists a solution of $P_{II}(\mu)$ whose asymptotic behaviour as $|x| \rightarrow \infty$ in \mathcal{S} is given by the asymptotic series (3.2).*

Proof. Let $y(x)$ be a solution of $P_{II}(\mu)$. We use the change of variables given in definition 3.1 and further define

$$w(z) = u(z) - a_0 = u(z) - i/\sqrt{2}.$$

Then a straightforward calculation shows that the function $w(z)$ satisfies

$$w''(z) = 2w(z)^3 + 3i\sqrt{2}w(z)^2 - 2w(z) - \frac{3\mu}{\mu+2} \frac{w'(z)}{z} - \frac{\mu(\mu-2)}{(\mu+2)^2} \frac{(w(z) + i/\sqrt{2})}{z^2}, \quad (3.3)$$

and this ODE admits the formal power-series solution

$$w_f(z) = \sum_{n=1}^{\infty} \frac{a_n}{z^{2n}}.$$

Now, by defining variables $w_1 = w$ and $w_2 = w'$, equation (3.3) becomes the system

$$\begin{aligned} w'_1 &= w_2, \\ w'_2 &= 2w_1^3 + 3i\sqrt{2}w_1^2 - 2w_1 - \frac{3\mu}{\mu+2} \frac{w_2}{z} - \frac{\mu(\mu-2)}{(\mu+2)^2} \frac{(w_1 + i/\sqrt{2})}{z^2}. \end{aligned}$$

Let $f_1 = w_2$ and f_2 be the right side of the second equation. This system is, therefore, in the form

$$z^{-q} \frac{d\mathbf{w}}{dz} = \mathbf{f}(z, w_1, w_2),$$

where $\mathbf{w} = (w_1, w_2)^T$, $\mathbf{f} = (f_1, f_2)^T$ and $q = 0$. The Jacobian of \mathbf{f} , evaluated at $\mathbf{w} = \mathbf{0}$ and $|z| \rightarrow \infty$, has nonzero eigenvalues $\lambda_{\pm} = \pm i\sqrt{2}$. These results allow us to apply Wasow's theorem [34, theorem 12.1], showing the existence of a true solution with the desired asymptotic behaviour. \square

Definition 3.5. The solutions defined by proposition 3.4 are called tronquée solutions.

Corollary 3.6. *Let $\mu \in \mathbb{Z}^+$ and $\mathcal{S} \in \mathbb{C}$ be a given sector of angle less than $2\pi/(\mu+2)$. Suppose $y(x)$ is a tronquée solution of $P_{II}(\mu)$ in \mathcal{S} . Then the k th derivative, $y^{(k)}(x)$, for $k \in \{0, 1, 2, \dots\}$, has asymptotic behaviour given by the k th term-by-term differentiation of series (3.2), i.e. $y_f^{(k)}(x)$, as $|x| \rightarrow \infty$ in \mathcal{S} .*

Proof. The asymptotic behaviour of $y(x)$ implies that it is holomorphic in \mathcal{S} for sufficiently large x . The proof follows from Wasow's theorem [34, theorem 8.8]. \square

3.2. Uniqueness

In the following proposition, we prove that there is a unique true solution asymptotic to the given power-series in $|\arg(x)| \leq 2\pi/(\mu+2)$.

Proposition 3.7 (tritronquée solution). *Given $\mu \in \mathbb{Z}^+$, there exists a unique solution $Y(x)$ of $P_{II}(\mu)$ that has the asymptotic expansion $y_f(x)$ in the sector $|\arg(x)| \leq 2\pi/(\mu+2)$ as $|x| \rightarrow \infty$.*

Proof. Let $0 < \epsilon \in \mathbb{R}$. Let $y_1(x)$ and $y_2(x)$ be tronquée solutions as defined by proposition 3.4 in the respective sectors

$$S_1 : -\epsilon/2 < \arg(x) < \theta - \epsilon \quad \text{and} \quad S_2 : -\theta + \epsilon < \arg(x) < \epsilon/2,$$

where $\theta = 2\pi/(\mu + 2)$. Consider solutions $y_1(x)$ and $y_2(x)$ in the overlapping sector of angular width ϵ , centred about the positive real axis. Since the two tronquée solutions have the same asymptotic expansion, we get

$$v(x) := y_1(x) - y_2(x) \underset{\substack{|x| \rightarrow \infty \\ |\arg(x)| < \epsilon/2}}{=} o\left(\frac{\sqrt{x^\mu}}{x^{(\mu+2)n}}\right), \quad (3.4)$$

for every $n \in \{0, 1, 2, \dots\}$. Furthermore, we have $v''(x) = y_1''(x) - y_2''(x)$, and by applying $P_{\Pi}(\mu)$, we see that $v(x)$ satisfies the linear ODE

$$v''(x) = (2y_1(x)^2 + 2y_1(x)y_2(x) + 2y_2(x)^2 + x^\mu)v = (-\psi(x) + \omega(x))v, \quad (3.5)$$

where

$$\psi(x) = 2x^\mu \quad \text{and} \quad \omega(x) = 2y_1(x)^2 + 2y_1(x)y_2(x) + 2y_2(x)^2 + 3x^\mu.$$

The functions ψ and ω are holomorphic for sufficiently large x in the sector $S_1 \cap S_2$. It follows from asymptotic series (3.2) that ω is $\mathcal{O}(1/x^2)$, and thus the integral

$$\int_x^\infty \left| \psi^{-1/4} \left(\psi^{-1/4} \right)'' - \omega \psi^{-1/2} \right| dx, \quad (3.6)$$

taken along the real axis converges.

The study of solutions of equation (3.5) in the limit $|x| \rightarrow \infty$ is the subject of theorems given by Olver [25] under the term *Liouville-Green approximations*. In this context, the integral (3.6) is called the *error-control function*. When this integral is bounded, equation (3.5) satisfies the hypotheses of Olver's theorems 2.1 and 2.2 (see [25, chapter 6, §2]). On the positive real x -axis, Olver's theorem [25, chapter 6, § 2, theorem 2.2] gives

$$\begin{aligned} v(x) &= c_1 \psi^{-1/4} \exp\left(i \int_x^\infty \psi^{1/2} dx\right) (1 + o(1)) \\ &\quad + c_2 \psi^{-1/4} \exp\left(-i \int_x^\infty \psi^{1/2} dx\right) (1 + o(1)), \end{aligned}$$

where c_1 and c_2 are some constants. For real x , this contradicts the asymptotic behaviour (3.4) unless $c_1 = c_2 = 0$. So it follows that $y_1(x) = y_2(x)$ for sufficiently large positive real x . \square

Definition 3.8. The solution defined by proposition 3.7 is called the tritronquée solution $Y(x)$.

Remark 3.9. Note that the sector in which Y is asymptotically free of poles is symmetric around the positive real semi-axis.

Given $Y(x)$, defined above, we may generate other tritronquée solutions by using the discrete symmetry of $P_{\Pi}(\mu)$, which is given by the transformation

$$Y_n(x) = \exp\left(-\frac{2\pi i n}{\mu + 2}\right) Y\left(\exp\left(-\frac{2\pi i n}{\mu + 2}\right)x\right), \quad (3.7)$$

for $n \in \mathbb{Z}$. $Y_n(x)$ also satisfies $P_{\Pi}(\mu)$ and is tritronquée in a rotated sector of width $4\pi/(\mu + 2)$, which is bisected by a ray of angle $2\pi n/(\mu + 2)$.

4. The Hastings–McLeod-type solution

In this section, we define the generalised Hastings–McLeod solution via a boundary-value problem on the real-line. We then prove that the existence of this solution fails when μ is even while it exists and is unique when μ is odd.

Definition 4.1 (Hastings–McLeod-type solution). Given $\mu \in \mathbb{Z}^+$, for $x \in \mathbb{R}$, we call a solution $y(x)$ of $P_{II}(\mu)$ a Hastings–McLeod-type solution if and only if $y(x) \sim y_f(x)$ as $x \rightarrow -\infty$ (see equation (3.2)), and $y(x) \sim kf(x)$ as $x \rightarrow +\infty$ (see equation (2.1)) for some $k \in \mathbb{R}^+$.

We now prove the main result of this section.

Theorem 4.2. *For every odd integer $\mu > 1$, there exists a unique Hastings–McLeod-type solution $y(x)$ of $P_{II}(\mu)$. On the other hand, for every even integer $\mu > 1$, no Hastings–McLeod-type solution of $P_{II}(\mu)$ exists.*

Remark 4.3. As discussed in remark 3.3, the formal solution $y_f(x)$, for real $x < 0$, may only be real-valued for odd μ . For even values of μ , the corresponding Hastings–McLeod-type solution would not be real-valued and thus not mirror the distinctive properties of the original Hastings–McLeod solution.

Hastings and McLeod originally tackled this problem while considering a different generalisation of P_{II} , that is

$$y''(x) = 2y(x)^{\alpha+1} + xy(x),$$

which coincides with $P_{II}(\mu)$ when $\alpha = 2$ and $\mu = 1$ (i.e. P_{II}). We will prove theorem 4.2 by extending the approach taken by Hastings and McLeod to our case with $\alpha = 2$ and more general μ .

4.1. Existence

The proof of the existence (and non-existence when μ is even) part of theorem 4.2 follows from a series of lemmas.

Lemma 4.4. *Given $\mu \in \mathbb{Z}^+$ and $k \in \mathbb{R}^+$, there exists a unique solution $y_k(x)$, $x \in \mathbb{R}$, of $P_{II}(\mu)$ which is asymptotic to $kf(x)$ as $x \rightarrow +\infty$. At each x for which the solution continues to exist as x decreases, the solution and its derivatives are continuous functions of k .*

Proof. By using $f(x)$ and $g(x)$ as integrating factors, we find that $y_k(x)$ must satisfy the integral equation

$$y_k(x) = kf(x) + \int_x^\infty (f(t)g(t) - f(t)g(x)) y_k(t)^3 dt, \quad (4.1)$$

where f and g are defined in equation (2.1). By application of a standard contraction mapping argument, in an appropriate space of decaying, continuous functions equipped with the uniform norm, equation (4.1) may be shown to have a unique solution, giving y_k and its continuous dependence on k . \square

Lemma 4.5. *For every even $\mu \in \mathbb{Z}^+$, and every $k \in \mathbb{R}^+$, $y_k(x)$ remains positive as x decreases and becomes singular at some finite value of x .*

Proof. Let $k \in \mathbb{R}^+$ and assume $\mu \in \mathbb{Z}^+$ is even. For sufficiently large $x \in \mathbb{R}^+$, $y_k(x)$ behaves like $kf(x)$. Therefore, there exists $x_0 \in \mathbb{R}^+$ so that $y_k(x) > 0$ and $y'_k(x) < 0$ for all $x \in [x_0, \infty)$.

Since μ is even, we immediately deduce from $P_{II}(\mu)$ that $y_k(x)$ is convex and positive for all $x \in \mathbb{R}$.

The remainder of the proof proceeds by contradiction. Assume that $y_k(x)$ remains bounded for all $x \in (-\infty, x_0)$, i.e. it is not singular in this interval. For all $x \in (-\infty, x_0)$, we have $y_k''(x) > y_k(x)^3$ and thus $y_k'(x)y_k''(x) < y_k'(x)y_k(x)^3$ (since y_k' is negative). We integrate the latter autonomous inequality between x and x_0 , which after rearranging terms gives

$$y_k'(x)^2 > \frac{1}{2}y_k(x)^4 + y_k'(x_0)^2 - \frac{1}{2}y_k(x_0)^4. \quad (4.2)$$

We now show that $y_k(x)$ is monotonically increasing to $+\infty$ as x decreases. Suppose instead that $y_k'(x_1)$ vanishes at some $x_1 < x_0$, while $y_k'(x) < 0$ for all $x \in (x_1, x_0]$. The inequality (4.2) would then imply that $y_k(x_1) < y_k(x_0)$, which contradicts the fact that $y_k(x)$ is still increasing as we move from x_0 to x_1 .

Therefore, we can choose $x_1 \in (-\infty, x_0)$ such that $\frac{1}{4}y_k(x_1)^4 + y_k'(x_0)^2 - \frac{1}{2}y_k(x_0)^4 > 0$, hence $y_k'(x)^2 > \frac{1}{4}y_k(x)^4$ for all $x \in (-\infty, x_1)$. Therefore, $y_k'(x) < -\frac{1}{2}y_k(x)^2$, and another integration gives

$$\frac{1}{y_k(x_1)} - \frac{1}{y_k(x)} > \frac{x_1 - x}{2},$$

for all $x < x_1$, which contradicts the assumption that $y_k(x)$ is bounded and non-singular on the interval. \square

As a direct result of lemma 4.5, we have proven the non-existence of the Hastings–McLeod-type solution in the case of even μ . We now continue our argument in the case of odd μ .

Lemma 4.6. *For every odd $\mu \in \mathbb{Z}^+$, the set $S_1 \subset \mathbb{R}^+$ of k values, for which $y_k(x)$ remains positive as x decreases and becomes singular at some finite value of x , is an open set.*

Proof. Suppose we have $k_0 \in \mathbb{R}^+$ and $x_0 \in \mathbb{R}$ such that $y_{k_0}(x)$ blows up at $x = x_0$. We now show that there exists a neighbourhood of k values (sufficiently close to k_0) such that the solution y_k still becomes singular. Since $y_{k_0}(x_0)$ is unbounded, we may choose $x_1 > x_0$ to make $y_{k_0}(x_1)$ arbitrarily large. We choose x_1 near x_0 such that

$$y_{k_0}(x_1)^2 > (|x_1| + 1)^\mu, \quad (4.3)$$

noting then that $y_{k_0}(x)$ remains convex on $(x_1 - 1, x_1)$ while it continues to exist. Based on the continuous dependence of $y_k(x)$ on k , we may replace k_0 with sufficiently close k so that the inequality (4.3) continues to hold. Furthermore, on the interval $(x_1 - 1, x_1)$, we have $|x| < |x_1| + 1$, i.e. $|x|^\mu < (|x_1| + 1)^\mu$. Hence

$$|x|^\mu < (|x_1| + 1)^\mu < y_k(x_1)^2 < y_k(x)^2.$$

So if $x < 0$, we have

$$y_k''(x) = 2y_k(x)^3 + x^\mu y_k(x) = 2y_k(x)^3 - |x|^\mu y_k(x) > 2y_k(x)^3 - y_k(x)^3 = y_k(x)^3. \quad (4.4)$$

Note that if $x \geq 0$ we still have $y_k''(x) > y_k(x)^3$.

We have therefore shown that $y_k''(x) > y_k(x)^3$ on $x \in (x_1 - 1, x_1)$. Then from the integration of this autonomous inequality, the proof of lemma 4.5 shows that $y_k(x)$ will become unbounded in this interval given sufficiently large $|y_k(x_1)|$ and $|y_k'(x_1)|$. As previously noted, we may choose x_1 so that these are arbitrarily large and so the desired result follows. \square

Lemma 4.7. *For every odd $\mu \in \mathbb{Z}^+$, the set $S_2 \subset \mathbb{R}^+$ of k values, for which $y_k(x)$ becomes negative at some point as x decreases, is an open set.*

Proof. Suppose there exist k_0 and x_0 such that $y_{k_0}(x_0) < 0$, then by the continuity of y_k in k , we may deduce that $y_k(x_0) < 0$ for all k in some neighbourhood of k_0 . \square

Lemma 4.8. *For every odd $\mu \in \mathbb{Z}^+$, the set S_1 is non-empty.*

Proof. It follows from equation (4.1) that $y_k(x) > kf(x)$ for sufficiently large x . Differentiation of equation (4.1) shows similarly that $y'_k(x) < kf'(x) < 0$. Using the same argument as in the proof of lemma 4.6, we may choose sufficiently large x_1 and k so that $y_k(x)$ blows up in $(x_1 - 1, x_1)$. \square

Lemma 4.9. *For every odd $\mu \in \mathbb{Z}^+$, the set S_2 is non-empty.*

Proof. Let $\mu > 1$ be an odd integer. Consider the graph of $p(x)$ given by the positive real root of $2p(x)^2 = -x^\mu$ on the interval $x \in (-\infty, 0]$. Let $x_0 < 0$. Recall that $y_0(x)$ corresponds to the $k=0$ case in lemma 4.4 and is the identically zero solution of $P_{\text{II}}(\mu)$. By the continuity of $y_k(x_0)$ in k , we may choose k sufficiently small so that $y_k(x_0) < p(x_0)$ and $p'(x_0) < y'_k(x_0) < 0$. Examination of $P_{\text{II}}(\mu)$ then shows that $y''_k(x) < 0$ for $x \leq x_0$. Therefore, $y_k(x)$ vanishes in this interval and becomes negative to the left of x_0 . \square

We now conclude the proof of existence. We have two non-empty, open sets S_1 and S_2 , which are disjoint. So there must exist at least one positive value of k , which lies in neither S_1 nor S_2 . The solution $y_k(x)$ corresponding to such a k is then finite and positive for all x . If such a solution were to decay to zero as $x \rightarrow -\infty$, then it would be oscillatory in this asymptotic limit (see equation (2.3)), thus vanishing at some point and contradicting the fact that $k \notin S_2$. Therefore, we conclude that $y(x) \sim y_f(x)$ as $x \rightarrow -\infty$. So $y_k(x)$ is a Hastings–McLeod-type solution.

4.2. Uniqueness

We now give the argument for uniqueness to complete the proof of theorem 4.2.

Lemma 4.10. *Let μ be an odd integer. Suppose $y_{k_1}(x)$ and $y_{k_2}(x)$ are Hastings–McLeod-type solutions with $k_1 > k_2$, then $y_{k_1}(x) > y_{k_2}(x)$ for all x .*

Proof. Note that the asymptotic behaviour as $x \rightarrow +\infty$ implies that $y_{k_1}(x) > y_{k_2}(x)$ for sufficiently large x . If the assertion of the lemma fails to hold, then there exists $x_0 \in \mathbb{R}$ such that $y_{k_1}(x_0) = y_{k_2}(x_0)$, while $y_{k_1}(x) > y_{k_2}(x)$ on (x_0, ∞) . For this to occur, we must have $y'_{k_1}(x_0)^2 \leq y'_{k_2}(x_0)^2$.

Define the energy function

$$V_k(x) = y'_k(x)^2 - x^\mu y_k(x)^2 - y_k(x)^4. \quad (4.5)$$

From $P_{\text{II}}(\mu)$, it follows that

$$V'_k(x) = -\mu x^{\mu-1} y_k(x)^2. \quad (4.6)$$

From equation (4.5), we have

$$V_{k_1}(x_0) - V_{k_2}(x_0) = y'_{k_1}(x_0)^2 - y'_{k_2}(x_0)^2 \leq 0, \quad (4.7)$$

which implies $V_{k_1}(x_0) \leq V_{k_2}(x_0)$.

Using equation (4.6), $y_{k_1}(x) > y_{k_2}(x)$ implies that $V'_{k_1}(x) < V'_{k_2}(x)$. We proceed to integrate the autonomous inequality $V'_{k_1}(x) - V'_{k_2}(x) < 0$ over the interval (x_0, N) , where $N \gg 1$, giving

$$V_{k_1}(N) - V_{k_2}(N) < V_{k_1}(x_0) - V_{k_2}(x_0). \quad (4.8)$$

However, for sufficiently large N we have $V_{k_1}(N) - V_{k_2}(N) > 0$, and we already deduced that $V_{k_1}(x_0) - V_{k_2}(x_0) \leq 0$. These observations contradict the inequality (4.8). Therefore, x_0 cannot exist and the desired result follows. \square

Lemma 4.11. *Let μ be an odd integer. Suppose $y_{k_1}(x)$ and $y_{k_2}(x)$ are Hastings–McLeod-type solutions. Then $k_1 = k_2$.*

Proof. Suppose $k_1 \neq k_2$. Without loss of generality, we assume $k_1 > k_2$. As in the proof of proposition 3.7, we can appeal to another theorem of Olver [25, theorem 2.1, p.193] to show that $p(x) = y_{k_1}(x) - y_{k_2}(x)$ satisfies

$$\begin{aligned} p(x) &= b_1 f^{-1/4} \exp \left(\int_{-\infty}^x f^{1/2} dx \right) (1 + o(1)) \\ &\quad + b_2 f^{-1/4} \exp \left(- \int_{-\infty}^x f^{1/2} dx \right) (1 + o(1)), \end{aligned} \quad (4.9)$$

for some constants b_1 and b_2 , where $f = -2x^\mu$, for sufficiently large $|x|$, $x < 0$. Olver’s theorem also states that the behaviour (4.9) is twice differentiable. However, this contradicts the assumed asymptotic behaviour $y_{k_j} \sim y_f$, $j = 1, 2$ given in (3.2) unless $b_1 = 0$.

Suppose $b_2 \neq 0$. Then we use the differentiability of the result (4.9) to obtain

$$\begin{aligned} V_{k_1}(x) - V_{k_2}(x) &= -2(2y_f(x)^3 + x^\mu y_f(x))p(x)(1 + o(1)) \\ &= -2y_f''(x)p(x)(1 + o(1)). \end{aligned}$$

For sufficiently large $|x|$, $x < 0$, recalling that $\mu > 1$ is an odd integer and $p(x) > 0$, this asymptotic behaviour is negative. However, we showed in the previous lemma that $V_{k_1}(x) - V_{k_2}(x)$ is positive for sufficiently large positive x and that $V'_{k_1}(x) - V'_{k_2}(x) < 0$ for all x , which is a contradiction. Therefore, we must have $b_2 = 0$.

It follows that $y_{k_1}(x) = y_{k_2}(x)$ for sufficiently large real $|x|$, with $x < 0$. This contradicts the result of lemma 4.10 unless $k_1 = k_2$. \square

This completes our proof of uniqueness of the Hastings–McLeod type solution.

Remark 4.12. Consider the behaviours of $y_k(x)$ depending on the parameter $k \in \mathbb{R}^+$. For odd μ , suppose k^* is the unique value for which $y_{k^*}(x)$ is the Hastings–McLeod-type solution. Using the same argument as in lemma 4.10, we may show that $0 < k < k^*$ implies $|y_k(x)| < y_{k^*}(x)$ for all x . So for $0 < k < k^*$, we have a family of bounded solutions on the real-line; these solutions have oscillatory asymptotic behaviour given by equation (2.3). On the other hand, when $k^* < k$, we have solutions which are positive and convex for all x , becoming singular at some finite point as x decreases.

5. Conclusion

In this paper, we investigated a family of nonlinear ODEs, parameterised by μ , which are perturbations of the second Painlevé equation. In the appendix, we show how such equations arise in an application, specifically, in a mathematical model of electro-diffusion. Our main results show that the perturbed equation possesses solutions that are natural generalisations of two celebrated solutions of P_{II} .

We found that the perturbed equation always has solutions which we call tritronqueé, first described by Boutroux for P_{II} . These solutions are asymptotic to a power-series expansion in a surprisingly broad annular region of the complex plane, near infinity. Furthermore, if our

integer parameter μ is odd, $P_{\text{II}}(\mu)$ has a solution analogous to the famous Hastings–McLeod solution of P_{II} . The Hastings–McLeod-type solution is holomorphic, real-valued and positive on the entire real-line, with known asymptotic behaviour in either direction.

These results naturally give rise to similar questions about perturbations of other Painlevé equations. They suggest that certain behaviours, important in applications, are preserved by classes of perturbations.

Data availability statement

No new data were created or analysed in this study.

Appendix. Physical application of the perturbed equation

This section shows how $P_{\text{II}}(\mu)$ arises in Bass' electro-diffusion model [4] under a small change in the physical setting. Specifically, we allow particle flux to be non-constant.

Assume that $c_+(x)$ and $c_-(x)$ represent the respective (dimensionless) concentrations of two ionic species with equal and opposite charge, while $E(x)$ represents the (dimensionless) induced electric field. The following coupled nonlinear ODEs govern the model of electro-diffusion:

$$c'_+(x) = E(x)c_+(x) + A_+, \quad (\text{A.1})$$

$$c'_-(x) = -E(x)c_-(x) + A_-, \quad (\text{A.2})$$

$$\lambda^2 E'(x) = c_+(x) - c_-(x), \quad (\text{A.3})$$

where λ is a constant, the terms A_{\pm} arise due to Fick's law's standard passive diffusion process, and we note that A_{\pm} is proportional to the ratio between the species flux in the x -direction, ϕ_{\pm} , and the species diffusivity, D_{\pm} .

In the original model, ϕ_{\pm} is constant in x due to the conservation of mass in a strictly one-dimensional setting. However, we consider the possibility that ϕ_{\pm} may undergo small amounts of variation, as the species may fluctuate in other spatial dimensions. However, we assume D_{\pm} is constant.

Consider A_{\pm} as a perturbative series, whereby A_{\pm} is polynomial in x and constant at leading order. For now, consider a linear correction term. Furthermore, we shall consider a somewhat simplified case whereby $A_+ = A_-$, so we have

$$A_+ = A_- = A_0 + \epsilon x. \quad (\text{A.4})$$

Then by adding equations (A.1) and (A.2), and applying equation (A.3), the result is

$$c'_+(x) + c'_-(x) = \lambda^2 E(x)E'(x) + 2A_0 + 2\epsilon x. \quad (\text{A.5})$$

Upon integrating equation (A.5) and disregarding the constant of integration, we have

$$c_+(x) + c_-(x) = \frac{\lambda^2}{2} E(x)^2 + \epsilon x^2 + 2A_0 x. \quad (\text{A.6})$$

Meanwhile, by differentiating equation (A.3), and then applying equations (A.1) and (A.2), we obtain

$$\lambda^2 E''(x) = E(x)(c_+(x) + c_-(x)). \quad (\text{A.7})$$

Hence, by combining equations (A.6) and (A.7), we obtain

$$\lambda^2 E''(x) = \frac{\lambda^2}{2} E(x)^3 + (\epsilon x^2 + 2A_0 x) E(x). \quad (\text{A.8})$$

Equation (A.8) is a nonlinear ODE which governs the induced electric field as a function of x and is similar in form to $P_{\text{II}}(\mu)$ with $\mu = 2$. Adding higher degree correction terms in equation (A.4) would correspond to a higher degree polynomial $p(x)$ in the coefficient of E , that is,

$$\lambda^2 E''(x) = \frac{\lambda^2}{2} E(x)^3 + p(x) E(x).$$

The presence of such higher degree (>1) terms in the coefficient function $p(x)$ motivates the study of our perturbed form of P_{II} .

ORCID iDs

Joshua Holroyd  <https://orcid.org/0000-0003-1354-1271>
Nalini Joshi  <https://orcid.org/0001-7504-4444>

References

- [1] Ablowitz M J and Segur H 1977 Exact linearization of a Painlevé transcendent *Phys. Rev. Lett.* **38** 1103
- [2] Airault H 1979 Rational solutions of Painlevé equations *Stud. Appl. Math.* **61** 31–53
- [3] Albrecht D W, Mansfield E L and Milne A E 1996 Algorithms for special integrals of ordinary differential equations *J. Phys. A: Math. Gen.* **29** 973
- [4] Bass L 1964 Irreversible interactions between metals and electrolytes *Proc. R. Soc. A* **277** 125–36
- [5] Bass L, Nimmo J, Rogers C and Schief W 2010 Electrical structures of interfaces: a Painlevé II model *Proc. R. Soc. A* **466** 2117–36
- [6] Bender C M and Orszag S A 1999 *Advanced Mathematical Methods for Scientists and Engineers I: Asymptotic Methods and Perturbation Theory* (New York: Springer Science & Business Media)
- [7] Boutroux P 1913 Recherches sur les transcendantes de M Painlevé et l'étude asymptotique des équations différentielles du second ordre *Ann. Sci. Normale Supérieure* **30** 255–375
- [8] Claeys T and Grava T 2010 Painlevé II asymptotics near the leading edge of the oscillatory zone for the Korteweg-de Vries equation in the small-dispersion limit *Commun. Pure Appl. Math. A* **63** 203–32
- [9] Clarkson P A 2006 Painlevé equations–nonlinear special functions *Orthogonal Polynomials and Special Functions* (Berlin: Springer) pp 331–411
- [10] Clarkson P A and Mansfield E L 2003 The second Painlevé equation, its hierarchy and associated special polynomials *Nonlinearity* **16** R1
- [11] Joshi N and Kruskal M 1994 A direct proof that solutions of the six Painlevé equations have no movable singularities except poles *Stud Appl. Math.* **93** 184–207
- [12] De Boer P and Ludford G 1975 Spherical electric probe in a continuum gas *Plasma Phys.* **17** 29
- [13] Deift P 2007 *Universality for Mathematical and Physical Systems* vol 1 pp 125–52 (Zurich: Eur. Math. Society)
- [14] Fokas A and Ablowitz M 1981 Linearization of the Korteweg-de Vries and Painlevé II equations *Phys. Rev. Lett.* **47** 1096
- [15] Forrester P J and Witte N S 2015 Painlevé II in random matrix theory and related fields *Constructive Approx.* **41** 589–613
- [16] Fukutani S, Okamoto K and Umemura H 2000 Special polynomials and the Hirota bilinear relations of the second and the fourth Painlevé equations *Nagoya Math. J.* **159** 179–200
- [17] Gambier B 1910 Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes *Acta Math.* **33** 1–55
- [18] Gromak V I, Laine I and Shimomura S 2008 *Painlevé Differential Equations in the Complex Plane* (Berlin: De Gruyter)

[19] Hastings S P and Mcleod J B 1980 A boundary value problem associated with the second Painlevé transcendent and the Korteweg-de Vries equation *Arch. Ration. Mech. Anal.* **73** 31–51

[20] Howes P and Joshi N 2014 Global asymptotics of the second Painlevé equation in Okamoto’s space *Constructive Approx.* **39** 11–41

[21] Ince E L and Equations O D 1956 *Ordinary Differential Equations* (New York: Dover Books on Mathematics. Dover Publications)

[22] Joshi N 2003 Tritronquée solutions of perturbed first Painlevé equations *Theor. Math. Phys.* **137** 1515–19

[23] Joshi N and Kitaev A 2001 On Boutroux’s tritonquée solutions of the first Painlevé equation *Stud. Appl. Math.* **107** 253–91

[24] Olde Daalhuis A 2022 Exponentially-improved asymptotics and numerics for the (un) perturbed first Painlevé equation *J. Phys. A: Math. Gen.* **55** 304004

[25] Olver F 1997 *Asymptotics and Special Functions* (Boca Raton, FL: CRC Press)

[26] Olver S 2011 Numerical solution of Riemann–Hilbert problems: Painlevé II *Found. Comput. Math.* **11** 153–79

[27] Olver W J, Olde Daalhuis A B, Lozier D W, Schneider B I, Boisvert R F, Clark C W, Miller B R, Saunders B V, Cohl H S and McClain M A (eds) NIST Digital Library of Mathematical Functions *Release 1.1.7* (available at: <http://dlmf.nist.gov/> and <http://dlmf.nist.gov/10.40.E2>) (Accessed 15 October 2022)

[28] Periwal V and Shevitz D 1990 Exactly solvable unitary matrix models: multicritical potentials and correlations *Nucl. Phys. B* **344** 731–46

[29] Schiappa R and Vaz R 2014 The resurgence of instantons: multi-cut Stokes phases and the Painlevé II equation *Commun. Math. Phys.* **330** 655–721

[30] Seiberg N and Shih D 2005 Flux vacua and branes of the minimal superstring *J. High Energy Phys.* **2005** 055

[31] Tracy C A and Widom H 1994 Level-spacing distributions and the Airy kernel *Commun. Math. Phys.* **159** 151–74

[32] Tracy C A and Widom H 1999 Random unitary matrices, permutations and Painlevé *Commun. Math. Phys.* **207** 665–85

[33] Umemura H 2007 Invitation to Galois theory Differential Equations and Quantum Groups, Andrey A. Bolibrukh Memorial Volume IRMA Lectures in Mathematics and Theoretical Physics vol 9, ed B Daniel, E Benjamin, M Claude, S Claude and S Reinhard (Zurich: EMS Press) pp 269–89

[34] Wasow W 2018 *Asymptotic Expansions for Ordinary Differential Equations* (Mineola NY: Dover Publications)