

KANTOWSKI-SACHS GRAVITATIONAL INSTANTON SOLUTION

BY D. LORENZ

Fakultät für Physik, Universität Konstanz*

(Received March 21, 1983)

We present a new self-dual solution to Euclidean gravity which is of the Kantowski-Sachs group type and may be considered as a gravitational instanton.

PACS numbers: 04.20.Jb

1. Introduction

The discovery of pseudoparticle (instanton) solutions to the Euclidean $SU(2)$ Yang-Mills theory (Belavin et al. 1975; Rajamaran 1982) has suggested the possibility that analogous solutions might occur in Einstein's theory of gravitation. Since the Yang-Mills pseudoparticles possess self-dual field strength, one likely possibility is that gravitational pseudoparticles are characterized by self-dual curvature.

Starting from the idea that the Yang-Mills potential is asymptotically a pure gauge (see Actor 1979 for a detailed description of Yang-Mills solutions), a gravitational instanton should have an asymptotically flat metric. Four types of non-compact instantons (a gravitational instanton with a cosmological constant $\Lambda > 0$ is necessarily compact) have been discovered (for recent reviews see Eguchi et al. 1980; Pope 1981; Perry 1982): asymptotically Euclidean (AE), asymptotically locally Euclidean (ALE), asymptotically flat (AF) and asymptotically locally flat (ALF). In general a gravitational instanton can be defined as a complete Riemannian manifold (with Euclidean signature) (M, g) of dimension four with finite action satisfying the Einstein Equations $R_{\mu\nu} = \Lambda g_{\mu\nu}$, where $R_{\mu\nu}$ is the Ricci tensor.

A variety of solutions of Einstein's field equations with instanton-like properties have been discovered. The corresponding metrics of many of these solutions may be cast into the Euclidean equivalent of the Bianchi type-IX space-time. Thus one is forced to consider the complete set of Bianchi types I-IX.

* Address: Fakultät für Physik, Universität Konstanz, Bücklestr. 13, D-7750 Konstanz, West Germany.

In papers published previously (Lorenz 1983a, b, c) we presented new self-dual solutions for the Bianchi types I–IX. In this paper we consider the Euclidean Kantowski-Sachs space-time (Kantowski 1966; Kantowski and Sachs 1966), which is closely related to the Bianchi type-III model.

2. Metric, field equations and solutions

The Kantowski-Sachs metric is a special case of the metrics

$$ds^2 = dt^2 + S^2(t)dr^2 + R^2(t)(d\theta^2 + f^2(\theta)d\phi^2), \quad (1)$$

where $f(\theta)$ is $\sinh \theta$, θ or $\sin \theta$, respectively, for Bianchi type-III, type-I or Kantowski-Sachs. By requiring that the Riemann tensor $R^\mu_{\nu\alpha\beta}$ is self-dual we obtain the following field equations

$$\ddot{S} = \ddot{R} = \dot{S}\dot{R} = 0, \quad \dot{R}^2 = -f''/f, \quad (2)$$

where $(\dot{}) = d/dt$ and $()' = d/d\theta$. We have $k = f''/f = 1, 0, -1$ in case of Bianchi type-III, type-I or K-S. The case $k = 0$ gives flat space solutions. According to the Positive Action Theorem (Hawking 1979; Schoen and Yau 1979) the only asymptotically Euclidean instanton is flat space. The case $k = 1$ has been already discussed by us (Lorenz 1983c) and gives only complex solutions. However, in the Euclidean Kantowski-Sachs space-time considered here we obtain the following real solution of Equations (2):

$$S = a, \quad R = b(t - t_1), \quad (3)$$

where a, b, t_1 are constants of integration obeying $b^2 = 1$.

Our solution possesses the interesting feature that the components of the curvature two-forms

$$\theta^\mu_{\nu} = \frac{1}{2} R^\mu_{\nu\alpha\beta} \sigma^\alpha \wedge \sigma^\beta \quad (4)$$

all vanish whereas the curvature

$$R^* = -f''/(fR^2) \quad (5)$$

of the two-dimensional surfaces $(t, r = \text{const})$ is always positive in a finite region but falls off as $1/t^2$ at infinity.

Defining the new coordinates (X, Y, Z, W) by

$$X = (t - t_1) \sin \theta \cos \phi, \quad Y = (t - t_1) \sin \theta \sin \phi, \quad Z = (t - t_1) \cos \theta, \quad W = ar, \quad (6)$$

the metric (1) can be transformed into flat space-time. In these new coordinates our solution turns out to be identical with the flat space solution obeying the Positive Action Theorem.

REFERENCES

- Actor, A., *Rev. Mod. Phys.* **51**, 461 (1979).
 Belavin, A. A., Polyakov, A. M., Schwarz, A. S., Tyupkin, Yu. S., *Phys. Lett.* **59B**, 85 (1975).
 Eguchi, T., Gilkey, P. B., Hanson, A. J., *Phys. Rep.* **66**, 213 (1980).

- Hawking, S. W., in *Recent Developments in Gravitation*, edited by M. Lévy and S. Deser, Plenum Press, New York-London 1979.
- Kantowski, R., Ph. D. Thesis, Univ. of Texas 1966.
- Kantowski, R., Sachs, R. K., *J. Math. Phys.* **7**, 443 (1966).
- Lorenz, D., *Acta. Phys. Pol.* **B14**, 479 (1983a).
- Lorenz, D., *Acta Phys. Pol.* **B14**, 723 (1983b).
- Lorenz, D., *Acta Phys. Pol.* **B14**, 791 (1983c).
- Perry, M., in *Seminar on Differential Geometry*, edited by S.-T. Yau, Ann. Math. Stud. N. 102, Princeton Univ. Press, Princeton, N. Y. 1982.
- Pope, C. N., in *Quantum Gravity 2*, edited by C. J. Isham, R. Penrose and D. W. Sciama, Clarendon Press, Oxford 1981.
- Rajaraman, R., *Solitons and Instantons*, North-Holland Publ. Comp., Amsterdam-New York-Oxford 1982.
- Schoen, R. M., Yau, S. T., *Phys. Rev. Lett.* **42**, 547 (1979).