



Constraining the in-medium nucleon-nucleon cross section from the width of nuclear giant dipole resonance

Rui Wang ^{a,b}, Zhen Zhang ^c, Lie-Wen Chen ^d, Che Ming Ko ^e, Yu-Gang Ma ^{a,b}



^a Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Institute of Modern Physics, Fudan University, Shanghai 200433, China

^b Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China

^c Sino-French Institute of Nuclear Engineering and Technology, Sun Yat-Sen University, Zhuhai 519082, China

^d School of Physics and Astronomy and Shanghai Key Laboratory for Particle Physics and Cosmology, Shanghai Jiao Tong University, Shanghai 200240, China

^e Cyclotron Institute and Department of Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA

ARTICLE INFO

Article history:

Received 23 March 2020

Received in revised form 27 May 2020

Accepted 29 May 2020

Available online 1 June 2020

Editor: W. Haxton

Keywords:

Heavy-ion collisions

Transport models

Nuclear giant dipole resonance width

In-medium nucleon-nucleon cross section

ABSTRACT

We develop a new lattice Hamiltonian method for solving the Boltzmann-Uehling-Uhlenbeck (BUU) equation. Adopting the stochastic approach to treat the collision term and using the GPU parallel computing to carry out the calculations allows for a rather high accuracy in evaluating the collision term, especially its Pauli blocking, leading thus to a new level of precision in solving the BUU equation. Applying this lattice BUU method to study the width of giant dipole resonance (GDR) in nuclei, where the accurate treatment of the collision term is crucial, we find that the obtained GDR width of ^{208}Pb shows a strong dependence on the in-medium nucleon-nucleon cross section σ_{NN}^* . A very large medium reduction of σ_{NN}^* is needed to reproduce the measured value of the GDR width of ^{208}Pb at the Research Center for Nuclear Physics in Osaka, Japan.

© 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The in-medium nucleon-nucleon (NN) cross section σ_{NN}^* has significant effects on the dynamics of heavy-ion collisions (HICs), and it thus plays a crucial role in understanding the reaction mechanisms as well as various phenomena and observables in these collisions [1–6]. The importance of σ_{NN}^* also lies in its intimate relation to the transport properties of nuclear matter [7,8] and the nucleon effective interactions [9]. Since a major goal of studying HICs is to extract the equation of state (EOS) of nuclear matter from experimental data [10–15], a thorough understanding of σ_{NN}^* helps reduce the uncertainties in transport models [16,17] that are used for describing these reactions. While the NN cross section in free space $\sigma_{\text{NN}}^{\text{free}}$ can be directly measured in experiments, the determination of the value of σ_{NN}^* in nuclear medium usually relies on theoretical investigations. These include calculations based on microscopic theories, such as the nonrelativistic and relativistic Brueckner theories [9,18–23] and the closed time path Green's function approach [24,25]. Also, there have been attempts to extract σ_{NN}^* from experiments by comparing results from transport

model calculations, where σ_{NN}^* is a crucial input, with observables measured in HICs that are sensitive to σ_{NN}^* , e.g., the collective flow and nuclear stopping [1,8,26–30]. Although these studies have reached the consensus that the NN cross section is suppressed in nuclear medium, the reduction factor is still far from certainty.

The transport model used in describing HICs is a straightforward tool for studying σ_{NN}^* because one of its main ingredients, the NN collision term, embodies the information of σ_{NN}^* . Since the mean field or the EOS of nuclear matter is another major ingredient of one-body transport models, finding the proper observables that depend on σ_{NN}^* rather than the nuclear EOS is essential for studying σ_{NN}^* . One such observable is the width of nuclear giant dipole resonance (GDR) as it is naturally related to σ_{NN}^* through the NN collision term in transport models. In general, the damping width of nuclear collective motion originates from three sources: 1) the escape width associated with particle emissions; 2) the fragmentation or the Landau damping width due to couplings between single particle states and the mean field; 3) the spreading or collisional damping width caused by the coupling to more complex states like the two-particle-two-hole ($2p-2h$), $3p-3h$, etc. For a heavy nucleus at zero temperature, the width of its GDR is mainly exhausted by collisional damping [31–33] before the contribution from deformation fluctuations appears as a result of the finite temperature effect [34]. In the transport model, the colli-

E-mail addresses: wangrui@sinap.ac.cn (R. Wang), zhangzh275@mail.sysu.edu.cn (Z. Zhang), lwchen@sjtu.edu.cn (L.-W. Chen), ko@comp.tamu.edu (C.M. Ko), mayugang@fudan.edu.cn (Y.-G. Ma).

sional damping is incorporated in the binary collisions of nucleons and thus depends directly on σ_{NN}^* . It is therefore expected that the GDR width of a heavy nucleus in studies based on the transport model depends strongly on σ_{NN}^* and weakly on the nuclear EOS.

The major obstacle that has so far prevented the use of transport models to accurately calculate the spreading width of GDR is due to the fermionic nature of nucleons. Specifically, the accurate treatment of Pauli blocking in transport models is challenging [16,17], especially for small amplitude nuclear collective motions. Both subtle implementations and advanced computing techniques are required for overcoming this difficulty. In the present work, we extend the previous study using the lattice Hamiltonian Vlasov method based on the next-to-next-to-next-to leading order (N3LO) Skyrme pseudopotential [35] to include a stochastic elastic NN collision term. Solving the resulting Boltzmann-Uehling-Uhlenbeck (BUU)-type one-body transport model with the high computation efficiency provided by GPU parallel computing [36], which enables the accurate treatment of Pauli blocking in the collision term of the BUU equation, allows us to calculate precisely the spreading width of the GDR in nuclei. We then obtain a stringent constraint on the in-medium NN cross section σ_{NN}^* by comparing the GDR width of ^{208}Pb from the present lattice BUU (LBUU) method with that measured from $^{208}\text{Pb}(\vec{p}, \vec{p}')$ reaction with polarized protons at the Research Center for Nuclear Physics (RCNP) in Osaka, Japan [37].

2. Model description

The BUU equation is a semi-classical approximation to the quantum transport equation [38,39]. For a momentum-dependent mean-field potential $U(\vec{r}, \vec{p})$, it reads as

$$\frac{\partial f}{\partial t} + \frac{\vec{p}}{E} \cdot \nabla_{\vec{r}} f + \nabla_{\vec{p}} U(\vec{r}, \vec{p}) \cdot \nabla_{\vec{r}} f - \nabla_{\vec{r}} U(\vec{r}, \vec{p}) \cdot \nabla_{\vec{p}} f = I_c, \quad (1)$$

where $f = f(\vec{r}, \vec{p})$ is the one-body phase-space distribution function of nucleons or their Wigner function. The r.h.s of Eq. (1) is the NN collision term including the Pauli blocking effect due to the Fermi statistics of nucleons, i.e.,

$$I_c = -g \int \frac{d\vec{p}_2}{(2\pi\hbar)^3} \frac{d\vec{p}_3}{(2\pi\hbar)^3} \frac{d\vec{p}_4}{(2\pi\hbar)^3} |\mathcal{M}_{12 \rightarrow 34}|^2 \times (2\pi)^4 \delta^4(p + p_2 - p_3 - p_4) \times [ff_2(1 - f_3)(1 - f_4) - f_3f_4(1 - f)(1 - f_2)], \quad (2)$$

where g is the degeneracy, $\mathcal{M}_{12 \rightarrow 34}$ is the in-medium transition matrix element, and $(1 - f_i)$ is the Pauli suppression factor. It is worth mentioning that higher-order quantum corrections to Eq. (1) can be added perturbatively [40].

In the present work, we solve the BUU equation by the lattice Hamiltonian (LH) method [41–43], which is a variant of the usual test particle method [44]. In the LH method, the total Hamiltonian H of the system is approximated by the lattice Hamiltonian H_L , i.e.,

$$H = \int \mathcal{H}(\vec{r}) d\vec{r} \approx l^3 \sum_{\alpha} \mathcal{H}(\vec{r}_{\alpha}) \equiv H_L, \quad (3)$$

where \mathcal{H} is the Hamiltonian density, \vec{r}_{α} represents the coordinate of certain lattice site α , and l is the lattice spacing. For the nucleon one-body phase-space distribution function $f_{\tau}(\vec{r}_{\alpha}, \vec{p})$, it is expressed as

$$f_{\tau}(\vec{r}_{\alpha}, \vec{p}, t) = \frac{(2\pi\hbar)^3}{gN_E} \sum_i^{\alpha, \tau} S[\vec{r}_i(t) - \vec{r}_{\alpha}] \delta[\vec{p}_i(t) - \vec{p}], \quad (4)$$

where S is the form factor and N_E is the number of ensembles or test particles used in the calculation. The sum in Eq. (4) runs over all test nucleons of isospin state τ that contribute to the lattice site α . In the present work, we adopt a triangular form factor S with the size of $4l$, and its detail can be found in Ref. [35]. The Hamiltonian in Eq. (3) contains both the Coulomb and the nuclear part [35] with the latter obtained from the N3LO Skyrme pseudopotential [45] SP6h, whose details can be found in Ref. [46].

In the present LBUU method, the ground state of a spherical nucleus at zero-temperature is obtained from the Thomas-Fermi approach [41,47–49] via the variation of the Hamiltonian with respect to the radial nucleon density $\rho_{\tau}(r)$. The obtained $\rho_{\tau}(r)$ is then used to determine the initial coordinates of test nucleons, while their initial momenta are generated according to zero-temperature Fermi distribution with local Fermi momentum given by $p_{\tau}^F(r) = \hbar[3\pi^2\rho_{\tau}(r)]^{1/3}$. This method for initialization ensures the stability of ground-state nuclei in BUU-like transport models [35,49].

For the collision term in the BUU equation, we implement it using the stochastic approach [50], which is more reliable than the commonly used geometric method when the mean free path λ_{MFP} of a test nucleon is not much larger than the interaction length between two test nucleons [51] or when the NN scattering cross section is very large. The collision probability P_{ij} of two test nucleons in the stochastic approach is determined from the NN collision term in Eq. (2), which is

$$P_{ij} = v_{\text{rel}} \sigma_{\text{NN}}^* S(\vec{r}_i - \vec{r}_{\alpha}) S(\vec{r}_j - \vec{r}_{\alpha}) l^3 \Delta t. \quad (5)$$

To reduce the statistical fluctuations of collision events and better reflect the nature of the BUU equation, we include collisions of test nucleons from different ensembles. In this case, the collision probability is reduced to P_{ij}/N_E , because of the scaling $\sigma_{\text{NN}}^* \rightarrow \sigma_{\text{NN}}^*/N_E$ of the in-medium NN cross section between test nucleons. Under such a scaling, the diluteness of the system, which is characterized by $\sqrt{\sigma_{\text{NN}}^*}/\lambda_{\text{MFP}}$, is reduced by the factor $\sqrt{N_E}$, and this makes it possible to solve the BUU equation almost exactly with a sufficiently large N_E achieved by adopting the GPU parallel computing.

For the i -th and j -th test nucleons colliding at the lattice site \vec{r}_{α} , the direction of their final momenta \vec{p}_3 and \vec{p}_4 are sampled according to the differential cross-section given in Ref. [52]. However, this collision can only happen if it is allowed by the Pauli principle via the factor $[1 - f(\vec{r}_{\alpha}, \vec{p}_3)] \times [1 - f(\vec{r}_{\alpha}, \vec{p}_4)]$. In the present LBUU method, the distribution function $f_{\tau}(\vec{r}_{\alpha}, \vec{p})$ is calculated from averaging its value in Eq. (4) over a given momentum-space sphere centered at \vec{p} with radius $R_{\tau}^p(\vec{r}_{\alpha}, \vec{p})$. In typical transport model calculations, $R_{\tau}^p(\vec{r}_{\alpha}, \vec{p})$ is taken to have a constant value of about one hundred MeV. In the present work, we use an improved form for $R_{\tau}^p(\vec{r}_{\alpha}, \vec{p})$ that is specifically proposed for small-amplitude nuclear collective dynamics near ground state [48], i.e., $R_{\tau}^p(\vec{r}_{\alpha}, \vec{p}) = \max[\Delta p, p_{\tau}^F(\vec{r}_{\alpha}) - |\vec{p}|]$, where $p_{\tau}^F = \hbar(3\pi^2\rho_{\tau})^{1/3}$ is the local nucleon Fermi momentum and Δp is a constant with the dimension of momentum that needs to be taken to be sufficiently small.

The treatment of Pauli blocking in transport models is crucial in calculating the width of nuclear collective excitations. At low incident energy or temperature, the Pauli blocking is notoriously difficult to handle in transport models [16,17]. This is mainly caused by the inaccuracy in calculating the local momentum distribution $f_{\tau}(\vec{r}_{\alpha}, \vec{p})$, which then leads to numerically spurious collisions and thus an overestimated GDR width as a result of the enhanced collisional damping. There are three main origins for the numerically spurious collisions in transport models: 1) fluctuations in calculating $f_{\tau}(\vec{r}_{\alpha}, \vec{p})$ caused by insufficiently large N_E ; 2) spurious thermal excitation caused by finite Δp in calculating $f_{\tau}(\vec{r}_{\alpha}, \vec{p})$ (also see Ref. [48]); and 3) diffusion in local momentum caused by finite

lattice spacing l when averaging over different local densities on the nuclear surface.

In choosing the parameter values in the LBUU calculations, we use the following criteria. For a given l and Δp , N_E should be large enough to eliminate the overwhelming majority of the spurious collisions caused by the first origin mentioned above, and at the same time l and Δp should be chosen to be sufficiently small to suppress the effects due to the second and third origins on the GDR width. After careful tests based on considerations of numerical accuracy and computation efficiency, we find the optimal values of $l = 0.5$ fm, $\Delta p = 0.05$ GeV and $N_E = 30000$. It is worth to mention that with the adoption of GPU parallel computing, it is possible to use a value for N_E that far exceeds those used in all previous calculations based on the BUU transport equation. Further reducing Δp and l and increasing N_E only leads to a negligible variation in the calculated GDR width.

We note that for the case of free NN cross section, an average of 97.93% of the attempted collisions in the ground state of ^{208}Pb are blocked by the Pauli principle, resulting in an average of 1.30 successful collisions of physical nucleons per fm/c during the time evolution of 0–500 fm/c. Also, the root mean square (rms) radius and the ground-state energy of ^{208}Pb vary by less than 3.6% (0.2 fm) and 3.2% (50 MeV), respectively, during this time evolution. With a reduced in-medium NN cross section, both the number of successful collisions and the change in the radius and binding energy are even smaller. Since the binding energy decreases monotonically with time without oscillations, it is not expected to have much effect on the calculated excitation energy of GDR. The energy violation, which is caused by our use of in-vacuum energy conservation in NN scatterings, instead of the in-medium energy conservation in the presence of the momentum-dependent potential, is not expected to affect the calculated width of GDR either. This is because the latter is controlled by the NN scattering rate, which depends on the NN scattering cross section and the Pauli blocking factor. We also note that although the radius and binding energy variations in LBUU are larger than those in the lattice Hamiltonian Vlasov approach of Ref. [35], where the rms radius and the binding energy almost do not change, and the radial density profile only changes slightly during the time evolution of 0–1000 fm/c, the LBUU method used in the present study is sufficiently accurate for investigating the GDR width.

3. Results and discussions

The collective excitation of a nucleus consisting of A nucleons can be induced by adding a perturbation to its Hamiltonian at the initial time t_0 , i.e., $\hat{H}_{ex}(t) = \lambda \hat{Q} \delta(t - t_0)$, where \hat{Q} is an appropriate excitation operator and λ is a small parameter. The width of a collective excitation is defined by the full width at half maximum (FWHM) of its strength function $S(E)$ as a function of the excitation energy E . In the linear response theory [53], the $S(E)$ is obtained from the Fourier integral

$$S(E) = -\frac{1}{\pi \lambda} \int_0^\infty dt \Delta(\hat{Q})(t) \sin \frac{Et}{\hbar}, \quad (6)$$

where $\Delta(\hat{Q})(t) = \langle 0' | \hat{Q} | 0' \rangle - \langle 0 | \hat{Q} | 0 \rangle$ is the time evolution of the response function of the nucleus to the excitation operator \hat{Q} with $|0\rangle$ and $|0'\rangle$ denoting the nuclear states before and after the perturbation, respectively. In terms of the Wigner transform $q(\vec{r}, \vec{p})$ of the one-body excitation operator \hat{q} , which is related to \hat{Q} by $\hat{Q} = \sum_i^A \hat{q}_i$, the expectation values in the above can be evaluated according to $\langle \hat{Q} \rangle(t) = \int f(\vec{r}, \vec{p}, t) q(\vec{r}, \vec{p}) d\vec{r} d\vec{p}$ using the nucleon phase-space distribution function $f(\vec{r}, \vec{p}, t)$. Details on the




Fig. 1. Time evolution of the isovector dipole response function $\Delta(\hat{Q}_{IVD})$ (left) and strength function $S(E)$ (right) of ^{208}Pb due to the perturbation of $\hat{H}_{ex} = \lambda \hat{Q}_{IVD} \delta(t - t_0)$ with $\lambda = 15$ MeV/c from the Vlasov calculation (solid lines) and the LBUU calculation (dashed lines) with $\sigma_{NN}^{\text{free}}$. The dotted cyan curve in the left window represents the expectation value of the \hat{Q}_{IVD} in the ground state of ^{208}Pb from the LBUU calculation with $\sigma_{NN}^{\text{free}}$.

single-particle operator used in exciting a ground state nucleus in transport models can be found in Ref. [35].

We first employ the present LBUU method to study the effect of NN scatterings on the isovector dipole response of ^{208}Pb using the excitation operator $\hat{Q}_{IVD} = \frac{N}{A} \sum_i^A \hat{z}_i - \frac{Z}{A} \sum_i^N \hat{z}_i$. In Fig. 1, we show the results obtained by using the free NN elastic scattering cross section taken from Ref. [52] with $\sigma_{NN}^{\text{free}}(p_{\text{lab}}) = \sigma_{NN}^{\text{free}}(0.1 \text{ GeV}/c)$ for neutron-neutron (nn) or proton-proton (pp) collisions at $p_{\text{lab}} \leq 0.1 \text{ GeV}/c$ and $\sigma_{NN}^{\text{free}}(p_{\text{lab}}) = \sigma_{NN}^{\text{free}}(0.05 \text{ GeV}/c)$ for neutron-proton (np) collisions at $p_{\text{lab}} \leq 0.05 \text{ GeV}/c$, as experimental data for lower incident momenta (p_{lab}) are unavailable. For comparison purpose, results from the LBUU calculation without NN scatterings, i.e., the Vlasov calculation, are also shown in Fig. 1. In both cases, we use in the initial perturbation the same parameter $\lambda = 15 \text{ MeV}/c$, which is also used in all the calculations in the present study, and we find that varying the value of λ by 2/3 almost has no effects on the value of the GDR width. As shown in the left window of Fig. 1 for the response function $\Delta(\hat{Q}_{IVD})(t)$, the inclusion of NN scatterings significantly enhances the damping of the oscillations. The dotted cyan curve in the left window of Fig. 1 represents the expectation value of the \hat{Q}_{IVD} in the ground state of ^{208}Pb from the LBUU calculation with $\sigma_{NN}^{\text{free}}$, which is negligible compared with that in the excited cases. To illustrate more clearly the effect of collisional damping, we show in the right window of Fig. 1 the GDR strength function $S(E)$ from the Fourier transformation of the response function. Note that the Vlasov calculation is carried out for a long evolution time of 1000 fm/c when the amplitude of the oscillation of $\Delta(\hat{Q})(t)$ almost vanishes so that the fluctuation in the calculated strength function from the Fourier transform of $\Delta(\hat{Q})(t)$ is negligible. We clearly see the large increase of GDR width after including NN scatterings, namely, the GDR width of ^{208}Pb are 6.5 MeV and 1.5 MeV in the LBUU calculations with and without NN scatterings, respectively.

Experimentally, the GDR width of ^{208}Pb has been well determined to be 4.0 MeV from the $^{208}\text{Pb}(\vec{p}, \vec{p}')$ reaction carried out at RCNP [37]. Our result from the LBUU calculation with $\sigma_{NN}^{\text{free}}$ thus significantly overestimates the GDR width of ^{208}Pb . This is understandable because of the absence of medium effect on the NN scattering in the calculation. Its inclusion is expected to reduce the NN cross section, weaken the collisional damping, and result in a smaller GDR width. The sensitivity of the GDR width to NN scatterings shown in Fig. 1 makes it possible to constrain the medium effect on the NN scattering cross section.

For σ_{NN}^* , we parameterize it by multiplying the free NN cross section with a medium-dependent correction factor. Specifically, we choose an exponential reduction factor as suggested by the T -matrix approach in Ref. [20], i.e.,




Fig. 2. The GDR width of ^{208}Pb from LBUU calculations for different values of α in σ_{NN}^* . The horizontal line represents the RCNP experimental value of 4.0 MeV. The inset shows the strength function with $\alpha = 1.8$ (solid line) and the shifted one (dash-dotted line) to match the experimental GDR peak energy. See text for details.




Fig. 3. Density dependence of the medium correction at different values for the total kinetic energy $T_{\text{c.m.}}$ of two scattering nucleons using NN cross sections from Eq. (7) with $\alpha = 1.8$, the Rostock cross section with $\alpha = 0.6$, and the FU4FP6 parameterization.

$$\sigma_{\text{NN}}^* = \sigma_{\text{NN}}^{\text{free}} \exp \left[-\alpha \frac{\rho / \rho_{\text{nuc}}}{1 + (T_{\text{c.m.}} / T_0)^2} \right]. \quad (7)$$

In the above, $T_{\text{c.m.}}$ is the total kinetic energy of two scattering test nucleons at the rest frame of the local medium or cell, $\rho_{\text{nuc}} = 0.16 \text{ fm}^{-3}$ is the nuclear normal density, and $T_0 = 0.015 \text{ GeV}$. For the parameter α , its original value in Ref. [20] is 0.6, which is called the Rostock cross section. In the present study, we treat it as a free parameter to control the strength of medium effect. Displayed in Fig. 2 is the GDR width Γ of ^{208}Pb obtained with different values of α . As expected, the GDR width decreases with increasing α . To reproduce the experimental value of $\Gamma = 4.0 \text{ MeV}$ measured at RCNP requires α to be as large as about 1.8, which indicates a very large medium reduction of the NN scattering cross section.

Although an early study on the balance energy, at which the nucleon direct flow in HICs vanishes, favors a small medium reduction of the NN cross section [1], more recent studies based on the analysis of the collective flow and nuclear stopping data [6,29] as well as the nucleon induced reaction cross section [30] require a large medium reduction. For comparisons, we also calculate the GDR width from the LBUU method with two different σ_{NN}^* , namely, the Fuchs cross section [23], which is obtained from the in-medium Dirac-Brueckner T matrix, and the FU4FP6 parameterization, which is preferred by the nucleon induced reaction cross section [30]. The value of GDR width of ^{208}Pb calculated using the FU4FP6 parameterization is 4.32 MeV, which is consistent with the experimental data. On the other hand, the values obtained with the Fuchs cross section and the Rostock cross section with $\alpha = 0.6$ based on microscopic calculations are 5.39 MeV and 5.59 MeV, respectively, which both overestimate the experimental value. In Fig. 3, we show the density dependence of the medium correction

$\sigma_{\text{NN}}^* / \sigma_{\text{NN}}^{\text{free}}$ at three different $T_{\text{c.m.}}$ values for the NN cross section in Eq. (7) with $\alpha = 1.8$, the Rostock cross section with $\alpha = 0.6$, and the FU4FP6 parameterization with the isospin asymmetry δ set to be 0.21 as in ^{208}Pb . The $T_{\text{c.m.}}^F \approx 0.073 \text{ GeV}$ in this figure represents the $T_{\text{c.m.}}$ of two nucleons at the Fermi surface of normal nuclear matter density ρ_{nuc} . It is seen that both the $\alpha = 1.8$ case and the FU4FP6 parameterization, which can describe the data of the GDR width of ^{208}Pb , show similar medium reductions, which are very large compared with that from the Rostock cross section.

In the inset of Fig. 2, we further show the strength function of the iso-vector excitation of ^{208}Pb from the LBUU calculation using the cross section in Eq. (7) with $\alpha = 1.8$ together with the experimental measurements at RCNP [37]. The LBUU result shifted to match the GDR peak energy is also included for comparison. As can be seen, our calculation nicely reproduces the shape of the experimental strength function, but overestimates the peak energy by about 1.4 MeV. A better agreement for the peak value could be achieved by varying the mean-field potential or nuclear EOS, which is known to significantly affect the peak energy of nuclear GDR [54]. Further tests show that the mean-field potential or nuclear EOS only weakly affects the obtained GDR width. As to the quantum corrections to the BUU equation [40], it is not expected to modify our results qualitatively as their effect on the damping of collective motion of heavy nucleus [55] is insignificant compared with that of NN scatterings.

4. Conclusions

We have used the LH method to solve the BUU transport equation with the binary collisions in the collision term treated via the stochastic approach. With the use of a sufficiently large number of test particles, the present LBUU method treats the Pauli blocking in the collision term of BUU equation with very high precision and thus significantly increases the accuracy in solving the BUU equation. From the accurately calculated GDR width of ^{208}Pb , we have found that it depends strongly on the magnitude of the in-medium NN cross section σ_{NN}^* , and the experimentally measured GDR width of ^{208}Pb from the $^{208}\text{Pb}(\vec{p}, \vec{p}')$ reaction at RCNP can only be reproduced with a NN cross section that is significantly reduced in nuclear medium. The large medium reduction of σ_{NN}^* raises challenges to microscopic calculations based on realistic NN interactions. Also, the effects of such a large medium reduction of σ_{NN}^* on the widths of other modes of giant resonances in nuclei and on the dynamics of HICs need to be studied as it may significantly affect the extracted information on the properties of nuclear matter at various densities.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Bao-An Li and Jun Su for useful discussions, and Meisen Gao, Jie Pu, Chen Zhong and Ying Zhou for the maintenance of the GPU servers. This work was partially supported by the National Natural Science Foundation of China under Contracts No. 11905302, No. 11947214, No. 11890714, No. 11625521 and No. 11421505, the Key Research Program of Frontier Sciences of the CAS under Grant No. QYZDJ-SSW-SLH002, the Strategic Priority Research Program of the CAS under Grants No. XDB16 and No. XDB34000000, the Major State Basic Research Development Program (973 Program) in China under Contract No. 2015CB856904, the US Department of Energy under Contract No. DE-SC0015266, and The Welch Foundation under Grant No. A-1358.

References

- [1] G.D. Westfall, W. Bauer, D. Craig, M. Cronqvist, E. Gaultier, S. Hannuschke, D. Klakow, T. Li, T. Reposeur, A.M. Vander Molen, W.K. Wilson, J.S. Winfield, J. Yee, S.J. Yennello, R. Lacey, A. Elmaani, J. Lauret, A. Nadasen, E. Norbeck, Mass dependence of the disappearance of flow in nuclear collisions, *Phys. Rev. Lett.* 71 (13) (1993) 1986–1989.
- [2] X. Cai, J. Feng, W. Shen, Y.-G. Ma, J. Wang, W. Ye, In-medium nucleon-nucleon cross section and its effect on total nuclear reaction cross section, *Phys. Rev. C* 58 (1) (1998) 572–575.
- [3] L.-W. Chen, F.-S. Zhang, G.-M. Jin, Z.-Y. Zhu, Isospin dependence of radial flow in heavy-ion collisions at intermediate energies, *Phys. Lett. B* 459 (1–3) (1999) 21–26.
- [4] J.-Y. Liu, W.-J. Guo, S.-J. Wang, W. Zuo, Q. Zhao, Y.-F. Yang, Nuclear stopping as a probe for in-medium nucleon-nucleon cross sections in intermediate energy heavy ion collisions, *Phys. Rev. Lett.* 86 (6) (2001) 975–978.
- [5] T.-T. Wang, Y.-G. Ma, C.-J. Zhang, Z.-Q. Zhang, Effect of in-medium nucleon-nucleon cross section on proton-proton momentum correlation in intermediate-energy heavy-ion collisions, *Phys. Rev. C* 97 (3) (2018) 034617.
- [6] P.-C. Li, Y.-J. Wang, Q.-F. Li, H.-F. Zhang, Collective flow and nuclear stopping in heavy ion collisions in Fermi energy domain, *Nucl. Sci. Tech.* 29 (12) (2018) 177.
- [7] H.L. Liu, Y.G. Ma, A. Bonasera, X.G. Deng, O. Lopez, M. Veselský, Mean free path and shear viscosity in central $^{129}\text{Xe}+^{119}\text{Sn}$ collisions below 100 MeV/nucleon, *Phys. Rev. C* 96 (6) (2017) 064604.
- [8] B. Barker, P. Danielewicz, Shear viscosity from nuclear stopping, *Phys. Rev. C* 99 (3) (2019) 034607.
- [9] F. Sammarruca, P. Krastev, Effective nucleon-nucleon cross sections in symmetric and asymmetric nuclear matter, *Phys. Rev. C* 73 (1) (2006) 014001.
- [10] B.-A. Li, C.M. Ko, Z. Ren, Equation of state of asymmetric nuclear matter and collisions of neutron-rich nuclei, *Phys. Rev. Lett.* 78 (9) (1997) 1644–1647.
- [11] P. Danielewicz, R. Lacey, W.G. Lynch, Determination of the equation of state of dense matter, *Science* 298 (5598) (2002) 1592–1596.
- [12] L.-W. Chen, C.M. Ko, B.-A. Li, Determination of the stiffness of the nuclear symmetry energy from isospin diffusion, *Phys. Rev. Lett.* 94 (3) (2005) 032701.
- [13] M.A. Famiano, T. Liu, W.G. Lynch, M. Mocko, A.M. Rogers, M.B. Tsang, M.S. Wallace, R.J. Charity, S. Komarov, D.G. Sarantites, L.G. Sobotka, G. Verde, Neutron and proton transverse emission ratio measurements and the density dependence of the asymmetry term of the nuclear equation of state, *Phys. Rev. Lett.* 97 (5) (2006) 052701.
- [14] M.B. Tsang, Y. Zhang, P. Danielewicz, M. Famiano, Z. Li, W.G. Lynch, A.W. Steiner, Constraints on the density dependence of the symmetry energy, *Phys. Rev. Lett.* 102 (12) (2009) 122701.
- [15] L. Ou, Z. Xiao, H. Yi, N. Wang, M. Liu, J. Tian, Dynamic isovector reorientation of deuteron as a probe to nuclear symmetry energy, *Phys. Rev. Lett.* 115 (21) (2015) 212501.
- [16] J. Xu, L.-W. Chen, M.B. Tsang, H. Wolter, Y.-X. Zhang, J. Aichelin, M. Colonna, D. Cozma, P. Danielewicz, Z.-Q. Feng, A. Le Fèvre, T. Gaitanos, C. Hartnack, K. Kim, Y. Kim, C.-M. Ko, B.-A. Li, Q.-F. Li, Z.-X. Li, P. Napolitani, A. Ono, M. Papa, T. Song, J. Su, J.-L. Tian, N. Wang, Y.-J. Wang, J. Weil, W.-J. Xie, F.-S. Zhang, G.-Q. Zhang, Understanding transport simulations of heavy-ion collisions at 100A and 400A MeV: comparison of heavy-ion transport codes under controlled conditions, *Phys. Rev. C* 93 (4) (2016) 044609.
- [17] Y.-X. Zhang, Y.-J. Wang, M. Colonna, P. Danielewicz, A. Ono, M.B. Tsang, H. Wolter, J. Xu, L.-W. Chen, D. Cozma, Z.-Q. Feng, S. Das Gupta, N. Ikeno, C.-M. Ko, B.-A. Li, Q.-F. Li, Z.-X. Li, S. Mallik, Y. Nara, T. Ogawa, A. Ohnishi, D. Oliinychenko, M. Papa, H. Petersen, J. Su, T. Song, J. Weil, N. Wang, F.-S. Zhang, Z. Zhang, Comparison of heavy-ion transport simulations: collision integral in a box, *Phys. Rev. C* 97 (3) (2018) 034625.
- [18] G.Q. Li, R. Machleidt, Microscopic calculation of in-medium nucleon-nucleon cross sections, *Phys. Rev. C* 48 (4) (1993) 1702–1712.
- [19] G.Q. Li, R. Machleidt, Microscopic calculation of in-medium proton-proton cross sections, *Phys. Rev. C* 49 (1) (1994) 566–569.
- [20] T. Alm, G. Röpke, M. Schmidt, Critical enhancement of the in-medium nucleon-nucleon cross section at low temperatures, *Phys. Rev. C* 50 (1) (1994) 31–37.
- [21] H.-J. Schulze, A. Schnell, G. Röpke, U. Lombardo, Nucleon-nucleon cross sections in nuclear matter, *Phys. Rev. C* 55 (6) (1997) 3006–3014.
- [22] M. Kohno, M. Higashi, Y. Watanabe, M. Kawai, In-medium nucleon-nucleon cross sections from nonrelativistic reaction matrices in nuclear matter, *Phys. Rev. C* 57 (6) (1998) 3495–3498.
- [23] C. Fuchs, A. Faessler, M. El-Shabshiry, Off-shell behavior of the in-medium nucleon-nucleon cross section, *Phys. Rev. C* 64 (2) (2001) 024003.
- [24] G. Mao, Z. Li, Y. Zhuo, Y. Han, Z. Yu, Study of in-medium NN inelastic cross section from relativistic Boltzmann-Uehling-Uhlenbeck approach, *Phys. Rev. C* 49 (6) (1994) 3137–3146.
- [25] Q. Li, Z. Li, E. Zhao, Density and temperature dependence of nucleon-nucleon elastic cross section, *Phys. Rev. C* 69 (1) (2004) 017601.
- [26] B.-A. Li, P. Danielewicz, W.G. Lynch, Probing the isospin dependence of the in-medium nucleon-nucleon cross sections with radioactive beams, *Phys. Rev. C* 71 (5) (2005) 054603.
- [27] B.-A. Li, L.-W. Chen, Nucleon-nucleon cross sections in neutron-rich matter and isospin transport in heavy-ion reactions at intermediate energies, *Phys. Rev. C* 72 (6) (2005) 064611.
- [28] Y. Zhang, Z. Li, P. Danielewicz, In-medium NN cross sections determined from the nuclear stopping and collective flow in heavy-ion collisions at intermediate energies, *Phys. Rev. C* 75 (3) (2007) 034615.
- [29] O. Lopez, D. Durand, G. Lehaut, B. Borderie, J.D. Frankland, M.F. Rivet, R. Bougault, A. Chbihi, E. Galichet, D. Guinet, M. La Commara, N. Le Neindre, I. Lombardo, L. Manduci, P. Marini, P. Napolitani, M. Pärlog, E. Rosato, G. Spadaccini, E. Vient, M. Vigilante, In-medium effects for nuclear matter in the Fermi-energy domain, *Phys. Rev. C* 90 (6) (2014) 064602.
- [30] L. Ou, X.-y. He, In-medium nucleon-nucleon elastic cross-sections determined from the nucleon induced reaction cross-section data, *Chin. Phys. C* 43 (4) (2019) 044103.
- [31] V.M. Kolomietz, V.A. Plujko, S. Shlomo, Interplay between one-body and collisional damping of collective motion in nuclei, *Phys. Rev. C* 54 (6) (1996) 3014–3024.
- [32] N.D. Dang, Shear-viscosity to entropy-density ratio from giant dipole resonances in hot nuclei, *Phys. Rev. C* 84 (3) (2011) 034309.
- [33] U. Garg, G. Colò, The compression-mode giant resonances and nuclear incompressibility, *Prog. Part. Nucl. Phys.* 101 (2018) 55–95.
- [34] A. Bracco, F. Camera, M. Mattiuzzi, B. Million, M. Pignanelli, J.J. Gaardhøje, A. Maj, T. Ramsøy, T. Tveter, Z. Želazny, Increase in width of the giant dipole resonance in hot nuclei: shape change or collisional damping?, *Phys. Rev. Lett.* 74 (19) (1995) 3748–3751.
- [35] R. Wang, L.-W. Chen, Z. Zhang, Nuclear collective dynamics in the lattice Hamiltonian Vlasov method, *Phys. Rev. C* 99 (4) (2019) 044609.
- [36] G. Ruetsch, M. Fatica, CUDA Fortran for Scientists and Engineers: Best Practices for Efficient CUDA Fortran Programming, Morgan Kaufmann, Waltham, MA, 2013.
- [37] A. Tamii, I. Poltoratska, P. von Neumann-Cosel, Y. Fujita, T. Adachi, C.A. Bertulani, J. Carter, M. Dozono, H. Fujita, K. Fujita, K. Hatanaka, D. Ishikawa, M. Itoh, T. Kawabata, Y. Kalmykov, A.M. Krumbholz, E. Litvinova, H. Matsubara, K. Nakanishi, R. Neveling, H. Okamura, H.J. Ong, B. Öznel-Tashenov, V.Y. Ponomarev, A. Richter, B. Rubio, H. Sakaguchi, Y. Sakemi, Y. Sasamoto, Y. Shimbara, Y. Shimizu, F.D. Smit, T. Suzuki, Y. Tameshige, J. Wambach, R. Yamada, M. Yosoi, J. Zenihiro, Complete electric dipole response and the neutron skin in ^{208}Pb , *Phys. Rev. Lett.* 107 (6) (2011) 062502.
- [38] P. Carruthers, F. Zachariasen, Quantum collision theory with phase-space distributions, *Rev. Mod. Phys.* 55 (1) (1983) 245–285.
- [39] G. Bertsch, S. Das Gupta, A guide to microscopic models for intermediate energy heavy in collisions, *Phys. Rep.* 160 (4) (1988) 189–233.
- [40] A. Bonasera, V.N. Kondratyev, A. Smerzi, E.A. Remler, Nuclear dynamics in the Wigner representation, *Phys. Rev. Lett.* 71 (4) (1993) 505–508.
- [41] R.J. Lenk, V.R. Pandharipande, Nuclear mean field dynamics in the lattice Hamiltonian Vlasov method, *Phys. Rev. C* 39 (6) (1989) 2242–2249.
- [42] H.M. Xu, W.G. Lynch, P. Danielewicz, G.F. Bertsch, Disappearance of fusion-like residues and the nuclear equation of state, *Phys. Rev. Lett.* 65 (7) (1990) 843–846.
- [43] H.M. Xu, Disappearance of flow in intermediate-energy nucleus-nucleus collisions, *Phys. Rev. Lett.* 67 (20) (1991) 2769–2772.
- [44] C.-Y. Wong, Dynamics of nuclear fluid. VIII. Time-dependent Hartree-Fock approximation from a classical point of view, *Phys. Rev. C* 25 (3) (1982) 1460–1475.
- [45] F. Raimondi, B.G. Carlsson, J. Dobaczewski, Effective pseudopotential for energy density functionals with higher-order derivatives, *Phys. Rev. C* 83 (5) (2011) 054311.
- [46] R. Wang, L.-W. Chen, Y. Zhou, Extended Skyrme interactions for transport model simulations of heavy-ion collisions, *Phys. Rev. C* 98 (5) (2018) 054618.
- [47] P. Danielewicz, Determination of the mean-field momentum-dependence using elliptic flow, *Nucl. Phys. A* 673 (1–4) (2000) 375–410.
- [48] T. Gaitanos, A.B. Larionov, H. Lenske, U. Mosel, Breathing mode in an improved transport approach, *Phys. Rev. C* 81 (5) (2010) 054316.
- [49] H. Lin, P. Danielewicz, One-body Langevin dynamics in heavy-ion collisions at intermediate energies, *Phys. Rev. C* 99 (2) (2019) 024612.
- [50] P. Danielewicz, G.F. Bertsch, Production of deuterons and pions in a transport model of energetic heavy-ion reactions, *Nucl. Phys. A* 533 (1991) 712–748.
- [51] Z. Xu, C. Greiner, Thermalization of gluons in ultrarelativistic heavy ion collisions by including three-body interactions in a parton cascade, *Phys. Rev. C* 71 (6) (2005) 064901.
- [52] J. Cugnon, D. L'Hôte, J. Vandermeulen, Simple parametrization of cross-sections for nuclear transport studies up to the GeV range, *Nucl. Instrum. Methods Phys. Res., Sect. B, Beam Interact. Mater. Atoms* 111 (3–4) (1996) 215–220.
- [53] A. Fetter, J.D. Walecka, Quantum Theory of Many-Particle Systems, McGraw-Hill, New York, 1971.
- [54] L. Trippa, G. Colò, E. Vigezzi, Giant dipole resonance as a quantitative constraint on the symmetry energy, *Phys. Rev. C* 77 (6) (2008) 061304(R).
- [55] V. Kondratyev, A. Smerzi, A. Bonasera, Dynamics of a quantal system, *Nucl. Phys. A* 577 (3–4) (1994) 813–828.