

Elastic scattering of alpha particles from ^{28}Si target

*Aparajita Dey¹, S. Ganguly², V. Srivastava¹, T.K. Rana¹, S. Kundu¹, K. Banerjee¹, P. Roy¹, H. Pai¹, C. Bhattacharya¹, T.K. Ghosh¹, R. Pandey¹, G. Mukherjee¹, J.K. Meena¹, M.R. Gohil¹, and S. Bhattacharya¹

¹Variable Energy Cyclotron Centre, 1/AF, Bidhan Nagar, Kolkata – 700 064, INDIA

²Department of Physics, Bethune College, 181, Bidhan Sarani, Kolkata – 700 006, INDIA

*email: aparajita@vecc.gov.in

At energies of a few tens of MeV α -particle elastic scattering angular distribution exhibits pronounced structure which usually varies smoothly with target mass and incident energy. The elastic scattering has been generally well described by the optical model potential in which the parameters of the potential have been found to vary smoothly with target mass (A) and bombarding energy (E_{lab}). However, there are cases of low energy α -scattering from light targets which show irregular variations with A and/or E_{lab} which may be due to compound nucleus or nuclear structure effects. This is particularly true for targets in the s-d shell.

Here we report the α -particle elastic scattering angular distributions from ^{28}Si targets at various incident energies. The α -particle elastic scattering was studied earlier at 14.47 to 40 MeV [1], 41 MeV [2], 45 MeV [3] and 57.8 MeV [4]. In the present experiment, we study the $\alpha + ^{28}\text{Si}$ reaction at 50 MeV.

The experiment was performed using α -ion beam of energy 50 MeV from the Variable Energy Cyclotron at VECC, Kolkata. The target was self-supporting ^{28}Si foil ($\sim 740 \mu\text{g/cm}^2$). Both elastic and transfer channels were detected using two three element (Si-Si-CsI) solid state telescopes mounted on two arms of the scattering chamber which could move independently. Each telescope consists of a $50 \mu\text{m}$ ΔE single-sided Si-strip detector (16 horizontal strips of 3 mm width), $500 \mu\text{m}$ E/ ΔE double-sided Si-strip detector (16 strips, width 3 mm, both side mutually orthogonal to each other) and two 6 cm CsI(Tl) detectors. A 6 mm slit was placed in front of each telescope and the data was collected only at the reaction plane. Well separated ridges corresponding to different particles as well as excited states are clearly seen in ΔE -E scatter plot (Fig. 1). The solid angle subtended by each strip is $\sim 5 \text{ msr}$. The detectors

were calibrated using elastically scattered α -ion from Au target and a Th- α source.




Fig. 1: Two dimensional ΔE -E plot using Si (500 μm) – CsI(Tl) combination for the α (50 MeV) + ^{28}Si reaction at the angle (a) $\theta_{\text{lab}} = 16.5^\circ$ and (b) $\theta_{\text{lab}} = 87^\circ$.

The measured elastic scattering angular distributions exhibit pronounced oscillations, as shown in Fig. 2. The optical model analysis of the data was carried out using the parametric Woods-Saxon forms for both the real and imaginary potentials. The optical model potential, used to

describe the elastic angular distributions at each energy, had the following form

$$U(r) = V^m(r) + iW_F^n(r) + iW_D(r) + V_C(r)$$

where $V(r)$ denotes the volume type Woods-Saxon real potential, $W_F(r)$ is a volume type Woods-Saxon imaginary potential to simulate the fusion after penetration of the barrier and $W_D(r)$ is a derivative type Woods-Saxon imaginary potential to account for the absorption due to reactions occurring at the surface.

$$V(r) = -V_o [1 + \exp(r - R_o)/a_o]^{-1}$$

$$W_F(r) = -W_v [1 + \exp(r - R_v)/a_v]^{-1}$$

$$W_D(r) = -W_s [1 + \exp(r - R_s)/a_s]^{-1}$$

$V_C(r)$ is the Coulomb potential due to a uniformly charged sphere of radius $R_C = 1.3A_T^{1/3}$ fm. The form factor powers m and n are integers, and the values are 1 or 2. The optical model calculations were carried out using different combinations of the values of m and n .

The search code ECIS94 [5] has been used to perform the model calculations. In the present work, we will perform the optical model fitting for the energy 14.5 to 58 MeV for the $\alpha + {}^{28}\text{Si}$ reaction.

In the present calculation, we would take nine variable parameters described earlier. The optical model potential fitting for different energies have been shown in Fig. 2 by solid curves.

Further analysis is in progress.

The authors thank the cyclotron operating crew for the smooth running of the machine. This work is a part of the SERB project No. SR/FTP/PS067/2012 dated 10/06/2012. A.D acknowledges with thanks the financial support provided by the Science and Engineering Research Board, Department of Science and Technology, Government of India.




Fig. 2: Elastic scattering angular distributions for the $\alpha + {}^{28}\text{Si}$ reactions.

References

- [1] A.S.B. Tariq, *et al.*, Phys. Rev. C **59** 2558 (1999).
- [2] C.B. Fulmer, *et al.*, Phys. Rev. C **18** 621 (1978).
- [3] V.P. Darshan, *et al.*, J. Phys. G: Nucl. Part. Phys. **21** 385 (1995).
- [4] P. Demetrian, *et al.*, Phys. Rev. C **72** 034607 (2005).
- [5] J. Raynal, ECIS94, NEA 0850/16.