


From the “Nuclear Mill” to the Large Hadron Collider ... and Beyond

Chris Quigg


Fermilab Colloquium · October 18, 2023



Shoe-Fitting Fluoroscope (50-kV)



Rolf Widerøe (1943), Tuddal (Telemark) NO:
colliding clouds inspire colliding beams
Kernmühle German patent issued 1953



XVIII. THE DEPARTMENT OF PHYSICS.

OFFICERS OF INSTRUCTION.

ALBERT A. MICHELSON, PH.D., *Head Professor of Physics.*

SAMUEL W. STRATTON, S.B., *Assistant Professor of Physics.*

GLEN M. HOBBS, S.B., *Assistant in Laboratory.*


INTRODUCTION.

Within the last twenty years the teaching and practice of Physics, have undergone a revolution more complete than in any of its sister sciences. This result may be attributed, to a very great extent, to the enormous development of its applications to electrical industries. No other industrial application since the invention of the steam-engine has so enhanced the appreciation of the importance of exact knowledge, or given a greater impetus to the search for new truths in the unexplored regions on the borderland of science.

So closely interwoven are the advances in pure science and its applications that it is difficult to say which has been of greater service to the other, but it is evident that it is as ill-advised to ignore the powerful stimulus furnished by the practical development of scientific ideas, as it is to belittle the influence which theoretical and experimental science have had on the world's material prosperity.

While it is never safe to affirm that the future of Physical Science has no marvels in store even more astonishing than those of the past, it seems probable that most of the grand underlying principles have been firmly established and that further advances are to be sought chiefly in the rigorous application of these principles to all the phenomena which come under our notice.

It is here that the science of measurement shows its importance—where quantitative results are more to be desired than qualitative work. An eminent physicist has remarked that the future truths of Physical Science are to be looked for in the sixth place of decimals.




Wilhelm Conrad Röntgen (Würzburg, 1895–1896)

RADIOACTIVITY, NEW PROPERTY OF MATTER


40 - 1922 - 92. - Jules Vaill Jarry L d. Dr. Polakowski
Papier noir. Cuir de cuir noir.
Estampe bleue le 27. de cette année. Le 16 -
Vendredi le 17. -



Henri Becquerel (Paris, 1896)




J. J. Thomson (Cambridge, 1897)




Thorium C = ^{212}Bi ; 6.208 MeV; C' = ^{212}Po ; 8.784 MeV


Maria Skłodowska-Curie & Pierre Curie (Paris, 1898–1902)



Max Planck (Berlin, 1900); Quanta

h... blackbody spectrum





Einstein (Bern, 1905) photoelectric effect / Brownian motion

20

ALBERT EINSTEIN

No. 18 of 22 Famous Americans

One of the great men of modern science, Albert Einstein was driven out of Germany by Adolf Hitler and came to the United States in 1933. He is a man of very simple tastes . . . even refusing to wear socks except in winter. His study is furnished with only an unpainted desk, a few unpainted shelves, a pencil and paper! He likes to wear old clothes . . . refusing to change from his baggy pants even when distinguished visitors are coming!


LOOK 'n see

HOW MANY PEOPLE
UNDERSTAND EINSTEIN'S
THEORY?



PLACE THE RED PAPER OVER THIS
CARD AND SEE THE ANSWER.

I.C.T.C.G.

PRINTED IN U.S.A.





Jean Perrin (Paris, 1908)



Heike Kamerlingh Onnes (Leiden, 1908)
Door meten tot weten



First superconducting magnet: Pb wire, 600 gauss (< 7.2 K)
Museum Boerhaave, Leiden



Hans Geiger, Ernest Marsden, *Ernest Rutherford* (Manchester, 1908–1913)



18,000,000 Volts

Cable, Strung between Mountains, Acts



Dr. Kurt Urban, most shocked man in the world, on roof of Mt. Generoso lightning-proof cabin.

AT THE peril of their lives, three young Germans have just succeeded in drawing electric currents of 18,000,000 volts from the skies during a violent display of lightning. On the slopes of Mt. Generoso in Switzerland they duplicated Benjamin Franklin's famous experiment of catching atmospheric electricity with a kite and key, on a scale which that pioneer never dreamed of.

From the side of Generoso's bony-ridged slope, they stretched, as an aerial, a metallic cable across a chasm to a neighboring peak. Threaded through cylinders of galvanized metal, this antenna resembled a string of beads for a giantess. Its knobs were designed to keep the currents from leaping from the ends. Instead, an escape was provided in an adjustable spark gap, from which the electricity could be carried to a lightning-proof cabin sheathed with metal beneath the brow of the mountain. Here were meters and other instruments to gage the force of the electricity.

ONE day not long ago the daring three, Drs. A. Brasch, F. Lange, and Kurt Urban, of the University of Berlin, saw a storm of unusual intensity brewing. The sky became pitch dark. Peals of thunder hurried the scientists as they threw switches and tested instruments. Then with a rush of wind up the valley that almost swept away the aerial, the fury of the storm broke.

Tongues of electric flame played about the rocks of the mountain's face near the summit. Filled with metallic ore, it was a natural lightning rod. Great yellow sparks snapped every second across the spark gap. The pointers of the voltmeters within the cabin were doing a dance. There came a brief lull, ominous in its calm.

Suddenly a terrific thunderclap seemed



Workmen installs metal "corona cylinders" in the Mt. Generoso cable to keep the high-voltage current from leaking into the air.



High-voltage currents, knowing no bonds, do strange things. Here is an arc that is leaping between two high-tension wires.

forces up to 2,600,000 volts—by a considerable margin the most powerful ever made.

The rays from this tube far exceed in power those of all the radium in the world. They easily penetrate a wall of lead a yard thick, an impassable barrier to all ordinary X-rays. Next the experimenters plan to build a 7,000,000-volt tube of similar

Captured from the Sky

as Antenna to Gather up Lightning

design. Only lightning can run it, for the greatest electrical tension ever produced in a laboratory is 5,000,000 volts.

How can such staggering forces be put to use? No one knows exactly, yet. But there are several long-awaited tests for which the power of lightning may be tried.

One of these is the dream of alchemists of old—the "transmutation of elements," such as turning base metals into gold. In recent years science has conceded that metals and other substances once held unchangeable can actually be transformed into entirely different ones under certain circumstances, requiring tremendous forces.

So far the only proved cases of transmutation have been done with the mysterious power of radium; thus, minute quantities of aluminum, phosphorus, and other things have been

This is not lightning, but a discharge of laboratory electricity leaping from insulators. It shows why extraordinary apparatus is required in harnessing 18,000,000 volts.


turned into hydrogen. Perhaps lightning's power may permit more useful transformations of other elements, when harnessed in a vacuum tube.

Then there is the question of unlocking the power of the atom. Some scientists hold that the energy contained in a single spoonful of water, for example, is sufficient to drive a modern steamship across the ocean. But this energy, if it does exist, is so securely locked up that no way has yet been found to release it. Should lightning's force break an entrance into this stronghold of power, the world might see a new era of industrial greatness based upon free atomic energy.

LASTLY, the rays from an electric tube operated by lightning's power may well have the most profound effect upon human beings. Whether beneficial or not, it is too early to say. Perhaps the audacious experimenters will find a curative ray more effective against cancer than any in sight!



Here is part of the apparatus that was used in the Mt. Generoso tests. This strange network was designed to carry the enormous voltage to the laboratory where the scientists waited in danger of instant death.



This 55-foot man-made arc was drawn out in Westinghouse laboratory in recent test.


instrument hitherto known to science. On the other hand it may be a "death ray."

Such fascinating possibilities spur man's advance in his attempt to harness stupendous electric forces. Only a few years ago the pinnacle of these efforts

was a crackling spark of 3,600,000 volts produced at the General Electric Company's laboratory in Pittsfield, Mass.

Then experts at the Carnegie Institution in Washington, D. C., built a mighty machine that could command 5,000,000 volts. Meanwhile the German experimenters had already installed a preliminary apparatus on Mt. Generoso, a peak famed for the frequency and violence of its electric storms, and were drawing two-million-volt sparks (P. S. M., Jan. '29, p. 23). A higher, rebuilt antenna made their recent feat possible. One of the experimenters, Dr. Urban, has acquired the sobriquet of "most shocked man in the world" from being knocked unconscious by sky currents.

IT MUST be understood that their antenna is not directly struck by lightning, for if it were, despite an electrical "safety valve" they have provided, they would probably all be killed. The aerial takes current from the clouds in two ways. Electricity in the air itself leaks down the cables in steady sparks. But the greatest voltages are obtained when a lightning bolt passes close to the aerial and a sympathetic surge of electricity is induced in the wire. In this way they have now captured 18,000,000 volts, and even greater voltages, up to thirty million, are in sight!




Walton · Rutherford · Cockcroft (Cambridge, 1932)



Fermilab's Cockcroft–Walton Machine: 750 keV, 1972–2012

Lawrence & Livingston Cyclotron (Berkeley, 1931–2)




60" cyclotron magnet frame (1938)



16-MeV deuterons from Lawrence's 60" cyclotron

SCIENCE



LIFE, February 5, 1940


THE BEAM OF PURPLE LIGHT IS AIR IONIZED BY FUSILLADE OF SUB-ATOMIC BULLETS FROM CYCLOTRON AT RIGHT.

THOUGH THE BEAM ITSELF IS NOT VERY HOT, IT WOULD

BURN YOUR HAND TO A CRISP IN AN INSTANT OR EXPLODE WATER INTO SUPERHEATED STEAM.

TO PROTECT OPERATORS, CYCLOTRON IS ENCLOSED IN THICK WATER





THE UNIVERSITY OF CHICAGO
CHICAGO 37 - ILLINOIS
THE CENTRAL ADMINISTRATION

ROBERT M. HUTCHINS - Chancellor
ERNEST CADMAN COLWELL - President
R.W. HARRISON - Vice-President
and Dean of Faculties
J. A. CUNNINGHAM - Vice-President
LYNN A. WILLIAMS, JR. - Vice-President

November 5, 1948

Dear Mr. Grace:

The fabrication and erection by your company of the magnet frame for the University's 170" synchrocyclotron has now been completed. Although the synchrocyclotron will not be in operation until early in 1950, the erection of the magnet frame represents a major step toward completion.

The 2070 ton magnet frame was fabricated from low-carbon steel forgings in accordance with our specifications. The magnet is an assembly of some twenty-eight units, weighing up to 81 tons each. Dr. Samuel K. Allison, Director of our Institute for Nuclear Studies, has just advised me that in all of the vital dimensions, the deviations are smaller than 10/1000 of an inch. In view of the fact that such accuracy has been obtained in a length of 250 inches, we are impressed by the mechanical precision achieved.


The representatives of your company have, without exception, been most cooperative and helpful in all phases of the work. We are extremely grateful to you and your associates for the splendid results which have been achieved.

Sincerely yours,
Ernest Cadman Colwell


E. G. Grace, chairman of Bethlehem Steel, reads letter (reproduced above) acknowledging Bethlehem's part in building the cyclotron for the University of Chicago. Left to right, A. B. Homer, president of Bethlehem Steel; E. C. Colwell, president of the University; Mr. Grace; and Dr. S. K. Allison, Director of Nuclear Studies at Chicago.

8


Shaping the ingot on 7500-ton press forge.



Machining a member of the magnet yoke.



Pole pieces are also intricately shaped.



Magnets for Giant Cyclotrons Made by Bethlehem

BIG ATOM-SMASHERS TO DELVE INTO SECRETS OF NUCLEAR STRUCTURE

Two of the nation's chief scientific and educational institutions, Columbia University and the University of Chicago, both actively engaged in nuclear research, are now constructing laboratories to be used exclusively for such studies. They will have the most powerful cyclotrons, popularly known as atom-smashers, which the world has ever seen. The steel for the 2500-ton magnets for both of these cyclotrons was produced at the Bethlehem plant.

Recent studies of the structure of matter have revealed that enormous forces are locked up in the atoms, or rather the nuclei—the small units from which matter is supposedly built up. These forces, so dramatically demonstrated to a startled world at Hiroshima and Nagasaki, will, when directed into peacetime channels, become a real boon to mankind.

Due to the magnitude of the forces involved, the equipment used in their release must of necessity be massive.

The cyclotrons at the Columbia University and the University of Chicago laboratories are designed to fire protons, which are the bullets used in breaking up the nuclei, at the unprecedented energy of 400,000,000 electron volts, to give the protons a speed of over 150,000 miles per second, very close to that of light.


The central part of each of these giant cyclotrons, or synchro-cyclotrons as they are also called, is a huge electro-magnet which weighs about 2500 tons. In fact, each magnet is so big that it was necessary to erect it on the site and then build the laboratory around it. The Columbia magnet, for instance, consists of a rectangular yoke assembly, 33 ft. long, 21 ft. high, and 14 ft. wide, shaped as a window with an opening 22½ ft. x 10½ ft. in which the magnet pole pieces are placed. The University of Chicago magnet is a similar structure.

Bethlehem's contribution to this

branch of science in the production of the steel for these magnets was made possible because of the exceptional facilities available at the Bethlehem, Pa., plant and the accumulated experience and skill dating back to the very beginning of the company. These huge magnets cannot be made in one piece, but must be assembled from a number of large sections bolted together to

form the yoke and the poles. In the Columbia magnet this assembly is made up of 38 pieces, the largest of which weighs 60 tons. The Chicago unit consists of 28 pieces, the largest of which weighs over 80 tons. Both have a diameter of 170 inches.


The magnet members were shaped from ingots weighing as much as 465,000 pounds, among the largest ever made.




465,000-lb. ingot for the University of Chicago cyclotron.

CCM @ FNAL


The Chicago cyclotron during erection.



170-inch diameter pole pieces in place.



TRIUMF Cyclotron: 18-m diameter, 520 MeV, 10^{15} p/sec




vacuum chamber 46 cm high

The 1953 Cosmic Ray Conference at Bagnères de Bigorre: the Birth of Sub Atomic Physics

J.W. Cronin^a

Department of Astronomy and Astrophysics, Enrico Fermi Institute, University of Chicago,
5640 South Ellis Ave., Chicago, IL 60637, USA




Received 2 March 2011

Published online 29 August 2011


© EDP Sciences, Springer-Verlag 2011

Abstract. The cosmic ray conference at Bagnères de Bigorre in July, 1953 organized by Patrick Blackett and Louis Leprince-Ringuet was a seminal one. It marked the beginning of sub atomic physics and its shift from cosmic ray research to research at the new high energy accelerators. The knowledge of the heavy unstable particles found in the cosmic rays was essentially correct in fact and interpretation and defined the experiments that needed to be carried out with the new accelerators. A large fraction of the physicists who had been using cosmic rays for their research moved to the accelerators. This conference can be placed in importance in the same category as two other famous conferences, the Solvay congress of 1927 and the Shelter Island Conference of 1948.




Louis Leprince-Ringuet
Bagnères-de-Bigorre (1953)
« Mais nous devons aller vite, nous
devons courir sans ralentir notre
cadence : nous sommes poursuivis
... nous sommes poursuivis par les
machines ! »

Synchrotrons




Brookhaven Cosmotron, 3+ GeV (1952–4)



Berkeley Bevatron, 6.2 GeV (1954)

Fermi's Dream Machine (1954): 5000-TeV $p \sim E_{\text{cm}} = 3 \text{ TeV}$

2-tesla magnets at radius $\approx 8000 \text{ km}$




Projected operation 1994
cost \$170 Billion
1954 inflation rate 0.75%



Why must accelerators be so large?



$$\text{radius} \approx \frac{10 \text{ km}}{3} \cdot \left(\frac{\text{Energy}}{1 \text{ TeV}} \right) \left/ \left(\frac{\text{Magnetic Field}}{1 \text{ tesla}} \right) \right.$$

+ aperture to contain unruly beam



Bevatron Magnet Frame (1950)

Alternating-gradient (strong) focusing



Christofilos (Athens, 1949)
Courant, Livingston, Snyder (Brookhaven, 1952)

Impact of alternating-gradient (strong) focusing

Synchrotron (circumference, E)	Beam Tube	Magnet Size
Cosmotron (230 ft, 3 GeV)	9 in x 3 ft	8 ft x 8 ft
Bevatron (400 ft, 6.2 GeV)	1 ft x 4 ft	9-1/2 ft x 20-1/2 ft
Brookhaven AGS (0.5 mi, 30 GeV)	2.7 in x 6 in	33 in x 39 in
Fermilab Main Ring (2π km, 400 GeV)	2 in x 4 in	14 in x 25 in
Fermilab Tevatron (\approx 1 TeV)	70 mm	4.3 T (SC, 4.2 K)
CERN LHC (27 km, \rightarrow 7 TeV)	56 mm	8.3 T (SC, 1.3 K)



Main Ring volume under vacuum < Bevatron

Attainment of Very High Energy by Means of Intersecting Beams of Particles

D. W. KERST,* F. T. COLE,† H. R. CRANE,‡ L. W. JONES,‡ L. J. LASLETT,§ T. OHKAWA,|| A. M. SESSLER,¶ K. R. SYMON,** K. M. TERWILLIGER,‡ AND NILS VOGT NILSEN††

Midwestern Universities Research Association,‡‡ University of Illinois, Champaign, Illinois

(Received January 23, 1956)




O'Neill (1956)
25⊗25-GeV $p\bar{p}$
~1.3 TeV fixed target

Carl Barber, Bernard Gittelman, G. K. O'Neill, Burton Richter: CBX
300-MeV electrons in collision \Rightarrow 350 GeV fixed-target

CERN Intersecting Storage Rings (1971)—Kjell Johnsen: “the finest instrument one can imagine for research in accelerator physics”

0.3 π km circumference
two independent rings
1.33-T dipoles
 pp @ $E_{cm} \rightarrow 62$ GeV
 ~ 2 TeV fixed-target
millions events/sec

also stored d, a, \bar{p}





Underinstrumented!
discovered rising σ_t
hints of qq scattering
Missed $J/\psi, \Upsilon$
A learning experience


Superconducting accelerator magnets based on Type-II superconductors



John Hulm



Bernd Matthias



"Yep, with them new superconductors, they built that
little SSC ring right there beneath your feet"

BOOTH

How can accelerators be so small?



Development of large-scale **cryogenic technology**,
to maintain many km of magnets at a few kelvins.

Active optics to achieve real-time corrections of orbits
enables reliable, highly tuned accelerators with small-aperture magnets.
Also “cooling,” or phase-space compaction, of stored (anti)protons.

Evolution of **vacuum technology**.
Beams stored for approximately 20 hours travel
 $\sim 2 \times 10^{10}$ km, about 150 times the Earth–Sun distance,
without encountering a stray air molecule. **LHC: 10^{-13} atm.**

Improvements in accelerating gradient
through efficient **(superconducting)** RF cavities

Simon van der Meer, Carlo Rubbia: $S\bar{p}\bar{p}S$ Collider (1981)




G. I. Budker, A. N. Skrinsky (Novosibirsk, 1970s)

UA1: $Z^0 \rightarrow e^+e^-$


EVENT 7433. 1001.

a)



EVENT 7433. 1001.

b)



Tevatron proton-antiproton collider







WH15E

Tevatron Collider Pioneers



Alvin Tollestrup (Magnets)




Helen Edwards (Machine)



John Peoples (Antiprotons)



DIPOLE




QUADRUPOLE



NO
FISHING IN
A-Q-L
ADDITION, MAIN RING
PARKING
←





315 Physicists Report Failure In Search for Supersymmetry

The negative result illustrates the risks of Big Science, and its often sparse pickings.

By MALCOLM W. BROWNE

THREE HUNDRED AND FIFTEEN physicists worked on the experiment.

Their apparatus included the Tevatron, the world's most powerful particle accelerator, as well as a \$65 million detector weighing as much as a warship, an advanced new computing system and a host of other innovative gadgets.

But despite this arsenal of brains and technological brawn assembled at the Fermilab accelerator laboratory, the participants have failed to find their quarry, a disagreeable reminder that as science gets harder, even Herculean efforts do not guarantee success.


In Centennial of One of Its Biggest Failures, Science Rejoices



Clark University Archives
Albert A. Michelson (left) and Edward W. Morley.



Case Western Reserve University Archives



The New York Times/Horley Trust
Prof. Donald E. Schuele with a replica of the 1887 interferometer at Case Western Reserve University. It uses a laser beam.

By MALCOLM W. BROWNE

ONE hundred years ago this month, two experimenters in an Ohio basement gently rotated a carousel of mirrors floating in a tub of mercury and inadvertently smashed the reassuring vision of a clockwork cosmos that had prevailed since the time of Sir Isaac Newton.

Physicists and philosophers have agonized over the experiment ever since, but one by one, nearly all have been compelled to abandon cherished theory in the face of its evidence.

The experimenters, Albert Abraham Michelson and Edward William Morley, set out to prove the existence of

"ether" — an intangible and invisible fluid thought by most 19th-century scientists to permeate the entire universe. Michelson and Morley expected to demonstrate that ether sped or slowed light waves; if this were the case, the ether would represent a universal standard against which the positions and motions of everything in the universe could be measured.

Much to their consternation, they failed. But their failure was pivotal. A new generation of scientists, struggling to explain the shocking experiment, eventually demonstrated that nothing in the universe has absolute reality, and that causality itself ceases to operate at the microscopic level of existence.

"It came at a time of transition from a period when science was dominated by wealthy amateurs to the era of true scientific professionals," added Dr. Taylor, a professor of physics at Case Western Reserve University in Cleveland. "Michelson, one of the first of these professionals, was also the first American to win a Nobel prize,

yardstick for space and time. So last week, scientists, educators and artists gathered in Cleveland to begin a six-month observance of what many scientists regard as science's most crucial failure.

"I think it was the most important experiment ever performed," said Dr. Philip L. Taylor, an organizer of the centennial. "If Michelson and Morley had obtained a different result, our whole view of space and time today would have been different."

Many experimenters before Michelson and Morley had demonstrated that light behaves in some ways like the waves that traverse bodies of water or the sound waves that travel through air. Reasoning by analogy,

Continued on Page C3


in 1907, for scientific achievement." (Only one other American, Theodore Roosevelt, had won a Nobel prize in any category. President Roosevelt received the peace prize in 1906.)

Dr. Taylor said the experiment marked "the birth of modern physics in America." But Michelson and Morley themselves were deeply troubled by its implications, and never fully accepted the radical interpretation developed for it by Einstein and other leading theorists.

Many experimenters before Michelson and Morley had demonstrated that light behaves in some ways like the waves that traverse bodies of water or the sound waves that travel through air. Reasoning by analogy,

NYT, April 28, 1987

D0



Diverse searches for new phenomena

Limits on
supersymmetric particles
extra spatial dimensions
signs of new strong dynamics
leptoquarks
new gauge bosons
magnetic monopoles

...

*Tevatron experiments did not find
what is not there!*

CDF/D0 (1995)
top quark:
mass ≈ 173 GeV ≈ 186 u
lifetime ≈ 0.4 ys



The Tevatron Collider Physics Legacy

Paul D. Grannis¹ and Melvyn J. Shochet²

¹Department of Physics and Astronomy, Stony Brook University, Stony Brook, New York 11794; email: pgrannis@sunysb.edu

²Enrico Fermi Institute and Department of Physics, University of Chicago, Chicago, Illinois 60637; email: shochet@hep.uchicago.edu

Annu. Rev. Nucl. Part. Sci. 2013. 63:467–502

First published online as a Review in Advance on July 29, 2013

The *Annual Review of Nuclear and Particle Science* is online at nucl.annualreviews.org

This article's doi:
10.1146/annurev-nucl-102212-170621

Copyright © 2013 by Annual Reviews.
All rights reserved

Keywords

Tevatron, CDF, D0

Abstract

The proton–antiproton Tevatron Collider at Fermilab began physics operation in 1988, and by its shutdown in 2011 it had delivered more than 10 fb^{-1} of data to the two general-purpose detectors CDF and D0. Thus far, these experiments have published more than 800 papers studying the character of the strong nuclear force, measuring the properties of hadrons containing heavy quarks, elucidating the nature of the electroweak force, discovering and measuring the properties of the top quark, seeking evidence for the Higgs boson, and searching for new phenomena beyond the standard paradigm of particle physics. We summarize the results that define the physics legacy of the Tevatron program.

Contents

1. INTRODUCTION	468
2. QCD STUDIES	470
2.1. Jet Production	470
2.2. Prompt Photon Production	471
2.3. Associated Production of Gauge Bosons and Jets	472
2.4. Correlations Within Events	474
2.5. Diffraction and Elastic Scattering	474
3. HEAVY-FLAVOR PHYSICS	475
3.1. Heavy-Flavor Production	475
3.2. Heavy-Flavor Particle Properties	476
3.3. Mixing Phenomena	477
3.4. CP-Violation Studies	478
4. ELECTROWEAK PHYSICS	479
4.1. Vector Boson Production	480
4.2. Asymmetry Measurements	481
4.3. Diboson Production and Trilinear Couplings	482
4.4. W Boson Mass and Width	483
5. TOP QUARK STUDIES	485
5.1. Top Quark Discovery	485
5.2. Single Top Quark Production	486
5.3. Top Quark Properties	488
5.4. Top Quark Production	488
6. HIGGS BOSON SEARCH	490
6.1. $H \rightarrow WW$	490
6.2. $H \rightarrow b\bar{b}$	491
6.3. Searches for the Higgs Boson in Supersymmetry and Other Models	492
7. SEARCHES FOR NEW PHENOMENA	493
7.1. Searches for Supersymmetric Partners of Standard Model Particles	494
7.2. Searches for New Gauge Bosons	495
7.3. Searches for Leptoquarks	495
7.4. Searches for Extra Dimensions of Space	497
7.5. Searches for Other New Particles	497
7.6. Signature-Based Searches	497
8. A FINAL NOTE	498

1. INTRODUCTION

The Tevatron was conceived in the early 1970s to double the beam energy of the original Fermilab Main Ring accelerator to approximately 1 TeV through the first use of superconducting magnets in an accelerator (1). Beams were commissioned in 1983. The proposal for a $p\bar{p}$ collider in 1976 (2) provided antiprotons from interactions between Main Ring protons in a heavy-metal target and antiproton accumulation rings employing stochastic cooling that could supply nearly 10^{11} antiprotons per hour. The antiprotons would be accelerated and stored in the same Tevatron magnet ring as the protons and brought into collision at up to six interaction regions. The first

Tevatron Collider Highlights beyond the Top Quark

CP violation in B mesons

Mesons with beauty and charm

Heavy-baryon spectroscopy

Higgs-boson searches

Hadron collider makes precision measurements!

Jets; W, Z , properties; B_s – \bar{B}_s oscillation frequency

Global experimental collaborations



The Tevatron (1983–) gave us courage to think bigger

Thought experiment (1977) considering WW scattering had revealed

$$\left(\frac{8\pi\sqrt{2}}{3G_F} \right)^{1/2} \approx 1 \text{ TeV}$$

energy scale to discover the mechanism
of electroweak symmetry breaking

Large Hadron Collider
 $pp, E_{cm} = 1 \text{ TeV} = 13.6 \text{ TeV}$


CMS

LHCb

ALICE


ATLAS


CERN







CMS










$H(125 \text{ GeV}) \rightarrow ZZ^* \rightarrow \mu^+ \mu^- \mu^+ \mu^-$ candidate



Run: 189280

Event: 143576946

2011-09-14 12:37:11 CEST



July 4, 2012

Higgs discovery follow-ups

Refine Higgs-boson properties: (HL)-LHC, “Higgs Factory”
 $e^+e^- \rightarrow H + Z$

Does H have partners?

How does H interact with itself?



Does H regulate WW scattering?

Vacuum-energy problem

→ Perspectives & (120+) Questions


→ Gedanken Worlds without Higgs

“sixth place of the decimal” and beyond ...



e magnetic dipole moment measured to 0.13 ppt

$$-\mu_e/\mu_B = 1.00115965218059(13)$$



Harvard-NU

(Anti)proton magnetic moments: CPT test

$$\mu_{\bar{p}} = -2.792\,847\,344\,1(42) \mu_N$$

vs.

$$\mu_p = +2.792\,847\,344\,62(82) \mu_N$$

BASE Collaboration @CERN Antiproton Decelerator



fine structure constant

Berkeley: $\alpha^{-1}(Cs) = 137.035\,999\,046(27)$

Paris: $\alpha^{-1}(Rb) = 137.035\,999\,206(11)$

(differ by 5.4 s.d.)

High-luminosity e^+e^- , up to $t\bar{t}$ threshold



A 10-TeV $\mu^+\mu^-$ Collider for the Nuclear Mill Centennial?

