

# Black hole entropy for the general area spectrum

Tomo Tanaka<sup>1</sup> and Takashi Tamaki<sup>2</sup>

<sup>1</sup>*Department of Physics, Waseda University, Okubo 3-4-1, Tokyo 169-8555, Japan*

<sup>2</sup>*Department of Physics, Waseda University, Okubo 3-4-1, Tokyo 169-8555, Japan and Department of Physics, Rikkyo University, Toshima, Tokyo 171-8501, Japan*

## Abstract

We consider the possibility that the horizon area is expressed by the general area spectrum in loop quantum gravity and calculate the black hole entropy by counting the degrees of freedom in spin-network states related to its area. Although the general area spectrum has a complex expression, we succeeded in obtaining the result that the black hole entropy is proportional to its area as in previous works where the simplified area formula has been used. This gives new values for the Barbero-Immirzi parameter ( $\gamma = 0.5802 \dots$  or  $0.7847 \dots$ ) which are larger than that of previous works.

## 1 Introduction

Statistical mechanics in a self-gravitating system is quite different from that without gravity. For example, particles in the box have maximal entropy when they spread out uniformly in the box if gravity is not taken into account. On the other hand, if particles are self-gravitating, we can suppose that clusters appear as an entropically favorable state. Then, if the pressure of particles can be neglected, it is likely that a black hole appears as a maximal entropy state. Thus, black hole entropy would be the key for understanding statistics in a self-gravitating system.

One of the most mysterious things about black holes is their entropy  $S$  which is *not* proportional to its volume *but* to its horizon area  $A$ . This was first pointed out related to the first law of black hole thermodynamics. The famous relation  $S = A/4$  has been established by the discovery of the Hawking radiation. Recently, its statistical origin has been discussed in the candidate theories of quantum gravity, such as string theory, or loop quantum gravity (LQG) [1], etc. It has been discussed that LQG can describe its statistical origin independent of black hole species because of its background independent formulation [2]. For this reason, we concentrate on LQG here.

Quantum states in LQG are described by spin-network [3], and basic ingredients of the spin-network are edges, which are lines labeled by spin  $j$  ( $j = 0, 1/2, 1, 3/2, \dots$ ) reflecting the  $SU(2)$  nature of the gauge group, and vertices which are intersections between edges. For three edges having spin  $j_1, j_2$ , and  $j_3$  that merge at an arbitrary vertex, we have following conditions.

$$j_1 + j_2 + j_3 \in \mathbb{N}, \quad (1)$$

$$j_i \leq j_j + j_k, \quad (i, j, k \text{ different from each other}). \quad (2)$$

These conditions guarantee the gauge invariance of the spin-network.

Using this formalism, general expressions for the spectrum of the area and the volume operators can be derived [4, 5]. For example, the area spectrum  $A_j$  is

$$A_j = 4\pi\gamma \sum_i \sqrt{2j_i^u(j_i^u + 1) + 2j_i^d(j_i^d + 1) - j_i^t(j_i^t + 1)}, \quad (3)$$

where  $\gamma$  is the Barbero-Immirzi parameter related to an ambiguity in the choice of canonically conjugate variables. The sum is added up all intersections between a surface and edges. Here, the indices  $u$ ,  $d$ , and  $t$  mean edges upper side, down side, and tangential to the surface, respectively (We can determine which side is upper or down side arbitrarily).

---

<sup>1</sup>E-mail:tomo@gravity.phys.waseda.ac.jp

<sup>2</sup>E-mail:tamaki@gravity.phys.waseda.ac.jp

In [1], it was proposed that black hole entropy is obtained by counting the number of degrees of freedom about  $j$  when we fix the horizon area where a simplified area formula is used. This simplified area formula is obtained by assuming that there are no tangential edges on black hole horizon, that is  $j_i^t = 0$ . We obtain  $j_i^u = j_i^d := j_i$  by using the condition (2). Then, we consider the degrees of freedom about  $j$  satisfying

$$A_j = 8\pi\gamma \sum_i \sqrt{j_i(j_i + 1)} = A. \quad (4)$$

The standard procedure is to impose the Bekenstein-Hawking entropy-area law  $S = A/4$  for large black holes in order to fix the value of  $\gamma$ . Ashtekar et al. in [2] extended this idea using the isolated horizon framework (ABCK framework) [6]. Error in counting in this original work has been corrected in [7, 8]. Similar works appear related to how to count the number of freedom in [9, 10].

However, should we restrict to the simplified area spectrum (4) ? Thiemann in [11] used the boundary condition that there is no other side of the horizon, i.e.,  $j_i^d = 0$ . Then, by using (2), we obtain  $j_i^u = j_i^t := j_i$  which gives  $A_j = 4\pi\gamma \sum_i \sqrt{j_i(j_i + 1)}$ . Based on this proposal, the number counting has been performed in [12] which gives  $\gamma = 0.323 \dots$ .

Another interesting possibility is to use (3) motivated by the hypothesis that a black hole is a maximal entropy state in a self-gravitating system. If we agree that the origin of the black hole entropy is related to degrees of freedom in  $j$  (or  $m = -j, -j + 1, \dots, j$  considered in [2]), it is evident that (3) can gain larger number of states than (4) for the fixed area. See, also [13] which also discuss using (3) as expressing the horizon area. Of course, it is speculative and the typical objection to the idea is that since the black hole evaporates, it is not the maximal entropy state. However, the black hole we consider is the limit  $A \rightarrow \infty$  where the evaporation process can be negligible. The second objection is that if we require the entropy-area law  $S = A/4$ , the black hole entropy does not depend on what types of area formula we use, so it is not relevant to the above hypothesis. This is a delicate question to be answered carefully. From the view point that the Barbero-Immirzi parameter is determined *a priori*, the formula  $S = A/4$  only provides us the method to *know* the value of  $\gamma$ . If this is the case, using (3) would enhance the entropy. Therefore, we concentrate on evaluating the number of states using (3) by adopting this view point. To answer whether this view point is true or not, we need independent discussion to *know* the value of  $\gamma$  through, e.g., cosmology [14] or quasinormal modes of black holes [15, 16, 17].

Our strategy is as follows [18]. Based on the observation that the value of  $\gamma$  in [2] is qualitatively same as that inferred in [1] which counts the degrees of freedom of  $j$  without imposing the horizon conditions for the case (4), we restrict counting the corresponding  $j$  freedom for (3) as a first step. We can perform it by carefully reanalyzing the case (4). This paper is organized as follows. In section 2, we count the degrees of freedom for (3). In section 3, we mention concluding remarks.

## 2 Consideration of the general area spectrum

In the case for (4), horizon conditions does not affect entropy formula. Therefore in the case for (3) we consider only degrees of freedom about area. In this case, we also denote number of states as  $N(A)$  which is defined as

$$N(A) := \left\{ (j_1^u, j_1^d, j_1^t, \dots, j_n^u, j_n^d, j_n^t) \mid 0 \neq j_i^u, j_i^d \in \frac{\mathbb{N}}{2}, \quad 0 \neq j_i^t \in \mathbb{N}, \quad j_i^u, j_i^d, j_i^t \text{ should satisfy (1) and (2)} \right. \\ \left. \sum_i \sqrt{2j_i^u(j_i^u + 1) + 2j_i^d(j_i^d + 1) - j_i^t(j_i^t + 1)} = \frac{A}{4\pi\gamma} \right\}. \quad (5)$$

We adopt the condition  $j^t \in \mathbb{N}$  motivated by the ABCK framework where the “classical horizon” is described by  $U(1)$  connection. This is, of course, not verified in the present situation and should be reconsidered in future.

Then, we perform counting as follows. If we use the condition  $j^t \in \mathbb{N}$ , we have  $j^u + j^d := n \in \mathbb{N}$  by (1). If we fix  $n$ , we can classify the possible  $j^u, j^d, j^t$  as follows, which is one of the most important parts in this paper. First, we have  $(j^u, j^d) = (\frac{n}{2} \pm \frac{s}{2}, \frac{n}{2} \mp \frac{s}{2})$  (double-sign corresponds) for  $0 \leq s \leq n, s \in \mathbb{N}$  to

satisfy (2). Then, for each  $s$ , possible value of  $j^t$  is  $j^t = s, s+1, \dots, n$  to satisfy (2). For any eigenvalue  $x := 4\pi\gamma\sqrt{2j_x^u(j_x^u+1) + 2j_x^d(j_x^d+1) - j_x^t(j_x^t+1)}$  ( $0 < x \leq A$ ) of the area operator, we have

$$(\mathbf{j}_1, \dots, \mathbf{j}_n) \in N(A-x) \Rightarrow (\mathbf{j}_1, \dots, \mathbf{j}_n, \mathbf{j}_x) \in N(A), \quad (6)$$

where we used the abbreviation as  $\mathbf{j}_i = (j_i^u, j_i^d, j_i^t)$ . We have  $(\mathbf{j}_1, \dots, \mathbf{j}_n, \mathbf{j}_x) \neq (\mathbf{j}_1, \dots, \mathbf{j}_n, \mathbf{j}_{x'})$ , if  $\mathbf{j}_x \neq \mathbf{j}_{x'}$ . Therefore, as for the case in (4), if we consider all  $0 < x \leq A$  and  $(\mathbf{j}_1, \dots, \mathbf{j}_n) \in N(A-x)$ ,  $(\mathbf{j}_1, \dots, \mathbf{j}_n, \mathbf{j}_x)$  form the entire set  $N(A)$ .

Then, if we use the notation  $j^u = \frac{n}{2} + \frac{s}{2}, j^d = \frac{n}{2} - \frac{s}{2}, j^t = t$ , we have  $x(n, s, t) = 4\pi\gamma\sqrt{n^2 + 2n + s^2 - t(t+1)}$  and

$$N(A) = \sum_{n=1}^{\infty} \left[ \sum_{s=1}^n \sum_{t=s}^n 2N(A-x(n, s, t)) + \sum_{t=0}^n N(A-x(n, s=0, t)) \right], \quad (7)$$

where the factor 2 in front of  $N(A-x(n, s, t))$  for  $s \neq 0$  corresponds to the fact that same  $x(n, s, t)$  appears twice for the exchange of  $j^u$  and  $j^d$ . For  $A \rightarrow \infty$ , by assuming  $N(A) = Ce^{\frac{A\gamma_M}{4\gamma}}$ , where  $C$  is a constant and substituting to the recursion relation (7), we obtain the beautiful formula as a generalization of the case (4) as,

$$1 = \sum_{n=1}^{\infty} \left[ \sum_{s=1}^n \sum_{t=s}^n 2 \exp(-\gamma_M x(n, s, t)/4\gamma) + \sum_{t=0}^n \exp(-\gamma_M x(n, s=0, t)/4\gamma) \right]. \quad (8)$$

If we require  $S = A/4$ , we have  $\gamma = \gamma_M = 0.5802 \dots$ . This means that even if we use (3) as the horizon spectrum, we can reproduce the entropy formula  $S = A/4$  by adjusting the Barbero-Immirzi parameter. This is nontrivial and is our main result in this paper.

Let us turn back to our assumptions. Although we obtained  $\gamma$  satisfying  $S = A/4$  for the case (3), there may be a criticism that the result is underestimated by only counting  $j$  freedom. To answer it, we consider the following counting. When the simplified area formula was used, there is a proposal that we should count not only  $j$  but also  $m = -j, -j+1, \dots, j$  freedom based on the ABCK framework [9]. Although it is nontrivial whether this framework can be extended to the general area formula, let us count also the  $m$  freedom for each  $j^u$  to maximize the estimate. Counting only  $m$  related to  $j^u$  is reasonable from the point of view of the entanglement entropy or the holography principle.

If we notice that there are  $(n+s+1)$  and  $(n-s+1)$  freedoms for  $m$  (total  $2(n+1)$ ) corresponding to  $(j^u, j^d) = (\frac{n}{2} + \frac{s}{2}, \frac{n}{2} - \frac{s}{2})$  and  $(\frac{n}{2} - \frac{s}{2}, \frac{n}{2} + \frac{s}{2})$ , respectively, the factor 2 in the first term of the right-hand side of (8) is replaced by  $2(n+1)$  in this case. For  $s=0$ , the factor 1 in the second term is replaced by  $(n+1)$ . Then, we obtain

$$1 = \sum_{n=1}^{\infty} \left[ \sum_{s=1}^n \sum_{t=s}^n 2(n+1) \exp(-\gamma_M x(n, s, t)/4\gamma) + \sum_{t=0}^n (n+1) \exp(-\gamma_M x(n, s=0, t)/4\gamma) \right], \quad (9)$$

which gives  $\gamma = \gamma_M = 0.7847 \dots$ . Thus, we confirm that the black hole entropy is proportional to the area again. Naively speaking, we expect that there is no qualitative deviation from these two values of  $\gamma$  even if we take into account the ABCK framework for (3) appropriately.

### 3 Conclusion and Discussion

In this paper, we obtained the black hole entropy by considering the general area formula. It is surprising that we succeeded in obtaining the black hole entropy proportional to the horizon area even in this case. Then, it is natural to ask what the area formula should be in describing the horizon area. There are many possibilities examining the area spectrum. For example, we have not yet established the black hole thermodynamics in LQG which is one of the most important topics to be investigated. There is an idea that black hole evaporation process should also be described by using the general area formula [13]. Therefore, whether we can establish the generalized second law of black hole thermodynamics might be one of the criteria in judging which area formula is appropriate. For this purpose, it is desirable to extend

the ABCK framework for the general area formula since the exact counting is required. Though we do not take care of the topology of the horizon, discussing the difference caused by the topology is important as considered for the simplified area formula [19].

Of course, as we mentioned in the introduction, we should check the value of the Barbero-Immirzi parameter in several independent discussions. Therefore, we should also take care of cosmology [14] and quasinormal modes of black holes [15, 16, 17] in determining the Barbero-Immirzi parameter. Confirming LQG in many independent methods would be the holy grail of the theory.

## Acknowledgements

We would like to thank Tomohiro Harada and Kei-ichi Maeda for useful comments and continuous encouragement.

## References

- [1] C. Rovelli, Phys. Rev. Lett. **77**, 3288 (1996).
- [2] A. Ashtekar, J. Baez, A. Corichi, and K. Krasnov, Phys. Rev. Lett. **80**, 904 (1998); A. Ashtekar, J. Baez, and K. Krasnov, Adv. Theor. Math. Phys. **4**, 1 (2000).
- [3] C. Rovelli and L. Smolin, Phys. Rev. D **52**, 5743 (1995).
- [4] C. Rovelli and L. Smolin, Nucl. Phys. B **442**, 593 (1995); Erratum, *ibid.*, **456**, 753 (1995).
- [5] A. Ashtekar and J. Lewandowski, Class. Quantum Grav. **14**, A55 (1997).
- [6] A. Ashtekar, A. Corichi, and K. Krasnov, Adv. Theor. Math. Phys. **3**, 419 (1999).
- [7] M. Domagala and J. Lewandowski, Class. Quant. Grav. **21**, 5233 (2004).
- [8] K. A. Meissner, Class. Quant. Grav. **21**, 5245 (2004).
- [9] A. Ghosh and P. Mitra, Phys. Lett. B **616**, 114 (2005); Int. J. Mod. Phys. **80**, 867 (2006); Phys. Rev. D **74**, 064026 (2006).
- [10] T. Tamaki and H. Nomura, Phys. Rev. D **72**, 107501 (2005).
- [11] T. Thiemann, gr-qc/0110034, Modern Canonical Quantum General Relativity (Cambridge University Press, Cambridge, 2007).
- [12] T. Tamaki, Class. Quant. Grav. **24**, 3837 (2007).
- [13] M. H. Ansari, Nucl. Phys. B **783**, 179 (2007); *ibid.*, **795**, 635 (2008).
- [14] M. Bojowald, Living. Rev. Rel. **8**, 11 (2005); A. Ashtekar, T. Pawłowski, P. Singh, Phys. Rev. D **74**, 084003, (2006).
- [15] For review, see, e.g., J. Natario and R. Schiappa, hep-th/0411267.
- [16] O. Dreyer, Phys. Rev. Lett. **90**, 081301 (2003).
- [17] S. Hod, Phys. Rev. Lett. **81**, 4293 (1998).
- [18] T. Tanaka and T. Tamaki, arXiv:0808.4056.
- [19] S. Kloster, J. Brannlund, and A. DeBenedictis, Class. Quant. Grav. **25**, 065008 (2008).