


Superconducting Super Collider Laboratory



Quantum Variances for Transverse SSC Injection Dynamics

T. Garavaglia

May 1991

Quantum Variances for Transverse SSC Injection Dynamics*

T. Garavaglia

Superconducting Super Collider Laboratory[†]
2550 Beckleymeade Avenue
Dallas, TX 75237

May 1991

* Presented at the 1991 IEEE Particle Accelerator Conference, San Francisco, CA, May 6-9, 1991.

[†] Operated by the Universities Research Association, Inc., for the U.S. Department of Energy under Contract No. DE-AC02-89ER40486.

QUANTUM VARIANCES FOR TRANSVERSE SSC INJECTION DYNAMICS

T. Garavaglia
Superconducting Super Collider Laboratory*
2550 Beckleymeade Avenue
Dallas, TX 75237

Abstract

Quantum variances for the transverse dynamics associated with the injection phase of the Superconducting Super Collider are derived using squeezed state methods.

1 Introduction

During the injection phase into the SSC particles near the edge of the dynamic aperture perform chaotic motion [1]. With an injection amplitude difference as small as 10^{-10} mm chaotic divergences for the x and y transverse betatron motions begin to occur as soon as 15×10^3 turns around the SSC. It is of interest to see what quantum uncertainties are associated with transverse amplitudes. These limits are indeed significant for the SSC, and they are in fact larger than the small differences which lead to chaotic dynamics.

2 Betatron Equations

The equations of motion for transverse betatron oscillations can be found from a Lorentz invariant Hamiltonian [2]. This Hamiltonian which is equal to the rest energy of the proton is invariant for curvilinear coordinate transformations $q^i \rightarrow \bar{q}^i$. This allows one to represent equations for betatron motion in the coordinate system related to ideal orbit of a proton and to see the various approximation made to obtain the usual Courant-Snyder equations. For this invariant Hamiltonian, contravariant coordinates are defined as the four-vector $q^i = (q^0, q^1, q^2, q^3) = (ct, x, y, z) \rightarrow (ct, x, y, \bar{s})$ where c is the speed of light and where $\bar{s}$ is the arc length along the ideal orbit. The contravariant components of the four-velocity and four-momentum are

$$u^i = dq^i/ds \quad \text{and} \quad p^i = mc u^i \quad (1)$$

where $ds = \sqrt{dq_i dq^i}$. The invariant Hamiltonian is defined as

$$H(p^i, q^i)/c = p_i u^i - L(u^i, q^i) + mc \quad (2)$$

where the invariant Lagrangian for the motion of a particle of mass m and electric charge e in an electromagnetic field

with four-potential $A(q)^i$ is

$$L(u^i, q^i) = mc\sqrt{u_i u^i} + (e/c)A(q)_i u^i. \quad (3)$$

In terms of the conjugate four-momentum, the Lagrangian becomes $L(u^i, q^i) = p_i u^i$, and the invariant action is

$$S = \int p_i u^i ds = \int \bar{p}_i \bar{u}^i d\bar{s}. \quad (4)$$

Treating $\bar{s}$ as the time coordinate, $\bar{p}_i$ as the Hamiltonian, and using the method of stationary action, one finds equations which describe transverse betatron motion.

3 Quantum Limits on Injection

As a first approximation for finding the quantum limits, one considers a time-dependent linear oscillator Hamiltonian for an on momentum particle

$$H(\bar{s}) = \frac{p^2 + K(\bar{s})q^2}{2}. \quad (5)$$

With the conditions

$$\begin{aligned} w'' + K(\bar{s})w - \frac{1}{w^3} &= 0 \\ q'' + K(\bar{s})q &= 0 \end{aligned} \quad (6)$$

and with $q = we^{\pm i\psi}$ and $\psi' = \frac{1}{w^2}$, one finds the invariant

$$2I = \gamma(\bar{s})q^2 + \alpha(\bar{s})(q p + p q) + \beta(\bar{s})p^2 \quad (7)$$

with Courant-Snyder parameters

$$\begin{aligned} \alpha(\bar{s}) &= -ww', \quad \beta(\bar{s}) = w^2, \quad \text{and} \\ \gamma(\bar{s}) &= \frac{1 + \alpha^2(\bar{s})}{\beta(\bar{s})}. \end{aligned} \quad (8)$$

The quantum states for this system can be constructed with the aid of the squeezing operator defined as

$$S = \exp((\xi^* a^2 - \xi a^{\dagger 2})/2). \quad (9)$$

The time-independent rationalized Hamiltonian, H_o , is found from (5) with $K(\bar{s}) = 1$ where the boson operators are found from $\sqrt{2}q = a + a^\dagger$ and $\sqrt{2}ip = a - a^\dagger$. The invariant (7) can be found from H_o as

$$2I(\bar{s}) = S H_o S^\dagger = \left(b^\dagger b + \frac{1}{2} \right) \quad (10)$$

*Operated by the Universities Research Association, Inc. for the U. S. Department of Energy under Contract No. DE-AC02-89ER40486.

where

$$\begin{aligned} \mathbf{b}(\bar{s}) &= \mathbf{S} e^{i\theta} \mathbf{a} \mathbf{S}^\dagger = \frac{1}{2} \left(\frac{1}{w} + w - iw' \right) \mathbf{a} \\ &+ \frac{1}{2} \left(\frac{1}{w} - w - iw' \right) \mathbf{a}^\dagger \end{aligned} \quad (11)$$

with $\theta(w, w')$. The eigenstates of $I(\bar{s})$ satisfy the eigenvalue equation

$$\begin{aligned} \mathbf{I}(\bar{s})|n, \bar{s}\rangle &= \left(n + \frac{1}{2} \right) |n, \bar{s}\rangle \\ |n, \bar{s}\rangle &= \frac{(\mathbf{b}^\dagger)^n}{\sqrt{n!}} |0\rangle, \end{aligned} \quad (12)$$

and the units of $I(\bar{s})$ are $(\hbar/|\bar{p}|) \text{rad}$. The Schrödinger states [3] are

$$|n, \bar{s}\rangle_s = e^{i\alpha_n(\bar{s})} |n, \bar{s}\rangle \quad (13)$$

where the phase is

$$\alpha_n(\bar{s}) = - \left(n + \frac{1}{2} \right) \int^{\bar{s}} \frac{d\bar{s}'}{w^2(\bar{s}')} \quad (14)$$

The coherent state for a time-dependent linear oscillator used to evaluate variances is generated from the squeezed ground state as

$$|\beta, \bar{s}\rangle_s = e^{\beta \mathbf{b}^\dagger(\bar{s}) - \beta^* \mathbf{b}(\bar{s})} |0, \bar{s}\rangle_s \quad (15)$$

where β , related to the classical value of the invariant $I(\bar{s})$, is the eigenvalue of the operator $\mathbf{b}(\bar{s})$.

One can now use the states (15) to obtain results appropriate for the SSC. Using appropriate scaling transformations

$$P \rightarrow \frac{p}{|\bar{p}|}, \quad H \rightarrow \frac{H}{|\bar{p}|}, \quad \text{and } \hbar \rightarrow \frac{\hbar}{|\bar{p}|} \quad (16)$$

with the momentum $|\bar{p}| \approx \mathcal{E}/c$ for a proton of energy $\mathcal{E}$, one finds the variances and the uncertainty product represented in terms of the Courant-Snyder parameters (8) are

$$\begin{aligned} \sigma(q) &= \sqrt{\frac{\hbar \beta(\bar{s})}{2|\bar{p}|}}, \quad \sigma\left(\frac{p}{\bar{p}}\right) = \sqrt{\frac{\hbar \gamma(\bar{s})}{2|\bar{p}|}}, \\ \frac{p}{|\bar{p}|} &= \frac{dq}{d\bar{s}}, \quad \text{and } \sigma(q)\sigma\left(\frac{dq}{d\bar{s}}\right) = \frac{\hbar \sqrt{\beta(\bar{s})\gamma(\bar{s})}}{2|\bar{p}|}. \end{aligned} \quad (17)$$

Writing the amplitude as $q_{amp} = \sqrt{\epsilon/\pi\beta(\bar{s})}$ with the emittance $\epsilon = 2\pi I(\bar{s})$, one finds the results

$$\begin{aligned} \frac{\sigma(q_{amp})}{q_{amp}} &= \left(\frac{\epsilon_q}{\epsilon} \right)^{1/2} \\ \epsilon_q/\pi &= \frac{\hbar}{2|\bar{p}|} \approx \frac{\hbar c}{2\mathcal{E}} \approx \frac{\lambda_{proton}}{2} \end{aligned} \quad (18)$$

where ϵ_q/π , the quantum emittance, represents half the resolution length of a proton in the beam. With the approximations $\hbar c \approx 2 \times 10^{-19} \text{TeV} \cdot \text{m}$, and $\mathcal{E} \approx 2 \text{TeV}$, one finds

$$\epsilon_q/\pi \approx 5 \times 10^{-20} \text{m}. \quad (19)$$

For a typical SSCTRK tracking result showing chaotic motion with $\beta(\bar{s}) \approx 300 \text{m}$ and with $x_{amp} \approx 3.5 \text{mm}$, one finds

$$\epsilon/\pi \approx 4 \times 10^{-8} \text{m}$$

and

$$\sigma(x_{amp}) \approx 3.9 \times 10^{-6} \text{mm}. \quad (20)$$

Similarly, the angular uncertainty is

$$\sigma\left(\frac{dx}{d\bar{s}}\right) \approx 1.3 \times 10^{-11} \text{rad}. \quad (21)$$

V. Conclusions

The quantized invariant of the linearized betatron equations which are obtained from an invariant action leads to the time-dependent coherent state used to evaluate the uncertainties of the position and momentum operators. The maximum value of $\beta(\bar{s})$ in the SSC arcs during injection is 300m , the value used to obtain the variances (20) and (21). However, larger variances can occur during the injection phase in other regions where $\beta(\bar{s}) > 1800 \text{m}$. It is of interest to note that the emittance defined as $\epsilon = 2\pi I(\bar{s})$ which is the area of the (q, q') phase space ellipse is quantized and that the variance is

$$\sigma(\epsilon) = \pi |\beta| (\hbar/|\bar{p}|) \text{rad}.$$

A more complete account may be found in [4].

References

- [1] T. Garavaglia, S. K. Kauffmann, R. Stiening, and D. Ritson, "Numerical Studies of the SSC Injection Process: Long Term Tracking," in *Supercollider 2.*, edited by M. McAshan, New York: Plenum Press, 1990, pp. 39-58.
- [2] C. Garrod, "Covariant Hamiltonian Dynamics with Interactions," *Phys. Rev.* **167**, pp. 1143-1145, March 1968.
- [3] H.R. Lewis, Jr., and W.B. Riesenfeld, "An Exact Quantum Theory of the Time-Dependent Harmonic Oscillator," *J. Math. Phys.* **10**, pp. 1458-1473, August 1969.
- [4] T. Garavaglia, "Quantum Variances for Transverse Betatron Dynamics," SSCL Report, in press, 1991.

Disclaimer Notice

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government or any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Superconducting Super Collider Laboratory is an equal opportunity employer.