

PARTICLES AND FIELDS • OPEN ACCESS

## On the nature of $\chi_{c2}(2P)$ : two-gluon decay

To cite this article: N.N. Achasov and Xian-Wei Kang 2017 *Chinese Phys. C* **41** 123102

View the [article online](#) for updates and enhancements.

### Related content

- [Analysis of Vector Meson Transitions Among Heavy Quarkonium States](#)  
Wang Zhi-Gang

- [Heavy quarkonia spectra and their decays in a relativistic quark model](#)  
Bhagya K. B. Vijaya Kumar and Antony Prakash Monteiro

- [Some topics on charmonium decays at BESIII experiment](#)  
Chen Hai-Xuan

# On the nature of $\chi_{c2}(2P)$ : two-gluon decay<sup>\*</sup>

N.N. Achasov<sup>2;1)</sup> Xian-Wei Kang(康现伟)<sup>1,3;2)</sup>

<sup>1</sup> College of Nuclear Science and Technology, Beijing Normal University, Beijing 100875

<sup>2</sup> Laboratory of Theoretical Physics, Sobolev Institute for Mathematics, 630090, Novosibirsk

<sup>3</sup> Institute of Physics, Academia Sinica, 115 Taipei

**Abstract:** We expect that  $BR(\chi_{c2}(2P) \rightarrow \text{gluon gluon}) \gtrsim 2\%$  if the Particle Data Group as well as the BaBar and Belle collaborations have correctly identified the state. In reality, this branching ratio corresponds to the one for  $\chi_{c2}(2P)$  decaying into light hadrons. We also discuss the detection possibilities of these decays.

**Keywords:** charmonium, branching ratio, two-gluon decay mode

**PACS:** 13.25.Gv, 14.40.Pq      **DOI:** 10.1088/1674-1137/41/12/123102

## 1 Introduction

More and more  $XYZ$  states have been observed in experiments, and their interpretation is still challenging to the community. Some of these states, however, may just be quarkonium states.

In 2006, the Belle collaboration reported a resonance with  $5.3\sigma$  statistical significance of the signal via the  $\gamma\gamma \rightarrow D\bar{D}$  process [1]. The properties of the mass, angular distributions, and  $\Gamma_{\gamma\gamma}\Gamma_{D\bar{D}}/\Gamma$  (see also Eq. (6) below) are all consistent with the  $2^3P_2$  charmonium state, now identified by the Particle Data Group (PDG) as  $\chi_{c2}(2P)$  [2]. Later, the BaBar collaboration [3] confirmed this observation using data samples of comparable magnitude with those of Belle. Its mass and width are  $M = 3927.2 \pm 2.6$  MeV and  $\Gamma = 24 \pm 6$  MeV, respectively [2]. To date, only the quantity  $\Gamma_{\gamma\gamma}\Gamma_{D\bar{D}}/\Gamma$  has been determined, except for the constrained upper limit for the product of the branching ratio for  $\gamma\gamma$  and some selected hadronic states [4–6]. Little is known beyond them, especially for the branching ratio of hadronic decays. In this article,

we will predict the branching ratio of the decay  $\chi_{c2} \rightarrow 2g$  ( $g$  denoting gluon) in a reliable way, by combining the observation of the known experimental facts and the successful application of the charmonium model [7]. In practice, that branching ratio corresponds to summing over those for light hadronic decay.

## 2 Discussion on branching ratios

It is natural that in the nonrelativistic potential model of charmonium, the ratio of the two-photon and two-gluon widths of the charmonium decays does not depend on the wave function and slowly grows with increasing charmonium mass, because of the proportionality to  $1/\alpha_s^2$ . See, for example, Ref. [7]. The well established states [2] confirm this consideration:

$$\begin{aligned} BR(\eta_c(1S) \rightarrow 2\gamma) &= \frac{\Gamma(\eta_c(1S) \rightarrow 2\gamma)}{\Gamma(\eta_c(1S))} \\ &= \frac{\Gamma(\eta_c(1S) \rightarrow 2\gamma)}{\Gamma(\eta_c(1S) \rightarrow 2g)} \approx 1.59 \times 10^{-4}, \end{aligned} \quad (1)$$

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow 2\gamma)}{\Gamma(\chi_{c0}(1P) \rightarrow 2g)} = \frac{\Gamma(\chi_{c0}(1P) \rightarrow 2\gamma)}{\Gamma(\chi_{c0}(1P)) - \Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))} = \frac{BR(\chi_{c0}(1P) \rightarrow 2\gamma)}{1 - BR(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))} \approx 2.26 \times 10^{-4}, \quad (2)$$

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow 2\gamma)}{\Gamma(\chi_{c2}(1P) \rightarrow 2g)} = \frac{\Gamma(\chi_{c2}(1P) \rightarrow 2\gamma)}{\Gamma(\chi_{c2}(1P)) - \Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))} = \frac{BR(\chi_{c2}(1P) \rightarrow 2\gamma)}{1 - BR(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))} \approx 3.39 \times 10^{-4}, \quad (3)$$

Received 10 August 2017

\* N.N. Achasov was supported in part by RFBR, (16-02-00065), and by Presidium of the Russian Academy of Sciences, Project (0314-2015-0011), XWK's work is supported by MOST, Taiwan, (104-2112-M-001-022)

1) E-mail: achasov@math.nsc.ru

2) E-mail: kangxianwei1@gmail.com

 Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP<sup>3</sup> and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

where we have used [2]

$$\begin{aligned} BR(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) &= (1.27 \pm 0.06)\%, \\ BR(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) &= (19.2 \pm 0.7)\%. \end{aligned} \quad (4)$$

According to QCD, the decay of charmonium is due to the annihilation of the  $c\bar{c}$  pair. The mass of  $c\bar{c}$  is large and  $c\bar{c} \rightarrow$  gluons are perturbative, so two-gluon decay mode is dominant. In the above equations, we did not use  $\eta_c(2S)$  as an argument. Its hadronic decay channels are not well determined yet, and also the only measured radiative channel,  $\eta_c(2S) \rightarrow \gamma\gamma$ , suffers from very large uncertainty.

We know that

$$\begin{aligned} \Gamma(\chi_{c2}(2P) \rightarrow \gamma\gamma) BR(\chi_{c2}(2P) \rightarrow D\bar{D}) \\ = (0.24 \pm 0.05 \pm 0.04) \text{ keV} \quad [3] \\ \Gamma(\chi_{c2}(2P) \rightarrow \gamma\gamma) BR(\chi_{c2}(2P) \rightarrow D\bar{D}) \\ = (0.18 \pm 0.05 \pm 0.03) \text{ keV} \quad [1]. \end{aligned} \quad (5)$$

The PDG average gives  $0.21 \pm 0.04$  keV [2].

Taking into account  $\Gamma(\chi_{c2}(2P)) \approx 24$  MeV [2], we find

$$\begin{aligned} BR(\chi_{c2}(2P) \rightarrow 2\gamma) BR(\chi_{c2}(2P) \rightarrow D\bar{D}) &\approx 10^{-5} \text{ or} \\ BR(\chi_{c2}(2P) \rightarrow 2\gamma) BR(\chi_{c2}(2P) \rightarrow D\bar{D}) &\approx 0.75 \times 10^{-5}. \end{aligned} \quad (6)$$

Conservatively selecting from Eq. (3) the ratio of the two-gluon to two-photon widths of the charmonium decays to be around  $(1/4) \times 10^4$ , we obtain

$$\begin{aligned} BR(\chi_{c2}(2P) \rightarrow 2g) BR(\chi_{c2}(2P) \rightarrow D\bar{D}) &\approx 0.025 \text{ or} \\ BR(\chi_{c2}(2P) \rightarrow 2g) BR(\chi_{c2}(2P) \rightarrow D\bar{D}) &\approx 0.019. \end{aligned} \quad (7)$$

So we expect  $BR(\chi_{c2}(2P) \rightarrow 2g) \gtrsim (2 \pm 0.4)\%$  if the PDG has correctly identified the state.

The hadron channels of the two-gluon decays of  $\chi_{c2}(2P)$  could be the same as in the  $\chi_{c2}(1P)$  case, that is, there are a few tens of such channels. It is expected that the difference in the radial wave functions of  $\chi_{c2}(1P)$  and  $\chi_{c2}(2P)$  does not lead to a significant difference in  $\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma)$  and  $\Gamma(\chi_{c2}(2P) \rightarrow \gamma\gamma)$ . Indeed,  $\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma) \approx 0.5$  keV [2] and  $\Gamma(\chi_{c2}(2P) \rightarrow \gamma\gamma) \gtrsim 0.24$  keV or  $0.18$  keV, cf. Eq. (5). That is to say, it is possible that  $\Gamma(\chi_{c2}(2P) \rightarrow \gamma\gamma) \approx 0.5$  keV because

of the  $D\bar{D}^* + \bar{D}\bar{D}^*$  channel, which can be essential. For example, assuming  $BR(\chi_{c2}(2P) \rightarrow D\bar{D}) \approx BR(\chi_{c2}(2P) \rightarrow \bar{D}\bar{D}^* + \bar{D}\bar{D}^*)$ , then  $\Gamma(\chi_{c2}(2P) \rightarrow \gamma\gamma) \gtrsim 0.48$  keV or  $0.36$  keV. It is also clear that such a consideration may take place for  $\Gamma(\chi_{c2}(1P) \rightarrow 2g)$  and  $\Gamma(\chi_{c2}(2P) \rightarrow 2g)$ , which results in  $\Gamma(\chi_{c2}(2P) \rightarrow 2g) \approx \Gamma(\chi_{c2}(1P) \rightarrow 2g) = \Gamma(\chi_{c2}(1P))(1 - BR(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))) \approx 1.56$  MeV. We then obtain  $BR(\chi_{c2}(2P) \rightarrow 2g) \approx 6.5\%$ . In fact, the mass difference for  $\chi_{c2}(2P)$  and  $\chi_{c2}(1P)$  can be considered. In Ref. [8] the heavy quark mass  $m_Q$  is used in the non-relativistic limit, and instead, the meson mass  $M$  is adopted in Ref. [7], which leads to the difference of 30%. Guided by this estimate, we will write  $\Gamma(\chi_{c2}(2P) \rightarrow 2g) \approx (0.5 \pm 0.2)$  keV, and  $BR(\chi_{c2}(2P) \rightarrow 2g) \approx (6.5 \pm 2.0)\%$ . The current measurements have uncertainties with similar size. Then we note that our such observations and results agree very well with an explicit calculation from a heavy quark potential derived from the instanton vacuum along with the Coulomb and linear confinement potential [9].

### 3 Conclusion and outlook

Once the Particle Data Group as well as the BaBar and Belle collaborations have correctly identified the  $\chi_{c2}(2P)$  state, there will be a very clean and reliable constraint of  $BR(\chi_{c2}(2P) \rightarrow \text{gluon gluon}) \gtrsim 2\%$  considering the leading-order property of the Charmonium decay. It is likely that  $BR(\chi_{c2}(2P) \rightarrow 2g) \approx (6.5 \pm 2.0)\%$ .

The confirmation of the  $\chi_{c2}(2P)$  state can be tested by BESIII, for example, through the process  $e^+e^- \rightarrow \psi(4040) \rightarrow \gamma\chi_{c2}(2P)$  using their data above the center of mass of 4 GeV, as in their detector the two  $D$ s can be clearly reconstructed. The search for two-gluon decays of the  $\chi_{c2}(2P)$  state is feasible for BESIII as well as other super factories such as BaBar and Belle.

*The author XWK thanks Prof. Xiang Liu for stimulating discussion. He also acknowledges helpful discussion with Sadaharu Uehara about the Belle measurements, and with Hai-Bo Li about the BESIII measurements.*

### References

- 1 S. Uehara et al (Belle Collaboration), Phys. Rev. Lett., **99**: 082003 (2006)
- 2 C. Patrignani et al (Particle Data Group), Chin. Phys. C, **40**: 100001 (2016) and 2017 update
- 3 B. Aubert et al (BABAR Collaboration), Phys. Rev. D, **81**: 092003 (2010)
- 4 P. del Amo Sanchez et al (BaBar Collaboration), Phys. Rev. D, **84**: 012004 (2011)
- 5 J. P. Lees et al (BaBar Collaboration), Phys. Rev. D, **86**: 092005 (2012)
- 6 S. Uehara et al (Belle Collaboration), PTEP, **2013**(12): 123C01 (2013)
- 7 V. A. Novikov, L. B. Okun, M. A. Shifman, A. I. Vainshtein, M. B. Voloshin and V. I. Zakharov, Phys. Rept., **41**: 1 (1978)
- 8 W. Kwong, P. B. Mackenzie, R. Rosenfeld, and J. L. Rosner, Phys. Rev. D, **37**: 3210 (1988)
- 9 P. P. D'Souza, M. Bhat, A. P. Monteiro, and K. B. Vijaya Kumar, arXiv:1703.10413 [hep-ph]