

## Status of KAGRA

---

**Yousuke Itoh<sup>a,\*</sup> for the KAGRA collaboration**

<sup>a</sup>*Osaka Metropolitan University,  
Sumiyoshi 3-3-138, Osaka, Japan*

*E-mail:* [yousuke.itoh@omu.ac.jp](mailto:yousuke.itoh@omu.ac.jp)

The gravitational wave detectors LIGO and Virgo have so far discovered gravitational wave signals from 90 compact binary coalescences including two neutron star binary coalescences. Those discoveries are revolutionizing our understanding of the universe. This ongoing revolution will be accelerated by more accurate localizations of the gravitational wave sources and by longer coincident observation by multi-detectors, which will be made possible using as many detectors as possible. KAGRA is a gravitational wave detector built at Kamioka in Gifu prefecture, Japan, well away from the LIGO and Virgo detectors, with unique features of having cryogenic mirrors and being built underground. KAGRA can be a prototype for next-generation gravitational wave detectors that would have those features to achieve further sensitivity to gravitational waves. In this presentation, the status of KAGRA is given.

POS ( ICRC2023 ) 1555

38th International Cosmic Ray Conference (ICRC2023)  
26 July - 3 August, 2023  
Nagoya, Japan



---

\*Speaker

## 1. Introduction

The Laser Interferometer Gravitational-wave Observatory (LIGO) [1] which comprises two detectors at Hanford and Livingston in the US discovered gravitational waves (GWs) from a black hole binary merger in 2015, and this epoch-making detection has opened up a new field in astronomy by means of gravitational wave observation. In August 2017, the Virgo gravitational wave detector [2] joined the second observing run conducted by LIGO and three detectors soon discovered GWs from a black hole binary on August 14. It was just three days later that the two LIGO detectors and Virgo discovered GWs from a binary neutron star (BNS) merger named GW170817, where associated gamma-ray emission was observed 1.7 seconds later from GWs. The worldwide follow-up campaign was conducted with electromagnetic telescopes for years, giving us a clue into the long-standing mystery of the origin of gamma-ray bursts. This event then was the beginning of multimessenger astronomy.

It is noteworthy that the signal-to-noise ratio for GW170817 of the Virgo was just 2, yet that smallness allows for better localization of the GW source than that could be achieved by two LIGO detectors. In fact, observing GW signals with multi-detectors is essential to localize GW sources, thereby making multi-messenger astronomy possible [3]. Also interesting to note is that finding the number of the polarization states of GWs, as a test of general relativity, is possible with multi-detector observation [4].


As such, the fourth GW detector KAGRA [5, 6] is long-awaited in the GW and multimessenger astronomy communities. KAGRA has two key features as a GW detector: it has cryogenic mirrors and is built underground to mitigate two of the most troublesome obstacles against GW detections. LIGO, Virgo, and KAGRA are all laser interferometers. As typical GW amplitudes are extremely small, efforts to detect GWs have mostly been those to mitigate noise from various origins. Those include the seismic motion of the ground, thermal fluctuations of the mirrors and their suspensions in the interferometer, radiation pressure and shot noise of the laser, interference by scattering and stray light, frequency and intensity fluctuation of the laser, acoustic vibration, and so on. In fact, even LIGO took 13 years for its first detection since its first observing run in 2002 with its BNS range of  $\sim 200$ kpc, which tells how far on average over the sky the detector can see if a source is BNS and is a figure of merit of the detector sensitivity. Being built underground, we see the reduction of the noise due to the seismic motion in the KAGRA data, while we mitigate thermal noise by using cryogenic mirrors. A lower seismic noise level is also important for a stable operation. These two features would be adopted to some extent in designs of next-generation GW detectors like Einstein Telescope [7]. KAGRA provides valuable information on the pros and cons of those features as a prototype for advanced detector plans [8].

The KAGRA project was approved in 2010. The tunnel excavation started in 2012 and finished in 2014. In 2019, KAGRA finished all installations with the designed configuration, which we call the baseline KAGRA. We conducted the test operations using room temperature mirrors in 2016 (the iKAGRA operation) and with the cryogenic mirrors in 2018 (the bKAGRA phase-I operation), and finally conducted the joint observing run with the GEO600 GW detector [9] in Germany in 2020 April, called O3GK [10]. At that time the BNS range was approximately 500kpc on average. Since then, we have been working on improving both sensitivity and stability of the detector [11]. While Virgo was still under commissioning, the two LIGO detectors and KAGRA started the fourth

observation run called O4 on May 24, 2023. The BNS ranges are 150Mpc for the LIGO detectors and 1.3Mpc for KAGRA. The observing run will last 20 calendar months but KAGRA stopped the observation as previously planned, and we work for further improvements in sensitivity and plan to rejoin the run in 2024 spring. This contribution reviews our efforts before O4 starts and the recent detector status.


## 2. KAGRA and its O4a run

The O4 run is split into at least two segments, O4a and O4b. We have realized a power-recycling Fabry-Pérot Michelson interferometer (PRFPMI) with a DC-readout configuration in O4a. We dedicated a week to calibration and detector characterization work from May 17 to 24. The calibration and detector characterization in KAGRA are described in [12, 13]. Our observation started on May 24 and ended on June 21, while LIGO continues its observation. The duty cycle (the ratio of the time the detector is capable of taking observing data to the total calendar days) is 80% compared with 53% in O3GK. It was found that the main sources of lock-loss events, that is, out of operation were earthquakes. With the BNS range of 1.3 Mpc on average, no GW candidate has so far been detected by KAGRA. The sensitivity curve during the O3GK run and various noise contributions are shown in Fig. 1.



**Figure 1:** The KAGRA sensitivity curve during O3GK and various noise sources. The Type-A control noise had a large impact at lower frequencies and thereby the BNS range. In the KAGRA O4a run, we succeeded in reducing the Type-A control noise in lower frequencies and laser noises in higher frequencies.

We plan to adopt the resonant-sideband-extraction (RSE) technique to achieve the design sensitivity. Fig. 2 shows the optical configuration of the interferometer. We plan to cool the four sapphire mirrors at ETMX(Y) and ITMX(Y) to cryogenic temperature in an observation run in the near future but at this moment only ETMX has been cooled down to around 87K while ITMX, ETMY, and ITMY are at 250K by radiation cooling. In fact, we have succeeded in a continuous cryogenic operation of over a year at the cryogenic x-arm end test mass tank (EXC). The finesse values are 1420 and 1350 for the X-arm and Y-arm FP cavities, respectively.



**Figure 2:** KAGRA schematic optical layout. KAGRA is a resonant sideband extraction (RSE) detector with the Fabry-Pérot, power-recycling, and signal-recycling cavities. All mirrors with labels are suspended inside the vacuum tanks with four types of vibration isolation systems, e.g., test masses (TMs) by the Type-A suspension system. TMX (Y): input test mass X(Y), ETMX (Y): end test mass X (Y), BS: beam splitter, PRM: power recycling mirror, SRM: signal recycling mirror, IMMT (OMMT): input (output) mode-matching telescope, IFI (OFI): input (output) Faraday isolator

### 3. Recent efforts and achievements for sensitivity improvements and better stability

We worked on various things and have achieved improved performance of the detector toward our participation in the O4a run and after. Here we briefly explain some of the major improvements and achievements. Parts of the efforts and achievements described here and other notable works shall be presented in more detail by other contributions at this conference.

#### 3.1 Cryogenic system

We had seen frosting on sapphire mirrors when those were cooled down, resulting in unacceptably low finesse. For a resolution, we have taken the following measures. We have set a new regulation on acceptable vacuum leak level from  $10^{-9}$  [Pa m<sup>3</sup>/sec] to  $10^{-10}$  [Pa m<sup>3</sup>/sec]. Mass spectrometers were introduced in each cryostat for monitoring N<sub>2</sub>, O<sub>2</sub>, and H<sub>2</sub>O molecules. We have established a sophisticated multi-step cooling procedure to reduce the amount of frost. We also installed heaters at the intermediate mass to enhance the speed of the heating, preparing for emergencies where we may have to heat the mirrors to room temperature quickly. As a result of all the measures above, we see no serious reduction in finesse until now.

Frosting on the windows of the cryo-chambers where the optical lever light passes through was also a problem, giving us unreliable information on mirror alignment. This in turn resulted in

unreliable pitch and yaw damping control for mirrors. We thus have installed heaters at the inner and outer radiation shields, seeing no serious frosting now, although we need to watch the shields carefully when those will be cooled down at and below 40K.

When the cryo-payload was cooled down, the physical stiffness of the suspensions was enhanced and the suspended mirrors were lifted up, troubling the interferometer alignment control. We have implemented a new system to measure the heights of the mirrors using already installed cameras called TCam.

The birefringence of the sapphire mirrors is one of the biggest and long-standing problems in KAGRA. We have developed an optical simulator for the interferometer that enables us to study the effects of the sapphire birefringence

See also [14, 15] for the sapphire mirrors and [16, 17] for vibrations of the cooling system.

### 3.2 High-power laser

We have installed a new high-power laser (HPL) in the pre-stabilized laser (PSL) room with the maximum power being 60W (40W at the time of O3GK). Locks of the input mode cleaner (IMC) with this HPL were realized. The laser intensity stabilization system has been improved and we have achieved relative intensity noise (RIN) of  $10^{-8}$ , which was required for the O4 run. Indeed, this is 1-1.5 orders of magnitude better RIN compared with the system during the O3GK run. Further stability shall be aimed for future runs.

### 3.3 Alignment sensing and control

We have successfully implemented the alignment sensing and control (ASC) system including the wave form sensing (WFS) and the beam position control (BPC), except for the power-recycling mirror 3 (PR3) tank and input test masses (ITMs). Drastic enhancement of the intra-cavity power and its stability has been observed. The resulting smaller contrast fluctuation at the anti-symmetric port allows for injecting more laser power into the interferometer. We expect a noticeable reduction of shot noise in the O4b run.

### 3.4 Vibration Isolation System, Baffles, and others

Other notable improvements for the vibration isolation system (VIS), baffles, and some others are described below.

We found some faults in all the type-A suspensions almost 2 years ago after the O3GK run. The repair works and health checks on the type-A suspensions were completed before pumping the vacuum tanks. The checks were on sensor spectra, transfer functions, actuator performance and balance, low actuator coupling, securing proper positions of the instruments and sensors, and resonance damping systems among others. We verified that there were no unintentional contacts among the instruments and no disorders inside the suspensions, except for one geometric-anti-spring (GAS) filter. In addition, we installed ribbon heaters and introduced a temperature control system to stabilize the GAS filters in the suspensions. For VIS control, low-noise linear variable differential transformers (LVDT) have been introduced for IX, EX, IY, and EY. The control scheme will be refined with these new LVDTs. With all the efforts above, we have successfully reduced the control noise to a quite low level in the type-A/B/Bp suspensions except for the type-A suspension for the ETMX. See [18–21] for pre-O4a works in more details.

Mid-size baffles have been installed in the MICH area and optical dumpers have been installed in the tanks housing the input Faraday isolator (IFI), input mode matching telescope (IMMT), output mode cleaner (OMC), output mode matching telescope (OMMT) to mitigate the scattered or stray light. In fact, stray light causes non-stationary noise, and mitigation of it is key to cleaning the sensitivity and enhancing the data quality. We will make a quantitative study on the effects of these baffles for sensitivity improvements. As for OMC, repair work was completed and lower loss was realized.

We have installed gate valves (GV) to separate the MICH area from the housings for the signal recycling mirrors and the power recycling mirrors. Ion pumps are operated for all the vacuum systems. These additional GVs saved time for the repair works for the IFI and OMC vacuum tanks. We have now realized the leak level of  $10^{-10}$  [Pa m<sup>3</sup>/sec], although serious leaks at the hermetic pins at the beam splitter and around the Y-end are found.

### 3.5 Outlook

Approved two decades later than the LIGO and Virgo projects, the KAGRA project has been showing remarkably rapid progress. Being built underground and the use of the cryogenic mirrors are two key features that would be adopted by next-generation GW detectors. After the successful participation in the early O4a run, KAGRA is now in commissioning. We plan to rejoin the O4 run with improved sensitivity with the BNS range of 3~10 Mpc hopefully. In the meantime, investigation of the effects of the environmental disturbances on the detector performance and post-data-taking noise reduction techniques have been studied [22–25].

After the O4 run, the international GW detector network would conduct O5 run [26]<sup>1</sup>. Toward future runs, we are working on the possibilities of introducing a frequency-dependent squeezing technique to tackle the standard quantum noise [27, 28], obtaining larger sapphire mirrors, a higher power laser, and so on. KAGRA will contribute to the international GW detector network in the near future. KAGRA is expected to provide invaluable information about cryogenic interferometer construction and operation for future GW detector plans such as the Einstein Telescope.

### Acknowledgements

This work was supported by MEXT, JSPS Leading-edge Research Infrastructure Program, JSPS Grant-in-Aid for Specially Promoted Research 26000005, JSPS Grant-inAid for Scientific Research on Innovative Areas 2905: JP17H06358, JP17H06361 and JP17H06364, JSPS Core-to-Core Program A. Advanced Research Networks, JSPS Grantin-Aid for Scientific Research (S) 17H06133 and 20H05639 , JSPS Grant-in-Aid for Transformative Research Areas (A) 20A203: JP20H05854, the joint research program of the Institute for Cosmic Ray Research, University of Tokyo, National Research Foundation (NRF), Computing Infrastructure Project of Global Science experimental Data hub Center (GSDC) at KISTI, Korea Astronomy and Space Science Institute (KASI), and Ministry of Science and ICT (MSIT) in Korea, Academia Sinica (AS), AS Grid Center (ASGC) and the National Science and Technology Council (NSTC) in Taiwan under grants including the Rising Star Program and Science Vanguard Research Program, Advanced Technology Center (ATC) of NAOJ, and Mechanical Engineering Center of KEK.

<sup>1</sup>See <https://observing.docs.ligo.org/plan/> for the latest information.

## References

- [1] C. Cahillane and G. Mansell. Review of the Advanced LIGO Gravitational Wave Observatories Leading to Observing Run Four. *Galaxies*, 10(1):36, February 2022.
- [2] I. Nardecchia. Detecting Gravitational Waves with Advanced Virgo. *Galaxies*, 10(1):28, February 2022.
- [3] LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration. Prospects for observing and localizing gravitational-wave transients with Advanced LIGO, Advanced Virgo and KAGRA. *Living Reviews in Relativity*, 23(1):3, September 2020.
- [4] T. Akutsu et al. Overview of KAGRA: KAGRA science. *Progress of Theoretical and Experimental Physics*, 2021(5):05A103, May 2021.
- [5] T. Akutsu et al. Overview of KAGRA: Detector design and construction history. *Progress of Theoretical and Experimental Physics*, 2021(5):05A101, May 2021.
- [6] H. Abe et al. The Current Status and Future Prospects of KAGRA, the Large-Scale Cryogenic Gravitational Wave Telescope Built in the Kamioka Underground. *Galaxies*, 10(3):63, April 2022.
- [7] Michele Maggiore et al. Science case for the Einstein telescope. *Journal of Cosmology and Astroparticle Physics*, 2020(3):050, March 2020.
- [8] F. Badaracco et al. KAGRA underground environment and lessons for the Einstein Telescope. *Physical Review D*, 104(4):042006, August 2021.
- [9] Fabio Bergamin et al. Characterization and evasion of backscattered light in the squeezed-light enhanced gravitational wave interferometer GEO 600. *arXiv e-prints*, page arXiv:2305.18284, May 2023.
- [10] LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration. First joint observation by the underground gravitational-wave detector KAGRA with GEO 600. *Progress of Theoretical and Experimental Physics*, 2022(6):063F01, June 2022.
- [11] Y. Michimura et al. Prospects for improving the sensitivity of the cryogenic gravitational wave detector KAGRA. *Physical Review D*, 102(2):022008, July 2020.
- [12] T. Akutsu et al. Overview of KAGRA: Calibration, detector characterization, physical environmental monitors, and the geophysics interferometer. *Progress of Theoretical and Experimental Physics*, 2021(5):05A102, May 2021.
- [13] Y. Inoue et al. Development of Advanced Photon Calibrator for KAGRA. *arXiv e-prints*, page arXiv:2302.12180, February 2023.
- [14] E. Hirose et al. Characterization of Core Optics in Gravitational-Wave Detectors: Case Study of KAGRA Sapphire Mirrors. *Physical Review Applied*, 14(1):014021, July 2020.

[15] P. B. Li and othres. Optical Characterization of Sapphire Test Mirror in KAGRA Gravitational Wave Detector. *Acta Astronomica Sinica*, 62(6):62, November 2021.

[16] R. Bajpai et al. Vibration analysis of KAGRA cryostat at cryogenic temperature. *Classical and Quantum Gravity*, 39(16):165004, August 2022.

[17] R. Bajpai et al. Estimation of Newtonian noise from the KAGRA cooling system. *Physical Review D*, 107(4):042001, February 2023.

[18] T. Akutsu et al. Compact integrated optical sensors and electromagnetic actuators for vibration isolation systems in the gravitational-wave detector KAGRA. *Review of Scientific Instruments*, 91(11):115001, November 2020.

[19] T. Akutsu et al. An arm length stabilization system for KAGRA and future gravitational-wave detectors. *Classical and Quantum Gravity*, 37(3):035004, February 2020.

[20] T. Akutsu et al. Vibration isolation systems for the beam splitter and signal recycling mirrors of the KAGRA gravitational wave detector. *Classical and Quantum Gravity*, 38(6):065011, March 2021.

[21] Y. Zhao et al. Investigation of crackling noise in the vibration isolation systems of the KAGRA gravitational wave detector. *Physics Letters A*, 416:127664, November 2021.

[22] A. Longo et al. Local Hurst Exponent Computation of Data from Triaxial Seismometers Monitoring KAGRA. *Pure and Applied Geophysics*, 178(9):3461–3470, September 2021.

[23] T. Washimi et al. Response of the underground environment of the KAGRA observatory against the air pressure disturbance from the Tonga volcanic eruption on January 15, 2022. *Progress of Theoretical and Experimental Physics*, 2022(11):113H02, November 2022.

[24] H. Abe et al. Noise subtraction from KAGRA O3GK data using Independent Component Analysis. *Classical and Quantum Gravity*, 40(8):085015, April 2023.

[25] S. Hoshino et al. Evaluation of the microseismic motion at the KAGRA site based on the ocean wave data. *arXiv e-prints*, page arXiv:2306.12437, June 2023.

[26] P. Brady, G. Losurdo, and H. Shinkai. LIGO, VIRGO, and KAGRA as the International Gravitational Wave Network. In *Handbook of Gravitational Wave Astronomy*. Edited by C. Bambi, page 51. 2022.

[27] K. Somiya. Quantum noise reduction techniques in KAGRA. *European Physical Journal D*, 74(1):10, January 2020.

[28] Yuhang Zhao et al. Improving the stability of frequency-dependent squeezing with bichromatic control of filter cavity length, alignment, and incident beam pointing. *Physical Review D*, 105(8):082003, April 2022.

## Full Authors List: KAGRA collaboration

**Note comment afterwards:** Collaborations have the possibility to provide an authors list in xml format which will be used while generating the DOI entries making the full authors list searchable in databases like Inspire HEP.

H. Abe<sup>1</sup>, T. Akutsu<sup>2,3</sup>, M. Ando<sup>4,5</sup>, M. Aoumi<sup>6</sup>, A. Araya<sup>7</sup>, N. Aritomi<sup>8</sup>, Y. Aso<sup>2,9</sup>, S. Bae<sup>10</sup>, R. Bajpai<sup>2</sup>, K. Cannon<sup>5</sup>, Z. Cao<sup>11</sup>, R.-J. Chang<sup>12</sup>, A. H.-Y. Chen<sup>13</sup>, D. Chen<sup>14</sup>, H. Chen<sup>15</sup>, Y. Chen<sup>15</sup>, A. Chiba<sup>16</sup>, R. Chiba<sup>17</sup>, C. Chou<sup>18</sup>, M. Eisenmann<sup>2</sup>, S. Fujii<sup>17</sup>, I. Fukunaga<sup>19</sup>, D. Haba<sup>1</sup>, S. Haino<sup>20</sup>, W.-B. Han<sup>21</sup>, H. Hayakawa<sup>6</sup>, K. Hayama<sup>22</sup>, Y. Himemoto<sup>23</sup>, N. Hirata<sup>2</sup>, C. Hirose<sup>24</sup>, S. Hoshino<sup>24</sup>, H.-F. Hsieh<sup>25</sup>, C. Hsiung<sup>26</sup>, S.-C. Hsu<sup>27,25</sup>, D. C. Y. Hui<sup>28</sup>, K. Inayoshi<sup>29</sup>, Y. Itoh<sup>19,30</sup>, M. Iwaya<sup>17</sup>, H.-B. Jin<sup>31,32</sup>, K. Jung<sup>33</sup>, T. Kajita<sup>34</sup>, M. Kamiizumi<sup>6</sup>, N. Kanda<sup>30,19</sup>, J. Kato<sup>16</sup>, T. Kato<sup>17</sup>, S. Kim<sup>28</sup>, N. Kimura<sup>6</sup>, T. Kiyota<sup>19</sup>, K. Kohri<sup>35</sup>, K. Kokeyama<sup>36</sup>, K. Komori<sup>5,4</sup>, A. K. H. Kong<sup>25</sup>, N. Koyama<sup>24</sup>, J. Kume<sup>5</sup>, S. Kuroyanagi<sup>38,37</sup>, S. Kuwahara<sup>5</sup>, K. Kwak<sup>33</sup>, S. Lai<sup>18</sup>, H. W. Lee<sup>39</sup>, R. Lee<sup>15</sup>, S. Lee<sup>40</sup>, M. Leonardi<sup>41,2</sup>, K. L. Li<sup>12</sup>, L. C.-C. Lin<sup>12</sup>, C.-Y. Lin<sup>42</sup>, E. T. Lin<sup>25</sup>, G. C. Liu<sup>26</sup>, L.-T. Ma<sup>25</sup>, K. Maeda<sup>16</sup>, M. Matsuyama<sup>19</sup>, M. Meyer-Conde<sup>19</sup>, Y. Michimura<sup>43,5</sup>, N. Mio<sup>44</sup>, O. Miyakawa<sup>6</sup>, S. Miyamoto<sup>17</sup>, S. Miyoki<sup>6</sup>, S. Morisaki<sup>17</sup>, Y. Moriwaki<sup>16</sup>, M. Murakoshi<sup>45</sup>, K. Nakamura<sup>2</sup>, H. Nakano<sup>46</sup>, T. Narikawa<sup>17</sup>, L. Nguyen Quynh<sup>47</sup>, Y. Nishino<sup>2,48</sup>, A. Nishizawa<sup>5</sup>, K. Obayashi<sup>45</sup>, J. J. Oh<sup>49</sup>, K. Oh<sup>28</sup>, M. Ohashi<sup>6</sup>, M. Ohkawa<sup>24</sup>, K. Oohara<sup>50,51</sup>, Y. Oshima<sup>4</sup>, S. Oshino<sup>6</sup>, M. A. Page<sup>2</sup>, K.-C. Pan<sup>15,25</sup>, J. Park<sup>40</sup>, F. E. Peña Arellano<sup>6</sup>, S. Saha<sup>25</sup>, K. Sakai<sup>52</sup>, T. Sako<sup>16</sup>, R. Sato<sup>24</sup>, S. Sato<sup>16</sup>, Y. Sato<sup>16</sup>, T. Sawada<sup>6</sup>, Y. Sekiguchi<sup>53</sup>, L. Shao<sup>29</sup>, Y. Shikano<sup>54,55</sup>, K. Shimode<sup>6</sup>, H. Shinkai<sup>56</sup>, J. Shiota<sup>45</sup>, K. Somiya<sup>1</sup>, T. Suzuki<sup>24</sup>, T. Suzuki<sup>1</sup>, H. Tagoshi<sup>17</sup>, H. Takahashi<sup>57</sup>, R. Takahashi<sup>2</sup>, A. Takamori<sup>7</sup>, K. Takatani<sup>19</sup>, H. Takeda<sup>58</sup>, M. Takeda<sup>19</sup>, M. Tamaki<sup>17</sup>, K. Tanaka<sup>59</sup>, S. J. Tanaka<sup>45</sup>, T. Tanaka<sup>58</sup>, A. Taruya<sup>60</sup>, T. Tomaru<sup>2</sup>, K. Tomita<sup>19</sup>, T. Tomura<sup>6</sup>, A. Toriyama<sup>45</sup>, A. A. Trani<sup>5</sup>, S. Tsuchida<sup>61</sup>, N. Uchikata<sup>17</sup>, T. Uchiyama<sup>6</sup>, T. Uehara<sup>62</sup>, K. Ueno<sup>5</sup>, T. Ushiba<sup>6</sup>, M. H. P. M. van Putten<sup>63</sup>, H. Wang<sup>4</sup>, T. Washimi<sup>2</sup>, C. Wu<sup>15</sup>, H. Wu<sup>15</sup>, K. Yamamoto<sup>16</sup>, M. Yamamoto<sup>16</sup>, T. Yamamoto<sup>6</sup>, T. S. Yamamoto<sup>37</sup>, S. Yamamura<sup>17</sup>, R. Yamazaki<sup>45</sup>, L.-C. Yang<sup>18</sup>, Y. Yang<sup>18</sup>, S.-W. Yeh<sup>15</sup>, J. Yokoyama<sup>5,4</sup>, T. Yokozawa<sup>6</sup>, H. Yuzurihara<sup>6</sup>, Y. Zhao<sup>17,2</sup>, Z.-H. Zhu<sup>11,64</sup>  
(The KAGRA Collaboration)

<sup>1</sup>Graduate School of Science, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan

<sup>2</sup>Gravitational Wave Science Project, National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan

<sup>3</sup>Advanced Technology Center, National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan

<sup>4</sup>Department of Physics, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>5</sup>Research Center for the Early Universe (RESCEU), The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>6</sup>Institute for Cosmic Ray Research, KAGRA Observatory, The University of Tokyo, 238 Higashi-Mozumi, Kamioka-cho, Hida City, Gifu 506-1205, Japan

<sup>7</sup>Earthquake Research Institute, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-0032, Japan

<sup>8</sup>LIGO Hanford Observatory, Richland, WA 99352, USA

<sup>9</sup>The Graduate University for Advanced Studies (SOKENDAI), 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan

<sup>10</sup>Korea Institute of Science and Technology Information (KISTI), 245 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea

<sup>11</sup>Department of Astronomy, Beijing Normal University, Xinjiekouwai Street 19, Haidian District, Beijing 100875, China

<sup>12</sup>Department of Physics, National Cheng Kung University, No.1, University Road, Tainan City 701, Taiwan

<sup>13</sup>Institute of Physics, National Yang Ming Chiao Tung University, 101 Univ. Street, Hsinchu, Taiwan

<sup>14</sup>Kamioka Branch, National Astronomical Observatory of Japan, 238 Higashi-Mozumi, Kamioka-cho, Hida City, Gifu 506-1205, Japan

<sup>15</sup>Department of Physics, National Tsing Hua University, No. 101 Section 2, Kuang-Fu Road, Hsinchu 30013, Taiwan

<sup>16</sup>Faculty of Science, University of Toyama, 3190 Gofuku, Toyama City, Toyama 930-8555, Japan

<sup>17</sup>Institute for Cosmic Ray Research, KAGRA Observatory, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8582, Japan

<sup>18</sup>Department of Electrophysics, National Yang Ming Chiao Tung University, 101 Univ. Street, Hsinchu, Taiwan

<sup>19</sup>Department of Physics, Graduate School of Science, Osaka Metropolitan University, 3-3-138 Sugimoto-cho, Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan

<sup>20</sup>Institute of Physics, Academia Sinica, 128 Sec. 2, Academia Rd., Nankang, Taipei 11529, Taiwan

<sup>21</sup>Shanghai Astronomical Observatory, Chinese Academy of Sciences, 80 Nandan Road, Shanghai 200030, China

<sup>22</sup>Department of Applied Physics, Fukuoka University, 8-19-1 Nanakuma, Jonan, Fukuoka City, Fukuoka 814-0180, Japan

<sup>23</sup>College of Industrial Technology, Nihon University, 1-2-1 Izumi, Narashino City, Chiba 275-8575, Japan

<sup>24</sup>Faculty of Engineering, Niigata University, 8050 Ikarashi-2-no-cho, Nishi-ku, Niigata City, Niigata 950-2181, Japan

<sup>25</sup>Institute of Astronomy, National Tsing Hua University, No. 101 Section 2, Kuang-Fu Road, Hsinchu 30013, Taiwan

<sup>26</sup>Department of Physics, Tamkang University, No. 151, Yingzhan Rd., Danshui Dist., New Taipei City 25137, Taiwan

<sup>27</sup>Department of Physics, University of Washington, 3910 15th Ave NE, Seattle, WA 98195, USA

<sup>28</sup>Department of Astronomy and Space Science, Chungnam National University, 9 Daehak-ro, Yuseong-gu, Daejeon 34134, Republic of Korea

<sup>29</sup>Kavli Institute for Astronomy and Astrophysics, Peking University, Yiheyuan Road 5, Haidian District, Beijing 100871, China

<sup>30</sup>Nambu Yoichiro Institute of Theoretical and Experimental Physics (NITEP), Osaka Metropolitan University, 3-3-138 Sugimoto-cho, Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan

<sup>31</sup>National Astronomical Observatories, Chinese Academic of Sciences, 20A Datun Road, Chaoyang District, Beijing, China

<sup>32</sup>School of Astronomy and Space Science, University of Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing, China

<sup>33</sup>Department of Physics, Ulsan National Institute of Science and Technology (UNIST), 50 UNIST-gil, Ulju-gun, Ulsan 44919, Republic of Korea

<sup>34</sup>Institute for Cosmic Ray Research, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8582, Japan

<sup>35</sup>Institute of Particle and Nuclear Studies (IPNS), High Energy Accelerator Research Organization (KEK), 1-1 Oho, Tsukuba City, Ibaraki 305-0801, Japan

<sup>36</sup>School of Physics and Astronomy, Cardiff University, The Parade, Cardiff, CF24 3AA, UK

<sup>37</sup>Department of Physics, Nagoya University, ES building, Furocho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan

<sup>38</sup>Instituto de Fisica Teorica UAM-CSIC, Universidad Autonoma de Madrid, 28049 Madrid, Spain

<sup>39</sup>Department of Computer Simulation, Inje University, 197 Inje-ro, Gimhae, Gyeongsangnam-do 50834, Republic of Korea

<sup>40</sup>Technology Center for Astronomy and Space Science, Korea Astronomy and Space Science Institute (KASI), 776 Daedeokdae-ro, Yuseong-gu, Daejeon 34055, Republic of Korea

<sup>41</sup>Department of Physics, University of Trento, via Sommarive 14, Povo, 38123 TN, Italy

<sup>42</sup>National Center for High-performance computing, National Applied Research Laboratories, No. 7, R&D 6th Rd., Hsinchu Science Park, Hsinchu City 30076, Taiwan

<sup>43</sup>LIGO Laboratory , California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125, USA

<sup>44</sup>Institute for Photon Science and Technology, The University of Tokyo, 2-11-16 Yayoi, Bunkyo-ku, Tokyo 113-8656, Japan

<sup>45</sup>Department of Physical Sciences, Aoyama Gakuin University, 5-10-1 Fuchinobe, Sagamihara City, Kanagawa 252-5258, Japan

<sup>46</sup>Faculty of Law, Ryukoku University, 67 Fukakusa Tsukamoto-cho, Fushimi-ku, Kyoto City, Kyoto 612-8577, Japan

<sup>47</sup>Department of Physics and Astronomy, University of Notre Dame, 225 Nieuwland Science Hall, Notre Dame, IN 46556, USA

<sup>48</sup>Department of Astronomy, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>49</sup>National Institute for Mathematical Sciences, 70 Yuseong-daero, 1689 Beon-gil, Yuseong-gu, Daejeon 34047, Republic of Korea

<sup>50</sup>Graduate School of Science and Technology, Niigata University, 8050 Ikarashi-2-no-cho, Nishi-ku, Niigata City, Niigata 950-2181, Japan

<sup>51</sup>Niigata Study Center, The Open University of Japan, 754 Ichibancho, Asahimachi-dori, Chuo-ku, Niigata City, Niigata 951-8122, Japan

<sup>52</sup>Department of Electronic Control Engineering, National Institute of Technology, Nagaoka College, 888 Nishikatai, Nagaoka City, Niigata 940-8532, Japan

<sup>53</sup>Faculty of Science, Toho University, 2-2-1 Miyama, Funabashi City, Chiba 274-8510, Japan

<sup>54</sup>Graduate School of Science and Technology, Gunma University, 4-2 Aramaki, Maebashi, Gunma 371-8510, Japan

<sup>55</sup>Institute for Quantum Studies, Chapman University, 1 University Dr., Orange, CA 92866, USA

<sup>56</sup>Faculty of Information Science and Technology, Osaka Institute of Technology, 1-79-1 Kitayama, Hirakata City, Osaka 573-0196, Japan

<sup>57</sup>Research Center for Space Science, Advanced Research Laboratories, Tokyo City University, 8-15-1 Todoroki, Setagaya, Tokyo 158-0082, Japan

<sup>58</sup>Department of Physics, Kyoto University, Kita-Shirakawa Oiwake-cho, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan

<sup>59</sup>Institute for Cosmic Ray Research, Research Center for Cosmic Neutrinos, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8582, Japan

<sup>60</sup>Yukawa Institute for Theoretical Physics (YITP), Kyoto University, Kita-Shirakawa Oiwake-cho, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan

<sup>61</sup>National Institute of Technology, Fukui College, Geshi-cho, Sabae-shi, Fukui 916-8507, Japan

<sup>62</sup>Department of Communications Engineering, National Defense Academy of Japan, 1-10-20 Hashirimizu, Yokosuka City, Kanagawa 239-8686, Japan

<sup>63</sup>Department of Physics and Astronomy, Sejong University, 209 Neungdong-ro, Gwangjin-gu, Seoul 143-747, Republic of Korea

<sup>64</sup>School of Physics and Technology, Wuhan University, Bayi Road 299, Wuchang District, Wuhan, Hubei, 430072, China