



*universe*



Article

---

# Octonion Internal Space Algebra for the Standard Model

---

Ivan Todorov



<https://doi.org/10.3390/universe9050222>

Article

# Octonion Internal Space Algebra for the Standard Model <sup>†</sup>

Ivan Todorov <sup>‡</sup>

Institut des Hautes Études Scientifiques, 35 Route de Chartres, 91440 Bures-sur-Yvette, France;  
ivbortodorov@gmail.com

<sup>†</sup> Extended version of a lecture presented at the Workshop *Octonions and the Standard Model*, Perimeter Institute, Waterloo, ON, Canada, 8 February–17 May 2021, and at the 14th International Workshop *Lie Theory and Its Applications to Physics* (LT 14), Sofia, Bulgaria, 20–26 June 2021.

<sup>‡</sup> Permanent address: Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Tsarigradsko Chaussee 72, 1784 Sofia, Bulgaria.

**Abstract:** This paper surveys recent progress in our search for an appropriate internal space algebra for the standard model (SM) of particle physics. After a brief review of the existing approaches, we start with the Clifford algebras involving operators of left multiplication by octonions. A central role is played by a distinguished complex structure that implements the splitting of the octonions  $\mathbb{O} = \mathbb{C} \oplus \mathbb{C}^3$ , which reflect the lepton-quark symmetry. Such a complex structure on the 32-dimensional space  $\mathcal{S}$  of  $\mathcal{Cl}_{10}$  Majorana spinors is generated by the  $\mathcal{Cl}_6$  ( $\subset \mathcal{Cl}_{10}$ ) volume form,  $\omega_6 = \gamma_1 \cdots \gamma_6$ , and is left invariant by the Pati–Salam subgroup of  $Spin(10)$ ,  $G_{PS} = Spin(4) \times Spin(6) / \mathbb{Z}_2$ . While the  $Spin(10)$  invariant volume form  $\omega_{10} = \gamma_1 \cdots \gamma_{10}$  of  $\mathcal{Cl}_{10}$  is known to split  $\mathcal{S}$  on a complex basis into left and right chiral (semi)spinors,  $\mathcal{P} = \frac{1}{2}(1 - i\omega_6)$  is interpreted as the projector on the 16-dimensional *particle subspace* (which annihilates the antiparticles). The standard model gauge group appears as the subgroup of  $G_{PS}$  that preserves the sterile neutrino (which is identified with the Fock vacuum). The  $\mathbb{Z}_2$ -graded internal space algebra  $\mathcal{A}$  is then included in the projected tensor product  $\mathcal{A} \subset \mathcal{P} \mathcal{Cl}_{10} \mathcal{P} = \mathcal{Cl}_4 \otimes \mathcal{Cl}_6^0$ . The Higgs field appears as the scalar term of a superconnection, an element of the odd part  $\mathcal{Cl}_4^1$  of the first factor. The fact that the projection of  $\mathcal{Cl}_{10}$  only involves the even part  $\mathcal{Cl}_6^0$  of the second factor guarantees that the color symmetry remains unbroken. As an application, we express the ratio  $\frac{m_H}{m_W}$  of the Higgs to the W boson masses in terms of the cosine of the *theoretical* Weinberg angle.



**Citation:** Todorov, I. Octonion Internal Space Algebra for the Standard Model. *Universe* **2023**, *9*, 222. <https://doi.org/10.3390/universe9050222>

Academic Editor: Lorenzo Iorio

Received: 17 February 2023

Revised: 7 April 2023

Accepted: 20 April 2023

Published: 6 May 2023



**Copyright:** © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The elaboration of the standard model (SM) of particle physics was completed in the early 1970s. To quote John Baez [1], “years trying to go beyond the Standard Model hasn’t yet led to any clear success”. The present survey belongs to an equally long—albeit less fashionable—effort to clarify the algebraic (or geometric) roots of the SM or, more specifically, to find a natural framework featuring its internal space properties. After discussing some old explorations, we provide an updated exposition of recent developments (particularly, of [2]) while clarifying the meaning and role of complex structures, and we concentrate on one structure associated with a Clifford algebra (in our case,  $\mathcal{Cl}_6$ ) pseudoscalar.

Most ideas on the natural framework of the SM originate in the 1970s, the first decade of its existence (There are two exceptions: the Jordan algebras were introduced and classified in the 1930s [3,4]; and the non-commutative geometry approach was born in the late 1980s [5–7] and is still vigorously developed by Connes and collaborators and followers [8–12]).

First, early in 1973, the ultimate division algebra, the octonions<sup>1</sup>, were introduced by Gürsey<sup>2</sup> and his student Günaydin [14,15] for the description of quarks and their  $SU(3)$  color symmetry. The idea was taken up and extended to incorporate all four division algebras by G. Dixon (see [16,17] and the earlier work cited there) and was further developed by Furey [18–24]. Dubois-Violette (D-V) [25,26] emphasizes the lepton-quark correspondence and unimodularity of the color group  $SU(3)_c$  as a physical motivation for introducing the octonions; they come equipped with a complex structure preserved by subgroup  $SU(3)$  of the automorphism group  $G_2$  of  $\mathbb{O}$ :

$$\mathbb{O} = \mathbb{C} \oplus \mathbb{C}^3. \quad (1)$$

### 1.1. (Split) Octonions as Composition Algebras

One can in fact provide a basis-free definition of the octonions, starting with the splitting (1). To this end, one uses the skew symmetric vector product and standard inner product on  $\mathbb{C}^3$  to define a non-commutative and non-associative distributive product  $xy$  on  $\mathbb{O}$  and a real-valued, non-degenerate symmetric bilinear form  $\langle x, y \rangle = \langle y, x \rangle$ , such that the quadratic norm  $N(x) = \langle x, x \rangle$  is multiplicative:

$$N(xy) = N(x)N(y) \quad \text{for} \quad N(x) = \langle x, x \rangle \quad (2)$$

(cf. [25,27,28]). Furthermore, by defining the real part of  $x \in \mathbb{O}$  by  $\text{Re } x = \langle x, 1 \rangle$  and the octonionic conjugation by  $x \rightarrow x^* = 2\langle x, 1 \rangle - x$ , we have

$$xx^* = N(x)\mathbb{I} \Leftrightarrow x^2 - 2\langle x, 1 \rangle x + N(x)\mathbb{I} = 0. \quad (3)$$

A unital algebra with a non-degenerate quadratic norm obeying (2) is called a *composition algebra*.

Another basis-free definition of the octonions  $\mathbb{O}$  and their split version  $\tilde{\mathbb{O}}$  can be provided in terms of quaternions using the Cayley–Dickson construction. We represent the quaternion as scalars plus vectors

$$\begin{aligned} \mathbb{H} &= \mathbb{R} \oplus \mathbb{R}^3, \quad x = u + U, \quad y = v + V, \quad u, v \in \mathbb{R}, \quad U, V \in \mathbb{R}^3, \\ xy &= uv - \langle U, V \rangle + uV + Uv + U \times V \end{aligned} \quad (4)$$

with the vector product  $U \times V \in \mathbb{R}^3$  satisfying

$$U \times V = -V \times U, \quad (U \times V) \times W = \langle U, W \rangle V - \langle V, W \rangle U. \quad (5)$$

The product (4) is clearly non-commutative, but one verifies that it is associative. The Cayley–Dickson construction defines the octonions  $\mathbb{O}$  and split octonions  $\tilde{\mathbb{O}}$  in terms of a pair of quaternions and a new “imaginary unit”  $\ell$  is defined as:

$$\begin{aligned} x &= u + U + \ell(v + V), \quad \ell(v + V) = (v - V)\ell, \\ \ell^2 &= \begin{cases} -1 & \Rightarrow \quad x \in \mathbb{O} \\ 1 & \Rightarrow \quad x \in \tilde{\mathbb{O}}. \end{cases} \end{aligned} \quad (6)$$

We shall encounter the split octonions as generators of  $C\ell(4, 2)$  in Section 3.1 below.

### 1.2. Jordan Algebras; Guts; Clifford Algebras

Studying quantum field theory, it appears natural to replace classical observables (real-valued functions) by an algebra of functions on space-time with values in a *finite dimensional Euclidean Jordan algebra*<sup>3</sup>. As a particularly attractive choice that incorporates

the idea of lepton-quark symmetry, D-V proposes [25] the exceptional Jordan or *Albert* algebra of  $3 \times 3$  Hermitian octonionic matrices,

$$J_3^8 = \mathcal{H}_3(\mathbb{O}), \quad (7)$$

which is the only irreducible one that does not admit an associative envelope [32]. Further progress was achieved in [27,28,30,33–35] by considering the Clifford algebra envelope of its non-exceptional subalgebra  $J_2^8$ , which fits one generation of fundamental fermions. In these papers, as well as in the present one, we are effectively working with associative algebras that should be viewed as an internal space counterpart of Haag’s *field algebra* (see [36]).

A second development, *grand unified theory* (GUT), anticipated again in 1973 by Pati and Salam [37], became mainstream for a time<sup>4</sup>. Fundamental chiral fermions fit the complex spinor representation of  $Spin(10)$ , which was introduced as a GUT group by Fritzsch and Minkowski and by Georgi. A preferred symmetry breaking yields the maximal rank semisimple Pati–Salam subgroup

$$G_{PS} = \frac{Spin(4) \times Spin(6)}{\mathbb{Z}_2} \subset Spin(10),$$

$$Spin(4) = SU(2)_L \times SU(2)_R, \quad Spin(6) = SU(4). \quad (8)$$

We note that  $G_{PS}$  is the only GUT group that does not predict a gauge triggered proton decay; it is also encountered in the non-commutative geometry approach to the SM [8,10]. In general, GUTs provide a nice home for the fundamental fermions, as displayed by the two 16-dimensional complex conjugate “Weyl spinors” of  $Spin(10)$ . Their other representations, however, such as the 45-dimensional adjoint representation of  $Spin(10)$ , are much too big and involve hypothetical particles such as lepto-quarks which cause difficulties.

The Clifford algebra<sup>5</sup>  $\mathcal{C}\ell_{10}$ , on the other hand, like the Clifford algebra of any even-dimensional Euclidean vector space, has a unique irreducible representation (IR); in the case of<sup>6</sup>  $\mathcal{C}\ell_{10} \cong \mathbb{R}[2^5]$ , it is the 32-component real (Majorana) spinor. Viewed as a representation of  $Spin(10)$ , it splits upon complexification into two 16-dimensional (complex) IRs that can be naturally associated to the left and right chiral fundamental (anti)fermions of one generation:

$$32 = \mathbf{16}_L + \mathbf{16}_R. \quad (9)$$

Clifford algebras were also applied to the SM in the 1970s—see [40] and the references therein. An essential difference in our approach is the use of octonions with a preferred complex structure in  $\mathcal{C}\ell_{8+\nu}$ ,  $\nu = 0, 1, 2$  to restrict the corresponding gauge group (another new point, the use of the  $\mathbb{Z}_2$  grading of  $\mathcal{C}\ell_{10}$  to define the Higgs field, will be discussed in Section 1.3 below).

The Pati–Salam subgroup of  $Spin(10)$  is singled out as the stabilizer of the  $\mathcal{C}\ell_6 (\subset \mathcal{C}\ell_8 \subset \mathcal{C}\ell_{10})$  pseudoscalar

$$\omega_6 = \gamma_1 \dots \gamma_6 \quad \text{for} \quad \gamma_\alpha = \sigma_0 \otimes \epsilon \otimes L_\alpha, \quad \gamma_8 = \sigma_0 \otimes \sigma_1 \otimes \mathbb{1}_8 (\in \mathcal{C}\ell_{10}),$$

$$\sigma_0 = \mathbb{1}_2, \quad \epsilon = i\sigma_2, \quad L_\alpha = L_{e_\alpha}, \quad [L_\alpha, L_\beta]_+ = -2\delta_{\alpha\beta}\mathbb{1}_8, \quad \alpha, \beta = 1, \dots, 7. \quad (10)$$

Here,  $L_x$  is the operator of left multiplication in the eight-dimensional real vector space of the octonions,  $L_x y = xy$  for  $x, y \in \mathbb{O}$ . The action of the operators  $L_\alpha \in \mathbb{R}[8]$  on the octonion units will be made explicit in Section 2.2 (Equation (22)). The group  $G_{PS}$  (8) in fact preserves each factor in the graded tensor product representation of  $\mathcal{C}\ell_{10}$ :

$$\mathcal{C}\ell_{10} = \mathcal{C}\ell_4 \hat{\otimes} \mathcal{C}\ell_6 \quad (11)$$

Introduced earlier by Furey [20,21] and exploited in [2], the complex structure  $J \in SO(10)$  generated by  $\omega_6$  will be displayed, and the physical interpretation of the mutually orthogonal projection operators

$$\mathcal{P} = \frac{1}{2}(1 - i\omega_6), \mathcal{P}' = \frac{1}{2}(1 + i\omega_6) \quad (12)$$

will be revealed in Section 2.3.

### 1.3. Main Message and Organization of the Paper

The present survey focuses on ongoing attempts to answer two questions:

(1) Why is the arbitrarily looking gauge group of the SM

$$G_{\text{SM}} = S(U(2) \times U(3)) = \frac{SU(2) \times SU(3) \times U(1)}{\mathbb{Z}_6}, \quad (13)$$

and what dictates its highly reducible representation for fundamental fermions?

(2) How do we put together the Higgs field with the gauge bosons? Can we explain their mass ratios?

1. Most physicists accept GUT as an answer to the first question. One has the intriguing result of Baez and Huerta [38], where  $G_{\text{SM}}$  appears as the intersection of two popular GUT subgroups of  $Spin(10)$ :

$$G_{\text{SM}} = SU(5) \cap G_{\text{PS}} \subset Spin(10).$$

A top-down approach starting with  $Spin(10)$ , however, should involve the maximal rank subgroup  $U(5)$  instead of  $SU(5)$ , in line with the philosophy of Borel-de Siebenthal [41], yielding an extra  $U(1)$  factor in the intersection.

The minority that are trying to go further includes, besides the fans of octonions and the already cited enthusiasts of almost commutative real spectral triples, Holger Nielsen, whose more than two decades of musing over the problem are reviewed in [42]. Our approach exploits the complex structure and particle projector  $\mathcal{P}$  (12) associated with the Clifford pseudoscalar  $\omega_6$  (10); it permeates the entire paper (Sections 2.3, 3.2–3.4 and 5.1...).

2. The second problem has been universally recognized (see, e.g., the popular account [43]). We follow the superconnection approach anticipated by Ne’eman and Fairlie—for a concise review and references, see Section 4.1. We exploit the restricted particle projector  $\mathcal{P}_r$ , which annihilates the sterile neutrino (Section 3.3) to deform the Fermi oscillators in the lepton sector into the odd generators of the simple Lie superalgebra postulated in [44,45]. The resulting difference between the flavor spaces of leptons and colored quarks allows one to compute the mass ratio  $m_H/m_W$  in agreement with the experiment (Section 4.2).

The paper aims to be self-contained and combines our contribution in a single narrative with a review of the background material. Section 2.1 provides a summary of the known triality realization of  $Spin(8)$ . Section 2.2 and Appendix A spell out the relation between left and right multiplication using imaginary octonion units, which is applied in Section 3.4 to display the stabilizer of  $R_7$ . We would like to single out two messages from Section 2.3: (1) the indirect connection between the  $C\ell_6$  pseudoscalar and the complex structure  $J \in SO(8)$  (33) and (35); (2) the observation that the Lie subalgebra of  $so(8)$  that commutes with  $\omega_6$  and the electric charge operator  $Q$  (45) is the rank four subalgebra  $su(3)_c \oplus u(1)_Q \oplus u(1)_{B-L}$ . Section 3.1 contains, along with a glance on the equivalence class of Clifford algebras involving  $C\ell(3, 1)$ ,  $C\ell_{-6} (= C\ell(0, 6))$ ,  $C\ell_{10}$ , the observation that the conformal Clifford algebra  $C\ell(4, 2)$  of this class is generated by the split octonions and gives rise to their isometry group  $SO(4, 4)$ . Section 3.2 contains one of the main messages of the paper: the SM gauge group (13) is the subgroup of  $G_{\text{PS}}$  (8) that leaves the sterile neutrino invariant (Proposition 1). Section 3.3 discusses superselection rules and the superselection of the weak hypercharge. Section 3.4 reviews and comments on recent work [24,46] on the

complex structure associated with the right action of octonion units as well as the derivation of the gauge group for the SM [35] and its left-right symmetric extension [47].

The Dirac operator  $\gamma^\mu(\partial_\mu + A_\mu)$  anticommutes with the chirality  $\gamma_5$  and hence intertwines the left and right fermions; so does the Higgs field, which substitutes a mass term in the fermionic Lagrangian. This inspired Connes and coworkers [5,6,9] to identify the Higgs field with the internal space part of the Dirac operator. This idea finds a natural implementation in the Clifford algebra approach for the SM *superconnection* (reviewed in Section 4.1). The concise exposition in Section 4 emphasizes our assumptions and some delicate points, referring the reader for calculational details to the preceding publication [2].

We recapitulate our convoluted route to  $\mathcal{Cl}_{10}$  in Section 5.1. In Section 5.2, we compare our solution of the fermion doubling problem with the approach of [23]. A summary of the main results of the paper is given in Section 5.3, which also cites existing (inconclusive) attempts to understand why there are exactly three generations of fundamental fermions.

## 2. Triality Realization of $Spin(8)$ : $\mathcal{Cl}_{-6}$

### 2.1. The Action of Octonions on Themselves

The group  $Spin(8)$ , the double cover of the orthogonal group  $SO(8) = SO(\mathbb{O})$ , can be defined (see [48,49]) as the set of triples  $(g_1, g_2, g_3) \in SO(8) \times SO(8) \times SO(8)$  such that

$$g_2(xy) = g_1(x)g_3(y) \text{ for any } x, y \in \mathbb{O}. \quad (14)$$

If  $u$  is a unit octonion,  $u^*u = 1$ ; then, the left and right multiplications by  $u$  are examples of isometries of  $\mathbb{O}$

$$|L_u x|^2 = \langle ux, ux \rangle = \langle x, x \rangle, \quad |R_u x|^2 = \langle xu, xu \rangle = \langle x, x \rangle \text{ for } \langle u, u \rangle = 1. \quad (15)$$

Using the *Moufang identity*<sup>7</sup>

$$u(xy)u = (ux)(yu) \text{ for any } x, y, u \in \mathbb{O}, \quad (16)$$

One verifies that the triple  $g_1 = L_u$ ,  $g_2 = L_u R_u$ ,  $g_3 = R_u$  satisfies (14) and hence belongs to  $Spin(8)$ . It turns out that triples of this type generate  $Spin(8)$  (see [48] or Yokota's book [49] for a proof).

The mappings  $x \rightarrow L_x$  and  $x \rightarrow R_x$  are, of course, not algebra homomorphisms, as  $L_x$  and  $R_y$  each generate an associative algebra, while the algebra of octonions is non-associative. They do, however, preserve the quadratic relation  $xy^* + yx^* = 2\langle x, y \rangle 1$ :

$$L_x L_{y^*} + L_y L_{x^*} = 2\langle x, y \rangle \mathbb{1} = R_x R_{y^*} + R_y R_{x^*} \quad (L_x^* = L_{x^*}). \quad (17)$$

Equation (17) applied to the span of the first six imaginary octonion units  $e_j$ ,  $j = 1, \dots, 6$  and setting  $L_{e_j} =: L_j$ ,  $R_{e_j} =: R_j$  becomes the defining relation of the Clifford algebra  $\mathcal{Cl}_{-6}$ :

$$L_j L_k + L_k L_j = -2\delta_{jk} = R_j R_k + R_k R_j, \quad j, k = 1, \dots, 6. \quad (18)$$

### 2.2. $\mathcal{Cl}_{-6}$ as a Generating Algebra of $\mathbb{O}$ and $so(\mathbb{O})$

The octonions appear in any of the nested Clifford algebras  $\mathcal{Cl}_8 \subset \mathcal{Cl}_9 \subset \mathcal{Cl}_{10}$ . In fact, the minimal realization of  $\mathbb{O}$  is provided by  $\mathcal{Cl}_{-6}$ , generated by the left multiplication  $L_\alpha$  by six of the seven imaginary octonion units  $e_\alpha$ . In general,  $L_x L_y \neq L_{xy}$  (and similarly for  $R$ ), but remarkably, as noted in [20], the relation  $e_1(e_2(e_3(e_4(e_5(e_6 e_a)))))) = e_7 e_a$  is satisfied for all  $a = 1, \dots, 8$  ( $e_8 = 1$ ) so that

$$L_1 L_2 \cdots L_6 = L_{e_7} =: L_7, \quad R_1 R_2 \cdots R_6 = R_{e_7} =: R_7. \quad (19)$$

While  $L_x R_x = R_x L_x$  (for  $x \in \mathbb{O}$ ), the non-associativity of the algebra of octonions is reflected in the fact that for  $x \neq y$ ,  $L_x$  and  $R_y$  in general, do not commute. The Lie algebra

$so(8)$  is spanned by the elements of the negative square of  $C\ell_{-6}$ . If we denote the exterior algebra on the span of  $L_1, \dots, L_6$  by

$$\Lambda^* \equiv \Lambda^* C\ell_{-6} = \Lambda^0 + \Lambda^1 + \dots + \Lambda^6 \left( \Lambda^1 = \underset{1 \leq j \leq 6}{\text{Span}} L_j, \Lambda^6 = \{\mathbb{R} L_7\} \right)$$

then  $so(8) = \Lambda^1 + \Lambda^2 + \Lambda^5 + \Lambda^6$  (accordingly, the 28-dimensional adjoint representation of  $so(8)$  splits into four irreducible representations of  $so(6)$  :  $\mathbf{28} = \mathbf{6} + \mathbf{15} + \mathbf{6}^* + \mathbf{1}$ . In particular,  $\Lambda_5 = \text{Span}\{L_{\alpha 7}, \alpha = 1, \dots, 6\}$  for  $L_{\alpha\beta}$  defined below). A basis of the Lie algebra, given by

$$L_{\alpha 8} = \frac{1}{2} L_\alpha, L_{\alpha\beta} = -\frac{1}{4} [L_\alpha, L_\beta], \alpha, \beta = 1, \dots, 7 \quad (20)$$

obeys the standard commutation relations (CRs) for  $so(n)$  (herein  $n = 8$ ):

$$[L_{ab}, L_{cd}] = \delta_{bc} L_{ad} - \delta_{bd} L_{ac} + \delta_{ad} L_{bc} - \delta_{ac} L_{bd},$$

$$L_{ab} = \frac{1}{4} (L_a L_b^* - L_b L_a^*), a, b, c, d = 1, 2, \dots, 8 \quad (L_\alpha^* = -L_\alpha, L_8^* = L_8) \quad (21)$$

(and similarly for  $R_{ab}$ ). Each element of  $so(8)$  of square  $-1$  defines a *complex structure* (see Section 2.3). Following [24], we shall single out the *Clifford pseudoscalars*  $L_7$  and  $R_7$  (19) (called *volume forms* in the highly informative lectures [51] and Coxeter elements in [52]). We shall use the (mod 7) multiplication rules of [13] for the imaginary octonion units

$$L_i e_j (= e_i e_j) = -\delta_{ij} + f_{ijk} e_k, f_{ijk} = 1$$

$$\text{for } (i, j, k) = (1, 2, 4)(2, 3, 5)(3, 4, 6)(4, 5, 7)(5, 6, 1)(6, 7, 2)(7, 1, 3) \quad (22)$$

and  $f_{ijk}$  is fully antisymmetric within each of the above seven triples. The Clifford pseudoscalar is naturally associated with the Cartan subalgebra of  $so(6)$ , spanned by

$$(L_{13}, L_{26}, L_{45}) \text{ as } L_7(e_1, e_2, e_4) = (e_3, e_6, e_5). \quad (23)$$

We can write

$$L_7 = 2^3 L_{13} L_{26} L_{45} \text{ (as } 2L_{13} = L_1 L_3^* = -L_1 L_3 \text{ etc.).} \quad (24)$$

The infinitesimal counterpart of (14) reads

$$T_\alpha(x, y) = (L_\alpha x)y + x(R_\alpha y) \text{ for } \alpha, x, y \in \mathbb{O}, \alpha^* = -\alpha,$$

$$\text{i.e., } T_\alpha = L_\alpha + R_\alpha. \quad (25)$$

There is an involutive outer automorphism  $\pi$  of the Lie algebra  $so(8)$  such that

$$\pi(L_\alpha) = T_\alpha, \pi(R_\alpha) = -R_\alpha, \pi(T_\alpha) = L_\alpha \quad (\pi^2 = id). \quad (26)$$

As proven in Appendix A,

$$\pi(L_{ab}) = E_{ab}, \text{ where } E_{ab} e_c = \delta_{bc} e_a - \delta_{ac} e_b \quad (a, b, c = 1, 2, \dots, 8, e_8 = 1). \quad (27)$$

$(L_{ab})$ ,  $(E_{ab})$ , and  $(R_{ab})$  provide three bases of  $so(8)$ , each obeying the CRs (21). They are expressed by each other in terms of the involution  $\pi$ :

$$L_{ab} = \pi(E_{ab}), E_{\alpha 8} = L_{\alpha 8} + R_{\alpha 8}, \alpha = 1, \dots, 7. \quad (28)$$

We find, particularly (see Appendix A):

$$L_7 = 2L_{78} = E_{78} - E_{13} - E_{26} - E_{45} = 2E_{78} - R_7; 2L_{13} = E_{13} - E_{26} - E_{45} - E_{78},$$

$$2L_{26} = E_{26} - E_{13} - E_{45} - E_{78}, 2L_{45} = E_{45} - E_{13} - E_{26} - E_{78}. \quad (29)$$

While  $L_{78} = 4L_{13}L_{26}L_{45}$  (24) commutes with the entire Lie algebra  $spin(6) = su(4)$  and the  $u(1)$  generator (whose physical meaning is revealed by (45)).

$$C_1 = L_{13} + L_{26} + L_{45} \text{ centralizes } u(3) = u(1) \oplus su(3) \subset su(4) \quad (30)$$

where the second summand is the unbroken color Lie algebra  $su(3) = su(3)_c$ .

### 2.3. Complex Structure and Symmetry Breaking in $\mathcal{Cl}_n$

The algebra  $\mathcal{Cl}_8$  is generated by two-by-two Hermitian matrices whose elements involve the operators  $L_a$  of the left multiplication by octonion units:

$$\gamma_a = \begin{pmatrix} 0 & L_a \\ L_a^* & 0 \end{pmatrix}, L_a = L_{e_a}, L_a^* = L_{e_a^*}, a = 1, \dots, 8. \quad (31)$$

Here,  $e_8 = 1 (= e_8^*)$ ,  $L_8 = \mathbb{1}_8$  is the unit operator in  $\mathbb{R}^8$ ;  $L_\alpha^* = -L_\alpha$  for  $\alpha = 1, \dots, 7$  so that the  $\mathcal{Cl}_{10}$  generators  $\gamma_\alpha$  (10) are obtained from the above (for  $\alpha = 1, \dots, 7$ ) through tensoring with the  $2 \times 2$  unit matrix  $\sigma_0$ .

A compact way to identify the particle states in a Clifford algebra  $\mathcal{Cl}_{2n}$  is to introduce a complex structure which, as we shall demonstrate, gives rise to a fermionic Fock space in  $\mathcal{Cl}_{2n}$ .

$$(JX, JY) = (X, Y), (JX, Y) = -(X, JY), \forall X, Y \in E_{2n}. \quad (32)$$

For a non-zero vector  $X$  and a complex structure  $J$ , the vector  $Y = JX$  is orthogonal to  $X$  (and has the same norm):

$$Y = JX \Rightarrow (X, Y) = 0 \ ((X, X) = (Y, Y) > 0).$$

It follows that for each complex structure  $J$  in  $E_{2n}$ , there exists an orthonormal basis of the form  $(\gamma_1, \dots, \gamma_n, J\gamma_1, \dots, J\gamma_n)$  in  $\mathcal{Cl}_{2n}$ . Then,  $a_j = \frac{1}{2}(\gamma_j - iJ\gamma_j)$  and  $a_j^* = \frac{1}{2}(\gamma_j + iJ\gamma_j)$ ,  $j = 1, \dots, n$  each span the image in  $\mathcal{Cl}_{2n} (= \mathcal{Cl}_{2n}(\mathbb{C}))$  of a maximal isotropic subspace of the complexification of  $E_{2n}$ . Together, they yield a realization of the *canonical anticommutation relations* (CAR). Fermionic oscillators have been used in the present context in [21,53]. The complex structure in  $so(2n)$  involves a distinguished maximal (rank  $n$ ) Lie subalgebra (a notion studied in [41]),  $u(n) \subset so(2n)$ , which is generated by the products  $b_j b_k^*$ . It also selects two distinguished  $u(n)$  singlet states in  $\mathcal{Cl}_{2n}$ , the *vacuum*, which is annihilated by all  $b_j$  and the antipode, which is annihilated by the  $b_j^*$ . Both singlets are annihilated by the simple part  $su(n)$  of  $u(n)$ .

Complex structures have been studied in relation to spinors by Élie Cartan (since 1908), Veblen (1933), and Chevalley (1954). For a carefully written survey with historical highlights - see [54]. For a concise modern exposition that connects them to the states in a fermionic Fock space, see Dubois-Violette [55]. We have also been influenced by their use (in  $SO(9)$ ) by Krasnov [46] and by relating them to Clifford pseudoscalars in [24].

The pseudoscalar  $\omega_6$  of  $\mathcal{Cl}_6$  belongs to  $\mathcal{Cl}_8$  but only defines a complex structure through its action on the octonion units. More precisely, taking the basic relations (22) and the identity  $\epsilon^2 = -\sigma_0$  into account, we can write

$$\omega_6 = -\sigma_0 \otimes L_7, \text{ where } -L_7 e_a = \sum_b J_{ab} e_b, J_{\alpha\beta} = -f_{7\alpha\beta}, \alpha, \beta = 1, \dots, 6,$$

$$J_{ab} = -J_{ba}, J_{13} = J_{26} = J_{45} = -J_{78} = -1. \quad (33)$$

Warning: due to the non-associativity of  $\mathbb{O}$ ,  $-L_7 e_3 = e_1$  does not imply  $-L_7 L_3 = L_1$ , etc.

We shall see that each of the subgroups  $Spin(n)^{\omega_6}$  of the spin groups of  $C\ell_n$  for  $n = 8, 9, 10$  that leaves  $\omega_6$  invariant is relevant for the particle physics:

$$Spin(8)^{\omega_6} = U(4), \quad Spin(9)^{\omega_6} = SU(4) \times SU(2), \quad Spin(10)^{\omega_6} = G_{PS} \quad (34)$$

The  $U(1)$  factor in the  $U(4)$  of  $Spin(8)^{\omega_6}$  and the  $SU(2)$  in  $Spin(9)^{\omega_6}$  are generated by all three components of the “total weak isospin”  $\mathbf{I} = \mathbf{I}^L + \mathbf{I}^R$ , as will be made explicit in Section 3.2.

We shall define a complex structure  $J$  corresponding to  $\omega_6$  by extrapolating (33) to a transformation of the gamma matrices of  $C\ell_8$ :

$$J : \gamma_a \rightarrow \sum_b J_{ab} \gamma_b = \omega_J \gamma_a \omega_J^*, \quad \omega_J = \frac{1}{4}(1 + \gamma_{13})(1 + \gamma_{26})(1 + \gamma_{45})(1 - \gamma_{78}). \quad (35)$$

We can extend the basis (31) of  $C\ell_8$  to  $C\ell_{10}$ , setting (cf. (10)):

$$\gamma_\alpha = \sigma_0 \otimes \epsilon \otimes L_\alpha \text{ for } \alpha = 1, \dots, 7,$$

$$\gamma_8 = \sigma_0 \otimes \sigma_1 \otimes \mathbb{1}_8, \quad \gamma_9 = \sigma_2 \otimes \sigma_3 \otimes \mathbb{1}_8, \quad \gamma_{10} = \sigma_1 \otimes \sigma_3 \otimes \mathbb{1}_8. \quad (36)$$

In particular,  $C\ell_9 = \mathbb{R}^{16} \oplus \mathbb{R}^{16}$  is generated by the  $32 \times 32$  matrices  $\gamma_1, \dots, \gamma_9$ , which commute with  $\omega_9 = \gamma_1 \gamma_2 \dots \gamma_9 = \sigma_2 \otimes \mathbb{1}_{16}$ . The Lie subalgebra of  $so(n)$  of the derivations of  $C\ell_n$ ,  $n = 8, 9, 10$  that commute with  $\omega_6$  (10) is  $so(6) \oplus so(n-6)$ . For  $n = 10$ , it is the Lie algebra  $g_{PS}$  of the Pati–Salam group (8) that respects the tensor product representation (11) of  $C\ell_{10}$ .

We now proceed to give meaning to the projection operator

$$\mathcal{P} = \frac{1 - i\omega_6}{2} \quad (\mathcal{P}^2 = \mathcal{P}), \quad \text{tr } \mathcal{P} = \text{tr}(1 - \mathcal{P}) = 2^{\ell-1} \text{ for } n = 2\ell (= 6, 8, 10). \quad (37)$$

To begin with, we introduce the isotropic elements

$$2b_1 = (1 - iJ)\gamma_1 = \gamma_1 + i\gamma_3, \quad 2b_2 = (1 - iJ)\gamma_2 = \gamma_2 + i\gamma_6, \quad 2b_3 = (1 - iJ)\gamma_4 = \gamma_4 + i\gamma_5. \quad (38)$$

These correspond to the projected octonion units  $\frac{1}{2}(1 + iL_7)e_\ell$ ,  $\ell = 1, 2, 4$ . Together with their conjugates  $b_j^*(b_1^* = \frac{1}{2}(\gamma_1 - i\gamma_3)$ , etc.), they realize the CAR

$$[b_j, b_k]_+ = 0, \quad [b_j, b_k^*]_+ = \delta_{jk}, \quad j, k = 1, 2, 3. \quad (39)$$

The annihilation operators  $b_j$  span the (maximal) three-dimensional isotropic subspace  $\mathcal{H}^{(1,0)}$  of the six-dimensional complex vector space  $\mathbb{C}E_6$ , while  $b_j^*$  span its orthogonal complement  $\mathcal{H}^{(0,1)}$ ; we have:

$$J(1 \mp iJ)\gamma_\ell = \pm i(1 \mp iJ)\gamma_\ell \text{ for } \ell = 1, 2, 4. \quad (40)$$

The commuting Hermitian elements  $i\gamma_{\ell 3\ell \text{ mod } 7}$ ,  $\ell = 1, 2, 4$ , which span a Cartan subalgebra of the complexified  $so(6)$ , can be expressed as commutators of  $b_j^*$  and  $b_j$ ,  $j = 1, 2, 3$  or as differences of the associated projection operators  $p'_j - p_j$ :

$$i\gamma_{\ell 3\ell} = [b_\ell^*, b_\ell] = p'_\ell - p_\ell, \quad \ell = 1, 2, \quad i\gamma_{45} = [b_3^*, b_3] = p'_3 - p_3,$$

$$p_j = b_j b_j^*, \quad p'_j = b_j^* b_j = 1 - p_j, \quad p_j p'_j = 0, \quad j = 1, 2, 3. \quad (41)$$

In terms of these operators, the  $C\ell_6$  pseudoscalar and the projector  $\mathcal{P}$  assume the form:

$$i\omega_6 = (p'_1 - p_1)(p'_2 - p_2)(p'_3 - p_3) = \mathcal{P}' - \mathcal{P}, \mathcal{P}' = 1 - \mathcal{P},$$

$$\mathcal{P} = \ell + q, \ell = p_1 p_2 p_3, q = q_1 + q_2 + q_3, q_j = p_j p'_k p'_\ell. \quad (42)$$

The triple  $(j, k, \ell)$  being a permutation of  $(1, 2, 3)$ .

We shall identify the generators (of the complexification  $sl(3, \mathbb{C})$ ) of  $su(3)$  with the traceless part of the matrix  $(b_j b_k^*)$ , whose elements belong to  $\mathcal{H}^{(1,1)}$ . Then, the splitting (42) of  $\mathcal{P}$  into the  $su(3)$  singlet  $\ell$  and the triplet  $q$  implements the lepton-quark splitting anticipated by its image (1) on the octonions. We shall thus interpret the one-dimensional projectors  $\ell$  and  $q_j$  as describing the lepton and the colored quark states in  $C\ell_6$ . The states  $\ell$  and  $q_j$  are mutually orthogonal idempotents, with  $\ell$  playing the role of the Fock vacuum in  $C\ell_6$ :

$$\ell^2 = \ell, \ell q_j = 0, q_j q_k = \delta_{jk} q_j, b_j \ell = 0 = \ell b_k^*. \quad (43)$$

**Remark 1.** We shall argue in Section 3.3 that the identification of  $\mathcal{P}$  as a particle subspace projector (adopted in [34]) would only be justified if we have a clear distinction between particles and antiparticles. This can be claimed for the 30 fundamental (anti)fermions of the  $C\ell_{10}$  multiplet 32 (9), which have different quantum numbers with respect to the gauge Lie algebra  $\mathfrak{g}_{SM}$  of the SM. It fails in the two-dimensional subspace of sterile neutrinos annihilated by  $\mathfrak{g}_{SM}$ ;  $\nu_R$  and  $\bar{\nu}_L$  are allowed to form a coherent superposition—a Majorana spinor. In Sections 3.3 and 4, we shall adopt the restricted projector  $\ell_r$  on the (three- rather than four-dimensional) lepton subspace excluding the sterile neutrino.

In order to extend the Fock space picture to  $C\ell_8$ , we shall set

$$i\gamma_7 = a^* - a, \gamma_8 = a + a^* \Rightarrow i\gamma_{78} = [a^*, a] \quad (44)$$

where the pair  $(a^*, a)$  describes another Fermi oscillator ( $[a, a^*]_+ = 1$ ) anticommuting with  $b_j, b_k^*$ . We shall fix the physical interpretation of  $[a^*, a]$  by postulating that the electric charge operator is given by

$$Q := \frac{1}{3} \sum_{j=1}^3 b_j^* b_j - a^* a = \frac{1}{2} (B - L - [a^*, a]),$$

$$\text{where } B - L = \frac{2i}{3} (L_{13} + L_{26} + L_{45}) = \frac{1}{3} \sum_j [b_j^*, b_j], \quad (45)$$

and stands for the difference between the baryon and the lepton numbers.  $B - L$  takes the eigenvalues  $\pm\frac{1}{3}$  for (anti)quarks and  $\mp 1$  for (anti)leptons. Demanding that the gauge Lie algebra within  $so(8)$  commutes with both  $\omega_6$  and  $Q$ , we shall further reduce it from  $so(6) \oplus so(2)$  to the rank four Lie subalgebra

$$\mathfrak{g}_4 = su(3)_c \oplus u(1)_Q \oplus u(1)_{B-L} = \{X \in u(4), [X, Q] = 0\}. \quad (46)$$

The knowledge of the charges  $Q, B - L$  along with the color Lie algebra allows us to identify the primitive idempotents of  $C\ell_8$ , given by  $\ell, q_j$  and multiplied by  $aa^*$  or  $a^*a$ , with the fundamental fermions:

$$\ell aa^* = \nu, \ell a^* a = e, q_j aa^* = u_j, q_j a^* a = d_j. \quad (47)$$

The “isotopic doublets”  $(\nu, e)$  and  $(u_j, d_j)$  stand for neutrino/electron and up/down colored quarks. We see, in particular, that the Fock vacuum in  $C\ell_8$  that is associated with the complex structure (33) is identified with the neutrino (as it has no charge and  $a\nu = 0 = b_j\nu$ ). Note that the subalgebra of  $\mathfrak{g}_4$  that annihilates  $\nu$  is the known unbroken gauge Lie algebra  $u(3)$  of the SM:

$$u(3)_{SM} = su(3)_c \oplus u(1)_Q = \{X \in \mathfrak{g}_4; X\nu = 0\}. \quad (48)$$

This picture ignores chirality, which will find its place in  $\mathcal{C}\ell_{10}$  (Section 3.2).

### 3. The Internal Space Subalgebra of $\mathcal{C}\ell_{10}$

#### 3.1. Equivalence Class of Lorentz-like Clifford Algebras

Nature appears to select real Clifford algebras  $\mathcal{C}\ell(s, t)$  of the equivalence class of  $\mathcal{C}\ell(3, 1)$  (with a Lorentz signature in four dimensions) in Élie Cartan's classification (which involves<sup>8</sup> the signs,  $\omega^2(s, t)$  and  $(-1)^{s-t}$ ):

$$\mathcal{C}\ell(s, t) = \mathbb{R}[2^n], \text{ for } s - t = 2(\text{mod } 8), s + t = 2n. \quad (49)$$

They act on  $2^n$  dimensional *Majorana spinors* that irreducibly transform under the *real*  $2^n$  dimensional representation of the spin group  $Spin(s, t)$ . If  $\gamma_1, \dots, \gamma_{2n}$  is the image in  $\mathcal{C}\ell(s, t)$  of an orthonormal basis of the underlying vector space  $\mathbb{R}^{s, t}$ , then the Clifford pseudoscalar defines a complex structure

$$\omega_{s, t} = \gamma_1 \cdots \gamma_{2n}, 2n = s + t, \omega_{s, t}^2 = -1, \quad (50)$$

which commutes with the action of  $Spin(s, t)$ . Upon complexification, the resulting *Dirac spinor* splits into two *inequivalent*  $2^{n-1}$  dimensional complex *Weyl* (or *chiral*) *spinor* representations, which are irreducible over  $\mathbb{C}$  under  $Spin(s, t)$ . The corresponding projectors  $\Pi_L$  and  $\Pi_R$  on the left and right spinors are given in terms of the chirality  $\chi$ , which involves the imaginary unit  $i$ :

$$\begin{aligned} \Pi_L &= \frac{1}{2}(1 - \chi), \quad \Pi_R = \frac{1}{2}(1 + \chi), \quad \chi = i\omega_{s, t}, \\ \chi^2 &= \mathbb{1} \Leftrightarrow \Pi_L^2 = \Pi_L, \quad \Pi_R^2 = \Pi_R, \quad \Pi_L \Pi_R = 0, \quad \Pi_L + \Pi_R = \mathbb{1}. \end{aligned} \quad (51)$$

Another interesting example of the same equivalence class (also with indefinite metric) is the *conformal Clifford algebra*  $\mathcal{C}\ell(4, 2)$  (with isometry group  $O(4, 2)$ ). We shall demonstrate that just as  $\mathcal{C}\ell_{-6}$  was viewed (in Section 2.2) as the *Clifford algebra of the octonions*,  $\mathcal{C}\ell(4, 2)$  plays the role of the *Clifford algebra of the split octonions* (also appearing in bitwistor theory [56]):

$$x = v + V + l(w + W), \quad v, w \in \mathbb{R}, \quad V = iV_1 + jV_2 + kV_3, \quad W = iW_1 + jW_2 + kW_3$$

$$i^2 = j^2 = k^2 = ijk = -1, \quad l^2 = 1, \quad Vl = -lV. \quad (52)$$

Indeed, defining the mapping (cf. (6)),

$$i \rightarrow \gamma_{-1}, \quad j \rightarrow \gamma_0, \quad l \rightarrow \gamma_1, \quad jl \rightarrow \gamma_2, \quad \ell k \rightarrow \gamma_3, \quad \ell i \rightarrow \gamma_4$$

$$[\gamma_\mu, \gamma_\nu]_+ = 2\eta_{\mu\nu}\mathbb{1}, \quad \eta_{11} = \eta_{22} = \eta_{33} = \eta_{44} = 1 = -\eta_{-1,-1} = -\eta_{00} \quad (53)$$

we find that the missing split-octonion (originally, quaternion) imaginary unit  $k$  ( $= ij = -ji$ ) can be identified with the  $\mathcal{C}\ell(4, 2)$  pseudoscalar:

$$\omega_{4, 2} = \gamma_{-1} \gamma_0 \gamma_1 \gamma_2 \gamma_3 \gamma_4 \leftrightarrow k, \quad \omega_{4, 2}^2 = -1, \quad [w_{4, 2}, \gamma_\nu]_+ = 0. \quad (54)$$

The conjugate to the split octonion  $x$  (52) and its norm are

$$x^* = v - V - \ell(w + W), \quad N(x) = xx^* = v^2 + V^2 - w^2 - W^2,$$

so that the isometry group of  $\tilde{\mathbb{O}}$  is  $O(4, 4)$  (in particular, the maximal compact subalgebra  $so(4) \oplus so(4) \subset so(4, 4)$  is spanned by  $\gamma_{jk}, j, k = 1, \dots, 4$  and by  $\omega_{4, 2}, \gamma_\alpha, \alpha = -1, 0$ , and their commutators. The remaining 16 non-compact generators of  $so(4, 4)$  involve the square-one matrices  $\gamma_j, \gamma_\alpha \gamma_j, \gamma_j \omega_{4, 2}$ ).

As we are interested in the geometry of the internal space of the SM acted upon by a compact gauge group, we shall work with (positive or negative) definite Clifford algebras  $\mathcal{Cl}_{2\ell}$ ,  $\ell = 1(\text{mod } 4)$ . The algebra  $\mathcal{Cl}_{-6}$  considered in Section 2 belongs to this family (with  $\ell = -3$ ). For  $\ell = 1$ , we obtain the Clifford algebra of the two-dimensional conformal field theory; the one-dimensional Weyl spinors correspond to analytic and antianalytic functions. Here, we shall argue that for the next allowed value,  $\ell = 5$ , the algebra  $\mathcal{Cl}_{10} = \mathcal{Cl}_4 \hat{\otimes} \mathcal{Cl}_6$  (11) fits the internal space of the SM beautifully if we associate the two factors to the color and flavor degrees of freedom, respectively. We shall strongly restrict the physical interpretation of the generators  $\gamma_{ab}$  ( $= \frac{1}{2} [\gamma_a, \gamma_b]$ ,  $a, b = 1, \dots, 10$ ) of the derivations of  $\mathcal{Cl}_{10}$  by demanding that the splitting (11) of  $\mathcal{Cl}_{10}$  into  $\mathcal{Cl}_4$  and  $\mathcal{Cl}_6$  is preserved. This amounts to selecting a first step of symmetry breakings of the GUT group  $Spin(10)$ , which leads to the semi-simple Pati–Salam group  $(Spin(4) \times Spin(6)) / \mathbb{Z}_2$  (8). Each summand of  $\mathfrak{g}_{PS}$ ,  $so(4)$  and  $so(6)$ , expressed in terms of Fermi creation and annihilation operators, has a distinguished Lie subalgebra,  $u(2)$  and  $u(3)$ , that belongs to  $\mathcal{H}^{1,1}$ . We identify the leptons and quarks with  $u(3)$  singlets and triplets. This identification implements the lepton-quark symmetry alluded to by (1).

### 3.2. $\mathfrak{g}_{SM}$ as Annihilator of Sterile Neutrino

We proceed to extend the complex structure  $J$  (33) and (35) to  $\mathcal{Cl}_{10}$ , expressing, in particular, the electroweak gauge group generators in terms of the fermionic oscillators corresponding to the  $\mathcal{Cl}_4$  factor in (11). To this end, we complement the definition (38) of  $b_j$  by

$$2a_1 = (1 - iJ)\gamma_{10} = \gamma_{10} - i\gamma_9, 2a_2 = \gamma_8 - i\gamma_7 \Rightarrow i\gamma_{78} = [a_2^*, a_2], i\gamma_{910} = [a_1^*, a_1] \quad (55)$$

(where  $\gamma_a$  are given by (36)). In particular,  $(a_2, a_2^*)$  coincide with the unique flavor fermionic oscillator  $(a, a^*)$  (44) of  $\mathcal{Cl}_8$ . They allow us to define two pairs of complementary projectors.

$$\pi_\alpha = a_\alpha a_\alpha^*, \pi'_\alpha = a_\alpha^* a_\alpha = 1 - \pi_\alpha, \alpha = 1, 2, \pi_\alpha \pi'_\alpha = 0, \pi_\alpha + \pi'_\alpha = 1. \quad (56)$$

The three pairs of color  $(p_j, p'_j, j = 1, 2, 3)$  and two pairs of flavor  $(\pi_\alpha, \pi'_\alpha, \alpha = 1, 2)$  projectors give rise to a ( $2^5 = 32$ -dimensional) maximal abelian subalgebra of  $\mathcal{Cl}_{10}$  of the commuting observables. The flavor gauge Lie algebra generators, the left and right chiral isospin components, are expressed in terms of  $a_\alpha^{(*)}$ :

$$\begin{aligned} I_+^L &= a_1^* a_2, I_-^L = a_2^* a_1, [I_+^L, I_-^L] = 2I_3^L = \pi'_1 \pi_2 - \pi_1 \pi'_2 = \pi'_1 - \pi'_2; \\ I_+^R &= a_1 a_2, I_-^R = a_2 a_1^*, [I_+^R, I_-^R] = 2I_3^R = \pi_1 \pi_2 - \pi'_1 \pi'_2 = \pi_2 - \pi'_1. \end{aligned} \quad (57)$$

The *chirality operator*  $\chi = \Pi_R - \Pi_L$  is expressed in terms of the  $\mathcal{Cl}_{10}$  pseudoscalar (as implied by (51) for  $s = 10, t = 0$ ):

$$\chi = i\omega_{10} = i\omega_6 \gamma_{78} \gamma_{910} = (\mathcal{P}' - \mathcal{P})[a_1^*, a_1][a_2, a_2^*] = (\mathcal{P}' - \mathcal{P})(P_1 - P'_1),$$

$$P_1 = (2I_3^L)^2 = \pi'_1 \pi_2 + \pi_1 \pi'_2, P'_1 = (2I_3^R)^2 = \pi_1 \pi_2 + \pi'_1 \pi'_2 = 1 - P_1, \quad (58)$$

so that  $\Pi_L = \mathcal{P}P_1 + \mathcal{P}'P'_1$ ,  $\Pi_R = \mathcal{P}P'_1 + \mathcal{P}'P_1$ . Within the particle subspace  $\mathcal{P}$ , the operator  $P_1$  projects on the left chiral and  $P'_1$  on the right chiral fermions.

The sum  $I_3 = I_3^L + I_3^R$  coincides with the total isospin projection that generates the commutant  $u(1)$  of  $su(4)$  in  $u(4)$ —see the discussion after Equation (34) in Section 2.3. Conversely,  $I_3^L, I_3^R$  appear as chiral projections of  $I_3$ :

$$\begin{aligned} 2I_3 &= 2I_3^L + 2I_3^R = [a_2, a_2^*] = \pi_2 - \pi'_2 = 2Q - (B - L), \\ (2I_3)^2 &= 1, I_3^L = P_1 I_3 P_1, I_3^R = P'_1 I_3 P'_1 \quad (I_3^L I_3^R = 0). \end{aligned} \quad (59)$$

The identification of the vacuum vector  $a_1 a_2 b_1 b_2 b_3$  (annihilated by all  $a_\alpha, b_j$ ) becomes consequential if we demand that this ket-vector is a singlet with respect to the gauge group of the SM. The fact that the left and right isospin cannot vanish simultaneously (because  $(2(I_3^L + I_3^R))^2 = 1$ ) implies that the Lie algebra  $\mathfrak{g}_{SM}$  of the SM should be chiral:

$$\mathfrak{g}_{SM} \subset u(2) \oplus u(3), \text{ where } u(2) = su(2)_L \oplus u(1)_{I_3^R}, u(3) = su(3)_c \oplus u(1)_{B-L}. \quad (60)$$

It is therefore rewarding that we can identify the Fock space vacuum in  $C\ell_{10}$  (given by  $\nu$  of (47) for  $C\ell_8$ ) with the (right handed, hypothetical) *sterile neutrino* (in fact,  $\nu_R$  and its antipode  $\bar{\nu}_L$  do not interact with the gauge bosons):

$$\nu_R = \pi_1 \pi_2 \ell, a_\alpha \nu_R = 0 (= \nu_R a_\alpha^*), b_j \nu_R = 0 \Leftrightarrow \bar{\nu}_L = \pi'_1 \pi'_2 \ell', a_\alpha^* \bar{\nu}_L = 0 \text{ etc.} \quad (61)$$

The role of the electric charge  $Q$  (45) that breaks the  $u(4)$  symmetry of  $\omega_6$  in  $so(8)$  to  $u(3) \oplus u(1)_Q$  is played by the *weak hypercharge*  $Y$  in  $so(10)$ :

$$\frac{1}{2}Y = \frac{1}{3} \sum_{j=1}^3 b_j^* b_j - \frac{1}{2} \sum_{\alpha=1}^2 a_\alpha^* a_\alpha = \frac{1}{2} \sum_{\alpha=1}^2 a_\alpha a_\alpha^* - \frac{1}{3} \sum_{j=1}^3 b_j b_j^*. \quad (62)$$

They both annihilate the respective vacuum state as well as its antipode. This is made obvious by the two forms of  $Y$  in Equation (62) as sums of the normal and antinormal products. By definition,  $Y$  belongs to the center of the broken symmetry subalgebra of  $\mathfrak{g}_{PS}$ . As pointed out in [2] and as will be discussed below in Section 3.3, it gives rise to a superselection rule in the SM.

The significance of choosing the sterile neutrino as a Fock vacuum is summarized by the following:

**Proposition 1.** *The Lie subgroup of  $\mathfrak{g}_{PS}$  (8) that leaves the Fock vacuum  $\nu_R$  (61) invariant is the SM gauge group (13).*

**Proof.** We shall first complete the argument that the maximal Lie subalgebra of  $\mathfrak{g}_{PS}$  annihilating the sterile neutrino is  $\mathfrak{g}_{SM}$ . We have already noted that the Lie subalgebra of  $\mathfrak{g}_{PS}$  for which the vacuum transforms as a singlet is  $u(2) \oplus u(3)$  (60). This follows from the observation that generators involving  $a_1^* a_2^*$  and  $b_j^* b_k^*$  transform  $\nu_R$  into a right-handed electron  $e_R$  and an up quark  $u_R$ , respectively. It remains to analyze the two-dimensional center  $u(1)_{B-L} + u(1)_{I_3^R}$  of this extended algebra.  $\nu_R$  and  $\bar{\nu}_L$  are eigenvectors of both generators with eigenvalues of opposite sign; only multiples of  $Y$  annihilate the sterile neutrino:

$$(2I_3^R - 1)\nu_R = 0 = (B - L + 1)\nu_R, Y = B - L + 2I_3^R \Rightarrow Y\nu_R = 0 = Y\bar{\nu}_L. \quad (63)$$

This establishes the characterization of the Lie algebra  $\mathfrak{g}_{SM}$  as the annihilator of the sterile neutrino. It will be straightforward to extend the result to the SM gauge group (13) after displaying the quantum numbers of the fundamental fermions in the following subsection.  $\square$

### 3.3. Superselection Rules: Restricted Particle Subspace

The weak hypercharge (62) and (63) generates the  $u(1)$  center of the gauge Lie algebra of the SM and hence commutes with all gauge transformations. It is not only conserved in the observed micro processes but even in hypothetical ones, such as a possible proton decay (with a conserved  $B - L$ ), or in the presence of a Majorana neutrino (a coherent superposition of  $\nu_R$  and  $\bar{\nu}_L$ ) that would break  $B - L$  by two units. The weak hypercharge was proposed in [2] as a *superselection rule*, assuming that  $Y$  commutes with all observables. The Jordan algebra of the 32-dimensional space of internal observables of one generation splits into 11 superselection sectors corresponding to the 11 different eigenvalues of  $Y$  (see Appendix to [2]).

Superselection rules (SSR) were introduced by Wick, Wightman, and Wigner [57,58] in 1952. The superselection of the electric charge has been thoroughly discussed in [58] and the review [59]; for more references and a historical survey addressed to philosophers, see [60]. The charge  $Q$  (45) is superselected by the exact symmetry of the SM (otherwise,  $I_{\pm}^L$  do not commute with it). SSRs are also related to measurement theory [52]. SSR and superselection sectors are an essential part of the Doplicher–Haag–Roberts reconstruction of quantum fields from the algebra of observables—see [36].

For all we know, the exact symmetry of the SM is given by the rank four unbroken Lie algebra (obtained from  $\mathfrak{g}_4$  (46) by the substitution  $B - L \rightarrow Y$ ):

$$\mathfrak{a}_4 = su(3)_c \oplus u(1)_Y \oplus u(1)_Q, Q = \frac{1}{2}Y + I_3^L = \frac{1}{3} \sum_{j=1}^3 p'_j - \pi'_2. \quad (64)$$

The states of the fundamental (anti)fermions are given by the primitive idempotents of  $\mathcal{C}\ell_{10}$ , represented by the  $2^5 = 32$  different products of the five pairs of basic projectors  $\pi_{\alpha}^{(\prime)}, p_j^{(\prime)}$  (56) (41). The 16 particles can be labeled by the eigenvalues of the pair of superselected charges  $(Q, Y)$ :

$$\begin{aligned} (\nu_R) &= \ell \pi_1 \pi_2 = (0, 0) = |\nu_R\rangle \langle \nu_R|, (\nu_L) = \ell \pi'_1 \pi'_2 = (0, -1) = |\nu_L\rangle \langle \nu_L|, \\ (e_L) &= \ell \pi_1 \pi'_2 = (-1, -1) = |e_L\rangle \langle e_L|, (e_R) = \ell \pi'_1 \pi'_2 = (-1, -2) = |e_R\rangle \langle e_R|; \\ \ell &= (\nu_L) + (e_L) + (\nu_R) + (e_R) = p_1 p_2 p_3, \ell^2 = \ell, \text{tr } \ell = 4. \end{aligned} \quad (65)$$

$$\begin{aligned} (u_L^j) &= q_j \pi'_1 \pi'_2 = (\frac{2}{3}, \frac{1}{3}) = |u_L^j\rangle \langle u_L^j|, (d_L^j) = q_j \pi_1 \pi'_2 = (-\frac{1}{3}, \frac{1}{3}) = |d_L^j\rangle \langle d_L^j|, \\ (u_R^j) &= q_j \pi_1 \pi_2 = (\frac{2}{3}, \frac{4}{3}) = |u_R^j\rangle \langle u_R^j|, (d_R^j) = q_j \pi'_1 \pi'_2 = (-\frac{1}{3}, -\frac{2}{3}) = |d_R^j\rangle \langle d_R^j|; \\ q_j &= (u_L^j) + (d_L^j) + (u_R^j) + (d_R^j) = p_j p'_k p'_\ell, q_i q_j = \delta_{ij} q_j, \text{tr } q_j = 4 \end{aligned} \quad (66)$$

$(j, k, \ell) \in \text{Perm}(1, 2, 3)$ ,  $q = q_1 + q_2 + q_3 = q^2$ ,  $\text{tr } q = 12$  (as the color is unobservable, we do not bother to assign to it eigenvalues of the diagonal operators  $i\gamma_{13}, i\gamma_{26}, i\gamma_{45}$  that would replace the index  $j$ ). Note that chirality in the particle subspace  $\mathcal{P}\chi = \chi\mathcal{P}$  is determined by the hypercharge:

$$\mathcal{P}\chi = \mathcal{P}(\Pi_R - \Pi_L) = \mathcal{P}(-1)^{3Y}. \quad (67)$$

The charges  $(Q, Y)$  for the corresponding antiparticles have the opposite sign. The spectrum of  $Y$  and of  $2I_3^L = 2Q - Y$ , together with the analysis of [38], allow us to complete the group theoretic version of Proposition 1.

**Remark 2.** The factorization of the primitive idempotents (65) and (66) into bra and kets involves choices. We demand, following [2], that they are Hermitian conjugate elements of  $\mathcal{C}\ell_{10}$ , homogeneous in  $a_{\alpha}^{(*)}$  and  $b_j^{(*)}$  such that the kets corresponding to a left(right) chiral particle contains an odd (or even) number of factors. Choosing then  $|\nu_R\rangle = a_1 a_2 \ell, |\nu_L\rangle = a_1^* |\nu_R\rangle$ , we find:

$$\begin{aligned} \langle \nu_R | &= \ell a_2^* a_1^* \Rightarrow (\nu_R) = \pi_1 \pi_2 \ell, |\nu_L\rangle = \pi'_1 a_2 \ell, \\ |e_L\rangle &= I_-^L |\nu_L\rangle = -a_1 \pi'_2 \ell, |e_R\rangle = -a_1^* |e_L\rangle = \pi'_1 \pi'_2 \ell = I_-^R |\nu_R\rangle; \\ |d_L^j\rangle &= \pi_1 a_2^* q_j, |u_L^j\rangle = I_+^L |d_L^j\rangle = a_1^* \pi_2 q_j, \\ |d_R^j\rangle &= a_1^* |d_L^j\rangle = a_1^* a_2^* q_j, u_R^j = a_1 |u_L^j\rangle = \pi_1 \pi_2 q_j, \end{aligned} \quad (68)$$

$q_j = p_j p'_k p'_\ell, j, k, \ell \in \text{Perm}(1, 2, 3)$ , i.e.  $q_1 = p_1 p'_2 p'_3 = p_1 p'_3 p'_2$ , etc. We note that all above kets as well as all primitive idempotents (65) (66) obey a system of five equations (specific for each particle),  $a_{\alpha} |\nu_R\rangle = 0 = b_j |\nu_R\rangle, a_1^* |\nu_L\rangle = a_2 |\nu_L\rangle = 0 = b_j |\nu_L\rangle, \alpha = 1, 2, j = 1, 2, 3$ , etc., so that they are minimal right ideals in agreement with the philosophy of Furey [20].

The fact that  $\nu_R, \bar{\nu}_L$  are not distinguished by the superselected charges has a physical implication; one can consider their coherent superposition as in the now popular theory of a (hypothetical) Majorana neutrino. This suggests the introduction of a *restricted* 15-dimensional particle subspace with projector

$$\mathcal{P}_r = \mathcal{P} - (\nu_R) = q + \ell_r, \quad \ell_r = \ell(1 - \pi_1 \pi_2). \quad (69)$$

Theories whose field algebra is a tensor product of a Dirac spinor bundle on a space-time manifold with a finite dimensional internal space usually encounter the problem of fermion doubling [61] (still discussed over 20 years later, [62]). It was proposed in [34] as a remedy to consider the algebra  $\mathcal{P}C\ell_{10}\mathcal{P}$ , where  $\mathcal{P}$  is the projector (42) on the 16 dimensional particle subspace (including the hypothetical right-handed sterile neutrino). It is important—and will be essential in the treatment of the Higgs field (Section 4)—that the operators  $a_\alpha^{(*)}$  and  $b_j^{(*)}$  behave quite differently under particle projection. While  $a_\alpha^{(*)}$  commute with  $\mathcal{P}$  so that

$$\mathcal{P}a_\alpha^{(*)}\mathcal{P} = a_\alpha^{(*)}\mathcal{P} = \mathcal{P}a_\alpha^{(*)}, [\mathcal{P}a_\alpha^*, \mathcal{P}a_\beta]_+ = \mathcal{P}\delta_{\alpha\beta}, \quad (70)$$

the 2-sided particle projection of  $b_j^{(*)}$  vanishes:

$$\mathcal{P}b_j\mathcal{P} = 0 = \mathcal{P}b_j^*\mathcal{P}. \quad (71)$$

Accordingly, while the generators (57) of the (electroweak) flavor “left-right symmetry”  $su(2)_L \oplus su(2)_R$  just get multiplied by  $\mathcal{P}$ , the particle subspace projections of the  $su(3)_c$  generators take a modified form:

$$\begin{aligned} \mathcal{P}b_j b_k^* \mathcal{P} &= b_j b_k^* p'_\ell =: B_{jk} \text{ for } (j, k, \ell) \in \text{Perm}(1, 2, 3), B_{jj} - B_{kk} := q_j - q_k; \\ T_a &= \frac{1}{2} B_{jk} \lambda_a^{kj}, \lambda_a \in \mathcal{H}_3(\mathbb{C}), \text{tr } \lambda_a = 0, \text{tr } \lambda_a \lambda_b = 2\delta_{ab}, a, b = 1, \dots, 8, \end{aligned} \quad (72)$$

but still obey the same CR. It makes sense to separately consider the gauge Lie algebra in the lepton and quark sectors (or the factors  $C\ell_4$  and  $C\ell_6$  in  $C\ell_{10}$ ), noting that  $\mathcal{P}(B - L) = -1$  for leptons and  $\mathcal{P}(B - L) = \frac{1}{3}$  for quarks. It is particularly appropriate to treat the lepton sector by itself when using the restricted particle space as it is there that the flavor oscillators  $a_\alpha^{(*)}$  are also modified:

$$\ell_r a_1^{(*)} \ell_r = a_1^{(*)} \pi'_2 =: A_1^{(*)}, \ell_r a_2^{(*)} \ell_r = a_2^{(*)} \pi'_1 =: A_2^{(*)}. \quad (73)$$

The operators  $A_\alpha^{(*)}$  provide a realization of the four odd generators of the smallest simple Lie superalgebra,  $sl(2|1)$ , whose even part is  $su(2)_L \oplus u(1)_Y$  (for a detailed identification with the standard definition of  $sl(2|1)$ , see Section 3 of [2]). The non-vanishing anticommutators of  $A_\alpha^{(*)}$  are:

$$\begin{aligned} [A_1, A_1^*]_+ &= \pi'_2 = -Q, [A_2, A_2^*]_+ = \pi'_1 = Q - Y, \\ [A_1^*, A_2]_+ &= a_2 a_1^* = -I_+, [A_1, A_2^*]_+ = -I_-; [I_+, I_-] = 2I_3 = 2Q - Y \end{aligned} \quad (74)$$

(where we are omitting the superscript L on  $I_a$ ). We shall apply the odd generators  $A_\alpha^{(*)}$  in defining the Higgs part of a superconnection in Section 4. The *minimal associative envelope*  $\mathcal{A}_\ell$  of  $sl(2|1) \subset C\ell_4$  is nine-dimensional; it contains on top of  $A_\alpha^{(*)}$  and their anticommutators (74) the projector  $A_1^* A_1 = A_2^* A_2 = \pi'_1 \pi'_2 \in C\ell_4$ . The resulting *internal space algebra* that ignores the sterile neutrinos is the direct sum

$$\mathcal{A} = \mathcal{A}_\ell \otimes \ell \oplus C\ell_4 \otimes \mathcal{A}_q, \quad \mathcal{A}_q = q C\ell_6^0 q. \quad (75)$$

Here,  $\mathcal{A}_q$  is effectively the nine-dimensional associative envelope of  $u(3) \subset C\ell_6^0$ .

### 3.4. Complex Structure Associated with $R_7$ : A Comment

Inspired by [24,46,63], we shall display and discuss the symmetry subalgebras of  $\mathcal{C}\ell_n, n = 8, 9, 10$  of the complex structure generated by the Clifford pseudoscalar  $\omega_6^R$  corresponding to the right action of the octonions:

$$\omega_6^R = \gamma_1^R \cdots \gamma_6^R \text{ for } \gamma_\alpha^R = \epsilon \otimes R_\alpha \quad \alpha = 1, \dots, 7. \quad (76)$$

Written in terms of the color projectors  $p_j$  and  $p'_j$ , the Hermitian pseudoscalar  $i\omega_6^R$  assumes the form:

$$\begin{aligned} i\omega_6^R &= \frac{1}{2}(\mathcal{P}' - \mathcal{P} - 3(B - L)) = \ell + q' - \ell' - q, \\ \ell' &= p'_1 p'_2 p'_3, q' = \sum_{j=1}^3 q'_j, q'_1 = p'_1 p_2 p_3, q'_2 = p_1 p'_2 p_3, q'_3 = p_1 p_2 p'_3 \end{aligned} \quad (77)$$

where we have used

$$L = \ell - \ell', \quad 3B = q - q'. \quad (78)$$

While the term  $\mathcal{P}' - \mathcal{P}$  (42) commutes with the entire derivation algebra  $spin(6) = su(4)$  of  $\mathcal{C}\ell_6$ , the centralizer of  $B - L$  in  $su(4)$  is  $u(3)$ —see Proposition A2 in Appendix A. It follows that the commutant of  $\omega_6^R$  in  $so(8)$  is  $u(3) \oplus u(1)$ , while its centralizer in  $so(9)$  is the gauge Lie algebra  $\mathfrak{g}_{SM} = su(3) \oplus su(2) \oplus u(1)$  of the SM; finally, in  $so(10)$ ,  $\omega_6^R$  is invariant under the left-right symmetric extension of  $\mathfrak{g}_{SM}$  [24,63],

$$\mathfrak{g}_{LR} = su(3)_c \oplus su(2)_L \oplus su(2)_R \oplus u(1)_{B-L}. \quad (79)$$

Furthermore, as proven in [46], the subgroup of  $Spin(9)$  that leaves  $\omega_6^R$  invariant is precisely the gauge group<sup>9</sup>  $G_{SM} = S(U(2) \times U(3))$  (13) of the SM (with the appropriate  $\mathbb{Z}_6$  factored out). One is then tempted to assume that  $\mathcal{C}\ell_9$ , the associative envelope of the Jordan algebra  $J_2^8 = \mathcal{H}_2(\mathbb{O})$ , may play the role of the internal algebra of the SM, corresponding to one generation of fundamental fermions, with  $Spin(9)$  as a GUT group [27,28,33]. We shall demonstrate that although  $G_{SM}$  appears as a subgroup of  $Spin(9)$ , its representation, obtained by restricting the (unique) spinor IR 16 of  $Spin(9)$  to  $S(U(2) \times U(3))$  only involves  $SU(2)$  doublets, so it has no room for  $(e_R), (u_R), (d_R)$  (65) (66). We shall see how this comes about when restricting the realization (57) of  $\mathbf{I}^L$  and  $\mathbf{I}^R$  to  $Spin(9) \subset \mathcal{C}\ell_9$ . It is clear from (57) that only the sum  $a_1 + a_1^* = \gamma_9$  (not  $a_1$  and  $a_1^*$  separately) belongs to  $\mathcal{C}\ell_9$ . So, the  $su(2)$  subalgebra of  $spin(9)$  corresponds to the diagonal embedding  $su(2) \hookrightarrow su(2)_L \oplus su(2)_R$ :

$$\begin{aligned} I_+ &= I_+^L + I_+^R = (a_1^* + a_1) a_2 = \gamma_9 a_2, \quad I_- = I_-^L + I_-^R = a_2^* \gamma_9 \\ 2I_3 &= 2I_3^L + 2I_3^R = [a_2, a_2^*] = \pi_2 - \pi_2'. \end{aligned} \quad (80)$$

In other words, the spinorial IR 16 of  $Spin(9)$  is an eigensubspace of the projector  $P_1 = (2I_3^L)^2$ . It consists of four  $SU(2)_L$  particle doublets and their right chiral antiparticles. More generally, the only simple orthogonal groups with a pair of inequivalent complex that conjugate fundamental IRs are  $Spin(4n + 2)$  (see, e.g., [64], Proposition 5.2, p. 571). They include  $Spin(10)$  but not  $Spin(9)$ .

A direct description of the IR 16<sub>L</sub> of  $Spin(10)$  acting on  $\mathbb{CH} \otimes \mathbb{CO}$  is given in [23] (Here,  $\mathbb{CH}$  and  $\mathbb{CO}$  are a short hand for the complexified quaternions and octonions:  $\mathbb{CH} := \mathbb{C} \otimes_{\mathbb{R}} \mathbb{H}$ ). The right action of  $\mathbb{CH}$  on elements of  $\mathbb{CH} \otimes \mathbb{CO}$ , which commutes with the left-acting  $spin(10)$ , is interpreted in [23] as the Lorentz ( $SL(2, \mathbb{C})$ ) transformation of the (unconstrained) two-component Weyl spinors.

The left-right symmetric extension  $\mathfrak{g}_{LR}$  (78) of  $\mathfrak{g}_{SM}$  has a long history, starting with [65] and vividly (with an admitted bias) told in [66]; it has been recently invigorated in [67,68]. The group  $\mathfrak{g}_{LR}$  was derived by Boyle [47] starting with the group  $E_6$  of determinant

preserving linear automorphisms of the complexified Albert algebra  $\mathbb{C}J_3^8$  and following the procedure of [35].

#### 4. Particle Subspace and the Higgs Field

##### 4.1. The Higgs as a Scalar Part of a Superconnection

The space of differential forms  $\Lambda^* = \Lambda^0 + \Lambda^1 + \Lambda^2 + \dots$  can be viewed as a  $\mathbb{Z}_2$  graded setting  $\Lambda_{ev} = \Lambda_0 + \Lambda_2 + \dots$ ,  $\Lambda_{od} = \Lambda_1 + \Lambda_3 + \dots$ . Let  $M = M_0 + M_1$  be a  $\mathbb{Z}_2$  graded matrix algebra. A superconnection in the sense of Quillen [69,70] is an element of  $\Lambda_{ev} \otimes M_1 + \Lambda_{od} \otimes M_0$ , the odd part of the tensor product  $\Lambda^* \otimes M$ ; a critical review of the convoluted history of this notion and its physical implications is given in Section 4 of [71] (one should also mention the neat exposition of [72] in the context of the Weinberg–Salam model with two Higgs doublets).

Let  $D$  be the Yang–Mills connection one-form of the SM,

$$D = dx^\mu (\partial_\mu + A_\mu(x)),$$

$$iA_\mu = W_\mu^+ I_+^L + W_\mu^- I_-^L + W_\mu^3 I_3^L + \frac{N}{2} Y B_\mu + G_\mu^a T_a, \quad (81)$$

where  $Y$ ,  $I^L$ , and  $T_a$  are given by (62), (57), and (72), respectively;  $G_\mu^a$  is the gluon field, and  $W_\mu$  and  $B_\mu$  provide an orthonormal basis of electroweak gauge bosons; the normalization constant  $N$  will be fixed in Equation (93) below. Then, one defines a superconnection  $\mathbb{D}$  in [34] involving the chirality  $\chi$  (60) by

$$\mathbb{D} = \chi D + \Phi, \quad \Phi = \sum_\alpha (\phi_\alpha a_\alpha^* - \bar{\phi}_\alpha a_\alpha) \in \mathcal{P} \mathcal{C} \ell_{10}^1 \mathcal{P} = \mathcal{P} \mathcal{C} \ell_4^1 \quad (82)$$

(we omit, for the time being, the projector  $\mathcal{P}$  in  $A_\mu$  and  $\Phi$ ). The last equation follows from (71); the projection on the particle subspace kills the odd part of  $\mathcal{C} \ell_6$ , thus ensuring that the quarks' color symmetry remains unbroken. The factor  $\chi$  (first introduced in this context in [71]) ensures the anticommutativity of  $\Phi$  and  $\chi D$  without changing the Yang–Mills curvature  $D^2 = (\chi D)^2$ .

The projector  $\mathcal{P}$  (42) on the 16-dimensional particle subspace that includes the hypothetical right chiral neutrino (and is implicit in (82)) was adopted in [34]. By contrast, particles are only distinguished from antiparticles in [2] if they have different quantum numbers in the Lie algebra of the SM

$$\mathfrak{g}_{SM} = su(3)_c \oplus su(2)_L \oplus u(1)_Y. \quad (83)$$

Thus, in [2],  $\mathcal{P}$  is replaced by the 15-dimensional projector  $\mathcal{P}_r = q + \ell_r$  (69). We have seen that the projected odd operators  $A_\alpha^{(*)} = \ell_r a_\alpha^{(*)} \ell_r$  give rise to a realization of the four odd elements of the eight-dimensional simple Lie superalgebra  $sl(2|1)$  whose even part is the four-dimensional Lie algebra  $u(2)$  of the Weinberg–Salam model of the electroweak interactions. It is precisely the Lie superalgebra that was proposed in 1979 independently by Ne'eman and by Fairlie [44,45] (and denoted by them  $su(2|1)$ ) in their attempt to unify  $su(2)_L$  with  $u(1)_Y$  (and explain the spectrum of the weak hypercharge). Let us stress that the representation space of  $sl(2|1)$  consists of the observed left and right chiral leptons (rather than of bosons and fermions like in the popular speculative theories in which the superpartners are hypothetical). Note that the trace of  $Y$  on negative chirality leptons ( $\nu_L, e_L$ ) is equal to its eigenvalue on the unique positive chirality state ( $e_R$ ) (equal to  $-2$ ) so that only the supertrace of  $Y$  vanishes on the lepton (as well as on the quark) space. This observation is useful in the treatment of anomaly cancellation (cf. [73]).

We shall sketch the main steps in the application of the superconnection (82) to the bosonic sector of the SM, emphasizing specific additional hypotheses used on the way (for detailed calculations, see [2]).

The canonical curvature form

$$\mathbb{D}^2 = D^2 + \chi[D, \Phi] + \Phi^2, \quad [D, \Phi] = dx^\mu(\partial_\mu \Phi + [A_\mu, \Phi]) \quad (84)$$

satisfies the *Bianchi identity*

$$\mathbb{D}\mathbb{D}^2 = \mathbb{D}^2\mathbb{D} \quad (\Rightarrow \chi(d\Phi^2 + [A, \phi^2] + [\Phi, D\Phi]_+) = 0), \quad (85)$$

equivalent to the (super) Jacobi identity of our Lie superalgebra. It is important that the Bianchi identity, which is needed for the consistency of the theory, still holds if we add to  $\mathbb{D}^2$  a constant matrix term with a similar structure. Without such a term, the Higgs potential would be a multiple of  $\text{Tr } \Phi^4$  and would only have a trivial minimum at  $\Phi = 0$ , yielding no symmetry breaking. The projected form of  $\Phi$  (82) and hence the admissible constant matrix addition to  $\Phi^2$  depends on whether we use the projector  $\mathcal{P}$  (as in [34]) or  $P_r$  (as in [2]). In the first case we just replace  $a_\alpha^{(*)}$  with  $a_\alpha^{(*)}\mathcal{P}$ . In the second, however, the odd generators for leptons and quarks differ, and we set:

$$\Phi = \ell(\phi_1 A_1^* - \bar{\phi}_1 A_1 + \phi_2 A_2^* - \bar{\phi}_2 A_2) + \rho q \sum_{\alpha=1}^2 (\phi_\alpha a_\alpha^* - \bar{\phi}_\alpha a_\alpha), \quad (86)$$

where  $\rho$  (like  $N$  in (81)) is a normalization constant that will be fixed later. Recalling that  $\ell$  and  $q$  are mutually orthogonal ( $\ell q = 0 = q\ell$ ,  $\ell + q = \mathcal{P}$ ), we find

$$\begin{aligned} \Phi^2 &= \ell(\phi_1 \bar{\phi}_2 I_+^L + \bar{\phi}_1 \phi_2 I_-^L - \phi_1 \bar{\phi}_1 \pi'_2 - \phi_2 \bar{\phi}_2 \pi'_1) \\ &\quad - \rho^2 q(\phi_1 \bar{\phi}_1 + \phi_2 \bar{\phi}_2) \quad (\phi_\alpha = \phi_\alpha(x)). \end{aligned} \quad (87)$$

This suggests defining the SM field strength (the extended curvature form) as

$$\mathbb{F} = i(\mathbb{D}^2 + \hat{m}^2), \quad \hat{m}^2 = m^2(\ell(1 - \pi_1 \pi_2) + \rho^2 q) \quad (88)$$

(while  $\hat{m}^2 = m^2\mathcal{P}$  for the 16-dimensional particle subspace of [34]).

#### 4.2. Higgs Potential and Mass Formulae

This yields the bosonic Lagrangian (setting  $\text{Tr } X = \frac{1}{4}\text{tr } X$ —see [2])

$$\mathcal{L}(x) = \text{Tr} \left\{ \frac{1}{2} F_{\mu\nu} F^{\mu\nu} - (\partial_\mu \Phi + [A_\mu, \Phi])(\partial^\mu \Phi + [A^\mu, \Phi]) \right\} - V(\Phi) \quad (89)$$

where the Higgs potential  $V(\Phi)$  is given by (noting that  $\text{Tr } \ell_r = \frac{3}{4}$ ):

$$V(\Phi) = \text{Tr}(\hat{m}^2 + \Phi^2)^2 - \frac{1}{4}m^4 = \frac{1}{2}(1 + 6\rho^4)(\phi\bar{\phi} - m^2)^2. \quad (90)$$

Minimizing  $V(\Phi)$  gives the expectation value of the square of  $\phi = (\phi_1, \phi_2)$ :

$$\langle \phi \bar{\phi} \rangle = \phi_1^m \bar{\phi}_1^m + \phi_2^m \bar{\phi}_2^m = m^2, \quad \text{for } \Phi^m = \sum_{\alpha=1}^2 \phi_\alpha^m a_\alpha^* (\ell \pi'_{3-\alpha} + \rho q) + c \cdot c \quad (91)$$

(the superscript  $m$  indicates that  $\phi_\alpha$  takes a constant in  $x$  values depending on the mass parameter  $m$ ). The mass spectrum of the gauge bosons is determined by the term  $-\text{Tr}[A_\mu, \Phi][A^\mu, \Phi]$  of the Lagrangian (89), with  $A_\mu$  and  $\Phi$  given by (81) and (86) for  $\phi_\alpha = \phi_\alpha^m$ . The gluon field  $G_\mu$  does not contribute to the mass term as  $\mathcal{C}\ell_6^0$  commutes with  $\mathcal{C}\ell_4^1$ . The resulting quadratic form is generally not degenerate, so it does not yield a massless photon. It does so, however, if we assume that  $\Phi^m$  is electrically neutral (i.e., commutes with  $Q$  (64)):

$$[\Phi^m, Q] = 0 \Rightarrow \phi_2^m = 0 \quad (= \bar{\phi}_2^m). \quad (92)$$

The normalization constant  $N$  ( $=\tan \theta_w$ ) is fixed by assuming that  $2I_3^L$  and  $NY$  are equally normalized:

$$N^2 = \frac{\text{Tr}(2I_3^L)^2}{\text{Tr} Y^2} = \frac{3}{5} \left( = (\tan \theta_w)^2 \Leftrightarrow \sin^2 \theta_w = \frac{3}{8} \right). \quad (93)$$

As  $Y(v_R) = 0 = I_3^L(v_R)$ , this result for the “Weinberg angle at unification scale” is independent of whether we use  $\mathcal{P}$  or  $\mathcal{P}_r$ . If one takes the trace over the leptonic subspace, the result would have been  $(\tan \theta_w)^2 = \frac{1}{3} (\Rightarrow \sin \theta_w = \frac{1}{2}$ , [44]), which is closer to the measured low-energy value.

Demanding, similarly, that the leptonic contribution to  $\Phi^2$  is the same as that for a colored quark (which gives  $\rho = 1$  for the projector  $\mathcal{P}$ ), we find

$$\rho^2 = \frac{\text{Tr}(\ell(1 - \pi_1 \pi_2) \Phi^2)}{\text{Tr} q_j \Phi^2} = \frac{\text{Tr}(\pi'_1 \pi'_2 \phi \bar{\phi} + \pi'_1 \pi_2 \phi_2 \bar{\phi}_2 + \pi_1 \pi'_2 \phi_1 \bar{\phi}_1)}{4 \phi \bar{\phi}} = \frac{1}{2}. \quad (94)$$

The ratio  $\frac{m_H^2}{m_W^2}$ , on the other hand, is found to be

$$\frac{m_H^2}{m_W^2} = 4 \frac{1 + 6\rho^4}{1 + 6\rho^2} = \begin{cases} 4 \text{ for } \rho^2 = 1 \\ \frac{5}{2} \text{ for } \rho^2 = \frac{1}{2} \end{cases}. \quad (95)$$

The result of [2], which is much closer to the observed value, can also be written in the form  $m_H^2 = 4 \cos^2 \theta_W m_W^2$ , where  $\theta_W$  is the theoretical Weinberg angle (93).

## 5. Outlook

### 5.1. Coming to $\mathcal{Cl}_{10}$

The search for an appropriate choice of a finite dimensional algebra suited to represent the internal space  $\mathcal{F}$  of the SM is still ongoing. The road to the choice of  $\mathcal{Cl}_{10}$ , our first step to the restricted algebra  $\mathcal{A}$  (75), has been convoluted.

In view of the lepton-quark correspondence that is embodied in the splitting (1) of the normed division algebra  $\mathbb{O}$  of the octonions, the choice of Dubois-Violette [25] of the exceptional Jordan algebra  $\mathcal{F} = \mathcal{H}_3(\mathbb{O})$  (7) appeared to be particularly attractive. We realized [27,28,35] that the simpler to work with subalgebra

$$J_2^8 = \mathcal{H}_2(\mathbb{O}) \subset \mathcal{H}_3(\mathbb{O}) = J_3^8 \quad (96)$$

corresponds to the observables of one generation of fundamental fermions. The associative envelope of  $J_2^8$  is  $\mathcal{Cl}_9 = \mathbb{R}[16] \oplus \mathbb{R}[16]$  with the associated symmetry group  $Spin(9)$ . It was proven in [35] that the SM gauge group  $G_{SM}$  (13) is the intersection of  $Spin(9)$  with the subgroup of the automorphism group  $F_4$  of  $J_3^8$  that preserves the splitting (1); that is, the group  $\frac{SU(3) \times SU(3)}{\mathbb{Z}_3} \subset F_4$ .

We were thus inclined to identify  $Spin(9)$  as the most economic GUT group. As demonstrated in Section 3.4, however, the restriction of the spinor IR 16 of  $Spin(9)$  to its subgroup  $G_{SM}$  gives room to only half of the fundamental fermions, the  $SU(2)_L$  doublets; the right chiral singlets,  $e_R, u_R, d_R$ , are left out. It was then recognized that the (octonionic) Clifford algebra  $\mathcal{Cl}_{10}$  does the job. The particle interpretation of  $\mathcal{Cl}_{10}$  is dictated by the choice of a (maximal) set of five commuting operators in the Pati–Salam Lie subalgebra of  $so(10)$  that leaves our complex structure invariant. This led us to presenting all chiral leptons and quarks of one generation as mutually orthogonal idempotents (65) and (66).

Furey [21] arrived (back in 2018) at the tensor product  $\mathcal{Cl}_4 \hat{\otimes} \mathcal{Cl}_6$  (11) following the  $\mathbb{R} \otimes \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$  road. In fact, Clifford algebras have arisen as an outgrowth of Grassmann algebras and quaternions<sup>10</sup>. The 32 products  $e_a e_\nu (= \varepsilon_\nu e_a)$ ,  $a = 1, \dots, 8$  ( $e_8 = \mathbb{1}$ ),  $\nu = 0, 1, 2, 3$  of octonion and quaternion units may serve as components of a  $Spin(10)$  Dirac (bi)spinor, acted upon by  $\mathcal{Cl}_{10}$  (with generators (36) involving the operators  $L_\alpha$ )—cf. [23].

### 5.2. Two Ways to Avoid Fermion Doubling

There are two inequivalent possibilities to avoid fermion doubling within  $C\ell_{10}$ . One, which was adopted in [2,34] and in Section 3.3 of the present survey, consists in projecting on the particle subspace, which incorporates four  $SU(2)_L$  doublets and eight  $SU(2)_L$  (right chiral) singlets, with projector (42)

$$\mathcal{P} = \ell + q = \frac{1 - i\omega_6}{2}, \quad \ell = p_1 p_2 p_3, \quad q = q_1 + q_2 + q_3 \quad (97)$$

(see (70)–(72)). Here,  $\omega_6$  is the  $C\ell_6$  pseudoscalar, the distinguished complex structure used in [24] as a first step in the “cascade of symmetry breakings”. The particle projector (97) is only invariant under the Pati–Salam subgroup (8) of  $Spin(10)$ . The more popular alternative, which was adopted in [23], projects on the left chiral fermions (four particle doublets and eight antiparticle singlets) with projector  $\Pi_L$ , which is defined in terms of the  $C\ell_{10}$  chirality  $\chi = i\omega_{10}$ :

$$\Pi_L = \frac{1 - \chi}{2} = \mathcal{P}P_1 + \mathcal{P}'P'_1 \quad (\mathcal{P} + \mathcal{P}' = 1 = P_1 + P'_1), \quad (98)$$

invariant under the entire  $Spin(10)$ ; here,  $P_1$  projects on  $SU(2)_L$  doublets (cf. (58)). The components of the resulting  $\mathbf{16}_L$  are viewed in [23] as Weyl spinors; the right action of (complexified) quaternions (that commutes with the left  $Spin(10)$  action) is interpreted as an  $sl(2, \mathbb{C})$  (Lorentz) transformation.

The difference of the two approaches, which can be labeled by the projectors  $\mathcal{P}$  and  $\Pi_L$  (on the left and right particles and on the left particles and antiparticles, respectively), has implications in the treatment of the generalized connection (including the Higgs) and anomalies. Thus, for the  $\Pi_L$  (anti)leptons  $(\nu_L, e_L), \bar{e}_L, \bar{\nu}_L$  we have vanishing trace of the hypercharge,  $\text{tr } \Pi_L Y = 0$ . For  $\mathcal{P}$  leptons,  $(\nu_L, e_L), \nu_R, e_R$ , the traces of the left and right chiral hypercharge are equal:  $\text{tr}(\mathcal{P}\Pi_L Y) = -2 = \text{tr}(\mathcal{P}\Pi_R Y)$ , so that, as noted in Section 4.1, only the supertrace vanishes in this case. The associated Lie superalgebra fits Quillen’s notion of superconnection ideally. A real “physical difference” only appears under the assumption that the electroweak hypercharge is superselected and  $\mathcal{P}$  is replaced by the restricted projector  $\mathcal{P}_r$  on the 15-dimensional particle subspace (with the sterile neutrino  $\nu_R$  and with the vanishing hypercharge excluded). Then, the leptonic (electroweak) part of the SM is governed by the Lie superalgebra  $sl(2|1)$ , whose four odd generators are given by third-degree monomials in  $a_\alpha^{(*)}$ , the  $C\ell_4$  Fermi oscillators. The replacement of  $\ell$  by  $\ell_r$  breaks the quark-lepton symmetry; while each colored quark  $q_j$  appears in four flavors, the colorless leptons number just three. This yields a relative normalization factor between the quark and leptonic projection of the Higgs field and allows us to derive (in [2]) the relation

$$m_H^2 = \frac{5}{2} m_W^2 = 4 \cos^2 \theta_{\text{th}} m_W^2, \quad (99)$$

where  $\theta_{\text{th}}$  is the *theoretical* Weinberg angle, such that  $\text{tg}^2 \theta_W = \frac{3}{5}$ . The relation (99) is satisfied within 1% accuracy by the observed Higgs and  $W^\pm$  masses.

### 5.3. Summary and Discussion; a Challenge

After the pioneering work of Feza Gürsey and the collaborators in the 1970s, Geoffrey Dixon devoted over 30 years to division algebras, which is followed by Cohl Furey since the 2010s. The Clifford algebra approach to unification, coupled with fermionic creation and annihilation operators, has also been pursued since the late 1970s by the Italian group around Roberto Casalbuoni. The notion of superconnection was anticipated and applied to the Weinberg–Salam model during the first decade of the creation of the SM as well. Thus, the basic ingredients of our endeavor have been with us for some 50 years. The pretended new features of the present survey concern certain details. Here belong:

- The interpretation of the Clifford pseudoscalar  $\omega_6$  as  $i(\mathcal{P} - \mathcal{P}')$ , where  $\mathcal{P}$  and  $\mathcal{P}'$  are the particle and antiparticle projection operators, respectively.

- The realization that the projected Clifford algebra

$$\mathcal{P}C\ell_{10}\mathcal{P} = C\ell_4 \otimes C\ell_6^0, \quad (100)$$

only involves the even part  $C\ell_6^0$  of  $C\ell_6$ ; coupled with the assignment of the Higgs field to the odd part,  $C\ell_4^1$  of the first factor explains the symmetry breaking of the (electroweak) flavor symmetry while preserving the color gauge group.

- Exhibiting the role of the sterile neutrino (of the first generation of fundamental fermions) as the vacuum state of the theory. The gauge group of the SM is identified as the maximal subgroup of the Pati–Salam group  $G_{PS}$  (8) that leaves  $\nu_R$  invariant.

- Singling out the *reduced* 15-dimensional *particle subspace* yields a relation between the Higgs and the W boson masses and the theoretical Weinberg angle satisfied within 1% accuracy.

What is missing for completing the “Algebraic Design of Physics”—to quote from the title of the 1994 book by Geoffrey Dixon—is a true understanding of the *three generations* of fundamental fermions. None of the attempts in this direction [18,25,30,47,74] has brought a clear success. The exceptional Jordan algebra  $J_3^8 = \mathcal{H}_3(\mathbb{O})$  (7) with its built-in triality was first proposed to this end in [25] (continued in [33]); in its straightforward interpretation, however, it corresponds to the triple coupling of left and right chiral spinors with a vector in internal space rather than to three generations of fermions. As recalled in (Section 5.2 of) [30], any finite dimensional unital module over  $\mathcal{H}_3(\mathbb{O})$  has the (disappointingly unimaginative) form of a tensor product of  $\mathcal{H}_3(\mathbb{O})$  with a finite dimensional real vector space  $E$ . It was further suggested there that the dimension of  $E$  should be divisible by three, but the idea was not pursued any further. Boyle [47] proposed considering the complexified exceptional Jordan algebra, whose group of determinant preserving linear automorphisms is the compact form of  $E_6$ . This led to a promising left-right symmetric extension of the gauge group of the SM, but the discussion has not yet shed new light on the three generation problem. Yet another development based on the study of indecomposable representations of Lie superalgebras can be traced back from [75], but only the mathematical machinery has been discussed so far.

**Funding:** This research received no external funding.

**Data Availability Statement:** No new data were created or analyzed in this study. Data sharing is not applicable to this article.

**Acknowledgments:** Discussions with John Baez, Latham Boyle, Michel Dubois-Violette, Cohl Furey, and Kirill Krasnov are gratefully acknowledged. It is a pleasure to thank Cécile Gourges for her expert typing. The author thanks the IHES for their hospitality and support during the course of this work. The present exposition differs from the one in IHES/P/22/01; it owes much to the critical remarks on the earlier version made by Cohl Furey and especially by Kirill Krasnov. The author thanks the anonymous referees for their careful reading of the manuscript and constructive critical remarks that helped to improve the presentation in the final version of the paper. I thank Dimitar Nedanovski who helped me cope with the Publishers’ requirements.

**Conflicts of Interest:** The author declares no conflict of interest.

## Appendix A

### Appendix A.1. Notation for Clifford Algebras

$C\ell(s, t)$  stands for the Clifford algebra generated by  $\gamma_\alpha$  satisfying

$$[\gamma_\alpha, \gamma_\beta]_+ = 2\eta_{\alpha\beta}, \quad \eta_{\alpha\alpha} = 1 \text{ for } \alpha = 1, \dots, s, \quad \eta_{\alpha\alpha} = -1 \text{ for } \alpha = s+1, \dots, s+t$$

( $\eta_{\alpha\beta} = 0$  for  $\alpha \neq \beta$ ). Its automorphism group is the (non-compact for  $st \neq 0$ ) orthogonal group  $O(s, t) = O(t, s)$ . As internal symmetries correspond to compact gauge groups, we are mainly working with (positive or negative) definite forms and use the abbreviated notation  $C\ell_s = C\ell(s, 0)$  and  $C\ell_{-t} = C\ell(0, t)$  for the associated Clifford algebras. The even

subalgebra  $\mathcal{C}\ell^0(s, t)$  is defined as the (closed under multiplication) span of products of an even number of  $\gamma$  matrices; The odd subspace  $\mathcal{C}\ell^1(s, t)$  is defined as the (real) span of products of odd numbers of  $\gamma$ s (which is not closed under multiplication).

#### Appendix A.2. Interrelations between the $L$ , $E$ , and $R$ Bases of $so(8)$

The imaginary octonion units  $e_1, \dots, e_7$  obey the anticommutation relations of  $\mathcal{C}\ell_{-7}$

$$[e_\alpha, e_\beta]_+ := e_\alpha e_\beta + e_\beta e_\alpha = -2 \delta_{\alpha\beta}, \alpha, \beta = 1, \dots, 7 \quad (\text{A1})$$

and give rise to the seven generators  $L_\alpha = L_{e_\alpha}$  of the Lie algebra  $so(8)$ :

$$L_{\alpha 8} := \frac{1}{2} L_\alpha =: -L_{8\alpha}, L_{\alpha\beta} := [L_{\alpha 8}, L_{8\beta}] \in so(7) \subset so(8). \quad (\text{A2})$$

For  $\alpha \neq \beta$ , there is a unique  $\gamma$  such that

$$L_\alpha e_\beta = f_{\alpha\beta\gamma} e_\gamma = \pm e_\gamma, f_{\alpha\beta\gamma} = -f_{\beta\alpha\gamma} = f_{\gamma\alpha\beta}. \quad (\text{A3})$$

The structure constants  $f_{\alpha\beta\gamma}$  (22) (which only take the values  $0, \pm 1$ ) obey for different triples  $(\alpha, \beta, \gamma)$  the relations

$$f_{\alpha\beta\gamma} = f_{\alpha+1\beta+1\gamma+2} = f_{2\alpha, 2\beta, 2\gamma} \pmod{7}. \quad (\text{A4})$$

The list (22) follows from  $f_{124} = 1$  and the first Equation (A4), taking into account relations such as  $f_{679} \equiv f_{672} \pmod{7}$ , etc. Note that  $f_{\alpha\beta\gamma} \neq 0$   $f_{\alpha\beta\gamma}$  are the structure constants of a (quaternionic)  $su(2)$  Lie algebra; they are *not* the structure constants of  $so(7) \subset so(8)$ .

Define the involutive outer automorphism  $\pi$  of the Lie algebra  $so(8)$  by its action (26) on the left and right multiplication  $L_\alpha$  and  $R_\alpha$  of octonions by imaginary octonions  $\alpha = -\alpha^*$ :

$$\pi(L_\alpha) = L_\alpha + R_\alpha =: T_\alpha, \pi(R_\alpha) = -R_\alpha \Rightarrow \pi(T_\alpha) = L_\alpha. \quad (\text{A5})$$

In the basis (A1) and (A3) of imaginary octonion units  $e_\alpha$  ( $\alpha = 1, \dots, 7$ ), by setting  $e_8 = \mathbb{1}$  and  $L_{\alpha 8} = \frac{1}{2} L_\alpha$  (A2),  $R_{\alpha 8} = \frac{1}{2} R_\alpha = -R_{8\alpha}$ , we define  $E_{ab}$  by the second relation (27)

$$E_{ab} e_c := \delta_{bc} e_a - \delta_{ac} e_b, a, b, c = 1, \dots, 8 \quad (e_8 = 1). \quad (\text{A6})$$

**Proposition A1.** *Under the above assumptions/definitions, we have*

$$\pi(L_{ab}) = E_{ab} \quad (\text{for } L_{\alpha\beta} := [L_{\alpha 8}, L_{8\beta}], L_{\alpha 8} = \frac{1}{2} L_\alpha = -L_{8\alpha}). \quad (\text{A7})$$

**Proof.** From the first equation (A5) and from (A1), (A2), and (A6), it follows that

$$E_{\alpha 8} = L_{\alpha 8} + R_{\alpha 8} = \pi(L_{\alpha 8}). \quad (\text{A8})$$

The proposition then follows from the relations

$$L_{\alpha\beta} = [L_{\alpha 8}, L_{8\beta}], \quad E_{\alpha\beta} = [E_{\alpha 8}, E_{8\beta}] \quad (\text{A9})$$

and from the assumption that  $\pi$  is a Lie algebra homomorphism.  $\square$

**Corollary A1.** *From (A7) and the involutive character of  $\pi$ , it follows that, conversely,*

$$\pi(E_{ab}) = L_{ab}. \quad (\text{A10})$$

*To each  $\alpha = 1, \dots, 7$ , there are three pairs  $\beta\gamma$  such that  $L_{\beta\gamma}$  and  $E_{\beta\gamma}$  commute with  $L_\alpha$  and among themselves and allow for the expression  $L_\alpha = 2L_{\alpha 8}$  in terms of  $E_{\alpha 8}$  and the corresponding  $E_{\beta\gamma}$ :*

$$\begin{aligned}
L_1 &= 2L_{18} = E_{18} - E_{24} - E_{37} - E_{56}, \\
L_2 &= 2L_{28} = E_{28} + E_{14} - E_{35} - E_{67}, \\
L_3 &= 2L_{38} = E_{38} + E_{17} + E_{25} - E_{46}, \\
L_4 &= 2L_{48} = E_{48} - E_{12} + E_{36} - E_{57}, \\
L_5 &= 2L_{58} = E_{58} + E_{16} - E_{23} - E_{47}, \\
L_6 &= 2L_{68} = E_{68} - E_{15} + E_{27} - E_{34}, \\
L_7 &= 2L_{78} = E_{78} - E_{13} - E_{26} - E_{45}, \text{ or } L_\alpha = E_{\alpha 8} - \sum_{\beta < \gamma} f_{\alpha\beta\gamma} E_{\beta\gamma}. \tag{A11}
\end{aligned}$$

Recalling that  $E_{ab} = \pi(L_{ab})$  (A8) and the fact that  $\pi$  is involutive, so that  $\pi(E_{ab}) = L_{ab}$  (A10), we deduce, in particular,

$$2E_{78} = L_{78} - L_{13} - L_{26} - L_{45},$$

$$R_7 = 2E_{78} - 2L_{78} = -L_{78} - L_{13} - L_{26} - L_{45}, \tag{A12}$$

thus reproducing (29).

We now proceed to displaying the commutant of  $i\omega_6$  and  $i\omega_6^R$  in  $so(7+j)$ ,  $j = 1, 2, 3$ .

**Proposition A2.** While the Lie algebra  $spin(6) = su(4)$  commutes with  $L_7$ , the commutant of  $R_7$  (A12) in  $su(4) \subset sl(4, \mathbb{C})$  is  $u(3) (\subset sl(4, \mathbb{C}))$ , given by

$$u(3) = \left\{ \sum_{j,k=1}^3 C_{jk} [b_j^*, b_k]; C_{jk} \in \mathbb{C}, C_{kj} = -\overline{C_{jk}} \right\} \tag{A13}$$

in the fermionic oscillator realization of  $\mathcal{C}\ell_6(\mathbb{C})$  (the bar over  $C_{jk}$  standing for complex conjugation).

**Proof.** The fact that  $L_7 = 2L_{78}$  commutes with the generators  $L_{\alpha\beta}$  ( $\alpha, \beta = 1, \dots, 6$ ) of  $so(6)$  and follows from (21). To find the commutant of  $R_7$  (A12), it is convenient to use the fermionic realization of the complexification  $sl(4, \mathbb{C})$  of  $su(4)$ , which is spanned by the nine commutators  $[b_j^*, b_k]$  in (A13) and the six products

$$b_j b_k = -b_k b_j, b_j^* b_k^* = -b_k^* b_j^*, j, k = 1, 2, 3, j \neq k. \tag{A14}$$

The sum  $L_{13} + L_{26} + L_{45}$  in (A12) is a multiple of  $B - L$  (58), the Hermitian generator of the center of  $gl(3, \mathbb{C})$ ,

$$B - L \left( = \frac{i}{3} (\gamma_{13} + \gamma_{26} + \gamma_{45}) \right) = \frac{1}{3} \sum_{j=1}^3 [b_j^*, b_j]. \tag{A15}$$

The relations

$$\begin{aligned}
[B - L, b_j^* b_k^*] &= \frac{2}{3} b_j^* b_k^*, [B - L, b_j b_k] = -\frac{2}{3} b_j b_k, \\
[[B - L, [b_j^*, b_k]]] &= 0, j, k = 1, 2, 3, j \neq k, \tag{A16}
\end{aligned}$$

show that the commutant of  $B - L$  (and hence of  $R_7$ ) in  $su(4)$  is  $u(3)$ .  $\square$

**Corollary A2.** The commutant of  $\omega_6^R$  in  $so(8)$  is  $u(3) \oplus u(1)$ ; the commutant of  $\omega_6^R$  in  $spin(9)$  is the gauge Lie algebra of the SM:

$$\mathcal{G}_{\text{SM}} = \{a \in spin(9); [a, \omega_6^R] = 0\} = u(3) \oplus su(2). \tag{A17}$$

## Notes

- 1 For a pleasant-to-read review of octonions, their history, and applications, see [13].
- 2 I had the good fortune to know him personally. See Witten’s eloquent characterization of his personality and work in the Wikipedia entry on Feza Gürsey (1921–1991).
- 3 These algebras are defined and classified in [3,4]; for a concise review, see [29], Section 3.2 of [25], and Section 2 of [30]; concerning Pascual Jordan (1902–1980), see [31].
- 4 For an enlightening review of the algebra of GUTs and some 40 references, see [38].
- 5 Aptly called *geometric algebra* by its inventor—see [39].
- 6 For any associative ring  $\mathbb{K}$ , particularly for the division rings  $\mathbb{K} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ , we denote the algebra of  $m \times m$  matrices with entries in  $\mathbb{K}$  by  $\mathbb{K}[m]$ .
- 7 See [50] for a reader-friendly review of Moufang loops and for a glimpse of the personality of Ruth Moufang (1905–1971).
- 8 The 10-fold classification of  $\mathbb{Z}_2$  graded Clifford algebras also involves signs coming from squaring two antiunitary charge conjugation operators—see [51] Chapter 13, pp. 87–125.
- 9 The group  $G_{\text{SM}}$  was earlier obtained in [35], starting with the Albert algebra  $J_3^8$  (7).
- 10 The Dublin Professor of Astronomy William Rowan Hamilton (1805–1865) and the Stettin Gymnasium teacher Hermann Günther Grassmann (1809–1877) published their papers on quaternions and on “extensive algebras”, respectively, in the same year of 1844. William Kingdom Clifford (1845–1879) combined the two in a “geometric algebra” in 1878, a year before his death, aged 33, referring to both of them.

## References

1. Baez, J.C. Can we understand the Standard Model? talk at the Workshop. In *Octonions and the Standard Model*; Perimeter Institute: Waterloo, ON, Canada, 2021.
2. Todorov, I. Superselection of the weak hypercharge and the algebra of the Standard Model. *JHEP* **2021**, *4*, 164. [\[CrossRef\]](#)
3. Jordan, P. Über Verallgemeinerungsmöglichkeiten des Formalismus der Quantenmechanik. *Nachr. Akad. Wiss. Göttingen Math. Phys.* **1933**, *41*, 209217.
4. Jordan, P.; von Neumann, J.; Wigner, E. On an algebraic generalization of the quantum mechanical formalism. *Ann. Math.* **1934**, *36*, 29–64. [\[CrossRef\]](#)
5. Connes, A. Essay on physics and noncommutative geometry. In *The Interface of Mathematics and Particle Physics*; Clarendon Press: Oxford, UK, 1988; Volume 24, pp. 9–48.
6. Connes, A.; Lott, J. Particle models and noncommutative geometry. *Nucl. Phys. Proc. Suppl.* **1990**, *18*, 29–47. [\[CrossRef\]](#)
7. Dubois-Violette, M.; Kerner, R.; Madore, J. Non-commutative differential geometry and new models of gauge theory. *J. Math. Phys.* **1990**, *31*, 323–329.
8. Boyle, L.; Farnsworth, S. The standard model, the Pati-Salam model, and “Jordan geometry”. *New J. Phys.* **2020**, *22*, 073023. [\[CrossRef\]](#)
9. Chamseddine, A.H.; Connes, A. Noncommutative geometry as a framework for unification of all fundamental interactions including gravity. *Fortschr. Phys.* **2010**, *58*, 553–600. [\[CrossRef\]](#)
10. Chamseddine, A.H.; Connes, A.; van Suijlekom, W.D. Grand unification in the spectral Pati-Salam model. *JHEP* **2015**, *1511*, 011. [\[CrossRef\]](#)
11. Chamseddine, A.H.; Iliopoulos, J.; van Suijlekom, W.D. Spectral action in matrix form. *Eur. Phys. J.* **2020**, *80*, 1045. [\[CrossRef\]](#)
12. van Nuland, J.D.H.; van Suijlekom, W.D. One loop corrections to the spectral action. *arXiv* **2021**, arXiv:2107.08485.
13. Baez, J.C. The octonions. *Bull. Am. Math. Soc.* **2002**, *39*, 145–205; Erratum in *ibid* **2005**, *42*, 213. [\[CrossRef\]](#)
14. Günaydin, M.; Gürsey, F. Quark structure and octonions. *J. Math. Phys.* **1973**, *14*, 1651–1667. [\[CrossRef\]](#)
15. Gürsey, F. Color quarks and octonions, The Johns Hopkins Workshop on Current Problems in High Energy Theory, 1974 Octonionic structures in particle physics. In *Group Theoretical Methods in Physics*; LNP 94; Springer: Berlin/Heidelberg, Germany, 1979; pp. 508–521.
16. Dixon, G.M. Division algebras; spinors; idempotents; the algebraic structure of reality. *arXiv* **2010**, arXiv:1012.1304.
17. Dixon, G.M. Seeable matter; unseeable antimatter, *Comment. Math. Univ. Carolin.* **2014**, *55*, 381–386.
18. Furey, C. Generations: Three prints, in colour. *JHEP* **2014**, *10*, 46. [\[CrossRef\]](#)
19. Furey, C. Charge quantization from a number operator. *Phys. Lett.* **2015**, *742*, 195–199. [\[CrossRef\]](#)
20. Furey, C. Standard model physics from an algebra? *arXiv* **2016**, arXiv:1611.09182.
21. Furey, C.  $SU(3)c \times SU(2)L \times U(1)Y \times U(1)X$  as a symmetry of the division algebra ladder operators. *Eur. Phys. J.* **2018**, *78*, 375. [\[CrossRef\]](#)
22. Furey, C. Three generations, two unbroken gauge symmetries, and one eight-dimensional algebra. *Phys. Lett.* **2018**, *85*, 84–89. [\[CrossRef\]](#)
23. Furey, N.; Hughes, M.J. One generation of standard model Weyl spinors as a single copy of  $\mathbb{R} \otimes \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$ . *Phys. Lett.* **2022**, *827*, 136259. [\[CrossRef\]](#)
24. Furey, N.; Hughes, M.J. Division algebraic symmetry breaking. *Phys. Lett.* **2022**, *831*, 137186. [\[CrossRef\]](#)

25. Dubois-Violette, M. Exceptional quantum geometry and particle physics. *Nucl. Phys.* **2016**, *912*, 426–444. [\[CrossRef\]](#)

26. Dubois-Violette, M. Finite quantum geometry, octonions and the theory of fundamental particles, extended notes on talk at the Workshop. In Proceedings of the Octonions and the Standard Model, Paris, France, 8 February 2021.

27. Todorov, I.; Drenska, S. Composition algebras, exceptional Jordan algebra and related groups. *JGSP* **2017**, *46*, 59–93. [\[CrossRef\]](#)

28. Todorov, I.; Drenska, S. Octonions, exceptional Jordan algebra, and the role of the group  $F_4$  in particle physics. *Adv. in Appl. Clifford Alg.* **2018**, *28*, 82. [\[CrossRef\]](#)

29. Meng, G. Euclidean Jordan algebras, hidden actions, and J-Kepler problems. *J. Math. Phys.* **2011**, *52*, 112104. [\[CrossRef\]](#)

30. Todorov, I. Exceptional quantum algebra for the standard model of particle physics. In *Lie Theory and Its Applications in Physics*; Dobrev, V.K., Ed.; Chapter 3; Springer: Singapore, 2020.

31. Dahn, R.W. The Forgotten Founder of Quantum Mechanics: The Science and Politics of Physicist Pascual Jordan 1902–1980. Ph.D. Thesis, University of Chicago, Chicago, IL, USA, 2019.

32. Albert, A.A. On a certain algebra of quantum mechanics. *Ann. Math.* **1934**, *35*, 65–73. [\[CrossRef\]](#)

33. Dubois-Violette, M.; Todorov, I., Exceptional quantum geometry and particle physics II. *Nucl. Phys.* **2019**, *938*, 751–761. [\[CrossRef\]](#)

34. Dubois-Violette, M.; Todorov, I. Superconnection in the spinfactor approach to particle physics. *Nucl. Phys.* **2020**, *957*, 115065. [\[CrossRef\]](#)

35. Todorov, I.; Dubois-Violette, M. Deducing the symmetry of the standard model from the automorphism and structure groups of the exceptional Jordan algebra. *Int. J. Mod. Phys.* **2018**, *33*, 1850118. [\[CrossRef\]](#)

36. Haag, R. *Local Quantum Physics, Fields, Particles, Algebras*; Springer: Berlin, Germany, 1993.

37. Pati, J. Abdus Salam, Unified Lepton-Hadron Symmetry and a Gauge Theory of the Basic Interactions. *Phys. Rev.* **1973**, *8*, 1240–1251.

38. Baez, J.C.; Huerta, J. The algebra of grand unified theory. *Bull. Am. Math. Soc.* **2010**, *47*, 483–552. [\[CrossRef\]](#)

39. Doran, C.; Lasenby, A. *Geometric Algebra for Physicists*; Cambridge University Press: Cambridge, UK, 2003.

40. Casalbuoni, R.; Gato, R. Unified theory of quarks and leptons based on Clifford algebras. *Phys. Lett.* **1979**, *90*, 81–86. [\[CrossRef\]](#)

41. Borel, A.; de Siebenthal, J. Les sous-groupe fermés de rang maximum des groupes de Lie clos. *Comment. Math. Helv.* **1949**, *23*, 200–221. [\[CrossRef\]](#)

42. Nielsen, H.B.; Bennett, D. Seaking a game in which the standard model group shall win. *arXiv* **2013**, arXiv:1306.2668.

43. Maladacena, J. The symmetry and simplicity of the laws of physics and the Higgs boson. *Eur. J. Phys.* **2015**, *37*, 015802. [\[CrossRef\]](#)

44. Fairlie, D. Higgs field and the determination of the Weinberg angle. *Phys. Lett.* **1979**, *82*, 97–100. [\[CrossRef\]](#)

45. Neeman, Y. Internal subgroup prediction of the Goldstone-Higgs particle mass. *Phys. Lett.* **1979**, *81*, 309–310.

46. Krasnov, K.  $SO(9)$  characterization of the standard model gauge group. *J. Math. Phys.* **2021**, *62*, 021703. [\[CrossRef\]](#)

47. Boyle, L. The standard model, the exceptional Jordan algebra and triality. *arXiv* **2020**, arXiv:2006.16265.

48. Bryant, R.L. Notes on spinors in low dimensions. *arXiv* **2020**, arXiv:2011.05568.

49. Yokota, I. Exceptional Lie groups. *arXiv* **2009**, arXiv:0902.0431.

50. Stener, M. *Moufang Loops General Theory and Visualization of Non-Associative Moufang Loops of Order 16*. Ph.D. Thesis, Uppsala University, Uppsala, Sweden, 2016.

51. Moore, G.W. Quantum symmetries and compatible Hamiltonians. *Notes Phys.* **2013**, *695*, 267.

52. Tanimura, S. Superselection rules from measurement theory. *arXiv* **2011**, arXiv:1112.5701.

53. Barducci, A.; Buccella, F.; Casalbuoni, R.; Lusanna, L.; Sorace, E. Quantized Grassmann variables and unified theories. *Phys. Lett. B* **1977**, *67*, 344–346. [\[CrossRef\]](#)

54. Budinich, P.; Trautman, A. *The Spinorial Chessboard*; Trieste Notes in Physics; Springer: Berlin/Heidelberg, Germany, 1988; 128p.

55. Dubois-Violette, M. Complex structures and the Elie Cartan approach to the theory of spinors. In *Spinors, Twistors, Clifford Algebras and Quantum Deformations*; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1993; pp. 17–23.

56. Penrose, R. Basic twistor theory, bi-twistors and split octonions, talk at the online lecture series. In Proceedings of the Octonions, Standard Model and Unification, (OSMU23), Online, 24 February–15 December 2023.

57. Wick, G.C.; Wightman, A.S.; Wigner, E.P. The intrinsic parity of elementary particles. *Phys. Rev.* **1952**, *88*, 101–105. [\[CrossRef\]](#)

58. Wick, G.C.; Wightman, A.S.; Wigner, E.P. Superselection rule for charge. *Phys. Rev.* **1970**, *12*, 3267–3269. [\[CrossRef\]](#)

59. Giulini, D. Superselection rules, *arXiv* **2007**, arXiv:0710.1516v2.

60. Earman, J. Superselection rules for philosophers. *Erkenntnis* **2008**, *69*, 377–414. [\[CrossRef\]](#)

61. Gracia-Bondia, J.M.; Iochum, B.; Schucker, T. The Standard Model in noncommutative geometry and fermion doubling. *Phys. Lett.* **1998**, *416*, 123. [\[CrossRef\]](#)

62. Bochniak, A.; Sitarz, A.; Zalecki, P. Spectral action of the electroweak  $\theta$ -terms for the Standard Model without fermion doubling. *JHEP* **2021**, *12*, 142. [\[CrossRef\]](#)

63. Krasnov, K.  $Spin(11,3)$ , particles, and octonions. *J. Math. Phys.* **2022**, *63*, 031701. [\[CrossRef\]](#)

64. Strursberg, R.C.; De Traubenberg, M.R. *Group Theory in Physics, A Practitioner’s Guide*; World Scientific: Singapore, 2019.

65. Mahapatra, R.N.; Pati, J.C. “Natural” left-right symmetry. *Phys. Rev.* **1975**, *11*, 2558–2561. [\[CrossRef\]](#)

66. Senjanović, G. Is left-right symmetry the key? *Mod. Phys. Lett.* **2017**, *32*, 173004. [\[CrossRef\]](#)

67. Dunsky, D.; Hall, L.J.; Harigaya, K. Sterile neutrino dark matter and leptogenesis in left-right Higgs parity. *JHEP* **2021**, *1*, 125. [\[CrossRef\]](#)

68. Hall, L.J.; Harigaya, K. Implications of Higgs discovery to the strong CP problem and unification. *JHEP* **2018**, *10*, 130. [\[CrossRef\]](#)

69. Matthes, V.; Quillen, D. Superconnections, Thom classes, and covariant differential forms. *Topology* **1986**, *25*, 85–110. [[CrossRef](#)]
70. Quillen, D. Superconnections and the Chern character. *Topology* **1985**, *24*, 85–95. [[CrossRef](#)]
71. Thierry-Mieg, J. Chirality, the missing key to the definition of the connection and curvature of a Lie-Kac superalgebra. *JHEP* **2021**, *1*, 111. [[CrossRef](#)]
72. Roepstorff, G. Superconnections and the Higgs field. *J. Math. Phys.* **1999**, *40*, 2698–2715. [[CrossRef](#)]
73. Thierry-Mieg, J. Scalar anomaly cancellation reveals the hidden algebraic structure of the quantum chiral  $SU(2|1)$  model of leptons and quarks. *JHEP* **2020**, *10*, 167. [[CrossRef](#)] [[PubMed](#)]
74. Manogue, C.A.; Dray, T.; Wilson, R.A. Octons: An  $E_8$  description of the Standard Model. *J. Math. Phys.* **2022**, *63*, 081703. [[CrossRef](#)]
75. Thierry-Mieg, J.; Jarvis, P.; Germini, J.; Gorelik, M. Construction of matryoshka nested indecomposable N-replications of Kac modules of quasi-reductive Lie superalgebras, including the  $sl(m/n), sop(2/n)$  series. *arXiv* **2022**, arXiv:2207.06538v3.

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.