

NATURALNESS, CHIRAL SYMMETRY, AND SPONTANEOUS
CHIRAL SYMMETRY BREAKING

G. 't Hooft

Institute for Theoretical Physics

Utrecht, The Netherlands

ABSTRACT

A properly called "naturalness" is imposed on gauge theories. It is an order-of-magnitude restriction that must hold at all energy scales μ . To construct models with complete naturalness for elementary particles one needs more types of confining gauge theories besides quantum chromodynamics. We propose a search program for models with improved naturalness and concentrate on the possibility that presently elementary fermions can be considered as composite. Chiral symmetry must then be responsible for the masslessness of these fermions. Thus we search for QCD-like models where chiral symmetry is not or only partly broken spontaneously. They are restricted by index relations that often cannot be satisfied by other than unphysical fractional indices. This difficulty made the author's own search unsuccessful so far. As a by-product we find yet another reason why in ordinary QCD chiral symmetry must be broken spontaneously.

III.1. INTRODUCTION

The concept of causality requires that macroscopic phenomena follow from microscopic equations. Thus the properties of liquids and solids follow from the microscopic properties of molecules and atoms. One may either consider these microscopic properties to have been chosen at random by Nature, or attempt to deduce these from even more fundamental equations at still smaller length and time scales. In either case, it is unlikely that the microscopic equations contain various free parameters that are carefully adjusted by Nature to give cancelling effects such that the macroscopic systems have some special properties. This is a

philosophy which we would like to apply to the unified gauge theories: the effective interactions at a large length scale, corresponding to a low energy scale μ_1 , should follow from the properties at a much smaller length scale, or higher energy scale μ_2 , without the requirement that various different parameters at the energy scale μ_2 match with an accuracy of the order of μ_1/μ_2 . That would be unnatural. On the other hand, if at the energy scale μ_2 some parameters would be very small, say

$$\alpha(\mu_2) = \mathcal{O}(\mu_1/\mu_2) , \quad (\text{III1})$$

then this may still be natural, provided that this property would not be spoilt by any higher order effects. We now conjecture that the following dogma should be followed:

- at any energy scale μ , a physical parameter or set of physical parameters $\alpha_i(\mu)$ is allowed to be very small only if the replacement $\alpha_i(\mu) = 0$ would increase the symmetry of the system. - In what follows this is what we mean by naturalness. It is clearly a weaker requirement than that of P. Dirac¹⁾ who insists on having no small numbers at all. It is what one expects if at any mass scale $\mu > \mu_0$ some ununderstood theory with strong interactions determines a spectrum of particles with various good or bad symmetry properties. If at $\mu = \mu_0$ certain parameters come out to be small, say 10^{-5} , then that cannot be an accident; it must be the consequence of a near symmetry.

For instance, at a mass scale

$$\mu = 50 \text{ GeV},$$

the electron mass m_e is 10^{-5} . This is a small parameter. It is acceptable because $m_e = 0$ would imply an additional chiral symmetry corresponding to separate conservation of left handed and right handed electron-like leptons. This guarantees that all renormalizations of m_e are proportional to m_e itself. In sects. III2 and III3 we compare naturalness for quantum electrodynamics and ϕ^4 theory.

Gauge coupling constants and other (sets of) interaction constants may be small because putting them equal to zero would turn the gauge bosons or other particles into free particles so that they are separately conserved.

If within a set of small parameters one is several orders of magnitude smaller than another then the smallest must satisfy our "dogma" separately. As we will see, naturalness will put the severest restriction on the occurrence of scalar particles in renormalizable theories. In fact we conjecture that this is the reason why light, weakly interacting scalar particles are not seen.

It is our aim to use naturalness as a new guideline to construct models of elementary particles (sect. III4). In practice naturalness will be lost beyond a certain mass scale μ_0 , to be referred to as "Naturalness Breakdown Mass Scale" (NBMS). This simply means that unknown particles with masses beyond that scale are ignored in our model. The NBMS is only defined as an order of magnitude and can be obtained for each renormalizable field theory. For present "unified theories", including the existing grand unified schemes, it is only about 1000 GeV. In sect. 5 we attempt to construct realistic models with an NBMS some orders of magnitude higher.

One parameter in our world is unnatural, according to our definition, already at a very low mass scale ($\mu_0 \sim 10^{-2}$ eV). This is the cosmological constant. Putting it equal to zero does not seem to increase the symmetry. Apparently gravitational effects do not obey naturalness in our formulation. We have nothing to say about this fundamental problem, except to suggest that *only* gravitational effects violate naturalness. Quantum gravity is not understood anyhow so we exclude it from our naturalness requirements.

On the other hand it is quite remarkable that all other elementary particle interactions have a high degree of naturalness. No unnatural parameters occur in that energy range where our popular field theories could be checked experimentally. We consider this as important evidence in favor of the general hypothesis of naturalness. Pursuing naturalness beyond 1000 GeV will require theories that are immensely complex compared with some of the grand unified schemes.

A remarkable attempt towards a natural theory was made by Dimopoulos and Susskind²⁾. These authors employ various kinds of confining gauge forces to obtain scalar bound states which may substitute the Higgs fields in the conventional schemes. In their model the observed fermions are still considered to be elementary.

Most likely a complete model of this kind has to be constructed step by step. One starts with the experimentally accessible aspects of the Glashow-Weinberg-Salam-Ward model. This model is natural if one restricts oneself to mass-energy scales below 1000 GeV. Beyond 1000 GeV one has to assume, as Dimopoulos and Susskind do, that the Higgs field is actually a fermion-antifermion composite field. Coupling this field to quarks and leptons in order to produce their mass, requires new scalar fields that cause naturalness to break down at 30 TeV or so. Dimopoulos and Susskind speculate further on how to remedy this. To supplement such ideas, we toyed with the idea that (some of) the presently "elementary" fermions may turn out to be bound states of an odd number of fermions when considered beyond 30 TeV. The binding mechanism would be similar

to the one that keeps quarks inside the proton. However, the proton is not particularly light compared with the characteristic mass scale of quantum chromodynamics (QCD). Clearly our idea is only viable if something prevented our "baryons" from obtaining a mass (eventually a small mass may be due to some secondary perturbation).

The proton owes its mass to spontaneous breakdown of chiral symmetry, or so it seems according to a simple, fairly successful model of the mesonic and baryonic states in QCD: the Gell-Mann-Lévy sigma model³⁾. Is it possible then that in some variant of QCD chiral symmetry is not spontaneously broken, or only partly, so that at least some chiral symmetry remains in the spectrum of fermionic bound states? In this article we will see that in general in SU(N) binding theories this is not allowed to happen, i.e. chiral symmetry must be broken spontaneously.

III.2. NATURALNESS IN QUANTUM ELECTRODYNAMICS

Quantum Electrodynamics as a renormalizable model of electrons (and muons if desired) and photons is an example of a "natural" field theory. The parameters α , m_e (and m_μ) may be small independently. In particular m_e (and m_μ) are very small at large μ . The relevant symmetry here is chiral symmetry, for the electron and the muon separately. We need not be concerned about the Adler-Bell-Jackiw anomaly here because the photon field being Abelian cannot acquire non-trivial topological winding numbers⁴⁾.

There is a value of μ where Quantum Electrodynamics ceases to be useful, even as a model. The model is not asymptotically free, so there is an energy scale where all interactions become strong:

$$\mu_0 \simeq m_e \exp(6\pi^2/e^2 N_f) , \quad (\text{III.2})$$

where N_f is the number of light fermions. If some world would be described by such a theory at low energies, then a replacement of the theory would be necessary at or below energies of order μ_0 .

III.3. ϕ^4 -THEORY

A renormalizable scalar field theory is described by the Lagrangian

$$\mathcal{L} = -\frac{1}{2}(\partial_\mu \phi)^2 - \frac{1}{2}m^2 \phi^2 - \frac{1}{4!} \lambda \phi^4 . \quad (\text{III.3})$$

the interactions become strong at

$$\mu \simeq m \exp(16\pi^2/3\lambda) , \quad (\text{III.4})$$

but is it still natural there?

There are two parameters, λ and m . Of these, λ may be small because $\lambda = 0$ would correspond to a non-interacting theory with total number of ϕ particles conserved. But is small m allowed? If we put $m = 0$ in the Lagrangian (III3) then the symmetry is not enhanced*). However we can take both m and λ to be small, because if $\lambda = m = 0$ we have invariance under

$$\phi(x) \rightarrow \phi(x) + \Lambda . \quad (\text{III5})$$

This would be an approximate symmetry of a new underlying theory at energies of order μ_0 . Let the symmetry be broken by effects described by a dimensionless parameter ϵ . Both the mass term and the interaction term in the effective Lagrangian (III3) result from these symmetry breaking effects. Both are expected to be of order ϵ . Substituting the correct powers of μ_0 to account for the dimensions of these parameters we have

$$\begin{aligned} \lambda &= \mathcal{O}(\epsilon) , \\ m^2 &= \mathcal{O}(\epsilon \mu_0^2) . \end{aligned} \quad (\text{III6})$$

Therefore,

$$\mu_0 = \mathcal{O}(m/\sqrt{\lambda}) . \quad (\text{III7})$$

This value is much lower than eq. (III4). We now turn the argument around: if any "natural" underlying theory is to describe a scalar particle whose *effective* Lagrangian at low energies will be eq. (III3), then its energy scale cannot be given by (III4) but at best by (III7). We say that naturalness breaks down beyond $m/\sqrt{\lambda}$. It must be stressed that these are orders of magnitude. For instance one might prefer to consider λ/π^2 rather than λ to be the relevant parameter. μ_0 then has to be multiplied by π . Furthermore, λ could be much smaller than ϵ because $\lambda = 0$ separately also enhances the symmetry. Therefore, apart from factors π , eq. (III7) indicates a maximum value for μ_0 .

Another way of looking at the problem of naturalness is by comparing field theory with statistical physics. The parameter m/μ would correspond to $(T-T_c)/T$ in a statistical ensemble. Why would the temperature T chosen by Nature to describe the elementary particles be so close to a critical temperature T_c ? If $T_c \neq 0$ then T may not be close to T_c just by accident.

III4. NATURALNESS IN THE WEINBERG-SALAM-GIM MODEL

The difficulties with the unnatural mass parameters only occur in theories with scalar fields. The only fundamental scalar

*) Conformal symmetry is violated at the quantum level.

field that occurs in the presently fashionable models is the Higgs field in the extended Weinberg-Salam model. The Higgs mass-squared, m_H^2 , is up to a coefficient a fundamental parameter in the Lagrangian. It is small at energy scales $\mu \gg m_H$. Is there an approximate symmetry if $m_H \rightarrow 0$? With some stretch of imagination we might consider a Goldstone-type symmetry:

$$\phi(x) \rightarrow \phi(x) + \text{const.} \quad (\text{III8})$$

However we also had the local gauge transformations:

$$\phi(x) \rightarrow \Omega(x) \phi(x) . \quad (\text{III9})$$

The transformations (III8) and (III9) only form a closed group if we also have invariance under

$$\phi(x) \rightarrow \phi(x) + C(x) . \quad (\text{III10})$$

But then it becomes possible to transform ϕ away completely. The Higgs field would then become an unphysical field and that is not what we want. Alternatively, we could have that (III8) is an approximate symmetry only, and it is broken by all interactions that have to do with the symmetry (III9) which are the weak gauge field interactions. Their strength is $g^2/4\pi = \mathcal{O}(1/137)$. So at best we can have that the symmetry is broken by $\mathcal{O}(1/137)$ effects. Therefore

$$m_H^2/\mu^2 \gtrsim \mathcal{O}(1/137) .$$

Also the $\lambda\phi^4$ term in the Higgs field interactions breaks this symmetry. Therefore

$$m_H^2/\mu^2 \gtrsim \mathcal{O}(\lambda) \gtrsim \mathcal{O}(1/137) . \quad (\text{III11})$$

Now

$$m_H^2 = \mathcal{O}(\lambda F_H^2) , \quad (\text{III12})$$

where F_H is the vacuum expectation value of the Higgs field, known to be*)

$$F_H = (2G\sqrt{2})^{-1/2} = 174 \text{ GeV} . \quad (\text{III13})$$

We now read off that

$$\mu \lesssim \mathcal{O}(F_H) = \mathcal{O}(174 \text{ GeV}) . \quad (\text{III14})$$

*) Some numerical values given during the lecture were incorrect. I here give corrected values.

This means that at energy scales much beyond F_H our model becomes more and more unnatural. Actually, factors of π have been omitted. In practice one factor of 5 or 10 is still not totally unacceptable. Notice that the actual value of m_H dropped out, except that

$$m_H = \mathcal{O}\left(\frac{\sqrt{\lambda}}{g} M_W\right) \gtrsim \mathcal{O}(M_W) . \quad (\text{III15})$$

Values for m_H of just a few GeV are unnatural.

III5. EXTENDING NATURALNESS

Equation (III14) tells us that at energy scales much beyond 174 GeV the standard model becomes unnatural. As long as the Higgs field H remains a fundamental scalar nothing much can be done about that. We therefore conclude, with Dimopoulos and Susskind²⁾ that the "observed" Higgs field must be composite. A non-trivial strongly interacting field theory must be operative at 1000 GeV or so. An obvious and indeed likely possibility is that the Higgs field H can be written as

$$H = Z\bar{\psi}\psi , \quad (\text{III16})$$

where Z is a renormalization factor and ψ is a new quark-like object, a fermion with a new color-like interaction²⁾. We will refer to the object as meta-quark having meta-color. The theory will have all features of QCD so that we can copy the nomenclature of QCD with the prefix "meta-". The Higgs field is a meta-meson.

It is now tempting to assume that the meta-quarks transform the same way under weak $SU(2) \times U(1)$ as ordinary quarks. Take a doublet with left-handed components forming one gauge doublet and right handed components forming two gauge singlets. The meta-quarks are massless. Suppose that the meta-chiral symmetry is broken spontaneously just as in ordinary QCD. What would happen?

What happens is in ordinary QCD well described by the Gell-Mann-Lévy sigma model. The lightest mesons form a quartet of real fields, ϕ_{ij} , transforming as a

$$2^{\text{left}} \otimes 2^{\text{right}}$$

representation of

$$SU(2)^{\text{left}} \otimes SU(2)^{\text{right}} .$$

Since the weak interaction only deals with $SU(2)^{\text{left}}$ this quartet can also be considered as one complex doublet representation of weak $SU(2)$. In ordinary QCD we have

$$\phi_{ij} = \sigma \delta_{ij} + i \tau_{ij}^a \pi^a, \quad (\text{III17})$$

and

$$\langle \sigma \rangle_{\text{vacuum}} = \frac{1}{\sqrt{2}} f_{\pi} = 91 \text{ MeV}. \quad (\text{III18})$$

The complex doublet is then

$$\phi_i = \frac{1}{\sqrt{2}} \begin{pmatrix} \sigma + i\pi^3 \\ \pi^2 + i\pi^1 \end{pmatrix}, \quad (\text{III19})$$

and

$$\langle \phi_i \rangle_{\text{vacuum}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \times 64 \text{ MeV}. \quad (\text{III20})$$

We conclude that if we transplant this theory to the TeV range then we get a scalar doublet field with a non-vanishing vacuum expectation value for free. All we have to do now is to match the numbers. If we scale all QCD masses by a scaling factor κ then we match

$$\begin{aligned} F_H &= 174 \text{ GeV} = \kappa 64 \text{ MeV}; \\ \kappa &= 2700. \end{aligned} \quad (\text{III21})$$

Now the mesonic sector of QCD is usually assumed to be reproduced in the $1/N$ expansion⁵⁾ where N is the number of colors (in QCD we have $N = 3$). The 4-meson coupling constant goes like $1/N$. Then one would expect

$$f_{\pi} \propto \sqrt{N}. \quad (\text{III22})$$

Therefore

$$\kappa = 2700 \sqrt{\frac{3}{N}}, \quad (\text{III23})$$

if the metacolor group is $SU(N)$.

Thus we obtain a model that reproduces the W -mass and predicts the Higgs mass. The Higgs is the meta-sigma particle. The ordinary sigma is a wide resonance at about 700 MeV ³⁾, so that we predict

$$m_H = \kappa m_{\sigma} = 1900 \sqrt{\frac{3}{N}} \text{ GeV}, \quad (\text{III24})$$

and it will be extremely difficult to detect among other strongly interacting objects.

III6. WHAT NEXT?

The model of the previous section is to our mind nearly inevitable, but there are problems. These have to do with the observed fermion masses. All leptons and quarks owe their masses to an interaction term of the form

$$g \bar{\psi} H \psi \ ,$$

(III25)

where g is a coupling constant, ψ is the lepton or quark and H is the Higgs field. With (III16) this becomes a four-fermion interaction, a fundamental interaction in the new theory. Because it is non-renormalizable further structure is needed. In ref. 2 the obvious choice is made: a new "meta-weak interaction" gauge theory enters with new super-heavy intermediate vector bosons. But since H is a scalar this boson must be in the crossed channel, a rather awkward situation. (See option a in Figure 1.) A simpler theory is that a new scalar particle is exchanged in the direct channel. (See option b in Figure 1.)




Figure 1.

Notice that in both cases new scalar fields are needed because in case a) something must cause the "spontaneous breakdown" of the new gauge symmetries. Therefore choice b) is simpler. We removed a Higgs scalar and we get a scalar back. Does naturalness improve? The answer is yes. The coupling constant g in the interaction (III25) satisfies

$$g = g_1 g_2 / M_s^2 Z . \quad (\text{III26})$$

Here g_1 and g_2 are the couplings at the new vertices, M_s is the new scalar's mass, and Z is from (III16) and is of order

$$Z \sim \frac{1}{\sqrt{\frac{N}{3}} (\kappa m_\rho)^2} = \frac{\sqrt{N/3}}{(1800 \text{ GeV})^2} . \quad (\text{III27})$$

Suppose that the heaviest lepton or quark is about 10 GeV. For that fermion the coupling constant g is

$$g = \frac{m_f}{F} \simeq 1/20 .$$

We get

$$g_1 g_2 \simeq \left(\frac{M_s}{1800 \text{ GeV}} \right)^2 \sqrt{\frac{N}{3}} \cdot \frac{1}{20} .$$

Naturalness breaks down at

$$\mu = \mathcal{O} \left(\frac{M_s}{g_{1,2}} \right) = 8000 \sqrt[4]{\frac{3}{N}} \text{ GeV} ,$$

an improvement of about a factor 50 compared with the situation in sect. III4. Presumably we are again allowed to multiply by factors like 5 or 10, before getting into real trouble.

Before speculating on how to go on from here to improve naturalness still further we must assure ourselves that all other alleys are blind ones. An intriguing possibility is that the presently observed fermions are composite. We would get option c), Figure 2.




Fig. 2

The dotted line could be an ordinary weak interaction W or photon, that breaks an internal symmetry in the binding force for the new components. The new binding force could either act at the 1 TeV or at the 10-100 TeV range. It could either be an extension of meta-color or be a (color)" or paracolor force. Is such an idea viable?

Clearly, compared with the energy scale on which the binding forces take place, the composite fermions must be nearly massless. Again, this cannot be an accident. The chiral symmetry responsible for this must be present in the underlying theory. Apparently then, the underlying theory will possess a chiral symmetry which is not (or not completely) spontaneously broken, but reflected in the bound state spectrum in the Wigner mode: some massless chiral objects and parity doubled massive fermions. This possibility is most clearly described by the σ -model as a model for the lowest bound states occurring in ordinary quantum chromodynamics.

III7. THE σ MODEL

The fermion system in quantum chromodynamics shows an axial symmetry. To illuminate our problem let us consider the case of two flavors. The local color group is $SU(3)_c$. The subscript c here stands for color. The flavor symmetry group is $SU(2)_L \otimes SU(2)_R \otimes U(1)$ where the subscripts L and R stands for left and right and the group elements must be chosen to be space-time independent. We split the fermion fields ψ into left and right components:

$$\psi = \frac{1}{2}(1+\gamma_5)\psi_L + \frac{1}{2}(1-\gamma_5)\psi_R . \quad (\text{III28})$$

$$\psi_L \text{ transforms as a } 3_c \otimes 2_L \otimes 1_R \otimes 2_{\mathcal{L}} \quad (\text{III29})$$

$$\text{and } \psi_R \text{ transforms as a } 3_c \otimes 1_L \otimes 2_R \otimes \bar{2}_{\mathcal{L}} \quad (\text{III30})$$

where the indices refer to the various groups. \mathcal{L} stands for the Lorentzgroup $SO(3,1)$, locally equivalent to $SL(2,c)$ which has two

different complex doublet representations 2_L and $\bar{2}_L$ (corresponding to the transformation law for the neutrino and antineutrino, respectively). The fields ψ_L and ψ_R have the same charge under $U(1)$, whereas axial $U(1)$ group (under which they would have opposite charges) is absent because of instanton effects⁴⁾.

The effect of the color gauge fields is to bind these fermions into mesons and baryons all of which must be color singlets. It would be nice if one could describe these hadronic fields as representations of $SU(2)_L \otimes SU(2)_R \otimes U(1)$ and the Lorentz group, and then cast their mutual interactions in the form of an effective Lagrangian, invariant under the flavor symmetry group. In the case at hand this is possible and the resulting construction is a successful and one-time popular model for pions and nucleons: the σ model³⁾. We have a nucleon doublet

$$N = \frac{1}{2}(1+\gamma_5)N_L + \frac{1}{2}(1-\gamma_5)N_R, \quad (\text{III31})$$

where

$$N_L \text{ transforms as a } 1_c \otimes 2_L \otimes 1_R \otimes 2_L, \quad (\text{III32a})$$

$$\text{and } N_R \text{ transforms as a } 1_c \otimes 1_L \otimes 2_R \otimes \bar{2}_L. \quad (\text{III32b})$$

Further we have a quartet of real scalar fields $(\sigma, \vec{\pi})$ which transform as a $1_c \otimes 2_L \otimes 2_L \otimes 1_L$. The Lagrangian is

$$\mathcal{L} = -\bar{N}[\gamma\partial + g_0(\sigma + i\vec{\tau} \cdot \vec{\pi}\gamma_5)]N - \frac{1}{2}(\partial\pi)^2 - \frac{1}{2}(\partial\sigma)^2 - V(\sigma^2 + \vec{\pi}^2). \quad (\text{III33})$$

Here V must be a rotationally invariant function.

Usually V is chosen such that its absolute minimum is away from the origin. Let V be minimal at $\sigma = v$ and $\vec{\pi} = 0$. Here v is just a c-number. To obtain the physical particle spectrum we write

$$\sigma = v + s \quad (\text{III34})$$

and we find

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}\bar{N}(\gamma\partial + g_0 v)N - \frac{1}{2}(\partial\vec{\pi})^2 - \frac{1}{2}(\partial s)^2 - 2v^2 V''(v^2)s^2 \\ & + \text{interaction terms}. \end{aligned} \quad (\text{III35})$$

Clearly, in this case the nucleons acquire a mass term $m_s = g_0 v$ and the s particle has a mass $m_s^2 = 4v^2 V''(v^2)$, whereas the pion remains strictly massless. The entire mass of the pion must be due to effects that explicitly break $SU(2)_L \times SU(2)_R$, such as a small

mass term $m_q \bar{\psi} \psi$ for the quarks (III28). We say that in this case the flavor group $SU(2)_L \otimes SU(2)_R$ is spontaneously broken into the isospin group $SU(2)$.

Another possibility however, apparently not realised in ordinary quantum chromodynamics, would be that $SU(2)_L \otimes SU(2)_R$ is *not* spontaneously broken. We would read off from the Lagrangian (III33) that the nucleons N would form a massless doublet and that the four fields $(\sigma, \vec{\pi})$ could be heavy. The dynamics of other confining gauge theories could differ sufficiently from ordinary QCD so that, rather than a spontaneous symmetry breakdown, massless "baryons" develop. The principle question we will concentrate on is why do these massless baryons form the representation (III32), and how does this generalize to other systems. We would let future generations worry about the question where exactly the absolute minimum of the effective potential V will appear.

III8. INDICES

We now consider any color group G_c . The fundamental fermions in our system must be non-trivial representation of G_c and we assume "confinement" to occur: all physical particles are bound states that are singlets under G_c . Assume that the fermions are all massless (later mass terms can be considered as a perturbation). We will have automatically some global symmetry which we call the flavor group G_F . (We only consider exact flavor symmetries, not spoilt by instanton effects.) Assume that G_F is not spontaneously broken. Which and how many representations of G_F will occur in the massless fermion spectrum of the baryonic bound states? We must formulate the problem more precisely. The massless nucleons in (III33) being bound states, may have many massive excitations. However, massive Fermion fields cannot transform as a 2_L under Lorentz transformations; they must go as a $2_L \oplus \bar{2}_L$. That is because a mass term being a Lorentz invariant product of two fields at one point only links 2_L representations with $\bar{2}_L$ representations. Consider a given representation r of G_F . Let p be the number of field multiplets transforming as $r \otimes 2_L$ and q be the number of field multiplets $r \otimes \bar{2}_L$. Mass terms that link the 2_L with $\bar{2}_L$ fields are completely invariant and in general to be expected in the effective Lagrangian. But the absolute value of

$$\ell = p - q \tag{III36}$$

is the minimal number of surviving massless chiral field multiplets. We will call ℓ the index corresponding to the representation r of G_F . By definition this index must be a (positive or negative) integer. In the sigma model it is postulated that

$$\text{index } (2_L \otimes 1_R) = 1 \quad (\text{III37})$$

$$\text{index } (1_L \otimes 2_R) = -1$$

$\text{index } (r) = 0$ for all other representations r .

This tells us that if chiral symmetry is not broken spontaneously one massless nucleon doublet emerges. We wish to find out what massless fermionic bound states will come out in more general theories. Our problem is: how does (III37) generalize?

III9. ABSENCE OF MASSLESS BOUND STATES WITH SPIN 3/2 OR HIGHER

In the foregoing we only considered spin 0 and spin 1/2 bound states. Is it not possible that fundamentally massless bound states develop with higher spin? I believe to have strong arguments that this is indeed not possible. Let us consider the case of spin 3/2. Massive spin 3/2 fermions are described by a Lagrangian of the form

$$\mathcal{L} = \frac{1}{2} \bar{\psi}_\mu [\sigma_{\mu\nu} (\gamma \partial + m) + (\gamma \partial + m) \sigma_{\mu\nu}] \psi_\nu. \quad (\text{III38})$$

Just like spin-one particles, this has a gauge-invariance if $m \rightarrow 0$:

$$\psi_\mu \rightarrow \psi_\mu + \partial_\mu \eta(x), \quad (\text{III39})$$

where $\eta(x)$ is arbitrary. Indeed, massless spin 3/2 particles only occur in locally supersymmetric field theories. The field $\eta(x)$ is fundamentally unobservable.

Now in our model ψ_μ would be shorthand for some composite field: $\psi_\mu \rightarrow \psi \psi_\mu$. However, then all components of this, including η , would be^μobservables. If $m = 0$ we would be forced to add a gauge fixing term that would turn η into an unacceptable ghost particle*).

We believe, therefore, that unitarity and locality forbid the occurrence of massless bound states with spin 3/2. The case for higher spin will not be any better. And so we concentrate on a bound state spectrum of spin 1/2 particles only.

*) Note added: during the lectures it was suggested by one attendant to consider only gauge-invariant fields as $\Psi_{\mu\nu} = \partial_\mu \psi_\nu - \partial_\nu \psi_\mu$.

However, such fields must satisfy constraints: $\partial [\alpha \Psi_{\mu\nu}] = 0$.

Composite field will never automatically satisfy such constraints.

III10. SPECTATOR GAUGE FIELDS AND -FERMIONS

So far, our model consisted of a strong interaction color gauge theory with gauge group G_c , coupled to chiral fermions in various representations r of G_c but of course in such a way that the anomalies cancel. The fermions are all massless and form multiplets of a global symmetry group, called G_F . For QCD this would be the flavor group. In the metacolor theory G_F would include all other fermion symmetries besides metacolor.

In order to study the mathematical problem raised above we will add another gauge connection field that turns G_F into a local symmetry group. The associated coupling constants may all be arbitrarily small, so that the dynamics of the strong color gauge interactions is not much affected. In particular the massless bound state spectrum should not change. One may either think of this new gauge field as a completely quantized field or simply as an artificial background field with possibly non-trivial topology. We will study the behavior of our system in the presence of this "spectator gauge field". As stated, its gauge group is G_F .

Note however, that some flavor transformations could be associated with anomalies. There are two types of anomalies:

i) those associated with $G_c \times G_F$, only occurring where the color field has a winding number. Only $U(1)$ invariant subgroups of G_F contribute here. They simply correspond to small explicit violations of the G_F symmetry. From now on we will take as G_F only the anomaly-free part. Thus, for QCD with N flavors, G_F is not $U(N) \times U(N)$ but

$$G_F = SU(N) \otimes SU(N) \otimes U(1) .$$

ii) those associated with G_F alone. They only occur if the spectator gauge field is quantized. To remedy these we simply add "spectator fermions" coupled to G_F alone. Again, since these interactions are weak they should not influence the bound state spectrum.

Here, the spectator gauge fields and fermions are introduced as mathematical tools only. It just happens to be that they really do occur in Nature, for instance the weak and electromagnetic $SU(2) \times U(1)$ gauge fields coupled to quarks in QCD. The leptons then play the role of spectator fermions.

III11. ANOMALY CANCELLATION FOR THE BOUND STATE SPECTRUM

Let us now resume the particle content of our theory. At small distances we have a gauge group $G_c \otimes G_F$ with chiral fermions in several representations of this group. Those fermions which are

trivial under G_c are only coupled weakly and are called "spectator fermions". All anomalies cancel, by construction.

At low energies, much lower than the mass scale where color binding occurs, we see only the G_F gauge group with its gauge fields. Coupled to these gauge fields are the massless bound states, forming new representations r of G_F , with either left- or right handed chirality. The numbers of left minus right handed fermion fields in the representations r are given by the as yet unknown indices $\ell(r)$. And finally we have the spectator fermions which are unchanged.

We now expect these very light objects to be described by a new local field theory, that is, a theory local with respect to the large distance scale that we now use. The central theme of our reasoning is now that this new theory must again be anomaly free. We simply cannot allow the contradictions that would arise if this were not so. Nature must arrange its new particle spectrum in such a way that unitarity is obeyed, and because of the large distance scale used the effective interactions are either vanishingly small or renormalizable. The requirement of anomaly cancellation in the new particle spectrum gives us equations for the indices $\ell(r)$, as we will see.

The reason why these equations are sometimes difficult or impossible to solve is that the new representations r must be different from the old ones; if $G_c = SU(N)$ then r must also be faithful representations of $G_F/Z(N)$. For instance in QCD we only allow for octet or decuplet representations of $(SU(3))_{\text{flavor}}$, whereas the original quarks were triplets.

However, the anomaly cancellation requirement, restrictive as it may be, does not fix the values of $\ell(r)$ completely. We must look for additional limitations.

III12 APPELQUIST-CARAZZONE DECOUPLING AND N-INDEPENDENCE

A further limitation is found by the following argument. Suppose we add a mass term for one of the colored fermions.

$$\Delta\mathcal{L} = m \bar{\psi}_{1L} \psi_{1R} + \text{h.c.}$$

Clearly this links one of the left handed fermions with one of the right handed ones and thus reduces the flavor group G_F into $G_F' \subset G_F$. Now let us gradually vary m from 0 to infinity. A famous theorem 5) tells us that in the limit $m \rightarrow \infty$ all effects due to this massive quark disappear. All bound states containing this quark should also disappear which they can only do by becoming very heavy. And they can only become heavy if they form representations r' of G_F' with total index $\ell'(r') = 0$. Each representation r of G_F forms

an array of representations r' of G'_F . Therefore

$$\ell'(r') = \sum_{r \text{ with } r' \subset r} \ell(r) \quad . \tag{III40}$$

Apparently this expression must vanish.

Thus we found another requirement for the indices $\ell(r)$. The indices will be nearly but not quite uniquely determined now. Calculations show that this second requirement makes our indices $\ell(r)$ practically independent of the dimensions n_i of G_F . For instance, if $G_C = SU(3)$ and if we have left- and righthanded quarks forming triplets and sextets then

$$G_F = SU(n_1)_L \otimes SU(n_2)_R \otimes SU(n_3)_L \otimes SU(n_4)_R \otimes U(1)^3 \tag{III41}$$

where $n_{1,2}$ refer to the triplets and $n_{3,4}$ to the sextets. G_C is anomaly-free if

$$n_1 - n_2 + 7(n_3 - n_4) = 0 \quad . \tag{III42}$$

Here we have three independent numbers n_i . If we write the representations r as Young tableaux then $\ell(r)$ could still depend explicitly on n_i .

However, suppose that someone would start as approximation of Bethe-Salpeter type to discover the zero mass bound state spectrum. He would study diagrams such as Fig. 3




Fig. 3

The resulting indices $\ell(r)$ would follow from topological properties of the interactions represented by the blobs. It is unlikely that this topology would be seriously effected by details such as the contributions of diagrams containing additional closed fermion loops. However, that is the only way in which explicit n -dependence enters. It is therefore natural to assume $\ell(r)$ to be n -independent. This latter assumption fixes $\ell(r)$ completely. What is the result of these calculations?

III13. CALCULATIONS

Let G be any (reducible or irreducible) gauge group. Let chiral fermions in a representation r be coupled to the gauge fields by the covariant derivative

$$D_\mu = \partial_\mu + i \lambda^a(r) A_\mu^a, \quad (\text{III43})$$

where A_μ^a are the gauge fields and $\lambda^a(r)$ a set of matrices depending on the μ representation r . Let the left-handed fermions be in the representations r_L and the right-handed ones in r_R . Then the anomalies cancel if

$$\sum_L \text{Tr}\{\lambda^a(r_L), \lambda^b(r_L)\} \lambda^c(r_L) = \sum_R \text{Tr}\{\lambda^a(r_R), \lambda^b(r_R)\} \lambda^c(r_R). \quad (\text{III44})$$

The object $d^{abc}(r) = \text{Tr}\{\lambda^a(r), \lambda^b(r)\} \lambda^c(r)$ can be computed for any r . In table 1 we give some examples. The fundamental representation r_0 is represented by a Young tableau: \square . Let it have n components. We take the case that $\text{Tr} \lambda(r_0) = 0$. Write

$$\begin{aligned} \text{Tr} I(r_0) &= n, & \text{Tr} I(r) &= N(r), \\ \text{Tr} \lambda(r) &= 0, \\ \text{Tr} \lambda^a(r) \lambda^b(r) &= C(r) \text{Tr} \lambda^a(r_0) \lambda^b(r_0), \\ d^{abc}(r) &= K(r) d^{abc}(r_0). \end{aligned} \quad (\text{III45})$$

We read off C and K from table 1.








Now III44 must hold both in the high energy region and in the low energy region. The contribution of the spectator fermions in both regions is the same. Thus we get for the bound states

$$\left(\sum_L - \sum_R \right) d^{abc}(r) = n_c \left(d^{abc}(r_{oL}) - d^{abc}(r_{oR}) \right) \quad (\text{III46})$$

where a, b, c are indices of G_F and r_0 is the fundamental representation of G_F . We have the factor n_c written explicitly, being the number of color components.

Let us now consider the case $G_c = \text{SU}(3)$;
 $G_F = \text{SU}_L(n) \otimes \text{SU}_R(n) \otimes \text{U}(1)$. We have n "quarks" in the fundamental representations. The representations r of the bound states must be in $G_F/Z(3)$. They are assumed to be built from three quarks, but we are free to choose their chirality. The expected representations

Table 1

r	N(r)	C(r)	K(r)
	n	1	1
	n	1	-1
<div><div></div><div></div></div>	$\frac{n(n+1)}{2}$	$n+2$	$n+4$
<div><div></div><div></div></div>	$\frac{n(n+1)(n+2)}{6}$	$\frac{(n+2)(n+3)}{2}$	$\frac{(n+3)(n+6)}{2}$
	$\frac{n(n^2-1)}{3}$	n^2-3	n^2-9
A ⊗ B	N(A)N(B)	C(A)N(B) + C(B)N(A)	K(A)N(B) + K(B)N(A)

are given in table 2, where also their indices are defined. Because of left-right symmetry these numbers change sign under interchange of left \leftrightarrow right.

Table 2

representation	index	representation	index
$\begin{array}{ c c c } \hline L & L & L \\ \hline \end{array}$ $\left. \begin{array}{ c } \hline L \\ \hline L \\ \hline L \\ \hline \end{array} \right\}$	$\ell_{1\pm}$	$\begin{array}{ c c c } \hline R & R & R \\ \hline \end{array}$ $\left. \begin{array}{ c } \hline R \\ \hline R \\ \hline R \\ \hline \end{array} \right\}$	$-\ell_{1\pm}$
$\begin{array}{ c } \hline L \\ \hline \end{array} \otimes \begin{array}{ c c } \hline R & R \\ \hline \end{array}$ $\left. \begin{array}{ c } \hline L \\ \hline \end{array} \otimes \begin{array}{ c } \hline R \\ \hline R \\ \hline \end{array} \right\}$	$\ell_{2\pm}$	$\begin{array}{ c } \hline R \\ \hline \end{array} \otimes \begin{array}{ c c } \hline L & L \\ \hline \end{array}$ $\left. \begin{array}{ c } \hline R \\ \hline \end{array} \otimes \begin{array}{ c } \hline L \\ \hline L \\ \hline \end{array} \right\}$	$-\ell_{2\pm}$
$\begin{array}{ c c } \hline L & L \\ \hline L & \\ \hline \end{array}$	ℓ_3	$\begin{array}{ c c } \hline R & R \\ \hline R & \\ \hline \end{array}$	$-\ell_3$

For the time being we assume no other representations. In eq. III46 we may either choose a, b and c all to be $SU(n)_L$ indices, or choose a and b to be $SU(n)_L$ indices and c the $U(1)$ index. We get two independent equations:

$$\sum_{\pm} \frac{1}{2}(n \pm 3)(n \pm 6)\ell_{1\pm} - \sum_{\pm} \frac{1}{2}n(n \pm 7)\ell_{2\pm} + (n^2 - 9)\ell_3 = 3, \text{ if } n > 2,$$

and

$$\sum_{\pm} \frac{1}{2}(n \pm 2)(n \pm 3)\ell_{1\pm} - \sum_{\pm} \frac{1}{2}n(n \pm 3)\ell_{2\pm} + (n^2 - 3)\ell_3 = 1, \text{ if } n > 1.$$

(III47)

The Appelquist-Carazzone decoupling requirement, eq. (III40), gives us in addition two other equations:

$$\ell_{1+} - \ell_{2+} + \ell_3 = 0,$$

$$\ell_{1-} - \ell_{2-} + \ell_3 = 0, \text{ both if } n > 2.$$

(III48)

For $n > 2$ the general solution is

$$\begin{aligned}
\ell_{1+} &= \ell_{1-} = \ell, \\
\ell_{2+} &= \ell_{2-} = 3\ell - \frac{1}{3}, \\
\ell_3 &= 2\ell - \frac{1}{3}.
\end{aligned}
\tag{III49}$$

Here ℓ is still arbitrary. Clearly this result is unacceptable. We cannot allow any of the indices ℓ to be non-integer. Only for the case $n = 2$ (QCD with just two flavors) there is another solution. In that case ℓ_{2-} and ℓ_3 describe the same representation, and ℓ_{1-} an empty representation. We get

$$\ell_{2-} + \ell_3 = k = 1 - 10 \ell_{1+} + 5 \ell_{2+}. \tag{III50}$$

According to the σ -model, $\ell_{1+} = \ell_{2+} = 0$; $k = 1$. The σ -model is therefore a correct solution to our equations.

In the previous section we promised to determine the indices completely. This is done by imposing n -independence for the more general case including also other color representations such as sextets besides triplets. The resulting equations are not very illuminating, with rather ugly coefficients. One finds that in general no solution exists except when one assumes that all mixed representations have vanishing indices. With mixed representations we mean a product of two or more non-trivial representations of two or more non-Abelian invariant subgroups of G_F . If now we assume n -independence this must also hold if the number of sextets is zero. So ℓ_{2+} and ℓ_{2-} must vanish. We get

$$\begin{aligned}
\ell_{1+} &= \ell_{1-} = 1/9, \\
\ell_3 &= -1/9.
\end{aligned}
\tag{III51}$$

If all quarks were sextets, not triplets, we would get

$$\begin{aligned}
\ell_{1+} &= \ell_{1-} = 2/9, \\
\ell_3 &= -2/9.
\end{aligned}
\tag{III52}$$

In the case $G_C = SU(5)$ the indices were also found. See table 3.

Table 3
indices for $G_c = SU(5)$

$\begin{array}{ c c c c c } \hline L & L & L & L & L \\ \hline \end{array}$	$\ell_{1+} = 1/25$	
$\begin{array}{ c c c c } \hline L & L & L & L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$	$\ell_{2+} = -1/25$	
$\begin{array}{ c c c } \hline L & L & L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$	$\ell_3 = 1/25$	
$\begin{array}{ c c } \hline L & L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$	$\ell_{2-} = -1/25$	
$\begin{array}{ c } \hline L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$	$\ell_{1-} = 1/25$	
		$\left. \begin{array}{ c c c } \hline L & L & L \\ \hline \end{array} \right\} \ell_{4\pm} = 0$ $\left. \begin{array}{ c c } \hline L & L \\ \hline \end{array} \right\}$ $\begin{array}{ c } \hline L \\ \hline \end{array}$

This clearly suggests a general tendency for $SU(N)$ color groups to produce indices $\pm 1/N^2$ or 0.

III14. CONCLUSIONS

Our result that the indices we searched for are fractional is clearly absurd. We nevertheless pursued this calculation in order to exhibit the general philosophy of this approach and to find out what a possible cure might be. Our starting point was that chiral symmetry is not broken spontaneously. Most likely this is untenable, as several authors have argued⁶⁾. We find that explicit chiral symmetry in QCD leads to trouble in particular if the number of flavors is more than two. A daring conjecture is then that in QCD the strange quark, being rather light, is responsible for the spontaneous breakdown of chiral symmetry.

An interesting possibility is that in some generalized versions of QCD chiral symmetry is broken only partly, leaving a few massless chiral bound states. Indeed there are examples of models where our philosophy would then give integer indices, but since we must drop the requirement of n -dependence our result was not unique and it was always ugly. No such model seems to reproduce anything resembling the observed quark-lepton spectrum.

Finally there is the remote possibility that the paradoxes associated with higher spin massless bound states can be resolved. Perhaps the $\Delta(1236)$ plays a more subtle role in the σ -model than assumed so far (we took it to be a parity doublet).

We conclude that we are unable to construct a bound state theory for the presently fundamental fermions along the lines

suggested above.

We thank R. van Damme for a calculation yielding the indices in the case $G_c = \text{SU}(5)$.

REFERENCES

1. P.A.M. Dirac, *Nature* 139 (1937) 323, *Proc. Roy. Soc.* A165 (1938) 199, and in: *Current Trends in the Theory of Fields*, (Tallahassee 1978) AIP Conf. Proc. No 48, *Particles and Fields Subseries* No 15, ed. by Lannuti and Williams, p. 169.
2. S. Dimopoulos and L. Susskind, *Nucl. Phys.* B155 (1979) 237.
3. M. Gell-Mann and M. Lévy, *Nuovo Cim.* 16 (1960) 705.
B.W. Lee, *Chiral Dynamics*, Gordon and Breach, New York, London, Paris 1972.
4. G. 't Hooft, *Phys. Rev. Lett.* 37 (1976) 8; *Phys. Rev.* D14 (1976) 3432.
S. Coleman, "The Uses of Instantons", *Erice Lectures* 1977.
R. Jackiw and C. Rebbi, *Phys. Rev. Lett.* 37 (1976) 172.
C. Callan, R. Dashen and D. Gross, *Phys. Lett.* 63B (1976) 334.
5. G. 't Hooft, *Nucl. Phys.* B72 (1974) 461.
6. T. Appelquist and J. Carazzone, *Phys. Rev.* D11 (1975) 2856.
7. A. Casher, *Chiral Symmetry Breaking in Quark Confining Theories*, Tel Aviv preprint TAUP 734/79 (1979).