

Quantum Optics with Giant Atoms—the First Five Years



Anton Frisk Kockum

Abstract In quantum optics, it is common to assume that atoms can be approximated as point-like compared to the wavelength of the light they interact with. However, recent advances in experiments with artificial atoms built from superconducting circuits have shown that this assumption can be violated. Instead, these artificial atoms can couple to an electromagnetic field at multiple points, which are spaced wavelength distances apart. In this chapter, we present a survey of such systems, which we call *giant atoms*. The main novelty of giant atoms is that the multiple coupling points give rise to interference effects that are not present in quantum optics with ordinary, small atoms. We discuss both theoretical and experimental results for single and multiple giant atoms, and show how the interference effects can be used for interesting applications. We also give an outlook for this emerging field of quantum optics.

Keywords Quantum optics · Giant atoms · Waveguide QED · Relaxation rate · Lamb shift · Superconducting qubits · Surface acoustic waves · Cold atoms

1 Introduction

Natural atoms are so small (radius $r \approx 10^{-10}$ m) that they can be considered point-like when they interact with light at optical frequencies (wavelength $\lambda \approx 10^{-6}$ – 10^{-7} m) (Leibfried et al. 2003). If the atoms are excited to high Rydberg states, they can reach larger sizes ($r \approx 10^{-8}$ – 10^{-7} m), but quantum-optics experiments with such atoms have them interact with microwave radiation, which has much longer wavelength ($\lambda \approx 10^{-2}$ – 10^{-1} m) (Haroche 2013). It has thus been well justified in theoretical treatments of quantum optics to assume $r \ll \lambda$, called the *dipole approximation*,

A. Frisk Kockum (✉)
Wallenberg Centre for Quantum Technology, Chalmers University of Technology, 412 96
Gothenburg, Sweden
e-mail: anton.frisk.kockum@chalmers.se

which simplifies the description of the interaction between light and matter (Walls and Milburn 2008).

In recent years, experimental investigations of quantum optics have expanded to systems with *artificial atoms*, i.e., engineered quantum systems such as quantum dots (Hanson et al. 2007) and superconducting quantum bits (qubits) (You and Nori 2011; Xiang et al. 2013; Gu et al. 2017; Kockum and Nori 2019), which emulate essential aspects of natural atoms. The circuits making up superconducting qubits can be large, reaching sizes up to $r \approx 10^{-4}\text{--}10^{-3}$ m, but this is still small when compared with the wavelength of the microwave fields they interact with.

In 2014, one experiment (Gustafsson et al. 2014) forced quantum opticians to reconsider the dipole approximation. In that experiment, a superconducting transmon qubit (Koch et al. 2007) was coupled to surface acoustic waves (SAWs) (Datta 1986; Morgan 2007). Due to the low propagation velocity of SAWs, their wavelength was $\lambda \approx 10^{-6}$ m, and the qubit, due to its layout with an interdigitated capacitance, coupled to the SAWs at multiple points, which were spaced $\lambda/4$ apart.

Motivated by this experiment, theoretical investigations on *giant atoms* were initiated (Kockum et al. 2014). The main finding was that the multiple coupling points lead to interference effects, e.g., the coupling of the giant atom to its environment becomes frequency-dependent (Kockum et al. 2014).

These initial experimental and theoretical works on giant atoms were published 5 years ago, at the time of writing for this book chapter. In this chapter, we give a brief survey of the developments in the field of quantum optics with giant atoms that have followed since. We begin in Sect. 2 with theory for giant atoms, looking first at the properties of a single giant atom (Sect. 2.1), including what happens when the coupling points are extremely far apart (Sect. 2.2), and then at multiple giant atoms (Sect. 2.3). In Sect. 3, we survey the different experimental systems where giant atoms have been implemented or proposed. We conclude with an outlook (Sect. 4) for future work on giant atoms, pointing to several areas where interesting results can be expected.

2 Theory for Giant Atoms

The experimental setup where giant atoms were first implemented (Gustafsson et al. 2014) falls into the category of waveguide quantum electrodynamics (QED). In waveguide QED (Gu et al. 2017; Roy et al. 2017), a continuum of bosonic modes can propagate in a one-dimensional (1D) waveguide and interact with atoms coupled to this waveguide. As reviewed in Gu et al. (2017), Roy et al. (2017), there is an abundance of theoretical papers dealing with one, two, or more atoms coupled to a 1D waveguide, but they almost all assume that the dipole approximation is valid, or, in other words, that the atoms are “small”.

The difference between small and giant atoms is illustrated in Fig. 1. While a small atom, because of its diminutive extent, can be described as being connected to the waveguide at a single point, a giant atom couples to the waveguide at multiple




Fig. 1 The difference between a small atom and a giant atom. **a** A small atom (two levels) couples to the 1D waveguide (grey) at a single point (red, coordinate x_1). **b** A giant atom couples to the waveguide at multiple points (labelled k , coordinates x_k). The distance between two coupling points k and n , $|x_k - x_n|$, is *not* negligible compared to the wavelength of the modes in the waveguide that the atom interacts with

points, and the distance between these points *cannot* be neglected in comparison to the wavelength of the modes in the waveguide that couple to the atom. The relevant wavelength λ to compare with is set by the (angular) transition frequency ω_a of the atom and the propagation velocity v in the waveguide: $\lambda = 2\pi v/\omega_a$.

2.1 One Giant Atom

Quantum optics with a single giant atom was first studied theoretically in Kockum et al. (2014), prompted by the experiment in Gustafsson et al. (2014) (discussed in Sect. 3.1). For a small atom coupled to a continuum of modes, like in Fig. 1a, standard quantum-optics procedure is to derive a master equation by assuming that the coupling to the modes is relatively weak and tracing out the modes (Carmichael 1999; Gardiner and Zoller 2004; Walls and Milburn 2008). When considering whether the same procedure can be applied to a giant atom, there is a new timescale to take into account: the time it takes to travel in the waveguide between coupling points. In Kockum et al. (2014), this time was assumed small compared to the time it takes for an excitation in the atom to relax into the waveguide. With this assumption, the system is *Markovian*, i.e., the time evolution of the atom only depends on the present state of the system, not on the past (for the non-Markovian case, see Sect. 2.2). Thus, the standard master-equation derivation from quantum optics with small atoms can be applied here as well.

2.1.1 Master Equation for a Giant Atom

The derivation of a master equation for a giant atom starts from the total system Hamiltonian (we use units where $\hbar = 1$ throughout this chapter),

$$H = H_a + H_{wg} + H_I, \quad (1)$$

with the bare atomic Hamiltonian

$$H_a = \sum_m \omega_m |m\rangle\langle m|, \quad (2)$$

the bare waveguide Hamiltonian

$$H_{wg} = \sum_j \omega_j (a_{Rj}^\dagger a_{Rj} + a_{Lj}^\dagger a_{Lj}), \quad (3)$$

and the interaction Hamiltonian

$$H_I = \sum_{j,k,m} g_{jkm} \left(\sigma_-^{(m)} + \sigma_+^{(m)} \right) \times \left(a_{Rj} e^{-i\omega_j x_k/v} + a_{Lj} e^{i\omega_j x_k/v} + a_{Rj}^\dagger e^{i\omega_j x_k/v} + a_{Lj}^\dagger e^{-i\omega_j x_k/v} \right). \quad (4)$$

Here, the atomic levels are labelled $m = 0, 1, 2, \dots$, have energies ω_m , and are connected through lowering and raising operators $\sigma_-^{(m)} = |m\rangle\langle m+1|$ and $\sigma_+^{(m)} = |m+1\rangle\langle m|$. The bosonic modes in the waveguide are labelled with indices j and with an index R (L) for right-moving (left-moving) modes. The corresponding annihilation and creation operators are a and a^\dagger , respectively. The difference to the case of a small atom is the sum over coupling points labelled by k in Eq. (4). The phase factors $e^{\pm i\omega_j x_k/v}$ are not present for a small atom. These phase factors give rise to interference effects. Note that the coupling strengths g_{jkm} can depend on both j , k , and m .

Following the standard master-equation derivation using the Born-Markov approximation, the resulting master equation becomes

$$\dot{\rho} = -i \left[\sum_m (\omega_m + \Delta_m) |m\rangle\langle m|, \rho \right] + \sum_m \Gamma_{m+1,m} \mathcal{D} \left[\sigma_-^{(m)} \right] \rho, \quad (5)$$

where ρ is the density matrix for the atom, $\mathcal{D}[X]\rho = X\rho X^\dagger - \frac{1}{2}X^\dagger X\rho - \frac{1}{2}\rho X^\dagger X$ is the Lindblad superoperator describing relaxation (Lindblad 1976), and we have assumed negligible temperature T , i.e., $\omega_m \gg k_B T$. The relaxation rates for the atomic transitions $|m+1\rangle \rightarrow |m\rangle$ are

$$\Gamma_{m+1,m} = 4\pi J(\omega_{m+1,m}) |A_m(\omega_{m+1,m})|^2, \quad (6)$$

where $\omega_{a,b} = \omega_a - \omega_b$, $J(\omega)$ is the density of states at frequency ω in the waveguide, and we have defined

$$A_m(\omega_j) = \sum_k g_{jkm} e^{i\omega_j x_k/v}. \quad (7)$$

The frequency shifts Δ_m of the atomic energy levels are Lamb shifts (Lamb and Rutherford 1947; Bethe 1947) given by

$$\Delta_m = 2\mathcal{P} \int_0^\infty d\omega \frac{J(\omega)}{\omega} \left(\frac{|A_m(\omega)|^2 \omega_{m+1,m}}{\omega + \omega_{m+1,m}} - \frac{|A_{m-1}(\omega)|^2 \omega_{m,m-1}}{\omega - \omega_{m,m-1}} \right). \quad (8)$$

Both the relaxation rates and the Lamb shifts acquire a strong dependence on the atomic transition frequencies, encoded in the factor $A_m(\omega_j)$. For the case of a small atom, $A_m(\omega_j) = g_{jm}$, which is a constant provided that g_{jm} does not depend strongly on j . The effect of this frequency dependence for giant atoms can be seen clearly if one considers the simple case of an atom with two coupling points x_1 and x_2 [compare Fig. 1b] having equally strong coupling to the waveguide. If the two points are half a wavelength apart, i.e., $|x_1 - x_2| = \pi v / \omega_{m+1,m}$, there will be destructive interference between emission from the two points, and the relaxation for the corresponding atomic transition is completely suppressed: $\Gamma_{m+1,m} = 0$. If the two points are one wavelength apart, there is instead constructive interference and the relaxation rate is enhanced.

2.1.2 Frequency-Dependent Relaxation Rate

To further understand the frequency-dependence of the relaxation rates and the Lamb shifts, consider the case of a two-level atom coupled to the waveguide at N equidistant points with equal coupling strength at each point. In this case, introducing the notation $\varphi = \omega_{1,0}(x_2 - x_1)/v$, we obtain (Kockum et al. 2014)

$$\Gamma_{1,0} = \gamma \frac{\sin^2\left(\frac{N}{2}\varphi\right)}{\sin\left(\frac{1}{2}\varphi\right)} = \gamma \frac{1 - \cos(N\varphi)}{1 - \cos(\varphi)}, \quad (9)$$

$$\Delta_1 = \gamma \frac{N \sin(\varphi) - \sin(N\varphi)}{2[1 - \cos(\varphi)]}, \quad (10)$$

where γ is the relaxation rate that the atom would have had if it was coupled to the waveguide only at a single point. To obtain the Lamb shift, we have also made the simplifying assumption that $J(\omega)$ is constant, that the lower limit of the integral in Eq. (8) can be extended down to $-\infty$, and that only the dominating second term in that integral contributes. Since $\Delta_0 = 0$ with these assumptions, Eq. (10) gives the full frequency shift for the two-level atom. In fact, the relaxation rate and the Lamb shift are related through a Hilbert transform due to Kramers–Kronig relations (Cohen-Tannoudji et al. 1998).

The relaxation rates and Lamb shifts in Eqs. (9)–(10) are plotted for two values of N in Fig. 2. The central peak corresponds to the distance between neighbouring coupling points being one wavelength. Note that the frequency dependence becomes sharper when more coupling points are added; in frequency units, the width of the central peak is approximately $\omega_{1,0}/2\pi N$. This sharpness can be used to determine when the Markovian approximation underlying the master-equation derivation breaks down, which happens roughly when the relaxation rate changes noticeably within the linewidth of the atom, i.e., when $\Gamma_{1,0} \approx \omega_{1,0}/2\pi N$. Interestingly, this is




Fig. 2 Relaxation rates and Lamb shifts for a giant two-level atom with symmetrically spaced coupling points all having the same coupling strength. Red curves: $N = 3$ coupling points. Blue curves: $N = 10$ coupling points. Solid curves: Relaxation rates $\Gamma_{1,0}$. Dashed curves: Lamb shifts Δ_1 . The relaxation rates and Lamb shifts are scaled to the maximum relaxation rate Γ_{\max} for each N . Figure adapted from Kockum et al. (2014) with permission

approximately the same condition as when the travelling time between the outermost coupling points, $2\pi(N - 1)/\omega_{1,0}$, becomes comparable to the relaxation time $1/\Gamma_{1,0}$.

An attractive feature of giant atoms is that the frequency-dependence of their relaxation rates (and Lamb shifts) can be *designed* (Kockum et al. 2014). The frequency dependence is directly determined by Eq.(7), which simply is a discrete Fourier transform of the coupling-point coordinates, weighted by the coupling strength in each point. With N coupling points, an experimentalist thus has $2N - 1$ knobs to turn (the translational invariance of the setup removes one degree of freedom). With enough coupling points, the curves in Fig. 2 can be moulded into any shape. Note that although the coupling-point coordinates and coupling strengths will be fixed in an experiment, superconducting qubits offer the possibility to tune the atomic frequency widely *in situ* (Gu et al. 2017; Kockum and Nori 2019), making it possible to move between regions with high and low relaxation rates during an experiment.

If we consider more than two atomic levels, other interesting applications of the frequency-dependent relaxation rate open up. As illustrated in Fig. 3, if the atomic transition frequencies $\omega_{1,0} \neq \omega_{2,1}$, it is possible to engineer the relaxation rates such that $\Gamma_{2,1}$ is at a maximum when $\Gamma_{1,0}$ is at a minimum. At that point, one can then create population inversion, and thus lasing, by driving the transition from $|0\rangle$ to $|2\rangle$ (Kockum et al. 2014). Recent experiments have been making use of this possibility to control the ratio of relaxation rates to enable electromagnetically induced transparency (EIT) (Andersson et al. 2020; Vadiraj et al. 2020).




Fig. 3 Engineering population inversion in a giant atom. The blue curve and the red curve are the relaxation rates $\Gamma_{1,0}$ and $\Gamma_{2,1}$, respectively, as a function of transition frequency $\omega_{1,0}$. The plot assumes $N = 10$ equally spaced coupling points, with equal coupling strengths at all points, and an anharmonicity $\omega_{2,1} - \omega_{1,0} = -0.1 \times 2\pi\nu/(x_2 - x_1)$. The inset shows the level structure with the relaxation rates and a drive of strength Ω_d on the $|0\rangle \leftrightarrow |2\rangle$ transition. Figure adapted from Kockum et al. (2014) with permission

2.1.3 Comparison with an Atom in Front of a Mirror

It is possible to engineer frequency-dependent relaxation rates and Lamb shifts also for small atoms. This can be achieved by placing a small atom in front of a mirror instead of in an open waveguide, a setup which has been considered in several theoretical (Meschede et al. 1990; Dorner and Zoller 2002; Beige et al. 2002; Dong et al. 2009; Koshino and Nakamura 2012; Wang et al. 2012; Tufarelli et al. 2013; Fang and Baranger 2015; Shi et al. 2015; Pichler and Zoller 2016) and experimental works (Eschner et al. 2001; Wilson et al. 2003; Dubin et al. 2007; Hoi et al. 2015; Wen et al. 2018, 2019). Here, the atomic relaxation can be enhanced or suppressed by interference with the mirror image of the atom. This setup is equivalent to a giant atom with two coupling points in a unidirectional waveguide.

However, this is the limit with a small atom in front of a mirror. In such a setup, it is not possible to increase the number of coupling points, or to have different coupling strengths at different coupling points, which means that the frequency dependence cannot be designed like for a giant atom. Furthermore, since propagation is unidirectional, it is not possible to have more advanced scattering, possible with a giant atom, where both reflection and transmission are influenced by interference between coupling points.

2.1.4 Coupling a Giant Atom to a Cavity

By introducing reflective boundary conditions at both ends of the waveguide in Fig. 1, a multimode cavity will be formed. The coupling of a giant atom to such a cavity has yet to be explored as thoroughly as the open-waveguide case. We can see that similar interference effects as in the open waveguide will come into play. It will thus, for example, be possible to arrange the coupling points such that the giant atom couples strongly to some modes of the cavity and is decoupled from other modes. This can to some extent already be achieved with a small atom, whose single coupling point can be at a node for some modes and at an antinode for others. However, we note that a recent theory proposal (Ciani and DiVincenzo 2017) uses a superconducting qubit with tunable coupling connected at multiple points to two resonators to cancel certain unwanted interaction terms while keeping desired interaction terms; it is shown that this would not have been possible with a small atom.

2.2 One Giant Atom with Time Delay

Consider a giant atom with two coupling points spaced such that it takes a time τ for light (or sound) to travel between them. In the previous section, it was assumed that τ was small compared to the relaxation time $1/\Gamma$. When this no longer is the case, the giant atom enters the non-Markovian regime, where the time evolution of the system can depend on what the system state was at an earlier time. In a giant atom, this non-Markovianity can manifest itself in revivals of the atomic population if energy is sent out from the atom at one coupling point and later is reabsorbed at another coupling point.

Four theoretical studies (Guo et al. 2017; Ask et al. 2019a; Guo et al. 2019, 2020) have explored this regime (the latter three considering more than two coupling points). In Ask et al. (2019a), it was shown that $\Gamma\tau = 1$ constitutes a sharp border for when time-delay effects become visible. When the system transitions from $\Gamma\tau < 1$ to $\Gamma\tau > 1$, the response of the giant atom to a weak coherent probe goes from showing one resonance to showing two. This is similar to the appearance of a vacuum Rabi splitting when an atom becomes strongly coupled to a cavity (the mathematical condition for the appearance of the splitting is actually exactly the same as for an atom in a multimode cavity Ask et al. 2019a; Krimer et al. 2014). In the case of the giant atom, the multiple coupling points act as a cavity when the coupling becomes strong enough or the travelling time becomes long enough.

In Guo et al. (2017), the cases $\tau > \Gamma$ and $\tau \gg \Gamma$ were studied in more detail. As τ increases, an initially excited giant atom exhibits more and more revivals of its population. In the limit of large τ , it turns out that the total energy stored in the giant atom and between its coupling points no longer decays exponentially with time t , as for a small atom, but instead decays polynomially ($\propto 1/\sqrt{t}$). Furthermore, the timescale for this decay is no longer set by the decay rate Γ , but by the travel time τ . These predictions for a giant atom with time delay were recently confirmed in an experiment (Andersson et al. 2019) (see Sect. 3.1 for more on the experimental platform used).

In Guo et al. (2019), it was shown that extending the setup from Guo et al. (2017) to more three or more coupling points enables qualitatively different phenomena: oscillating bound states. These oscillating bound states do not decay into the waveguide, but the energy oscillates persistently between the atom and the waveguide modes in-between the outermost coupling points of the atom. This result appears connected to that of Ask et al. (2019a) discussed above, and similar results have been obtained in Guo et al. (2020).

There are similarities between a giant atom with time delay and the previously studied (Dorner and Zoller 2002; Tufarelli et al. 2013; Pichler and Zoller 2016) setup with a small atom placed far from a mirror. However, in the giant-atom case scattering processes will involve both reflection and transmission, and the second-order correlation functions for these signals, calculated in Guo et al. (2017), exhibit oscillations between bunching and anti-bunching on a timescale set by τ .

2.3 *Multiple Giant Atoms*

When multiple small atoms are coupled to a waveguide, they can be spaced wavelength distances apart, which leads to interference effects influencing the collective behaviour of the atoms (Gu et al. 2017; Roy et al. 2017; Lehmberg 1970b, a; Lalumière et al. 2013; Zheng and Baranger 2013). Well-known examples include super- and sub-radiance (Dicke 1954; Lalumière et al. 2013), i.e., increased and decreased emission rates due to collective decay, and an effective coupling (sometimes called collective Lamb shift) between pairs of atoms, mediated by virtual photons in the transmission line (Friedberg et al. 1973; Scully and Svidzinsky 2010; Wen et al. 2019). Given this, one might wonder whether there is something left to set multiple giant atoms apart from multiple small atoms. After all, it was mainly the interference effects that separated a single giant atom from a single small atom.

In Kockum et al. (2018), the properties of multiple giant atoms were studied thoroughly and compared to those of multiple small atoms. The simplest cases considered are pictured in Fig. 4. For each of these setups, a master equation of the same form can be derived, assuming again that the travel time between coupling points is negligible:

$$\begin{aligned} \dot{\rho} = & -i \left[\omega'_a \frac{\sigma_z^a}{2} + \omega'_b \frac{\sigma_z^b}{2} + g (\sigma_-^a \sigma_+^b + \sigma_+^a \sigma_-^b), \rho \right] \\ & + \Gamma_a \mathcal{D}[\sigma_-^a] \rho + \Gamma_b \mathcal{D}[\sigma_-^b] \rho + \Gamma_{\text{coll}} \left[\left(\sigma_-^a \rho \sigma_+^b - \frac{1}{2} \{ \sigma_+^a \sigma_-^b, \rho \} \right) + \text{H.c.} \right], \end{aligned} \quad (11)$$

where ω'_j is the transition frequency of atom j (we label the left atom a and the right atom b) including Lamb shifts, g is the strength of the exchange interaction mediated by the waveguide between the atoms, Γ_j is the individual relaxation rate of atom j , Γ_{coll} is the collective relaxation rate, and H.c. denotes Hermitian conjugate.




Fig. 4 Setups for two small and two giant atoms. **a** Two small atoms in an open waveguide. **b** Two small atoms in a waveguide terminated by a mirror on the left. **c** Two “separate” giant atoms, where the rightmost coupling point of the left atom is left of the leftmost coupling point of the right atom. **d** Two “braided” giant atoms, where each atom has a coupling point that lies in between the two coupling points of the other atom. **e** Two “nested” giant atoms, where the coupling points of one atom all lie in-between the coupling points of the other atom. Figure adapted from Kockum et al. (2018) with permission

Assuming that the atoms couple to the waveguide with equal strength at each coupling point, and that the distances between neighbouring coupling points are equal, corresponding to a phase shift φ , the coefficients g , Γ_j , and Γ_{coll} in Eq. (11) have simple expressions as functions of φ (Kockum et al. 2018). These functions are plotted in Fig. 5 for all the setups in Fig. 4. Looking at the individual relaxation rates (dashed curves), we see that they are always non-zero for small atoms in an open waveguide, but for setups with giant atoms there are points where $\Gamma_j = 0$, as we know from the discussion of single giant atoms in Sect. 2.1. Furthermore, at the points where $\Gamma_j = 0$, the collective relaxation rate Γ_{coll} also goes to zero. It is thus clear that setups with multiple giant atoms can be completely protected from relaxation into the waveguide.

The most remarkable feature in Fig. 5 is found when looking at the behaviour of the exchange interaction g at the points where the relaxation rates are zero. One might think that since interference effects at these points prevent the atoms from relaxing into the waveguide, it should not be possible for the waveguide to mediate interaction between the atoms. However, it turns out that g can be *non-zero* here for one of the three giant-atom setups: the braided giant atoms. This effect has recently been confirmed in experiment (Kannan et al. 2020) (see Sect. 3.2 for more on the experimental platform used).

One way to understand this protected interaction is to note that $\Gamma_j = 0$ when the phase between the coupling points of atom j is an odd integer multiple of π . The




Fig. 5 Exchange interaction g (solid curves), individual relaxation rates Γ_j (dashed curves), and collective relaxation rates Γ_{coll} (dotted curves) as a function of φ for the setups in Fig. 4. The colours of the curves denote the ordering of coupling points: ab [small atoms, Fig. 4a, black], $aabb$ [separate giant atoms, Fig. 4c, blue], $abab$ [braided giant atoms, Fig. 4d, green], and $abba$ [nested giant atoms, Fig. 4e, red]. The last case is qualitatively equivalent to small atoms in front of a mirror [Fig. 4b]. For this case, there are two dashed curves (red), one for Γ_a and one for Γ_b . Figure adapted from Kockum et al. (2018) with permission

collective relaxation is due to interference between emission from coupling points of different atoms, but the sum total of these contributions is zero if the emissions from the two coupling points of one of the atoms interfere destructively. The exchange interaction arises due to emission from coupling points of one atom being absorbed at coupling points of the other atom. If the giant atoms are in the separate or nested configurations, the emissions from the two coupling points of atom b cancel if $\Gamma_b = 0$, but in the case of *braided* giant atoms, the two inner coupling points are placed in-between the coupling points of the other atom, so there is no condition forcing the contributions from the two coupling points of the other atom to interfere destructively.

We note that the protected interaction with braided giant atoms is reminiscent of the interaction between two small atoms in a waveguide with a bandgap (Kurizki 1990; Lambropoulos et al. 2000; Sundaresan et al. 2019). In that case, a bound state of photons forms around each atom that has a frequency in the bandgap, where propagation in the waveguide is impossible. The extension of these bound states decays exponentially with distance, but if two bound states overlap, the atoms can interact without decaying into the waveguide.

It is shown in Kockum et al. (2018) that the above conclusions about relations between relaxation rates and exchange interactions in giant atoms remain true even for the most general setups, with an arbitrary number of giant atoms, each having an arbitrary number of coupling points at arbitrary coordinates and with different

coupling strength at each coupling point. This opens up interesting possibilities for constructing larger setups with protected exchange interaction between many giant atoms (Kockum et al. 2018).

It is also interesting that the case of two small atoms in front of a mirror, equivalent to nested giant atoms (red curves in Fig. 5), allows interaction even if one (but not both) of the atoms is prevented from relaxing into the waveguide. This has recently been confirmed in an experiment (Wen et al. 2019) with superconducting qubits in a transmission-line waveguide, and expanded upon in a connected theoretical study (Lin et al. 2019).

Finally, we note that a recent theoretical study (Karg et al. 2019) extended the treatment from giant atoms to arbitrary quantum systems, e.g., harmonic oscillators, interacting with a waveguide at multiple points. The study took into account losses in the waveguide and also considered the impact of time delays, and showed how these factors can affect the protected interaction that is possible with a nested setup.

3 Experiments with Giant Atoms

Waveguide QED can be implemented in several experimental systems (Gu et al. 2017; Roy et al. 2017), e.g., with quantum dots coupled to photonic crystal waveguides (Arcari et al. 2014), with quantum emitters coupled to plasmons in nanowires (Akimov et al. 2007; Huck and Andersen 2016), and with natural atoms coupled to optical fibres (Bajcsy et al. 2009), but the most versatile platform at the moment appears to be superconducting qubits coupled to transmission lines (Gu et al. 2017; Astafiev et al. 2010a, b; Hoi et al. 2011, 2012; van Loo et al. 2013; Hoi et al. 2013, 2015; Liu and Houck 2017; Forn-Díaz et al. 2017; Wen et al. 2018; Mirhosseini et al. 2018, 2019; Sundaresan et al. 2019; Wen et al. 2019). There are thus many systems where giant atoms could be implemented. So far, as reviewed in this section, experiments have been conducted exclusively with superconducting qubits, coupled to either surface acoustic waves (SAWs, Sect. 3.1) or transmission lines (Sect. 3.2). A theoretical proposal exists for an implementation with cold atoms in optical lattices (Sect. 3.3), and we expect that experiments will eventually be performed using more platforms.

3.1 Superconducting Qubits and Surface Acoustic Waves

Superconducting qubits (You and Nori 2011; Xiang et al. 2013; Gu et al. 2017; Kockum and Nori 2019) are electrical circuits with capacitances, inductances, and Josephson junctions (which function as non-linear inductances) that can emulate properties of natural atoms, e.g., energy-level structures and coupling to an electromagnetic field. These circuits usually have resonance frequencies ω on the order of




Fig. 6 Experimental implementation of a giant atom with a superconducting qubit coupled to SAWs. **a** Sketch of the experimental setup. The IDT on the left is used both to send out SAWs to the right towards the qubit and to convert reflected SAW signals from the qubit into a voltage signal that can be read out. The qubit on the right has its capacitance formed like an IDT to interact with the SAWs. The two islands of the capacitance are also connected through two Josephson junctions (boxes with crosses), which function as a non-linear inductance, making the qubit essentially an anharmonic LC oscillator. The qubit can also be driven electrically through a gate on the top. **b** False-colour image of the experimental sample. The blue parts are the IDT to the right and the qubit to the left. The yellow parts are ground planes and the electrodes connecting to the IDT. The aspect ratio of the IDT, with fingers being much longer than they are wide, collimates the SAW beam such that it travels straight towards the qubit (and also in the opposite direction). Figure from Aref et al. (2016) with permission

GHz and are cooled to low temperatures $T \ll \hbar\omega/k_B$ to prevent the thermal fluctuations interfering with quantum properties.

In 2014, an experiment (Gustafsson et al. 2014) managed to couple a superconducting qubit of the transmon type (Koch et al. 2007) to SAWs, which are vibrations that propagate on the surface of a substrate (Datta 1986; Morgan 2007). The experimental setup is shown in Fig. 6. The substrate on which the SAWs propagate is piezoelectric, which means that the vibrations acquire an electromagnetic component. Vibrations can be induced by applying an oscillating voltage across two electrodes, in the form of an interdigitated transducer (IDT), placed on the surface. If the spacing between fingers in the electrode matches the wavelength of SAWs at the frequency of the applied signal, the induced SAWs add up coherently. Conversely, propagating SAWs that arrive at the transducer induce charge on the fingers such that the vibrations are converted into a voltage signal. The crucial invention in Gustafsson et al. (2014) was to let the capacitance in the transmon qubit double as an IDT to mediate a direct coupling between qubit and SAWs. Because of the slow propagation speed of SAWs, $v \approx 3000$ m/s, the IDT finger spacing was on the order of $d \approx 1 \mu\text{m}$ to match the resonance frequency around $\omega \approx 5$ GHz. As can be seen in the figure, many fingers were used in the qubit IDT, which corresponded to tens of wavelengths, making this a truly giant atom.

This first experiment with a giant atom could only probe the atom around a single frequency, since the IDT used to convert signals had a narrow bandwidth. The frequency-dependence of the qubit coupling (see Sect. 2.1.2) could therefore not be tested. However, the experimental platform with SAWs and qubits, called cir-

cuit quantum acoustodynamics (QAD) (Gustafsson et al. 2014; Aref et al. 2016; Manenti et al. 2017), has been adopted in several research groups. In their experiments (Manenti et al. 2017; Noguchi et al. 2017; Moores et al. 2018; Satzinger et al. 2018; Bolgar et al. 2018; Sletten et al. 2019; Bienfait et al. 2019), the qubit is coupled to a resonator for the SAW modes. Since the resonator is long, it has a narrow free spectral range, and the frequency-dependent coupling of the qubit is evident from how it couples with different strength to different modes. This selective coupling to modes has been used in a clever way to read out the number of phonons in a mode via the qubit (Sletten et al. 2019).

A particular advantage of the SAWs is that their slow propagation speed makes it possible to engineer a giant atom with a very long distance between coupling points. In the experiment of Andersson et al. (2019), distances exceeding 400 wavelengths were realized, corresponding to $\Gamma\tau \approx 14$, i.e., well in the non-Markovian regime discussed in Sect. 2.2.

Another recent experiment (Andersson et al. 2020) with a superconducting transmon qubit and SAWs used the possibility to engineer the relaxation rates of the first two transitions of the transmon (see Sec. 2.1.2) to enable EIT. This appears to be the first time that EIT of a propagating mechanical mode has been demonstrated.

3.2 Superconducting Qubits and Microwave Transmission Lines

Superconducting qubits are usually coupled to microwave transmission lines, or *LC* resonators, instead of SAWs. Also the setup with a transmission line can be used to implement giant atoms, as proposed in Kockum et al. (2014). One simply couples the transmission line to the qubit at one point, meanders the transmission line back and forth on the chip until a wavelength distance has been reached, and then connects the transmission line to the qubit once more. Due to size limitations, this approach will not allow for distances between coupling points on the order of hundreds of wavelengths or more, as is possible with SAWs. However, with the transmission line it is possible to engineer the coupling at each point and the distance between coupling points with great precision, which can be crucial for demonstrating the interference effects that lie at the heart of giant atoms.

Two recent experiments have followed this approach to implement one (Vadiraj et al. 2020) and two (Kannan et al. 2020) giant atoms. In the experiment with one giant atom, the frequency-dependent coupling shown in Fig. 2 was measured and the ability to manipulate the relaxation rates in a multilevel atom as in Fig. 3 was shown. In the experiment with two giant atoms, the decoherence-free interaction discussed in Sect. 2.3 was demonstrated.

This opens up interesting possibilities for preparing entangled many-body states in waveguide QED with many atoms, which otherwise is difficult due to the dissipation into the waveguide which always is present for small atoms (Kannan et al. 2020).

3.3 *Cold Atoms in Optical Lattices*

All experiments with giant atoms so far have taken place in 1D geometries at microwave frequencies and used superconducting qubits. A recent theory proposal (González-Tudela et al. 2019) shows how giant atoms instead could be implemented in higher dimensions on another platform for quantum-optics simulation: cold atoms in optical lattices. Here, one would use atoms with two internal states, each of which couples to a different optical lattice, realized by counter-propagating lasers. In one state, the atom mimics a photon moving in a lattice; in the other state, the atom mimics an atom trapped in a specific site. By rapidly modulating the relative positions of the two lattices, it is possible to engineer an effective interaction where the atomic state couples to the photonic state at multiple points (González-Tudela et al. 2019). It may be possible to achieve a similar effect with superconducting qubits coupled to several sites in a 2D lattice of superconducting resonators. While such lattices have been analysed and realized previously (Koch et al. 2010; Houck et al. 2012; Underwood et al. 2016), to the best of our knowledge it has not been suggested previously to couple one qubit to several lattice sites in such a setup.

The proposed setup with cold atoms displays rich physics with the giant atoms coupled to 2D photonic environments that have a band structure. It is possible to construct interference such that a single giant atom relaxes by only emitting its energy in certain directions. It is also possible to decouple giant atoms completely from the environment, but still have them interact by exchange interactions, like in Sect. 2.3. While this interference was possible with just two coupling points per atom in 1D, the 2D case requires at least four coupling points.

4 Conclusion and Outlook

Giant atoms are emerging as a new, interesting field of quantum optics. Following the first experimental realization and theoretical study in 2014, the field has grown quickly in the past 5 years. Theoretical investigations have been extended from one to multiple giant atoms, from 1D to higher-dimensional environments coupling to the atoms, and from the Markovian to the non-Markovian regime, where time delays between coupling points matter. These investigations have revealed remarkable properties of giant atoms, including frequency-dependent couplings and decoherence-free interactions, which are hard or impossible to realize with small atoms.

In parallel, the experimental platform for giant atoms, with SAWs coupled to superconducting qubits, has been further developed. There are now also experiments with superconducting qubits coupled to microwave transmission lines, and an experimental platform with cold atoms in optical lattices has been proposed. The experiments have confirmed many of the theoretical predictions, and also contributed with new ideas for applications of giant atoms.

Looking towards the future, we can formulate a long research agenda for giant atoms. At the heart of this agenda is the fact that giant atoms mainly differ from small atoms by the interference effects introduced by the multiple coupling points, which already has been shown to lead to new effects. It therefore seems prudent to revisit many well-known quantum-optics phenomena to see if giant atoms can enhance them or enable new physics. Below, we give a list of such projects:

- **Superradiance:** For multiple small atoms coupled to light, it is well known that quantum interference effects can give rise to enhanced light emission, superradiance, where N atoms emit light at an increased rate, proportional to N^2 (Dicke 1954; Shammah et al. 2018). The reverse process, “superabsorption”, is also possible (Higgins et al. 2014; Yang et al. 2019), and may be of importance in photosynthesis and future solar cells. It is thus highly relevant to see if giant atoms can enhance superradiance and superabsorption.
- **Ultrastrong coupling:** When the strength of the coupling between light and matter starts to approach the bare resonance frequencies in the system, it is called ultrastrong (Kockum et al. 2019; Forn-Díaz et al. 2019). In this regime, the rotating-wave approximation breaks down and the number of excitations in the system is no longer conserved in the absence of drives. For a giant atom ultrastrongly coupled to an open waveguide, it would be interesting to map out the ground state of the system, since results for a small atom indicates that it should contain virtual photons clustered around each connection point (Sanchez-Burillo et al. 2014). However, ultrastrong coupling with giant atoms comes with several theoretical challenges, which make analytical results hard to achieve. For example, a giant atom with ultrastrong coupling will inevitably be in a regime where the travel time between coupling points is non-negligible (Ask et al. 2019a).
- **Generating non-classical light:** It has recently been shown that coherently driving a small atom in front of a mirror can lead to the generation of non-classical states of light with a negative Wigner function (Quijandría et al. 2018). Could a giant atom do the same?
- **Matryoshka atoms:** The topology in Fig. 4e, nested atoms, is reminiscent of a Russian matryoshka doll. Although it does not enable decoherence-free interaction like braided atoms do, it seems to have other interesting properties. If the distance between the coupling points of the outer atom is large, the outer atom could effectively act as a cavity (Ask et al. 2019b), similar to what two small atoms placed far away on either side of a central atom can do (Guimond et al. 2016). Also, preliminary results indicate that two nested giant atoms can emulate electromagnetically induced transparency in a Λ system without any external drive (Ask et al. 2019c). With many nested giant atoms, the situation is similar to having many atoms in front of a mirror. Thus, for certain inter-coupling-point distances, these giant atoms should be able to combine into fewer effective larger atoms, as can happen in the mirror case (Lin et al. 2019).
- **Chiral quantum optics:** In some waveguide-QED setups with small atoms, it is possible to realize chiral couplings, i.e., that the atoms only couple to one propagation direction in the waveguide (Lodahl et al. 2017). Although it is not

yet clear if this can be implemented in experiments with giant atoms, it seems interesting to study chiral quantum optics with giant atoms theoretically. A related question is whether interference between light propagating in a waveguide, and light taking the “shortcut” between two coupling points through a giant atom, can be used to realize an effective chiral coupling.

This was recently answered affirmatively for a setup with two atoms that are both directly coupled to each other and each coupled at its own single point to a waveguide ($\sim \lambda/4$ apart) (Guimond et al 2020).

Acknowledgements AFK acknowledges support from the Swedish Research Council (grant number 2019-03696), and from the Knut and Alice Wallenberg Foundation through the Wallenberg Centre for Quantum Technology (WACQT).

References

A.V. Akimov, A. Mukherjee, C.L. Yu, D.E. Chang, A.S. Zibrov, P.R. Hemmer, H. Park, M.D. Lukin, Generation of single optical plasmons in metallic nanowires coupled to quantum dots. *Nature* **450**, 402 (2007). <http://www.nature.com/doifinder/10.1038/nature06230>

G. Andersson, M. K. Ekström, P. Delsing, Electromagnetically induced transparency in a propagating mechanical mode. *Physical Review Letters* **124**, 240402 (2020)

G. Andersson, B. Suri, L. Guo, T. Aref, P. Delsing, Nonexponential decay of a giant artificial atom. *Nat. Phys.* **15**, 1123 (2019) <http://arxiv.org/abs/1812.01302>

M. Arcari, I. Söllner, A. Javadi, S. Lindskov Hansen, S. Mahmoodian, J. Liu, H. Thyrrstrup, E.H. Lee, J.D. Song, S. Stobbe, P. Lodahl, Near-unity coupling efficiency of a quantum emitter to a photonic crystal waveguide. *Phys. Rev. Lett.* **113**, 093603 (2014). <http://link.aps.org/doi/10.1103/PhysRevLett.113.093603>

T. Aref, P. Delsing, M.K. Ekström, A.F. Kockum, M.V. Gustafsson, G. Johansson, P.J. Leek, E. Magnusson, R. Manenti, Quantum acoustics with surface acoustic waves, in *Superconducting Devices in Quantum Optics*, eds. by R.H. Hadfield, G. Johansson (Springer, Berlin, 2016). http://dx.doi.org/10.1007/978-3-319-24091-6_9

A. Ask, et al., In preparation (2019b)

A. Ask, et al., In preparation (2019c)

A. Ask, M. Ekström, P. Delsing, G. Johansson, Cavity-free vacuum-Rabi splitting in circuit quantum acoustodynamics. *Phys. Rev. A* **99**, 013840 (2019a). <http://dx.doi.org/10.1103/PhysRevA.99.013840>

O.V. Astafiev, A.A. Abdumalikov, A.M. Zagoskin, Y.A. Pashkin, Y. Nakamura, J.S. Tsai, Ultimate on-chip quantum amplifier. *Phys. Rev. Lett.* **104**, 183603 (2010b). <http://link.aps.org/doi/10.1103/PhysRevLett.104.183603>

O. Astafiev, A.M. Zagoskin, A.A. Abdumalikov, Y.A. Pashkin, T. Yamamoto, K. Inomata, Y. Nakamura, J.S. Tsai, Resonance fluorescence of a single artificial atom. *Science* **327**, 840 (2010a). <http://www.sciencemag.org/cgi/doi/10.1126/science.1181918>

M. Bajcsy, S. Hofferberth, V. Balic, T. Peyronel, M. Hafezi, A.S. Zibrov, V. Vuletic, M.D. Lukin, Efficient all-optical switching using slow light within a hollow fiber. *Phys. Rev. Lett.* **102**, 203902 (2009). <http://link.aps.org/doi/10.1103/PhysRevLett.102.203902>

A. Beige, J. Pachos, H. Walther, Spontaneous emission of an atom in front of a mirror. *Phys. Rev. A* **66**, 063,801 (2002). <http://link.aps.org/doi/10.1103/PhysRevA.66.063801>

H.A. Bethe, The electromagnetic shift of energy levels. *Phys. Rev.* **72**, 339 (1947). <http://link.aps.org/doi/10.1103/PhysRev.72.339>

A. Bienfait, K.J. Satzinger, Y.P. Zhong, H.S. Chang, M.H. Chou, C.R. Conner, É. Dumur, J. Grebel, G.A. Peairs, R.G. Povey, A.N. Cleland, Phonon-mediated quantum state transfer and remote qubit entanglement. *Science* **364**, 368 (2019). <http://www.sciencemag.org/lookup/doi/10.1126/science.aaw8415>

A.N. Bolgar, J.I. Zotova, D.D. Kirichenko, I.S. Besedin, A.V. Semenov, R.S. Shaikhaidarov, O.V. Astafiev, Quantum regime of a two-dimensional phonon cavity. *Phys. Rev. Lett.* **120**, 223603 (2018). <https://link.aps.org/doi/10.1103/PhysRevLett.120.223603>

H.J. Carmichael, *Statistical Methods in Quantum Optics 1* (Springer, Berlin, 1999). <http://link.springer.com/10.1007/978-3-662-03875-8>

A. Ciani, D.P. DiVincenzo, Three-qubit direct dispersive parity measurement with tunable coupling qubits. *Phys. Rev. B* **96**, 214511 (2017). <http://dx.doi.org/10.1103/PhysRevB.96.214511>

C. Cohen-Tannoudji, J. Dupont-Roc, G. Grynberg, *Atom-Photon Interactions* (Wiley, Hoboken, 1998)

S. Datta, *Surface Acoustic Wave Devices* (Prentice Hall, Upper Saddle River, 1986)

R.H. Dicke, Coherence in spontaneous radiation processes. *Phys. Rev.* **93**, 99 (1954). <http://link.aps.org/doi/10.1103/PhysRev.93.99>

H. Dong, Z.R. Gong, H. Ian, L. Zhou, C.P. Sun, Intrinsic cavity QED and emergent quasinormal modes for a single photon. *Phys. Rev. A* **79**, 063847 (2009). <http://link.aps.org/doi/10.1103/PhysRevA.79.063847>

U. Dorner, P. Zoller, Laser-driven atoms in half-cavities. *Phys. Rev. A* **66**, 023816 (2002). <http://link.aps.org/doi/10.1103/PhysRevA.66.023816>

F. Dubin, D. Rotter, M. Mukherjee, C. Russo, J. Eschner, R. Blatt, Photon correlation versus interference of single-atom fluorescence in a half-cavity. *Phys. Rev. Lett.* **98**, 183003 (2007). <http://link.aps.org/doi/10.1103/PhysRevLett.98.183003>

J. Eschner, C. Raab, F. Schmidt-Kaler, R. Blatt, Light interference from single atoms and their mirror images. *Nature* **413**, 495 (2001). <http://www.nature.com/doifinder/10.1038/35097017>

Y.L.L. Fang, H.U. Baranger, Waveguide QED: Power spectra and correlations of two photons scattered off multiple distant qubits and a mirror. *Phys. Rev. A* **91**, 053845 (2015). <http://link.aps.org/doi/10.1103/PhysRevA.91.053845>

P. Forn-Díaz, J.J. García-Ripoll, B. Peropadre, J.L. Orgiazzi, M.A. Yurtalan, R. Belyansky, C.M. Wilson, A. Lupascu, Ultrastrong coupling of a single artificial atom to an electromagnetic continuum in the nonperturbative regime. *Nat. Phys.* **13**, 39 (2017). <http://www.nature.com/doifinder/10.1038/nphys3905>

P. Forn-Díaz, L. Lamata, E. Rico, J. Kono, E. Solano, Ultrastrong coupling regimes of light-matter interaction. *Rev. Mod. Phys.* **91**, 025005 (2019). <http://dx.doi.org/10.1103/RevModPhys.91.025005>

R. Friedberg, S.R. Hartmann, J.T. Manassah, Frequency shifts in emission and absorption by resonant systems of two-level atoms. *Phys. Rep.* **7**, 101 (1973). <http://linkinghub.elsevier.com/retrieve/pii/037015737390001X>

C.W. Gardiner, P. Zoller, *Quantum Noise*, 3rd edn. (Springer, Berlin, 2004)

A. González-Tudela, C. Sánchez Muñoz, J.I. Cirac, Engineering and harnessing giant atoms in high-dimensional baths: a proposal for implementation with cold atoms. *Phys. Rev. Lett.* **122**, 203603 (2019). <https://link.aps.org/doi/10.1103/PhysRevLett.122.203603>

X. Gu, A.F. Kockum, A. Miranowicz, Y.X. Liu, F. Nori, Microwave photonics with superconducting quantum circuits. *Phys. Rep.* **718–719**, 1–102 (2017). <https://doi.org/10.1016/j.physrep.2017.10.002>

P.O. Guimond, A. Roulet, H.N. Le, Scarani, V, Rabi oscillation in a quantum cavity: Markovian and non-Markovian dynamics. *Phys. Rev. A* **93**, 023808 (2016). <http://dx.doi.org/10.1103/PhysRevA.93.023808>

P. O. Guimond, B. Vermersch, M. L. Juan, A. Sharafiev, G. Kirchmair, P. Zoller, A unidirectional on-chip photonic interface for superconducting circuits. *npj Quantum Inf.* **6**, 32 (2020)

L. Guo, A.L. Grimsmo, A.F. Kockum, M. Pletyukhov, G. Johansson, Giant acoustic atom: a single quantum system with a deterministic time delay. *Phys. Rev. A* **95**, 053821 (2017). <http://link.aps.org/doi/10.1103/PhysRevA.95.053821>

L. Guo, A. F. Kockum, F. Marquardt, G. Johansson, Oscillating bound states for a giant atom. *Phys. Rev. Res.* [arXiv:1911.13028](https://arxiv.org/abs/1911.13028) (2019)

S. Guo, Y. Wang, T. Purdy, J. Taylor, Beyond Spontaneous Emission: Giant Atom Bounded in Continuum. *Phys. Rev. A* **102**, 033706 (2020). <https://doi.org/10.1103/PhysRevA.102.033706>

M.V. Gustafsson, T. Aref, A.F. Kockum, M.K. Ekström, G. Johansson, P. Delsing, Propagating phonons coupled to an artificial atom. *Science* **346**, 207 (2014). <http://www.sciencemag.org/cgi/doi/10.1126/science.1257219>

R. Hanson, L.P. Kouwenhoven, J.R. Petta, S. Tarucha, L.M.K. Vandersypen, Spins in few-electron quantum dots. *Rev. Mod. Phys.* **79**, 1217 (2007). <http://link.aps.org/doi/10.1103/RevModPhys.79.1217>

S. Haroche, Nobel Lecture: Controlling photons in a box and exploring the quantum to classical boundary. *Rev. Mod. Phys.* **85**, 1083 (2013). <https://doi.org/10.1103/RevModPhys.85.1083>

K.D.B. Higgins, S.C. Benjamin, T.M. Stace, G.J. Milburn, B.W. Lovett, E.M. Gauger, Superabsorption of light via quantum engineering. *Nat. Commun.* **5**, 4705 (2014). <http://www.nature.com/doifinder/10.1038/ncomms5705>

I.C. Hoi, A.F. Kockum, T. Palomaki, T.M. Stace, B. Fan, L. Tornberg, S.R. Sathyamoorthy, G. Johansson, P. Delsing, C.M. Wilson, Giant cross-Kerr effect for propagating microwaves induced by an artificial atom. *Phys. Rev. Lett.* **111**, 053601 (2013). <http://link.aps.org/doi/10.1103/PhysRevLett.111.053601>

I.C. Hoi, A.F. Kockum, L. Tornberg, A. Pourkabirian, G. Johansson, P. Delsing, C.M. Wilson, Probing the quantum vacuum with an artificial atom in front of a mirror. *Nat. Phys.* **11**, 1045 (2015). <http://www.nature.com/doifinder/10.1038/nphys3484>

I.C. Hoi, T. Palomaki, J. Lindkvist, G. Johansson, P. Delsing, C.M. Wilson, Generation of nonclassical microwave states using an artificial atom in 1D open space. *Phys. Rev. Lett.* **108**, 263601 (2012). <http://link.aps.org/doi/10.1103/PhysRevLett.108.263601>

I.C. Hoi, C.M. Wilson, G. Johansson, T. Palomaki, B. Peropadre, P. Delsing, Demonstration of a single-photon router in the microwave regime. *Phys. Rev. Lett.* **107**, 073601 (2011). <http://link.aps.org/doi/10.1103/PhysRevLett.107.073601>

A.A. Houck, H.E. Türeci, J. Koch, On-chip quantum simulation with superconducting circuits. *Nat. Phys.* **8**, 292 (2012). <http://www.nature.com/doifinder/10.1038/nphys2251>

A. Huck, U.L. Andersen, Coupling single emitters to quantum plasmonic circuits. *Nanophotonics* **5**, 1 (2016). <http://www.degruyter.com/view/j/nanoph.2016.5.issue-3/nanoph-2015-0153/nanoph-2015-0153.xml>

B. Kannan, M. Ruckriegel, D. Campbell, A. F. Kockum, J. Braumüller, D. Kim, M. Kjaergaard, P. Krantz, A. Melville, B. M. Niedzielski, A. Vepsäläinen, R. Winik, J. Yoder, F. Nori, T. P. Orlando, S. Gustavsson, W. D. Oliver, Waveguide quantum electrodynamics with giant superconducting artificial atoms. *Nat.* **583**, 775 (2020). <http://www.nature.com/articles/s41586-020-2529-9>

T.M. Karg, B. Gouraud, P. Treutlein, K. Hammerer, Remote Hamiltonian interactions mediated by light. *Phys. Rev. A* **99**, 063829 (2019). <https://link.aps.org/doi/10.1103/PhysRevA.99.063829>

J. Koch, A.A. Houck, K. Le Hur, S.M. Girvin, Time-reversal-symmetry breaking in circuit-QED-based photon lattices. *Phys. Rev. A* **82**, 043811 (2010). <http://link.aps.org/doi/10.1103/PhysRevA.82.043811>

J. Koch, T.M. Yu, J. Gambetta, A.A. Houck, D.I. Schuster, J. Majer, A. Blais, M.H. Devoret, S.M. Girvin, R.J. Schoelkopf, Charge-insensitive qubit design derived from the Cooper pair box. *Phys. Rev. A* **76**, 042319 (2007). <http://link.aps.org/doi/10.1103/PhysRevA.76.042319>

A.F. Kockum, P. Delsing, G. Johansson, Designing frequency-dependent relaxation rates and Lamb shifts for a giant artificial atom. *Phys. Rev. A* **90**, 013837 (2014). <http://link.aps.org/doi/10.1103/PhysRevA.90.013837>

A.F. Kockum, G. Johansson, F. Nori, Decoherence-Free Interaction between Giant Atoms in Waveguide Quantum Electrodynamics. *Phys. Rev. Lett.* **120**, 140404 (2018). <https://link.aps.org/doi/10.1103/PhysRevLett.120.140404>

A.F. Kockum, A. Miranowicz, S. De Liberato, S. Savasta, F. Nori, Ultrastrong coupling between light and matter. *Nat. Rev. Phys.* **1**, 19 (2019). <http://dx.doi.org/10.1038/s42254-018-0006-2>

A.F. Kockum, F. Nori, Quantum bits with Josephson junctions, *Fundamentals and Frontiers of the Josephson Effect*, ed. by F. Tafuri (Springer, Berlin, 2019). <https://doi.org/10.1007/978-3-030-20726-7>

K. Koshino, Y. Nakamura, Control of the radiative level shift and linewidth of a superconducting artificial atom through a variable boundary condition. *New J. Phys.* **14**, 043005 (2012). <http://stacks.iop.org/1367-2630/14/i=4/a=043005?key=crossref.8e8ee9c5cb468c374b4c2a56c13ab881>

D.O. Krimer, M. Liertzer, S. Rotter, H.E. Türeci, Route from spontaneous decay to complex multimode dynamics in cavity QED. *Phys. Rev. A* **89**, 033820 (2014). <https://link.aps.org/doi/10.1103/PhysRevA.89.033820>

G. Kurizki, Two-atom resonant radiative coupling in photonic band structures. *Phys. Rev. A* **42**, 2915 (1990). <https://link.aps.org/doi/10.1103/PhysRevA.42.2915>

K. Lalumi  re, B.C. Sanders, A.F. van Loo, A. Fedorov, A. Wallraff, A. Blais, Input-output theory for waveguide QED with an ensemble of inhomogeneous atoms. *Phys. Rev. A* **88**, 043806 (2013). <http://link.aps.org/doi/10.1103/PhysRevA.88.043806>

W.E. Lamb, R.C. Rutherford, Fine structure of the hydrogen atom by a microwave method. *Phys. Rev.* **72**, 241 (1947). <http://link.aps.org/doi/10.1103/PhysRev.72.241>

P. Lambropoulos, G.M. Nikolopoulos, T.R. Nielsen, S. Bay, Fundamental quantum optics in structured reservoirs. *Rep. Prog. Phys.* **63**, 455 (2000). <http://stacks.iop.org/0034-4885/63/i=4/a=201?key=crossref.4e89934b6d1170192620f7f10999fb37>

R.H. Lehmberg, Radiation from an N-atom system. II. Spontaneous emission from a pair of atoms. *Phys. Rev. A* **2**, 889 (1970b). <http://link.aps.org/doi/10.1103/PhysRevA.2.889>

R.H. Lehmberg, Radiation from an N-atom system. I. General formalism. *Phys. Rev. A* **2**, 883 (1970a). <http://link.aps.org/doi/10.1103/PhysRevA.2.883>

D. Leibfried, R. Blatt, C. Monroe, D. Wineland, Quantum dynamics of single trapped ions. *Rev. Mod. Phys.* **75**, 281 (2003). <http://link.aps.org/doi/10.1103/RevModPhys.75.281>

K.T. Lin, T. Hsu, C.Y. Lee, I.C. Hoi, G.D. Lin, Scalable collective Lamb shift of a 1D superconducting qubit array in front of a mirror. *Sci. Rep.* **9**, 19175 (2019). <https://doi.org/10.1038/s41598-019-55545-5>

G. Lindblad, On the generators of quantum dynamical semigroups. *Commun. Math. Phys.* **48**, 119 (1976). <https://doi.org/10.1007/BF01608499>

Y. Liu, A.A. Houck, Quantum electrodynamics near a photonic bandgap. *Nat. Phys.* **13**, 48 (2017). <http://www.nature.com/doifinder/10.1038/nphys3834>

P. Lodahl, S. Mahmoodian, S. Stobbe, A. Rauschenbeutel, P. Schneeweiss, J. Volz, H. Pichler, P. Zoller, Chiral quantum optics. *Nature* **541**, 473 (2017). <http://www.nature.com/doifinder/10.1038/nature21037>

R. Manenti, A.F. Kockum, A. Patterson, T. Behrle, J. Rahamim, G. Tancredi, F. Nori, P.J. Leek, Circuit quantum acoustodynamics with surface acoustic waves. *Nat. Commun.* **8**, 975 (2017). <http://www.nature.com/articles/s41467-017-01063-9>

D. Meschede, W. Jhe, E.A. Hinds, Radiative properties of atoms near a conducting plane: An old problem in a new light. *Phys. Rev. A* **41**, 1587 (1990). <http://link.aps.org/doi/10.1103/PhysRevA.41.1587>

M. Mirhosseini, E. Kim, V.S. Ferreira, M. Kalaei, A. Sipahigil, A.J. Keller, O. Painter, Superconducting metamaterials for waveguide quantum electrodynamics. *Nat. Commun.* **9**, 3706 (2018). <http://dx.doi.org/10.1038/s41467-018-06142-z>

M. Mirhosseini, E. Kim, X. Zhang, A. Sipahigil, P.B. Dieterle, A.J. Keller, A. Asenjo-Garcia, D.E. Chang, O. Painter, Cavity quantum electrodynamics with atom-like mirrors. *Nature* **569**, 692 (2019). <http://www.nature.com/articles/s41586-019-1196-1>

B.A. Moores, L.R. Sletten, J.J. Viennot, K.W. Lehnert, Cavity quantum acoustic device in the multimode strong coupling regime. *Phys. Rev. Lett.* **120**, 227701 (2018). <https://link.aps.org/doi/10.1103/PhysRevLett.120.227701>

D. Morgan, *Surface Acoustic Wave Filters*, 2nd edn. (Academic Press, Cambridge, 2007)

A. Noguchi, R. Yamazaki, Y. Tabuchi, Y. Nakamura, Qubit-assisted transduction for a detection of surface acoustic waves near the quantum limit. *Phys. Rev. Lett.* **119**, 180505 (2017). <https://link.aps.org/doi/10.1103/PhysRevLett.119.180505>

H. Pichler, P. Zoller, Photonic circuits with time delays and quantum feedback. *Phys. Rev. Lett.* **116**, 093601 (2016). <http://link.aps.org/doi/10.1103/PhysRevLett.116.093601>

F. Quijandría, I. Strandberg, G. Johansson, Steady-state generation of wigner-negative states in one-dimensional resonance fluorescence. *Phys. Rev. Lett.* **121**, 263603 (2018). <https://link.aps.org/doi/10.1103/PhysRevLett.121.263603>

D. Roy, C.M. Wilson, O. Firstenberg, Colloquium: strongly interacting photons in one-dimensional continuum. *Rev. Mod. Phys.* **89**, 021001 (2017). <http://link.aps.org/doi/10.1103/RevModPhys.89.021001>

E. Sanchez-Burillo, D. Zueco, J.J. Garcia-Ripoll, L. Martin-Moreno, Scattering in the ultrastrong regime: nonlinear optics with one photon. *Phys. Rev. Lett.* **113**, 263604 (2014). <http://link.aps.org/doi/10.1103/PhysRevLett.113.263604>

K.J. Satzinger, Y.P. Zhong, H.S. Chang, G.A. Pears, A. Bienfait, M.H. Chou, A.Y. Cleland, C.R. Conner, É. Dumur, J. Grebel, I. Gutierrez, B.H. November, R.G. Povey, S.J. Whiteley, D.D. Awschalom, D.I. Schuster, A.N. Cleland, Quantum control of surface acoustic-wave phonons. *Nature* **563**, 661 (2018). <http://www.nature.com/articles/s41586-018-0719-5>

M.O. Scully, A.A. Svidzinsky, The Lamb shift - yesterday, today, and tomorrow. *Science* **328**, 1239 (2010). <http://www.sciencemag.org/cgi/doi/10.1126/science.1190737>

N. Shammah, S. Ahmed, N. Lambert, S. De Liberato, F. Nori, Open quantum systems with local and collective incoherent processes: efficient numerical simulations using permutational invariance. *Phys. Rev. A* **98**, 063815 (2018). <http://arxiv.org/abs/1805.05129> <https://link.aps.org/doi/10.1103/PhysRevA.98.063815>

T. Shi, D.E. Chang, J.I. Cirac, Multiphoton-scattering theory and generalized master equations. *Phys. Rev. A* **92**, 053834 (2015). <http://link.aps.org/doi/10.1103/PhysRevA.92.053834>

L.R. Sletten, B.A. Moores, J.J. Viennot, K.W. Lehnert, Resolving phonon Fock states in a multimode cavity with a double-slit qubit. *Phys. Rev. X* **9**, 021056 (2019). <http://dx.doi.org/10.1103/PhysRevX.9.021056>

N.M. Sundaresan, R. Lundgren, G. Zhu, A.V. Gorshkov, A.A. Houck, Interacting qubit-photon bound states with superconducting circuits. *Phys. Rev. X* **9**, 011021 (2019). <http://dx.doi.org/10.1103/PhysRevX.9.011021>

T. Tufarelli, F. Ciccarello, M.S. Kim, Dynamics of spontaneous emission in a single-end photonic waveguide. *Phys. Rev. A* **87**, 013820 (2013). <http://link.aps.org/doi/10.1103/PhysRevA.87.013820>

D.L. Underwood, W.E. Shanks, A.C.Y. Li, L. Ateshian, J. Koch, A.A. Houck, Imaging photon lattice states by scanning defect microscopy. *Phys. Rev. X* **6**, 021044 (2016). <https://link.aps.org/doi/10.1103/PhysRevX.6.021044>

A. M. Vadiraj, A. Ask, T. G. McConkey, I. Nsanzineza, C. W. Sandbo Chang, A. F. Kockum, C. M. Wilson, Engineering the level structure of a giant artificial atom in waveguide quantum electrodynamics. [arXiv:2003.14167](https://arxiv.org/abs/2003.14167) (2020)

A.F. van Loo, A. Fedorov, K. Lalumiere, B.C. Sanders, A. Blais, A. Wallraff, Photon-mediated interactions between distant artificial atoms. *Science* **342**, 1494 (2013). <http://www.sciencemag.org/cgi/doi/10.1126/science.1244324>

D.F. Walls, G.J. Milburn, *Quantum Optics*, 2nd edn. (Springer, Berlin, 2008). <http://link.springer.com/10.1007/978-3-540-28574-8>

Y. Wang, J. Minár, G. Hétet, V. Scarani, Quantum memory with a single two-level atom in a half cavity. *Phys. Rev. A* **85**, 013823 (2012). <http://link.aps.org/doi/10.1103/PhysRevA.85.013823>

P.Y. Wen, A.F. Kockum, H. Ian, J.C. Chen, F. Nori, I.C. Hoi, Reflective amplification without population inversion from a strongly driven superconducting qubit. *Phys. Rev. Lett.* **120**, 063603 (2018). <https://link.aps.org/doi/10.1103/PhysRevLett.120.063603>

P.Y. Wen, K.T. Lin, A.F. Kockum, B. Suri, H. Ian, J.C. Chen, S.Y. Mao, C.C. Chiu, P. Delsing, F. Nori, G.D. Lin, I.C. Hoi, Large collective Lamb shift of two distant superconducting artificial atoms. *Phys. Rev. Lett.* **123**, 233602 (2019). <https://doi.org/10.1103/PhysRevLett.123.233602>

M.A. Wilson, P. Bushev, J. Eschner, F. Schmidt-Kaler, C. Becher, R. Blatt, U. Dorner, Vacuum-field level shifts in a single trapped ion mediated by a single distant mirror. *Phys. Rev. Lett.* **91**, 213602 (2003). <http://link.aps.org/doi/10.1103/PhysRevLett.91.213602>

Z.L. Xiang, S. Ashhab, J.Q. You, F. Nori, Hybrid quantum circuits: superconducting circuits interacting with other quantum systems. *Rev. Mod. Phys.* **85**, 623 (2013). <http://link.aps.org/doi/10.1103/RevModPhys.85.623>

D. Yang, S.H. Oh, J. Han, G. Son, J. Kim, J. Kim, K. An, Observation of superabsorption by correlated atoms (2019). <http://arxiv.org/abs/1906.06477>

J.Q. You, F. Nori, Atomic physics and quantum optics using superconducting circuits. *Nature* **474**, 589 (2011). <http://dx.doi.org/10.1038/nature10122>

H. Zheng, H.U. Baranger, Persistent quantum beats and long-distance entanglement from waveguide-mediated interactions. *Phys. Rev. Lett.* **110**, 113601 (2013). <http://link.aps.org/doi/10.1103/PhysRevLett.110.113601>

Open Access This chapter is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

