

Double neutral Higgs production in the two-Higgs doublet model at the LHC

Abdesslam Arhrib,^{a,b,c} Rachid Benbrik,^{c,d,e} Chuan-Hung Chen,^{d,e} Renato Guedes^f
and Rui Santos^f

^aNational Center for Theoretical Physics, National Taiwan University,
Taipei, Taiwan 10617, R.O.C.

^bDépartement de Mathématique, Faculté des Sciences and Techniques,
Université Abdelmalek Essaâdi,
B. 416, Tangier, Morocco

^cLPHEA, Faculté des Sciences-Semlalia,
B.P. 2390 Marrakesh, Morocco

^dDepartment of Physics, National Cheng Kung University,
Taiwan 701, Taiwan, R.O.C.

^eNational Center for Theoretical Physics,
Taiwan 701, R.O.C.

^fNExT Institute and School of Physics and Astronomy, University of Southampton Highfield,
Southampton SO17 1BJ, U.K.

E-mail: aarhrib@ictp.it, rbenbrik@ictp.it, physchen@mail.ncku.edu.tw,
R.B.Guedes@soton.ac.uk, rsantos@cii.fc.ul.pt

ABSTRACT: In this work we study the production and decay of two CP-even Higgs bosons in the general CP-conserving two-Higgs doublet model at the LHC. We also study the limiting case of the decoupling scenario. For each Yukawa version of the model, we look for the region of the parameter space where a signal would be seen at the LHC. Taking into account theoretical and experimental constraints on the two Higgs doublet model parameters, we show that the cross section can be two orders of magnitude above the corresponding Standard Model cross section. We have also studied in detail the decay profile of the Higgs bosons and showed that interesting signatures may emerge for some particular values of the parameters. There are scenarios where some of the triple Higgs couplings could be measured.

KEYWORDS: Higgs Physics, Beyond Standard Model

Contents

1	Introduction	1
2	The CP-conserving 2HDM	3
3	Theoretical and experimental bounds	5
4	Results	6
4.1	The calculation	6
4.2	The decoupling limit	8
4.3	The general case for $pp \rightarrow h^0 h^0$	10
4.4	The $H^0 H^0$ and $H^0 h^0$ production modes	14
5	Higgs signatures	15
6	Discussion and conclusions	18
A	Fermion-scalar vertices	19

1 Introduction

CERN's Large Hadron Collider (LHC) is due to start operation in 2009 with a center of mass energy of 10 TeV . Therefore, testing the spontaneous symmetry breaking (SSB) mechanism and the search for the scalar particle or particles associated to that process is most probably postponed for at least another year. On the other hand, the Tevatron [1] has recently excluded a 160-170 GeV Standard Model (SM) Higgs which decays to a pair of W gauge bosons. The mass region probed will obviously grow as data is being collected and a Higgs boson can still be discovered at the Tevatron.

If a Higgs boson is found, the next task will be to understand the underlying mechanism that gives mass to all known particles. Characterizing the Higgs potential is the first step towards inferring the nature of the SSB mechanism. There are several alternatives to the SM that can be tested at the LHC. The most popular are the Minimal Supersymmetric Model (MSSM) or some of its variants like two Higgs Doublet Models (2HDM), little Higgs models and extra dimensions models among others. It is well known that the CP-conserving 2HDM has in its spectrum 2 CP-even scalars, h^0 and H^0 , one CP-odd scalar A^0 and a pair of charged Higgs bosons, H^\pm . Regarding the Higgs sector, the 2HDM is a clear candidate to be probed at the LHC because it is the minimal version of the SM that gives rise to charged Higgs bosons while being consistent with all available experimental data.

The difference between all these models can in principle be investigated by examining the processes involving scalar triple and quartic vertices.

In the SM, the potential can be studied in processes which have at least two Higgs in the final state. The process $gg \rightarrow h^0 h^0$ was studied in the SM in [2, 3], in the SM with a sequential fourth generation in [4] and in the MSSM in [5]. The SM cross section is of the order of 20 fb for a Higgs mass of 115 GeV and it drops steeply with increasing Higgs mass as a consequence of the fast falloff of the gluon-gluon luminosity. A sequential fourth generation can enhance the double Higgs production rate to ≈ 300 fb for a Higgs mass of 115 GeV. In the MSSM, the triple coupling can be probed in a wide range of Higgs masses whenever the resonant decay $H^0 \rightarrow h^0 h^0$ is possible. The measurement will be easier for small $\tan\beta$, where the trilinear coupling is larger and the heavy Higgs mass is less than twice the light Higgs mass in major parts of the parameter space. More detailed studies at parton level were then performed for the SM in [6–8] and for the MSSM in [9, 10]. QCD corrections to Higgs pair production in gluon collisions have been calculated for the SM and for the MSSM in the heavy top quark limit in [11]. Other processes such as vector boson fusion have been proposed for the 2HDM and shown to be rather powerful in reconstructing the triple Higgs self coupling [12] at the LHC. Similar processes were proposed for a linear electron-positron collider [13].

The process $pp \rightarrow h^0 h^0$ is also possible in the general 2HDM. In order to be able to distinguish between models, such as 2HDM, MSSM, sequential fourth generation and little Higgs models, it is mandatory to perform a detailed study for the double Higgs production within the general 2HDM. In a previous study, we have considered double Higgs production at hadron colliders in the 2HDM fermiophobic limit [14] and we have shown that information on the triple Higgs couplings could be extracted. In this work, we extend the above study to the general 2HDM. We will consider all CP-even double Higgs production, that is, $pp \rightarrow h^0 h^0, h^0 H^0, H^0 H^0$. We will present the regions of the parameter space of a general CP-conserving 2HDM where this process is large enough to be probed at the LHC taking into account the various experimental constraints on the 2HDM parameters as well as the tree-level unitarity and vacuum stability constraints. We will show that in some cases, even after imposing all constraints, cross sections are still large enough to allow for a determination of 2HDM triple Higgs couplings. In other cases only a deviation from the SM can be measured and other processes would be needed to complete the picture. We will give particular attention to the decoupling limit where the lightest Higgs mimics the SM Higgs and look for the possible non-decoupling effects.

This paper is organized as follows. In the next section we review the 2HDM potential. We then present all experimental and theoretical constraints that bound the 2HDM parameter space. In section 4 we present our numerical results. In section 5 we discuss the final states in the different 2HDM scenarios. Finally, our findings are summarized and discussed in section 6.

2 The CP-conserving 2HDM

The most general two-Higgs doublet model has 14 independent parameters. It violates CP explicitly and it can break electric charge. Even if explicit CP-violating interactions are not present in the potential, spontaneous CP breaking can still occur. One way to force the CP minimum not to exist is to impose the exact Z_2 discrete symmetry $\Phi_1 \rightarrow \Phi_1$, $\Phi_2 \rightarrow -\Phi_2$ [15, 16]. This symmetry can be softly broken by dimension two terms we define as $[m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.}]$. In this case, we can build either a CP-conserving theory or one that breaks CP spontaneously [17, 18]. In this work the vacuum structure is chosen so that the potential is CP-conserving. The theory build on a minimum that does not break CP-invariance nor electric charge was shown to be stable at tree level [19, 20]. Under these constraints, the most general renormalizable potential which is invariant under $SU(2) \otimes U(1)$ can be written as

$$V(\Phi_1, \Phi_2) = m_1^2 \Phi_1^\dagger \Phi_1 + m_2^2 \Phi_2^\dagger \Phi_2 + (m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.}) + \frac{1}{2} \lambda_1 (\Phi_1^\dagger \Phi_1)^2 + \frac{1}{2} \lambda_2 (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) + \frac{1}{2} \lambda_5 \left[(\Phi_1^\dagger \Phi_2)^2 + \text{h.c.} \right], \quad (2.1)$$

where Φ_i , $i = 1, 2$ are complex $SU(2)$ doublets with 4 degrees of freedom each and all m_i^2 , λ_i and m_{12}^2 are real which follows from the hermiticity of the potential. From the initial eight degrees of freedom, if the $SU(2)$ symmetry is broken, we end up with two CP-even Higgs states usually denoted by h^0 and H^0 , one CP-odd state, A^0 , two charged Higgs boson, H^\pm and three Goldstone bosons.

The potential in eq. (2.1) has a total of 10 parameters if one includes the vacuum expectation values. In a CP-conserving minimum there are two minimization conditions that can be used to fix the tree-level value of the parameters m_1^2 and m_2^2 . The combination $v^2 = v_1^2 + v_2^2$ is fixed as usual by the electroweak breaking scale through $v^2 = (2\sqrt{2}G_F)^{-1}$. We are thus left with 7 independent parameters, namely $(\lambda_i)_{i=1,\dots,5}$, m_{12}^2 , and $\tan \beta \equiv v_2/v_1$. Equivalently, we can take instead the set, m_{h^0} , m_{H^0} , m_{A^0} , m_{H^\pm} , $\tan \beta$, α and m_{12}^2 , as the 7 independent parameters. The angle β is the rotation angle from the group eigenstates to the mass eigenstates in the CP-odd and charged sector. The angle α is the corresponding rotation angle for the CP-even sector. The parameter m_{12} is a measure of how the discrete symmetry is broken. As stated previously, the potential with $m_{12} = 0$ has an exact Z_2 symmetry and is always CP-conserving.

From the above potential, eq. (2.1), we can derive the triple Higgs couplings needed for the present study as a function of the 2HDM parameters m_{h^0} , m_{H^0} , m_{A^0} , m_{H^\pm} , $\tan \beta$, α and m_{12}^2 . They are the same for all different Yukawa versions of the 2HDM and are given by:

$$\lambda_{h^0 h^0 h^0}^{2HDM} = \frac{-3e}{m_W s_W s_{2\beta}^2} \left[(c_\beta c_\alpha^3 - s_\beta s_\alpha^3) s_{2\beta} m_{h^0}^2 + 2c_{\beta-\alpha}^2 c_{\beta+\alpha} m_{12}^2 \right] \quad (2.2)$$

$$\lambda_{H^0 H^0 H^0}^{2HDM} = \frac{-3e}{m_W s_W s_{2\beta}^2} \left[(c_\beta c_\alpha^3 - s_\beta s_\alpha^3) s_{2\beta} m_{H^0}^2 + 2s_{\beta-\alpha}^2 s_{\beta+\alpha} m_{12}^2 \right] \quad (2.3)$$

$$\lambda_{H^0 h^0 h^0}^{2HDM} = -\frac{1}{2} \frac{ec_{\beta-\alpha}}{m_W s_W s_{2\beta}^2} \left[(2m_{h^0}^2 + m_{H^0}^2) s_{2\alpha} s_{2\beta} - 2(3s_{2\alpha} - s_{2\beta}) m_{12}^2 \right] \quad (2.4)$$

$$\lambda_{H^0 H^0 h^0}^{2HDM} = \frac{1}{2} \frac{e s_{\beta-\alpha}}{m_W s_W s_{2\beta}^2} \left[(m_{h^0}^2 + 2m_{H^0}^2) s_{2\alpha} s_{2\beta} + 2(3s_{2\alpha} + s_{2\beta}) m_{12}^2 \right] \quad (2.5)$$

$$\lambda_{h^0 h^0 h^0}^{SM} = \frac{-3em_{h^0}^2}{2m_W s_W} \quad (2.6)$$

where $\lambda_{h^0 h^0 h^0}^{SM}$ is the triple Higgs SM coupling, e is the electric charge, m_W is the W boson mass, s_W is the sin of the Weinberg angle and s and c were used as shorthand notation for sin and cos. As we can see from eqs. (2.2)–(2.5), all triple Higgs couplings have a quadratic dependence of the physical masses of the fields present in the vertex and on the soft breaking term m_{12}^2 . These couplings also depend strongly on $\tan\beta$ and α .

The Yukawa Lagrangian is a straightforward generalization of the SM one. Its most general form in the 2HDM is

$$\mathcal{L}_Y = G_{ij}^k (\bar{u} \quad \bar{d})_L^i \phi_k d_R^j + \tilde{G}_{ij}^k (\bar{u} \quad \bar{d})_L^i \tilde{\phi}_k u_R^j + h.c. \quad (2.7)$$

where the G_{ij}^k and \tilde{G}_{ij}^k are arbitrary constants, i and j are generation indices and k is the doublet number. However, tree-level flavor changing neutral currents (FCNCs) are severely constrained by experiment. The symmetry imposed to the potential can be naturally extended to the Yukawa Lagrangian to guarantee that FCNCs are not present. It suffices that fermions of a given electric charge couple to no more than one Higgs doublet [21]. This can be accomplished naturally by imposing on all fields appropriate discrete symmetries that forbid the unwanted FCNC couplings. There are essentially four ways of doing this [22] and so there are four variations of the model. We define as Type I the model where only the doublet ϕ_2 couples to all fermions; Type II is the model where ϕ_2 couples to up-type quarks and ϕ_1 couples to down-type quarks and leptons; in a Type III model ϕ_2 couples to all quarks and ϕ_1 couples to all leptons; a Type IV model is instead built such that ϕ_2 couples to up-type quarks and to leptons and ϕ_1 couples to down-type quarks. Models III and IV have been explored in a number of papers [23] and more recently with a renewed interest in [24]. We present all Yukawa coupling in appendix A. Regarding the Higgs couplings to the quarks, the models can be grouped in pairs like (I, IV) and (II, III) as the only difference between these two groups is in the couplings to the leptons. Moreover, the way we have chosen to build the Yukawa Lagrangian is such that up type quarks have the same couplings to the Higgs bosons in all four models. Down type quarks have different couplings in the two groups defined above, (I, IV) and (II, III), as can be seen in the appendix. In the large $\tan\beta$ limit, in 2HDM-(II, III), the Higgs coupling to a pair of down quarks is enhanced by a factor of $1/\cos\beta \approx \tan\beta$ while for 2HDM-(I, IV) there is no such enhancement. The processes described in this work of neutral Higgs pair production are either one-loop and proceed via quark loops or tree-level in the case of $b\bar{b}$ fusion. We will show that the theoretical constraints force $\tan\beta$ to be small, let us say smaller than 5 (we will come back to this point later). Because the Yukawa couplings are proportional to the quark masses and some function of the angles, for such small values of $\tan\beta$ the bottom loop can not compete with the top loop and the results for $gg \rightarrow h^0 h^0$ proved to be almost Yukawa-model independent. We will show that the only difference comes from higher order corrections that enter the calculation via the width of the Higgs bosons.

3 Theoretical and experimental bounds

The parameter space of the scalar potential of the 2HDM is reduced both by theoretical constraints as well as by the results of experimental searches. From the theoretical constraints which the 2HDM is subjected to, the most important are the ones that insure tree-level unitarity and vacuum stability of the theory. We also force the potential to be perturbative by requiring that all quartic couplings of the scalar potential, eq. (2.1), obey $|\lambda_i| \leq 8\pi$ for all i . For the vacuum stability conditions that ensure that the potential is bounded from below, we use those from [25], which are given by

$$\begin{aligned} \lambda_1 &> 0, & \lambda_2 &> 0, \\ \sqrt{\lambda_1 \lambda_2} + \lambda_3 + \min(0, \lambda_4 - |\lambda_5|) &> 0. \end{aligned} \tag{3.1}$$

The most restrictive theoretical bound comes from the full set of unitarity constraints [26, 27] established using the high energy approximation as well as the equivalence theorem and which can be written as

$$|a_{\pm}|, |b_{\pm}|, |c_{\pm}|, |d_{\pm}|, |e_{1,2}|, |f_{\pm}|, |g_{1,2}| < 8\pi \tag{3.2}$$

with

$$a_{\pm} = \frac{3}{2} \left\{ (\lambda_1 + \lambda_2) \pm \sqrt{(\lambda_1 - \lambda_2)^2 + \frac{4}{9}(2\lambda_3 + \lambda_4)^2} \right\}, \tag{3.3}$$

$$b_{\pm} = \frac{1}{2} \left\{ (\lambda_1 + \lambda_2) \pm \sqrt{(\lambda_1 - \lambda_2)^2 + 4\lambda_4^2} \right\}, \tag{3.4}$$

$$c_{\pm} = d_{\pm} = \frac{1}{2} \left\{ (\lambda_1 + \lambda_2) \pm \sqrt{(\lambda_1 - \lambda_2)^2 + 4\lambda_5^2} \right\}, \tag{3.5}$$

$$e_1 = (\lambda_3 + 2\lambda_4 - 3\lambda_5), \quad e_2 = (\lambda_3 - \lambda_5), \tag{3.6}$$

$$f_+ = (\lambda_3 + 2\lambda_4 + 3\lambda_5), \quad f_- = (\lambda_3 + \lambda_5), \tag{3.7}$$

$$g_1 = g_2 = (\lambda_3 + \lambda_4). \tag{3.8}$$

Since all λ_i can be expressed in terms of masses and mixing angles so can the above expressions. Some of the λ_i have a strong dependence on $\tan\beta$ and grow with increasing $\tan\beta$. Therefore, they are very restrictive in the allowed values of $\tan\beta$, especially when the parameter m_{12}^2 is also large.

The 2HDM parameters are also constrained by direct experimental searches and by precision experimental data. First, the extra contributions to the $\delta\rho$ parameter from the extra Higgs scalars [28] should not exceed the current limits from precision measurements [29]: $|\delta\rho| \lesssim 10^{-3}$. Such an extra contribution to $\delta\rho$ vanishes in the limit $m_{H\pm} = m_{A^0}$. To ensure that $\delta\rho$ will be within the allowed range we allow only a small splitting between $m_{H\pm}$ and m_{A^0} . Second, it has been shown in ref. [30] that data from $B \rightarrow X_s \gamma$ impose a lower limit of $m_{H\pm} \gtrsim 290$ GeV in Models type II and III. This constraint no longer applies in the case of 2HDM-I and 2HDM-IV. Third, values of $\tan\beta$ smaller than ≈ 1 are disallowed both by the constraints coming from $Z \rightarrow b\bar{b}$ and from $B_q \bar{B}_q$ mixing [30] for all Yukawa versions of




Figure 1. Tree-level contribution to $b\bar{b} \rightarrow S_i S_j$.

the model. On the other hand, because $\tan\beta$ can not be too large due to the theoretical constraints, except in some very specific scenarios, there is no need to discuss experimental constraints for very large $\tan\beta$. Finally, searches for Higgs bosons at colliders directly constrains their masses. The constraints on neutral Higgs bosons depend on their decay modes. The scenario where the light CP-even Higgs h^0 is fermiophobic, has been discussed in [14]. If the neutral Higgs bosons decay mainly into fermions, the OPAL, DELPHI and L3 collaborations have set a limit on the h^0 and A^0 masses of the 2HDM [31–33]. OPAL concluded that the regions $1 \lesssim m_{h^0} \lesssim 55$ GeV and $3 \lesssim m_{A^0} \lesssim 63$ GeV are excluded at 95% CL independently of the values of α and $\tan\beta$ [31]. It should be noted however that this is true as long as we are away from the limit $\alpha - \beta \approx 0$. In this limit, the cross section for the process $e^+e^- \rightarrow Z \rightarrow Zh^0$ vanishes in the 2HDM and the above bound no longer applies. L3 set a lower limit on the CP-even mass of the order 110.3 GeV [32] if the Higgs-strahlung cross section is the SM one and the Higgs decays hadronically. The DELPHI collaboration also studied the decay $h^0 \rightarrow A^0 A^0$ in $e^+e^- \rightarrow h^0 Z$ and $e^+e^- \rightarrow h^0 A^0$ production and large portions of the (m_{h^0}, m_{A^0}) plane were excluded depending on the 2HDM suppression factor that enters the $e^+e^- \rightarrow h^0 Z$ and $e^+e^- \rightarrow h^0 A^0$ cross sections [33] with respect to the SM. In what follows, we will assume that all Higgs masses are above 100 GeV unless the ratio of couplings $g_{ZZh^0}^{2HDM}/g_{ZZh}^{SM}$ is suppressed down to $\sin(\beta - \alpha) \approx 0.1$ when no bound to m_{h^0} applies.

4 Results

4.1 The calculation

The double CP-even Higgs production processes $pp \rightarrow S_i S_j$ ($S_{i,j} = h^0, H^0$) have tree level contributions from $q\bar{q} \rightarrow S_i S_j$ with s-channel Higgs exchange,¹ t-channel quark exchange and one loop contributions from $gg \rightarrow S_i S_j$. The most relevant contribution for process $q\bar{q} \rightarrow S_i S_j$ is the one where $q = b$ because the Higgs couples proportionally to the quark masses. The tree level diagrams for double Higgs production via b-quark fusion are shown in figure 1.

The process $gg \rightarrow S_i S_j$ occurs only at the one-loop level and the four types of diagrams that participate in the process are depicted in figure 2. The triangle diagrams, as shown

¹Bose Symmetry forbids the couplings $Zh^0 h^0$, $Zh^0 H^0$ and $ZH^0 H^0$.




Figure 2. Triangle and box contributions to $gg \rightarrow S_i S_j$, where $S = h^0, H^0$ and F is a generic quark.

in figure 2 (a), represent the generic contribution to the process with a fermion F which can be either an up or a down quark. The second type of diagrams are the box contributions which come in three different topologies, with virtual up and down quarks being exchanged. Comparing the tree-level with the loop diagrams it is easy to understand that no complementarity is expected from these two contributions - if a given set of parameters originate a small value of $\sigma(b\bar{b} \rightarrow S_i S_j)$ it will forcibly lead to a small $\sigma(gg \rightarrow S_i S_j)$ as well. We have explicitly checked that the tree-level contribution is always well below the loop contribution - usually less than 5%. All other quark fusion processes are negligible when compared to $b\bar{b} \rightarrow S_i S_j$.

The tree and one-loop amplitudes were generated and calculated with the packages FeynArts [34] and FormCalc [35]. The scalar integrals were evaluated with LoopTools [36]. We have included the total width of the scalar particle $S_i = h^0, H^0$ in the calculation of the corresponding amplitude. Even if the dominant contribution to the one-loop amplitude comes from top and bottom diagrams, in our numerical results we include all quarks in the loops. In models I and IV the largest contribution is always the one with top quarks in the loops. In the remaining two Yukawa models the situation is not as clear - for instance, in the triangle loop diagram with a virtual light Higgs, $S = h^0$, when $m_t \approx \tan \alpha \tan \beta m_b$ the contribution from the bottom quark triangle loop is larger than the one with the top quark loop. However, the same is not true for the triangle diagram with the virtual H^0 . We will show that the difference between models arises mainly from the combined dependence of the two CP-even masses, m_{12} , $\tan \beta$ and $\sin \alpha$.

We will start our discussion with the very interesting scenario of the decoupling limit, where the light Higgs resembles the SM one. We will then move to the study of the more general case and try to identify the regions of the 2HDM parameter space that can be probed in this more general scenario. As stated in the introduction, NLO QCD corrections to double Higgs production were calculated in [11] for the SM and for the MSSM. The total K factor, in the large top mass limit, was shown to vary between 1.8 and 2 for the SM and for a Higgs mass between 70 and 200 GeV. As $\tan \beta$ is constrained to be small, the top loop always dominate in our analysis. This means that these results are to a good approximation the enhancement one would also expect in the 2HDM.

4.2 The decoupling limit

The decoupling limit of the 2HDM is usually defined as the limiting case where all scalar masses except the lighter CP-even Higgs become infinite and the effective theory is just the SM with one doublet (see [37] for a discussion). In this case, the CP-even h^0 is the lightest scalar particle while the other Higgs particles, H^0 , A^0 and H^\pm are heavy and mass-degenerate. Using purely algebraic arguments at the tree level, one can derive that the main consequences of the decoupling limit is that because $\cos(\beta - \alpha) \rightarrow 0$, the CP-even h^0 of the 2HDM and the SM Higgs h_{SM} have similar tree level couplings to the gauge bosons and to the fermions [37, 38]. Obviously, the decoupling limit does not rigorously apply to the more realistic world where the particle masses are finite. Actually, one may consider a less rigorous scenario, labeled as the decoupling regime [37], where the heavy Higgs particles have masses much larger than the Z boson mass and may escape detection in the planned experiments.

Several studies have been carried out looking for non-decoupling effects in Higgs boson decays and Higgs self-interactions. Large loop effects in $h^0 \rightarrow \gamma\gamma$, $h^0 \rightarrow \gamma Z$ and $h^0 \rightarrow b\bar{b}$ have been pointed out for the 2HDM [40, 41] and may give indirect information on Higgs masses and the involved triple Higgs couplings such as $h^0 H^+ H^-$, $h^0 H^0 H^0$, $h^0 A^0 A^0$ and $h^0 h^0 h^0$. The non-decoupling contributions to the triple Higgs self coupling $h^0 h^0 h^0$ has been investigated in the 2HDM in ref. [38], revealing large non-decoupling effects in the 2HDM. In this section we will show that the large non-decoupling effects in the $h^0 h^0 h^0$ coupling will modify the double Higgs pair production $pp \rightarrow h^0 h^0$ cross section making it larger than its SM counterpart. We will focus on the scenario where all Higgs particles of the 2HDM, except for the lightest CP-even Higgs, are heavy and can escape detection at the early stages of next generation of colliders.

It is easy to check that in the decoupling limit the relation $\beta - \alpha \rightarrow \pi/2$ holds; hence, the triple coupling $h^0 h^0 h^0$ given in eq. (2.2) is reduced to its SM value $g_{h^0 h^0 h^0} = -3m_{h^0}^2/v$. Furthermore, we have $g_{H^0 h^0 h^0} \rightarrow 0$ at tree level and so the $gg \rightarrow h^0 h^0$ and $b\bar{b} \rightarrow h^0 h^0$ amplitudes reduce exactly to the SM results. As shown in [38], the one-loop leading contributions from all heavy Higgs boson loops and also from top quark loops to the effective $h^0 h^0 h^0$ coupling, can be written as

$$\lambda_{hhh}^{\text{eff}}(2HDM) = -\frac{3m_{h^0}^2}{\sqrt{2}v} \left\{ 1 + \frac{m_{H^0}^4}{12\pi^2 m_{h^0}^2 v^2} \left(1 + \frac{M^2}{m_{H^0}^2} \right)^3 + \frac{m_{A^0}^4}{12\pi^2 m_{h^0}^2 v^2} \left(1 + \frac{M^2}{m_{A^0}^2} \right)^3 + \frac{m_{H^\pm}^4}{6\pi^2 m_{h^0}^2 v^2} \left(1 + \frac{M^2}{m_{H^\pm}^2} \right)^3 - \frac{N_c M_t^4}{3\pi^2 m_{h^0}^2 v^2} \right\} \quad (4.1)$$

where $M = m_{12}/\sqrt{\sin\beta \cos\beta}$, M_Φ and p_i represent the mass of the H^0 , A^0 or H^\pm bosons and the momenta of the external Higgs lines, respectively. We note that in eq. (4.1) m_{h^0} is the renormalized physical mass of the lightest CP-even Higgs boson h^0 . In our calculation of the cross section of $pp \rightarrow h^0 h^0$ in the decoupling limit, we ignore one-loop effects due to the $h^0 b\bar{b}$ coupling and replace the $h^0 h^0 h^0$ coupling by its effective coupling given in eq. (4.1). In figure 3 we present the double Higgs production process for the LHC in the decoupling limit of the 2HDM. In the left panel we show $\sigma_{gg \rightarrow h^0 h^0}$




Figure 3. $pp \rightarrow h^0 h^0$ in the decoupling limit. On the right we show the strength of the one-loop corrected $h^0 h^0 h^0$ coupling relative to SM, for several values of m_{12} . On the left plot we show the production cross section for $m_{h^0} = 115 \text{ GeV}$ as a function of M_Φ for the same values of m_{12} as well as the corresponding SM cross section.

for $m_{h^0} = 115 \text{ GeV}$ as a function of M_Φ for several values of m_{12} . In the right panel we show how the effective coupling behaves with respect to the corresponding SM coupling when we vary M_Φ and m_{12} . Let us first note that in the decoupling limit, the cross section depends only on these three parameters and very mildly on $\tan\beta$ - the sensitivity of the cross section to $\tan\beta$ is mild since $\tan\beta$ only enters the $h^0 h^0 h^0$ coupling through $M^2 = m_{12}^2 / (\sin\beta \cos\beta)$. Moreover, $\tan\beta$ is constrained by perturbative unitarity to be rather moderate. The abrupt cuts in the plots are due perturbative unitarity and vacuum stability constraints. We are considering the case $m_{12}^2 > 0$ where the non-decoupling effects are larger. It is interesting to note that in the decoupling limit the cross sections can be smaller than the SM one. This is because the box diagrams do not depend on m_{12} neither on m_Φ . For small m_Φ , there is a destructive interference between boxes and vertices but as m_Φ grows the total contribution becomes dominated by the vertex diagrams which explain why it is enhanced for large m_Φ . If m_{12} is very large (300 GeV or more) the box contributions become negligible and non-decoupling effects can be seen already for small M_Φ . For our choice of $m_{h^0} = 115 \text{ GeV}$, the SM cross section for the LHC is small, 0.02 pb . As is clear in figure 3, the 2HDM contributions can enhance the cross section by about one order of magnitude and reach 0.4 pb for $m_{12} = 300 \text{ GeV}$. Contrary to what happens in the corresponding process $\gamma\gamma \rightarrow h^0 h^0$ [39], we will not be able to test non-decoupling effects arising from the $h^0 H^+ H^-$ vertex at one loop level because charged Higgs bosons do not couple to gluons. Only non-decoupling effects from higher order corrections can be probed. In the right panel of figure 3, one can see that the coupling $h^0 h^0 h^0$ is enhanced by a factor of more than eight in some cases. This enhancement in the $h^0 h^0 h^0$ coupling is due

to the non decoupling effect shown in eq. (4.1). Finally we remark that the $b\bar{b}$ contribution is much smaller than the gluon fusion one and has a much higher uncertainty coming from the PDFs. We have also checked that $c\bar{c} \rightarrow h^0 h^0$ contribution is much smaller than the $b\bar{b} \rightarrow h^0 h^0$ one.

4.3 The general case for $pp \rightarrow h^0 h^0$

The allowed parameter space of the 2HDM is restricted after imposing experimental and theoretical constraints. Before presenting our results we will scan the parameter space imposing all theoretical constraints discussed before to search for allowed regions where the values of the cross section are above the SM ones. The most severe constraints are the 14 tree-level unitary conditions imposed on the scalar potential parameters.

As explained before, the 2HDM under study has 7 independent parameters. In this particular study, the mass on the pseudo-scalar, m_{A^0} and the mass of the charged Higgs, m_{H^\pm} , do not play any role in the value obtained for the cross section. The reason is that the pure scalar couplings $h^0 h^0 h^0$, $H^0 H^0 H^0$ and $H^0 h^0 h^0$, eqs. (2.2), (2.3) and (2.4), do not involve those parameters. The same is obviously true for the Yukawa couplings. The only possible contribution from those parameters occurs when one of the CP-even Higgs is allowed to decay to a pair of pseudo-scalars or/and to a pair of charged Higgs bosons. In this case, they will influence the total cross section via a change in the total width of the CP-even scalar in question. However, in order to check perturbative unitarity and vacuum stability constraints, m_{A^0} and m_{H^\pm} must be specified. The most relevant parameters we have to consider are therefore m_{h^0} , m_{H^0} , $\sin \alpha$, $\tan \beta$ and above all m_{12} . We have scanned the 2HDM parameter space to look for the regions where the cross sections are larger than the SM one. The results are shown in figures 4 and 5. The detailed parton level studies presented in [6–8] concluded that although hard there is a chance of measuring the SM triple Higgs couplings at the LHC. For some final states the measurement could only take place with the tenfold increase luminosity of the Super Large Hadron Collider. Although a more detailed analysis is needed we will take these studies as a benchmark and look for the parameter space where cross sections are at least larger than the SM double Higgs production.

In figure 4 we present a scan of in the $(m_{12}, \tan \beta)$ plane for $m_{h^0} = 115$ GeV, $m_{A^0} = 350$ GeV, $m_{H^\pm} = 300$ GeV, $m_{H^0} = 250$ GeV, $\tan \beta = 1$ and $\sin \alpha = -0.9$. With this choice of the charged Higgs boson mass, the $b \rightarrow s\gamma$ constraint is fulfilled for all 2HDM models. The scan shows the allowed region of the 2HDM parameter space where the cross section is larger than the SM cross section. Here we see that for reasonable values of the other parameters $\tan \beta$ has to be below 5. We can also witness a correlation between m_{12} and $\tan \beta$: large values of one of the variables force low values of the other variable. In the right panel of figure 4, we illustrate the size of the triple Higgs couplings $h^0 h^0 h^0$ and $H^0 h^0 h^0$ normalized to SM one as a function of $\tan \beta$ and scanned in m_{12} . For small $\tan \beta$ and large m_{12} the couplings can be larger than $8 \times \lambda_{h^0 h^0 h^0}^{\text{SM}}$.

In the left panel of figure 5 we again show the allowed region of the 2HDM parameter space where the cross section is larger than the SM cross section. It is clear that large values of m_{12} are preferred because the self-couplings have a strong dependence on this parameter. As expected, when $m_{H^0} \approx 2m_{h^0}$ the cross section reaches a maximum but




Figure 4. The allowed regions of the 2HDM where the $\sigma(gg \rightarrow h^0 h^0)$ is larger than the corresponding SM cross section (left) and the 2HDM triple coupling $h^0 h^0 h^0$ and $h^0 h^0 h^0$ normalized to SM couplings (right). We show a scan in the $(m_{12}, \tan\beta)$ plane for $m_{h^0} = 115$ GeV, $m_{A^0} = 350$ GeV, $m_{H^\pm} = 300$ GeV, $m_{H^0} = 250$ GeV and $\sin\alpha = -0.9$.




Figure 5. The allowed regions of the 2HDM where the $\sigma(gg \rightarrow h^0 h^0)$ is larger than the corresponding SM cross section. We have chosen $m_{h^0} = 115$ GeV, $m_{A^0} = 350$ GeV, $m_{H^\pm} = 300$ GeV and $\tan\beta = 1$. In the left panel we show a scan in the (m_{12}, m_{H^0}) plane with $\sin\alpha = -0.9$. In the right panel we show a scan in the $(m_{12}, \sin\alpha)$ plane for $m_{H^0} = 250$ GeV.




Figure 6. The cross section for $gg \rightarrow h^0 h^0$ with $m_{A^0} = 320 \text{ GeV}$, $m_{H^\pm} = 380 \text{ GeV}$, $\sin(\beta - \alpha) = 0.1$ and $\tan\beta = 2$. In the left panel we present the cross section as a function of m_{h^0} for $m_{H^0} = 200 \text{ GeV}$ while in the right panel we can see the variation with m_{H^0} for $m_{h^0} = 120 \text{ GeV}$.

all values above $2m_{h^0}$ can give rise to large cross sections depending on the remaining parameters. From the right panel of figure 5, where a scan in the $(m_{12}, \sin\alpha)$ plane is shown, we conclude that large values of $|\sin\alpha|$ maximize the cross section. It should be noted however that these plots would change if the values of the masses were modified.

Let us move to the discussion of double Higgs production $\sigma(gg \rightarrow h^0 h^0)$ at the LHC in the framework of the general 2HDM. The cases of $H^0 H^0$ and $H^0 h^0$ production will be discussed at the end of this section. We already know that the cross section does not depend on m_{A^0} nor on m_{H^\pm} except via the Higgs H^0 or h^0 width. We also know that, due to the theoretical constraints, $\tan\beta$ is constrained to be rather moderate. From various scans over the parameter space, we conclude that in order to have large enhancement with respect to SM cross section one needs to take small values of $\tan\beta$ together with large positive m_{12}^2 . We start by showing how the cross section $pp \rightarrow h^0 h^0$ behaves with the two CP-even Higgs masses. This is illustrated in figure 6, where the cross section $\sigma(gg \rightarrow h^0 h^0)$ at the LHC via gluon fusion is shown for three different values of m_{12} . We have chosen $\sin(\beta - \alpha) = 0.1$ which allow us to probe the very low mass region in m_{h^0} . However, we have checked that similar values for the cross section can be obtained for a very wide range of $\sin(\beta - \alpha)$. As expected, the cross section grows with m_{12} . In the left panel of figure 6 we show $\sigma(gg \rightarrow h^0 h^0)$ as a function of m_{h^0} for $m_{H^0} = 200 \text{ GeV}$. Before the threshold, $m_{H^0} = 2m_{h^0}$, the cross section is of the order of a few pb and does not depend much on the value of m_{12} . This is because once the channel $H^0 \rightarrow h^0 h^0$ is open it becomes dominant with a branching ratio close to 100%. On the other hand, there is another triangle diagram contribution, where a virtual h^0 is exchanged, with a much smaller coupling constant. Therefore, the dependence on m_{12} is mild. In this region, there is a clear chance of measuring the triple coupling $H^0 h^0 h^0$ as the cross section can easily be two orders of magnitude above the SM (note that we show the SM line for all values




Figure 7. The cross section for $gg \rightarrow h^0 h^0$ (left) and the contributing scalar couplings (right) as a function of $\sin \alpha$ for different values of m_{H^0} . The parameters are $m_{h^0} = 110 \text{ GeV}$, $m_{A^0} = 320 \text{ GeV}$, $m_{H^\pm} = 380 \text{ GeV}$, $m_{12} = 340 \text{ GeV}$, and $\tan \beta = 1$.

of the light Higgs mass to serve as a reference although masses below 115 GeV are already excluded). Below the threshold, in the region where $H^0 \rightarrow h^0 h^0$ is closed, all contributions are now on the same footing and depend much more on the set of parameters chosen. For large enough m_{12} the cross section is still above the SM one but as one can see in figure 6 it can also be well below the SM value for $m_{12} = 0$ in the large light Higgs mass region. In the right panel of figure 6 we show a plot for the same set of parameters but now as a function of m_{H^0} . This plot show us exactly what we have just discussed but from a different point of view - that of the heavy CP-even Higgs mass. Again we see that there is a limit for what the cross section can grow by increasing m_{12} and that limit is attained when $Br(H^0 \rightarrow h^0 h^0) \approx 100\%$. Finally let us comment that these cross sections do not vary much when we change the Yukawa couplings. There is only a small difference due to the Higgs widths which is however too small to be measured at the LHC.

In figure 7 (left panel) we show the cross section as a function of $\sin \alpha$ for different values of m_{H^0} , one below the $H^0 \rightarrow h^0 h^0$ threshold, one above but almost on the threshold and the other one above the threshold. In the right panel we present the value of the couplings $H^0 h^0 h^0$ and $h^0 h^0 h^0$ normalized to SM one which prove to be the main contributions by far to the value of the cross section. This was explicitly checked by repeating the calculation with only the triangle diagrams where either a virtual h^0 or a virtual H^0 was exchanged. Combining the information from the two figures we conclude that the triangle diagram with the virtual H^0 exchanged dominates for $-1 < \sin \alpha < -0.6$. Therefore in this range, and for this set of parameters the coupling $H^0 h^0 h^0$ could be measured because the cross section is much larger than the corresponding SM cross section. After crossing $\sin \alpha = -0.6$ both couplings grow and now the relative importance also depends on the value of m_{H^0} : if $H^0 \rightarrow h^0 h^0$ is open then the diagram with the virtual H^0 still dominates and there is a




Figure 8. $\sigma(gg \rightarrow h^0h^0, H^0h^0$ and $H^0H^0)$ as a function of $\sin\alpha$. The parameters chosen are $m_{h^0} = 120\text{ GeV}$, $m_{H^0} = 121\text{ GeV}$, $m_{A^0} = 200\text{ GeV}$, $m_{H^\pm} = 300\text{ GeV}$ and $\tan\beta = 1$. In the left panel $m_{12} = 100\text{ GeV}$ and in the right panel $m_{12} = 300\text{ GeV}$.

good chance of measuring it; below the threshold, and for this particular set of parameters, it is the diagram with the virtual h^0 exchange that dominates. We conclude by saying that depending on the parameters there is a good chance of measuring the coupling $H^0h^0h^0$ or $h^0h^0h^0$ and that this occurs in a large portion of the parameter space.

4.4 The H^0H^0 and H^0h^0 production modes

Processes $gg \rightarrow H^0H^0, H^0h^0$ differ from $gg \rightarrow h^0h^0$ mainly due to the fact that H^0 is heavier than h^0 . This means that there is no possibility of resonant behavior - the intermediate state is always off-shell - and that there is a reduction of the phase space available. We may ask ourselves what would the difference be if the two CP-even masses were of similar magnitude. In this case, it is the relation between $\sin\alpha$, m_{12} and $\tan\beta$ that determines the value of the cross section, that is, it is the way these three variables are combined that determines how large the cross section is. Because we know that $\tan\beta$ has to be small we have fixed $\tan\beta = 1$ and varied the remaining two parameters for $m_{h^0} = 120\text{ GeV}$ and $m_{H^0} = 121\text{ GeV}$. In figure 8 we present all three cross sections $\sigma(gg \rightarrow h^0h^0, H^0h^0$ and $H^0H^0)$ as a function of $\sin\alpha$ for $m_{h^0} = 120\text{ GeV}$, $m_{H^0} = 121\text{ GeV}$, $m_{A^0} = 200\text{ GeV}$, $m_{H^\pm} = 300\text{ GeV}$ and $\tan\beta = 1$. In the left panel we take $m_{12} = 100\text{ GeV}$ while we choose $m_{12} = 300\text{ GeV}$ for the right panel. We start by noting that we have checked that even if the value of m_{H^0} is shifted up several tenths of GeV the general trends do not change much. It is clear that in any case the cross section can reach several hundreds of fbarn. However, a large m_{12} is needed because for $m_{12} = 100\text{ GeV}$ the cross section barely reaches the SM value. We can see clearly two effects due to the couplings:




Figure 9. Light Higgs branching ratios in model I as a function of its mass (left panel) and as a function of $\sin \alpha$ (right panel). Although irrelevant in Model I we have chosen $\sin(\beta - \alpha) = 0.1$, $\tan \beta = 1$, $m_{H^0} = m_{A^0} = m_{H^\pm} = 300 \text{ GeV}$ and $m_{12} = 0 \text{ GeV}$.

the huge dependence on the value of $\sin \alpha$ due to the Yukawa couplings and the dependence on the relation between m_{12} and $\sin \alpha$ caused by the scalar couplings. We conclude that if the CP-even scalar masses are similar and not too large, the production of these particles is heavily conditioned by the values of m_{12} and $\sin \alpha$. Finally we should point out that in this particular limit, the two CP-even Higgs bosons would be undistinguishable experimentally and therefore the number of events would be obtained by summing the three contributions.

5 Higgs signatures

In most of the 2HDM parameter space, the dominant process in the production of a pair of neutral Higgs is the triangle diagram where $gg \rightarrow H^0 \rightarrow h^0 h^0$. As the mass of the heavy Higgs grows, the triangle diagram $gg \rightarrow h^0 \rightarrow h^0 h^0$ starts to play an important role. Therefore searches based on these topologies in the SM and in the MSSM should yield similar results. Searches based on $4b$ final state were performed in [7] for the SM and in [9, 10] for the MSSM. The final state $b\bar{b}\tau^+\tau^-$ was also explored in [7] for the SM. The search based on rare decays, $h^0 h^0 \rightarrow b\bar{b}\gamma\gamma$ both for the SM and for the MSSM was performed in [8]. In this section we will present the profile of the light Higgs boson in all four Yukawa type models. We will then comment on the results, showing that new rare decays can be used in the 2HDM case.

We will discuss a scenario where the light Higgs boson's mass is well below $\approx 200 \text{ GeV}$ and all decay channels that have at least one other Higgs in the final state are closed. Scenarios where the light Higgs boson decays to a pair of charged Higgs, a pair of CP-odd Higgs or a combination of Higgs and gauge bosons are obviously allowed in a general




Figure 10. Light Higgs branching ratios in models II (left panel) and III (right panel) as a function of $\sin \alpha$ for $\tan \beta = 1$. Again, although irrelevant, we have chosen the other parameters as $m_{H^0} = m_{A^0} = m_{H^\pm} = 300 \text{ GeV}$ and $m_{12} = 0 \text{ GeV}$.

2HDM. In the decoupling limit, the lightest CP-even Higgs has the SM signature. This means that close to the limit $(\beta - \alpha) \approx \pi/2$ no major differences should be expected in the light Higgs boson profile with respect to the SM Higgs. The same is true for model type I. In figure 9 we show the profile of the light Higgs boson in Model I as a function of its mass in the left panel and as a function of $\sin \alpha$ in the right panel. As in Model I only one Higgs couples to all fermions, the only dependence is in the mass of the light Higgs. The branching ratios do not depend on any other parameters of the model. It should be noted however that when $\sin \alpha = 1$ (the fermiophobic limit) the decay to gauge bosons become dominant. In this case the light Higgs will decay mainly to two photons or to two W bosons depending on the Higgs mass considered and this trend can be seen in the right panel of figure 9. The values presented in the right panel, $\sin(\beta - \alpha) = 0.1$ and $\tan \beta = 1$ are irrelevant for Model I for the channels presented. However, they are relevant for the other models and the left panel of figure 9 also shows the qualitative behavior of Models II, III and IV for the same set of parameters. Therefore, for the set of values presented this plot is the same in all four models. When $\tan \beta = 1$ and at the same time $\beta \approx \alpha$ the angular dependence almost cancels resulting in a SM-like scenario. As $\tan \beta$ grows this is no longer true: in Model II the coupling to up quarks is suppressed resulting again in a SM-like scenario with the $b\bar{b}$ and the $\tau^+\tau^-$ mode dominating; in Model III all couplings are now suppressed except the ones to the down quarks which means that the Higgs decays almost exclusively to b-pairs; finally in Model IV, and again for high $\tan \beta$ the Higgs decays almost exclusively to $\tau^+\tau^-$. We should add that, moving away from $\beta \approx \alpha$, as the mass of the light Higgs grows, the decays $h^0 \rightarrow WW^*$ and $h^0 \rightarrow ZZ^*$ become increasingly important although their branching ratios are always below the SM values.




Figure 11. Light Higgs branching ratios as a function of $\sin \alpha$ in model IV for $\tan \beta = 1$ (left panel) and $\tan \beta = 4$ (right panel). The other parameters are $m_{H^0} = m_{A^0} = m_{H^\pm} = 300 \text{ GeV}$ and $m_{12} = 0 \text{ GeV}$.

Most interesting is the dependence of the light Higgs branching ratio with $\sin \alpha$. In figure 10 we show the light Higgs branching ratios as a function of $\sin \alpha$ for Models II (left) and III (right) and for $\tan \beta = 1$. In Model II, when $\sin \alpha$ is close to zero, the dominant decays become $c\bar{c}$ and gg . Therefore, in this region a decay to non b-jets can easily reach 100% making the detection of Higgs pairs extremely hard due to the overwhelming QCD background. In Model III a more pleasant situation emerges. In the same $\sin \alpha$ region, the decay to $\tau^+\tau^-$ dominates reaching 50% for $\sin \alpha = 0$. At the same time the decay channel $h^0 \rightarrow \mu^+\mu^-$ is well above the 0.1% line turning the final state $\tau^+\tau^-\mu^+\mu^-$ into a very good candidate. We should point out that these final states occur in significant regions of the parameter space as can be seen in the figures. Finally, note that we don't show the respective plots for large $\tan \beta$ as we would recover the behavior described previously, that is, in Model II (left panel) the light Higgs would decay exclusively to $b\bar{b}$ and $\tau^+\tau^-$ and in Model III (right panel) $Br(h^0 \rightarrow b\bar{b}) \approx 100\%$.

In figure 11 we show the light Higgs branching ratios as a function of $\sin \alpha$ for Model IV and for $\tan \beta = 1$ (left panel) and $\tan \beta = 4$ (right panel). For $\tan \beta = 1$ the Higgs decays mainly to b pairs. Away from $\sin \alpha = 0$ the decay to τ pairs can reach interesting values and becomes the dominant decay mode when $|\sin \alpha| \approx 1$. For $\tan \beta = 4$ both modes almost share the total branching ratio between them. Once more we should point out that when $\sin \alpha$ moves away from zero, not only the decay to τ pairs become the dominant mode, but also the decay to μ pairs reach almost 0.4% for $|\sin \alpha| = 1$. Hence, new decay channels can be explored in the context of 2HDM.

The process $pp \rightarrow H^0 H^0$ gives rise to very similar signatures. The only disadvantage is that H^0 is in principle heavier than h^0 and phase space is lost. If the masses are of the

same order and below let us say 130 GeV , then the main decays are again $b\bar{b}$ and $\tau^+\tau^-$. For larger values of the mass again the decays $h^0 \rightarrow WW^*$ and $h^0 \rightarrow ZZ^*$ become dominant. In this limit they differ only in the behavior with $\sin\alpha$ and $\tan\beta$ in all Yukawa models except model I. Finally, process $pp \rightarrow h^0 H^0$, has in principle two drawbacks: first the phase space is again reduced due to the presence of a heavier Higgs in the final state; second we can not reduce the background by asking the two invariant masses from each pair that originates from a Higgs bosons (for instance two b-jets) to have a similar magnitude.

6 Discussion and conclusions

In this work we have calculated the cross section for the production processes $pp \rightarrow h^0 h^0, H^0 H^0, H^0 h^0$. We have shown that gluon fusion is always the dominant contribution. Regarding the different Yukawa models, little difference was found in the scanned parameter space - although the cross sections presented are for models II and III they are representative of all Yukawa models.

We started by assessing the possibility of distinguishing the lightest CP-even Higgs boson from the SM Higgs in the decoupling limit of the 2HDM. We concluded that the non-decoupling effects entering the amplitude via higher order corrections to the vertex $h^0 h^0 h^0$ can be seen for very large values of the soft breaking parameter m_{12} or for very large values of the common mass scale M_Φ . Taking for instance $m_{12} = 300 \text{ GeV}$ the cross section is more than one order of magnitude above the corresponding SM value.

The general 2HDM has too many parameters to allow for a detailed analysis taking all of them into account. However, for the processes studied the only relevant parameters are $m_{12}, \sin\alpha, \tan\beta, m_{H^0}$ and m_{h^0} . The remaining masses have only little influence via their contribution to the CP-even Higgs widths. We have scanned the parameter space to conclude that, as a general trend, the largest cross sections are obtained for large m_{12} and large $|\sin\alpha|$. The parameter $\tan\beta$ is constrained to be $O(1)$ except for very specific scenarios. Note however that we have showed that even for $m_{12} = 0$ cross sections can be 100 times above the SM cross section when $m_{H^0} > 2m_{h^0}$. The cross sections are larger for smaller values of the CP-even Higgs masses and can be two orders of magnitude above the SM cross section if the channel $H^0 \rightarrow h^0 h^0$ is open. We have shown that there are scenarios where the triple couplings could be measured.

We have then moved to the study of the other production modes $H^0 H^0$ and $H^0 h^0$. We have showed that all production modes can be large and that the main variation occurs with the combination of $\sin\alpha$ and m_{12} parameters. In fact, we have shown that when $m_{H^0} \approx m_{h^0}$ the different processes differ only by virtue of values of the parameters $\sin\alpha$ and m_{12} chosen as $\tan\beta$ is bounded to be small.

In the last section we have studied the possible final states for each production mode. We have shown that the main decays for a light CP-even Higgs are to b-quark pairs or to $\tau^+\tau^-$. The relative importance of each of these decays depend very much on the values of $\sin\alpha$ and $\tan\beta$. In [7] it was shown that the best channel to measure the SM triple Higgs coupling is $2b2\gamma$ for a Higgs mass below 140 GeV . In all Yukawa versions of the 2HDM, the branching ratio of Higgs to two photons does not differ much from the SM. For

	I	II	III	IV
α_{eh}	$-\frac{\cos \alpha}{\sin \beta}$	$\frac{\sin \alpha}{\cos \beta}$	$-\frac{\cos \alpha}{\sin \beta}$	$\frac{\sin \alpha}{\cos \beta}$
α_{dh}	$-\frac{\cos \alpha}{\sin \beta}$	$\frac{\sin \alpha}{\cos \beta}$	$\frac{\sin \alpha}{\cos \beta}$	$-\frac{\cos \alpha}{\sin \beta}$
α_{eH}	$-\frac{\sin \alpha}{\sin \beta}$	$-\frac{\cos \alpha}{\cos \beta}$	$-\frac{\sin \alpha}{\sin \beta}$	$-\frac{\cos \alpha}{\cos \beta}$
α_{dH}	$-\frac{\sin \alpha}{\sin \beta}$	$-\frac{\cos \alpha}{\cos \beta}$	$-\frac{\cos \alpha}{\cos \beta}$	$-\frac{\sin \alpha}{\sin \beta}$
β_e	$-\cot \beta$	$\tan \beta$	$-\cot \beta$	$\tan \beta$
β_d	$-\cot \beta$	$\tan \beta$	$\tan \beta$	$-\cot \beta$

Table 1. Coupling constants for the fermion-scalar interactions.

most of the parameter space, the decay to $b\bar{b}$ is either equal or enhanced relative to the SM. Therefore, this channel should again be one of the most promising to measure 2HDM triple couplings. The channel $2b\mu^+\mu^-$ was also studied and shown to be one order of magnitude below $2b2\gamma$. With larger values for the production cross section and the enhancement of the $h^0 \rightarrow \mu^+\mu^-$ branching ratio in some of the parameter space of Models III and IV, this is a channel worth exploring. Finally, other channels like $4b$, $2b\tau^+\tau^-$, and $2b\mu^+\mu^-$ could be reevaluated for the low mass region when the 2HDM cross section is well above the SM one. For heavier Higgs we would have two distinguish two scenarios. One, where decay channels with at least one other Higgs in the final state are open; in this scenario a new study would have to be performed [42]. The other where the decays to $WW^{(*)}$ and $ZZ^{(*)}$ become important, which means a larger h^0 mass and $\alpha - \beta$ away from zero; in this scenario, the analysis will heavily depend on the values of the branching ratios of the decays $h^0 \rightarrow WW^{(*)}$ and $h^0 \rightarrow ZZ^{(*)}$.

Acknowledgments

A. A. acknowledges the warm hospitality of the University of Southampton, where some of this research took place. A.A is supported by the National Science of Theoretical Studies-Taipei under contract # 980528731. R.B. is supported by National Cheng Kung University Grant No. HUA 97-03-02-063. C.C.H is supported by the National Science Council of R.O.C under Grant #s: NSC-97-2112-M-006-001-MY3. R.B.G. is supported by a Fundação para a Ciência e Tecnologia Grant SFRH/BPD/47348/2008. R.S. is supported by the FP7 via a Marie Curie Intra European Fellowship, contract number PIEF-GA-2008-221707.

A Fermion-scalar vertices

$$\begin{aligned}
 \bar{e}_i e_i h & \quad \frac{ig}{2M_W} \alpha_{eh} m_{e_i} & \quad \bar{u}_i u_i G_0 & \quad -\frac{g}{2M_W} m_{u_i} \gamma_5 \\
 \bar{u}_i u_i h & \quad -\frac{ig}{2M_W} \frac{\cos \alpha}{\sin \beta} m_{u_i} & \quad \bar{d}_i d_i G_0 & \quad \frac{g}{2M_W} m_{d_i} \gamma_5 \\
 \bar{d}_i d_i h & \quad \frac{ig}{2M_W} \alpha_{dh} m_{d_i} & \quad \bar{e}_i \nu_i H^+ & \quad \frac{ig}{2\sqrt{2}M_W} \beta_e m_{e_i} (1 + \gamma_5)
 \end{aligned}$$

$$\begin{array}{ll}
\bar{e}_i e_i H & \frac{ig}{2M_W} \alpha_{eH} m_{e_i} & \bar{u}_i d_j H^+ & \frac{ig}{2\sqrt{2}M_W} V_{ij} [\beta_d m_{d_j} (1 + \gamma_5) + \cot \beta m_{u_i} (1 - \gamma_5)] \\
\bar{u}_i u_i H & -\frac{ig}{2M_W} \frac{\sin \alpha}{\sin \beta} m_{u_i} & \bar{\nu}_i e_i H^- & \frac{ig}{2\sqrt{2}M_W} \beta_e m_{e_i} (1 - \gamma_5) \\
\bar{d}_i d_i H & \frac{ig}{2M_W} \alpha_{dH} m_{d_i} & \bar{d}_i u_j H^- & \frac{ig}{2\sqrt{2}M_W} V_{ij}^* [\beta_d m_{d_i} (1 - \gamma_5) + \cot \beta m_{u_j} (1 + \gamma_5)] \\
\bar{e}_i e_i A & -\frac{g}{2M_W} \beta_e m_{e_i} \gamma_5 & \bar{e}_i \nu_i G^+ & -\frac{ig}{2\sqrt{2}M_W} m_{e_i} (1 + \gamma_5) \\
\bar{u}_i u_i A & -\frac{g}{2M_W} \cot \beta m_{u_i} \gamma_5 & \bar{u}_i d_j G^+ & \frac{ig}{2\sqrt{2}M_W} V_{ij} [-m_{d_j} (1 + \gamma_5) + m_{u_i} (1 - \gamma_5)] \\
\bar{d}_i d_i A & -\frac{g}{2M_W} \beta_d m_{d_i} \gamma_5 & \bar{\nu}_i e_i G^- & -\frac{ig}{2\sqrt{2}M_W} m_{e_i} (1 - \gamma_5) \\
\bar{e}_i e_i G_0 & \frac{g}{2M_W} m_{e_i} \gamma_5 & \bar{d}_i u_j G^- & \frac{ig}{2\sqrt{2}M_W} V_{ij}^* [-m_{d_i} (1 - \gamma_5) + m_{u_j} (1 + \gamma_5)]
\end{array}$$

References

- [1] TEVATRON NEW PHENOMENA HIGGS WORKING GROUP collaboration, G. Bernardi et al., *Combined CDF and D0 upper limits on standard model Higgs boson production at high mass (155 – 200 GeV/c²) with 3fb⁻¹ of data*, [arXiv:0808.0534](#) [SPIRES].
- [2] E.W.N. Glover and J.J. van der Bij, *Higgs boson pair production via gluon fusion*, *Nucl. Phys. B* **309** (1988) 282 [SPIRES].
- [3] D.A. Dicus, C. Kao and S.S.D. Willenbrock, *Higgs boson pair production from gluon fusion*, *Phys. Lett. B* **203** (1988) 457 [SPIRES].
- [4] G.D. Kribs, T. Plehn, M. Spannowsky and T.M.P. Tait, *Four generations and Higgs physics*, *Phys. Rev. D* **76** (2007) 075016 [[arXiv:0706.3718](#)] [SPIRES].
- [5] T. Plehn, M. Spira and P.M. Zerwas, *Pair production of neutral Higgs particles in gluon-gluon collisions*, *Nucl. Phys. B* **479** (1996) 46 [Erratum *ibid.* **B 531** (1998) 655] [[hep-ph/9603205](#)] [SPIRES].
- [6] U. Baur, T. Plehn and D.L. Rainwater, *Determining the Higgs boson selfcoupling at hadron colliders*, *Phys. Rev. D* **67** (2003) 033003 [[hep-ph/0211224](#)] [SPIRES].
- [7] U. Baur, T. Plehn and D.L. Rainwater, *Examining the Higgs boson potential at lepton and hadron colliders: a comparative analysis*, *Phys. Rev. D* **68** (2003) 033001 [[hep-ph/0304015](#)] [SPIRES].
- [8] U. Baur, T. Plehn and D.L. Rainwater, *Probing the Higgs selfcoupling at hadron colliders using rare decays*, *Phys. Rev. D* **69** (2004) 053004 [[hep-ph/0310056](#)] [SPIRES].
- [9] J. Dai, J.F. Gunion and R. Vega, *Detection of the minimal supersymmetric model Higgs boson H⁰ in its h⁰h⁰ → 4b and A⁰A⁰ → 4b decay channels*, *Phys. Lett. B* **371** (1996) 71 [[hep-ph/9511319](#)] [SPIRES].
- [10] J. Dai, J.F. Gunion and R. Vega, *Detection of neutral MSSM Higgs bosons in four-b final states at the Tevatron and the LHC: an update*, *Phys. Lett. B* **387** (1996) 801 [[hep-ph/9607379](#)] [SPIRES].

- [11] S. Dawson, S. Dittmaier and M. Spira, *Neutral Higgs-boson pair production at hadron colliders: QCD corrections*, *Phys. Rev. D* **58** (1998) 115012 [[hep-ph/9805244](#)] [[SPIRES](#)].
- [12] M. Moretti, S. Moretti, F. Piccinini, R. Pittau and J. Rathsmann, *Vector-boson production of light Higgs pairs in 2-Higgs doublet models*, *JHEP* **12** (2007) 075 [[arXiv:0706.4117](#)] [[SPIRES](#)];
M. Moretti, S. Moretti, F. Piccinini, R. Pittau and A.D. Polosa, *Higgs boson self-couplings at the LHC as a probe of extended Higgs sectors*, *JHEP* **02** (2005) 024 [[hep-ph/0410334](#)] [[SPIRES](#)].
- [13] R.N. Hodgkinson, D. Lopez-Val and J. Solà, *Higgs boson pair production through gauge boson fusion at linear colliders within the general 2HDM*, *Phys. Lett. B* **673** (2009) 47 [[arXiv:0901.2257](#)] [[SPIRES](#)].
- [14] A. Arhrib, R. Benbrik, R.B. Guedes and R. Santos, *Search for a light fermiophobic Higgs boson produced via gluon fusion at hadron colliders*, *Phys. Rev. D* **78** (2008) 075002 [[arXiv:0805.1603](#)] [[SPIRES](#)].
- [15] J. Velhinho, R. Santos and A. Barroso, *Tree level vacuum stability in two-Higgs doublet models*, *Phys. Lett. B* **322** (1994) 213 [[SPIRES](#)].
- [16] G.C. Branco, *Spontaneous CP nonconservation and natural flavor conservation: a minimal model*, *Phys. Rev. D* **22** (1980) 2901 [[SPIRES](#)].
- [17] T.D. Lee, *A theory of spontaneous T violation*, *Phys. Rev. D* **8** (1973) 1226 [[SPIRES](#)].
- [18] G.C. Branco and M.N. Rebelo, *The Higgs mass in a model with two scalar doublets and spontaneous CP-violation*, *Phys. Lett. B* **160** (1985) 117 [[SPIRES](#)].
- [19] P.M. Ferreira, R. Santos and A. Barroso, *Stability of the tree-level vacuum in two Higgs doublet models against charge or CP spontaneous violation*, *Phys. Lett. B* **603** (2004) 219 [*Erratum ibid.* **B 629** (2005) 114] [[hep-ph/0406231](#)] [[SPIRES](#)].
- [20] I.P. Ivanov, *Minkowski space structure of the Higgs potential in 2HDM*, *Phys. Rev. D* **75** (2007) 035001 [*Erratum ibid.* **D 76** (2007) 039902] [[hep-ph/0609018](#)] [[SPIRES](#)].
- [21] S.L. Glashow and S. Weinberg, *Natural conservation laws for neutral currents*, *Phys. Rev. D* **15** (1977) 1958 [[SPIRES](#)].
- [22] V.D. Barger, J.L. Hewett and R.J.N. Phillips, *New constraints on the charged Higgs sector in two Higgs doublet models*, *Phys. Rev. D* **41** (1990) 3421 [[SPIRES](#)].
- [23] R.M. Barnett, G. Senjanovic, L. Wolfenstein and D. Wyler, *Implications of a light Higgs scalar*, *Phys. Lett. B* **136** (1984) 191 [[SPIRES](#)];
R.M. Barnett, G. Senjanovic and D. Wyler, *Tracking down Higgs scalars with enhanced couplings*, *Phys. Rev. D* **30** (1984) 1529 [[SPIRES](#)];
H.-S. Goh, L.J. Hall and P. Kumar, *The leptonic Higgs as a messenger of dark matter*, *JHEP* **05** (2009) 097 [[arXiv:0902.0814](#)] [[SPIRES](#)].
- [24] M. Aoki, S. Kanemura and O. Seto, *Neutrino mass, dark matter and baryon asymmetry via TeV-scale physics without fine-tuning*, *Phys. Rev. Lett.* **102** (2009) 051805 [[arXiv:0807.0361](#)] [[SPIRES](#)];
M. Aoki, S. Kanemura, K. Tsumura and K. Yagyu, *Models of Yukawa interaction in the two Higgs doublet model and their collider phenomenology*, [arXiv:0902.4665](#) [[SPIRES](#)];
S. Su and B. Thomas, *The LHC discovery potential of a leptophilic Higgs*, [arXiv:0903.0667](#) [[SPIRES](#)];

- H.E. Logan and D. MacLennan, *Charged Higgs phenomenology in the lepton-specific two Higgs doublet model*, *Phys. Rev. D* **79** (2009) 115022 [[arXiv:0903.2246](#)] [[SPIRES](#)];
- V. Barger, H.E. Logan and G. Shaughnessy, *Identifying extended Higgs models at the LHC*, *Phys. Rev. D* **79** (2009) 115018 [[arXiv:0902.0170](#)] [[SPIRES](#)];
- A.G. Akeroyd and W.J. Stirling, *Light charged Higgs scalars at high-energy e^+e^- colliders*, *Nucl. Phys. B* **447** (1995) 3 [[SPIRES](#)];
- A.G. Akeroyd, *Non-minimal neutral Higgs bosons at LEP2*, *Phys. Lett. B* **377** (1996) 95 [[hep-ph/9603445](#)] [[SPIRES](#)]; *Fermiophobic and other non-minimal neutral Higgs bosons at the LHC*, *J. Phys. G* **24** (1998) 1983 [[hep-ph/9803324](#)] [[SPIRES](#)].
- [25] N.G. Deshpande and E. Ma, *Pattern of symmetry breaking with two Higgs doublets*, *Phys. Rev. D* **18** (1978) 2574 [[SPIRES](#)].
- [26] S. Kanemura, T. Kubota and E. Takasugi, *Lee-Quigg-Thacker bounds for Higgs boson masses in a two doublet model*, *Phys. Lett. B* **313** (1993) 155 [[hep-ph/9303263](#)] [[SPIRES](#)].
- [27] A.G. Akeroyd, A. Arhrib and E.-M. Naimi, *Note on tree-level unitarity in the general two Higgs doublet model*, *Phys. Lett. B* **490** (2000) 119 [[hep-ph/0006035](#)] [[SPIRES](#)];
- A. Arhrib, *Unitarity constraints on scalar parameters of the standard and two Higgs doublets model*, [hep-ph/0012353](#) [[SPIRES](#)];
- J. Horejsi and M. Kladiva, *Tree-unitarity bounds for THDM Higgs masses revisited*, *Eur. Phys. J. C* **46** (2006) 81 [[hep-ph/0510154](#)] [[SPIRES](#)].
- [28] A. Denner, R.J. Guth, W. Hollik and J.H. Kuhn, *The Z width in the two Higgs doublet model*, *Z. Phys. C* **51** (1991) 695 [[SPIRES](#)].
- [29] PARTICLE DATA GROUP collaboration, S. Eidelman et al., *Review of particle physics*, *Phys. Lett. B* **592** (2004) 1 [[SPIRES](#)].
- [30] A.W. El Kaffas, O.M. Ogreid and P. Osland, *Profile of two-Higgs-doublet-model parameter space*, [arXiv:0709.4203](#) [[SPIRES](#)];
- A.W. El Kaffas, P. Osland and O.M. Ogreid, *Constraining the two-Higgs-doublet-model parameter space*, *Phys. Rev. D* **76** (2007) 095001 [[arXiv:0706.2997](#)] [[SPIRES](#)].
- [31] OPAL collaboration, G. Abbiendi et al., *Flavour independent $h^0 A^0$ search and two Higgs doublet model interpretation of neutral Higgs boson searches at LEP*, *Eur. Phys. J. C* **40** (2005) 317 [[hep-ex/0408097](#)] [[SPIRES](#)].
- [32] L3 collaboration, P. Achard et al., *Flavor independent search for neutral Higgs bosons at LEP*, *Phys. Lett. B* **583** (2004) 14 [[hep-ex/0402003](#)] [[SPIRES](#)].
- [33] DELPHI collaboration, J. Abdallah et al., *Searches for neutral Higgs bosons in extended models*, *Eur. Phys. J. C* **38** (2004) 1 [[hep-ex/0410017](#)] [[SPIRES](#)].
- [34] T. Hahn, *Generating Feynman diagrams and amplitudes with FeynArts 3*, *Comput. Phys. Commun.* **140** (2001) 418 [[hep-ph/0012260](#)] [[SPIRES](#)];
- T. Hahn and C. Schappacher, *The implementation of the minimal supersymmetric standard model in FeynArts and FormCalc*, *Comput. Phys. Commun.* **143** (2002) 54 [[hep-ph/0105349](#)] [[SPIRES](#)];
- J. Küblbeck, M. Böhm and A. Denner, *Feyn arts: computer algebraic generation of Feynman graphs and amplitudes*, *Comput. Phys. Commun.* **60** (1990) 165 [[SPIRES](#)].
- [35] T. Hahn and J.I. Illana, *Extensions in FormCalc 5.3*, [PoS\(ACAT\)074](#) [[arXiv:0708.3652](#)] [[SPIRES](#)]; *Excursions into FeynArts and FormCalc*, *Nucl. Phys. Proc. Suppl.* **160** (2006) 101 [[hep-ph/0607049](#)] [[SPIRES](#)];

- T. Hahn and M. Perez-Victoria, *Automatized one-loop calculations in four and D dimensions*, *Comput. Phys. Commun.* **118** (1999) 153 [[hep-ph/9807565](#)] [[SPIRES](#)].
- [36] G.J. van Oldenborgh, *FF: a package to evaluate one loop Feynman diagrams*, *Comput. Phys. Commun.* **66** (1991) 1 [[SPIRES](#)];
T. Hahn, *Loop calculations with FeynArts, FormCalc, and LoopTools*, *Acta Phys. Polon.* **B 30** (1999) 3469 [[hep-ph/9910227](#)] [[SPIRES](#)].
- [37] J.F. Gunion and H.E. Haber, *The CP-conserving two-Higgs-doublet model: the approach to the decoupling limit*, *Phys. Rev.* **D 67** (2003) 075019 [[hep-ph/0207010](#)] [[SPIRES](#)].
- [38] S. Kanemura, S. Kiyoura, Y. Okada, E. Senaha and C.P. Yuan, *New physics effect on the Higgs self-coupling*, *Phys. Lett.* **B 558** (2003) 157 [[hep-ph/0211308](#)] [[SPIRES](#)];
S. Kanemura, Y. Okada, E. Senaha and C.P. Yuan, *Higgs coupling constants as a probe of new physics*, *Phys. Rev.* **D 70** (2004) 115002 [[hep-ph/0408364](#)] [[SPIRES](#)].
- [39] A. Arhrib, R. Benbrik, C.-H. Chen and R. Santos, *Neutral Higgs boson pair production in photon-photon annihilation in the two Higgs doublet model*, [arXiv:0901.3380](#) [[SPIRES](#)];
F. Cornet and W. Hollik, *Pair production of two-Higgs-doublet-model light Higgs bosons in $\gamma\gamma$ collisions*, *Phys. Lett.* **B 669** (2008) 58 [[arXiv:0808.0719](#)] [[SPIRES](#)];
E. Asakawa, D. Harada, S. Kanemura, Y. Okada and K. Tsumura, *Higgs boson pair production at a photon-photon collision in the two Higgs doublet model*, *Phys. Lett.* **B 672** (2009) 354 [[arXiv:0809.0094](#)] [[SPIRES](#)].
- [40] I.F. Ginzburg, M. Krawczyk and P. Osland, *Potential of photon collider in resolving SM-like scenarios*, *Nucl. Instrum. Meth.* **A 472**, 149 (2001) [[hep-ph/0101229](#)] [[SPIRES](#)].
- [41] A. Arhrib, W. Hollik, S. Penaranda and M. Capdequi Peyranere, *Higgs decays in the two Higgs doublet model: large quantum effects in the decoupling regime*, *Phys. Lett.* **B 579** (2004) 361 [[SPIRES](#)].
- [42] S. Kanemura, S. Moretti, Y. Mukai, R. Santos and K. Yagyu, *Distinctive Higgs signals of a type II 2HDM at the LHC*, [arXiv:0901.0204](#) [[SPIRES](#)].