

Introduction to Little String Theory

David Kutasov*

Department of Physics, University of Chicago, Chicago, USA

*Lectures given at the
Spring School on Superstrings and Related Matters
Trieste, 2 – 10 April 2001*

LNS027004

* kutasov@theory.uchicago.edu

Abstract

These notes, based on lectures presented at the ICTP Spring School on Superstrings and Related Matters in April 2001, provide an introduction to Little String Theory.

Contents

| | | |
|----------|--|------------|
| 1 | Introduction | 167 |
| 2 | The decoupling limit of flat $NS5$-branes | 169 |
| 3 | A holographically dual description of LST | 172 |
| 3.1 | Example 1: Chiral operators in LST | 175 |
| 3.2 | Example 2: Normalizable states | 176 |
| 3.3 | The strong coupling problem | 177 |
| 4 | High energy thermodynamics of LST | 179 |
| 4.1 | Thermodynamics of near-extremal fivebranes | 179 |
| 4.2 | The leading $1/\mu$ correction to classical thermodynamics | 182 |
| 4.3 | Comments on the near-Hagedorn thermodynamics of LST | 189 |
| 5 | Weakly coupled LST | 191 |
| 6 | Other aspects of LST | 198 |
| 6.1 | Singular Calabi-Yau manifolds and lower dimensional vacua of LST | 198 |
| 6.2 | D-branes in the vicinity of $NS5$ -branes | 200 |
| 6.3 | Low dimensional toy models of LST | 201 |
| 7 | Some open problems in LST | 202 |
| | References | 204 |

1 Introduction

Much has been learned over the years by studying string dynamics near various kinds of “impurities.” Examples include string propagation on orbifolds [1], where one finds “twisted sectors” corresponding to fundamental strings trapped at the orbifold singularities, and vacua with D-branes which contain localized excitations corresponding to open strings ending on the branes.

In both of these examples, the states localized at the impurity couple to the bulk – *e.g.* two open strings ending on a D-brane can fuse into a closed string that can leave the brane. It is sometimes possible to decouple the physics of the localized modes from bulk dynamics by taking a low energy limit, $E \ll m_s$, where $m_s = 1/\sqrt{\alpha'}$ is the string scale, associated with the tension of the fundamental string $T = 1/2\pi\alpha'$.

Whenever this limit gives rise to an interacting theory, it corresponds to a local quantum field theory (QFT), such as the non-abelian gauge theories found on branes. This embedding of field theoretic dynamics into string theory led in recent years to many insights into field theory and string theory (see *e.g.* [2, 3] for reviews).

The purpose of these lectures is to describe another class of impurities – Neveu-Schwarz fivebranes [4], or equivalently singularities of Calabi-Yau manifolds and other spaces¹. One of the striking features of the dynamics of $NS5$ -branes is that it can be decoupled from the bulk without taking the low energy limit $\alpha' \rightarrow 0$. The decoupled theory of $NS5$ -branes is known as Little String Theory² (LST). It has the following properties:

- (1) The theory is non-local. In particular, upon compactification on tori, LST exhibits T-duality.
- (2) It has a Hagedorn density of states at high energies, $\rho(E) \sim E^\alpha \exp(\beta_H E)$.
- (3) The theory can be defined in six or fewer spacetime dimensions. It has super – Poincare invariant vacua with sixteen or fewer supercharges.
- (4) LST is a non-gravitational theory: there is no massless spin two particle in the spectrum.

¹Orbifolds are examples of such singularities, but in [1] they are in fact resolved by a finite expectation value of a modulus – the B field [5]. We will be interested below in situations where this v.e.v. is zero or at least very small.

²A name due to [6].

- (5) The theory appears to have well defined off-shell Green functions, unlike (closed) critical string theory, where it is believed that only on-shell observables can be studied.

Note that while properties (1) and (2) are reminiscent of critical string theory, properties (3), (4) and (5) are different in the two cases.

The main purpose of these lectures is to describe in more detail some of the above properties and the techniques that were used to study them. Most of these results were obtained by using holography, and this is the approach that will be followed here. In particular, We will not describe an alternative approach to LST based on a discrete light-cone quantization (DLCQ) of the theory, which utilizes a certain $1+1$ dimensional sigma model [7, 8, 9]. For a review of that approach and LST in general as of mid-1999, see [10].

There are several reasons why I think LST is of some interest. Among them:

- (1) In most (compactified) supersymmetric string theories one finds moduli spaces of vacua. For generic values of the moduli the perturbative description is non-singular, but one can often tune the moduli so that a singularity appears somewhere on the compact manifold. The dynamics near the singularity is described by LST. Thus LST is part of the dynamics of rather conventional looking string vacua at special points in the moduli space. Furthermore, when supersymmetry is broken, it is possible that the theory is dynamically driven to such singular points in moduli space.
- (2) LST is relevant for the study of strongly coupled gauge theories, which can be realized on $NS5$ -branes wrapped around Riemann surfaces or D-branes stretched between fivebranes (see [2] for a review). There are also applications to matrix theory [11], which in fact provided some of the original motivation for the construction of this theory [12, 13].
- (3) It was proposed that LST might be phenomenologically relevant for brane world scenarios with a relatively low string scale [14].

More generally, LST appears to be a structure that is intermediate in complexity between local QFT and critical string theory. It has the non-locality and Hagedorn spectrum characteristic of critical string theory, but not the complications associated with gravity. A better understanding of its structure might shed light on string theory, strongly coupled gauge theory (QCD strings), holography and other matters.

The plan of these lectures is as follows. We start in section 2 by describing the limit in which the dynamics of $NS5$ -branes decouples from bulk physics. In section 3 we discuss the holographic description of this limit and some of the properties of LST mentioned above. In particular, we exhibit some observables and physical states in the theory.

In section 4 we discuss the high energy thermodynamics of LST. We show that the spectrum has a Hagedorn growth and compute the Hagedorn temperature and the first subleading term in the entropy which shows that the thermodynamics is unstable. In section 5 we introduce and study a class of vacua of LST which can be analyzed in a controlled weak coupling expansion.

Section 6 contains some comments on aspects of LST that we cannot treat in detail due to lack of time, including singularities of Calabi-Yau manifolds which give rise to $d < 6$ dimensional vacua of LST and models with reduced supersymmetry, D-branes in the vicinity of $NS5$ -branes, and instabilities in LST. In section 7 we discuss some open problems.

2 The decoupling limit of flat $NS5$ -branes

Consider a vacuum of type II string theory which contains N parallel $NS5$ -branes³, which are extended in the directions (x^1, \dots, x^5) and are localized in (x^6, \dots, x^9) . We will initially take the fivebranes to be at the same point and will later examine the deformations that separate them in the directions $(6, 7, 8, 9)$.

The presence of the fivebranes breaks the Lorenz symmetry:

$$SO(9, 1) \rightarrow SO(5, 1) \times SO(4). \quad (2.1)$$

From the fivebrane worldvolume point of view, $SO(5, 1)$ is the Lorenz symmetry, while $SO(4)$ is an internal R -symmetry. The fivebranes also break half of the supersymmetry, reducing the number of unbroken supercharges from thirty two to sixteen. In terms of six dimensional supersymmetry along the fivebranes, IIA fivebranes preserve a chiral $(2, 0)$ supersymmetry⁴, while IIB fivebranes preserve $(1, 1)$ supersymmetry.

Since $NS5$ -branes are dynamical objects, like D-branes, one expects to find a rich spectrum of excitations on the branes. To decouple the dynamics

³Neveu-Schwarz fivebranes are magnetically charged under the Neveu-Schwarz $B_{\mu\nu}$ field. See *e.g.* [15] for a review of some of their properties.

⁴I.e. two complex supercharges in the $\mathbf{4}$ of $Spin(5, 1)$.

on the fivebranes from the bulk, consider the limit

$$g_s \rightarrow 0; \quad \frac{E}{m_s} = \text{fixed}. \quad (2.2)$$

Processes in which modes that live on the fivebranes are emitted into the bulk as closed strings are suppressed in this limit, since the corresponding amplitudes are proportional to g_s and thus go to zero. At the same time, the dynamics on the $NS5$ -branes does not become free in this limit. One way to see this is to consider the low energy limit of the resulting theory and to show that it is not free.

Consider first the low energy limit of N $NS5$ -branes in type IIB string theory. S-duality relates this to N $D5$ -branes; thus the low energy theory is a six dimensional gauge theory with $(1, 1)$ supersymmetry and gauge group $U(N)$. The gauge coupling of the theory on the $D5$ -branes is

$$\frac{1}{g_D^2} = \frac{m_s^2}{g_s}. \quad (2.3)$$

Using the transformation of g_s and m_s under S-duality one finds that the gauge coupling on the $NS5$ -branes is

$$\frac{1}{g_N^2} = m_s^2. \quad (2.4)$$

Thus in the limit (2.2) the gauge coupling remains fixed. Since the gauge theory in question is non-renormalizable, the gauge coupling g_N in fact changes with the scale, approaching zero at long distances and growing at short distances. At energies of order m_s the gauge theory description breaks down and more data needs to be supplied to define the theory. As we will see, there are in fact additional degrees of freedom in the theory at (roughly) that scale, and the full density of states is much larger than that in any local QFT. At any rate, since the dynamics at scales $E \simeq m_s$ is not free, the full theory must be interacting.

Note that the above arguments are only valid for $N > 1$ fivebranes. The low energy theory on a single $NS5$ -brane is free⁵. Indeed, we will see later that LST is interacting only for $N > 1$.

⁵In the IIA case it contains a self-dual $B_{\mu\nu}$ field, five massless scalars and fermions related to them by $(2, 0)$ supersymmetry. In the IIB theory one finds a gauge field, four scalars and fermions, related by $(1, 1)$ supersymmetry.

The infrared dynamics of N IIA $NS5$ -branes is more involved. One finds in this case a non-trivial IR fixed point with $(2, 0)$ superconformal symmetry [16]. To see that something special is happening in the IR imagine separating the fivebranes in the $(6, 7, 8, 9)$ directions. In the IIB theory, one then finds massive states corresponding to D-strings stretched between the fivebranes; their masses go to zero as the fivebranes approach each other. The resulting massless states are the off-diagonal $U(N)$ gauge bosons on the fivebranes.

The analogous process for IIA involves $D2$ -branes stretched between the fivebranes. The ends of the $D2$ -branes are strings bound to the fivebranes. Their tension goes to zero when the fivebranes coincide [17]. These tensionless strings signal the interacting nature of the low energy limit of the IIA fivebrane theory – the $(2, 0)$ superconformal field theory.

Thus, we conclude that the limit (2.2) corresponds to an interacting theory on the $NS5$ -branes decoupled from the bulk. What sort of theory is it? Already at the level of the present discussion there are a few hints of non-local/stringy behavior. Let us mention two:

- (1) T-duality: Compactify some or all of the dimensions $(1, 2, 3, 4, 5)$ on circles. $NS5$ -branes are known to transform to themselves under T-duality along their worldvolume. Since the limit (2.2) commutes with T-duality, inversion of the radius of a single circle ($R \rightarrow 1/m_s^2 R$) exchanges the IIA and IIB LST's, while inversion of an even number of radii is a symmetry of the theory.
- (2) The theory contains strings with tension $T = 1/2\pi\alpha'$, which can be interpreted as fundamental strings bound to the fivebranes. In the IIB case⁶, these strings can be constructed in the low energy gauge theory as instanton solutions, which are extended (say) in $(0, 1)$ and localized in $(2, 3, 4, 5)$. The tension of these strings is proportional to the instanton action, $1/g_N^2$, which using (2.4) is indeed tension of a fundamental string. Of course, this construction gives rise to long strings, and it is not clear what are the properties of short strings which actually govern the dynamics, but it suggests that LST is a theory of strings. Later we will see further evidence that supports this.

It is instructive to compare the decoupling limit (2.2) with the limits studied in D-brane physics. Usually, to decouple the physics of D-branes from the

⁶A similar construction can be performed in the IIA case.

bulk one considers the low energy limit

$$\frac{E}{m_s} \rightarrow 0; \quad g_s = \text{fixed}, \quad (2.5)$$

and the decoupling from the bulk is the standard low energy decoupling of QFT from gravity. In contrast, the limit (2.2) for D-branes gives rise in general to a free theory on the branes, since g_s determines both the open and the closed string couplings.

A limit for N D-branes which is more analogous to (2.2) is

$$N \rightarrow \infty; \quad g_s \rightarrow 0; \quad \lambda = g_s N = \text{fixed}; \quad \frac{E}{m_s} = \text{fixed}. \quad (2.6)$$

The open string coupling λ is fixed; hence the theory on the D-branes remains interacting. Since $g_s \rightarrow 0$, the closed string sector decouples, despite the fact that a low energy limit has not been taken. The resulting theory is an open string theory without closed strings; it has some things in common with LST although there are differences as well.

3 A holographically dual description of LST

The construction described in the previous section is useful for establishing the existence of LST, but it does not provide efficient techniques for studying the theory. To proceed, we will use a holographically dual description proposed in [18] (see also [19, 20]). This duality is a generalization of the AdS/CFT correspondence [3]; it postulates that LST is equivalent to ten dimensional string theory in the background of the fivebranes, in the limit (2.2). In this section we will describe the fivebrane geometry and will briefly discuss the duality of [18].

The metric, dilaton and NS B -field around N $NS5$ -branes in type II string theory are [4]:

$$\begin{aligned} ds^2 &= dx_\mu dx^\mu + (1 + \frac{N\alpha'}{r^2}) dx^i dx^i, \\ e^{2\Phi} &= g_s^2 (1 + \frac{N\alpha'}{r^2}), \\ H_{ijk} &= -\epsilon_{ijkl} \partial^l \Phi, \end{aligned} \quad (3.1)$$

where $\mu = 0, 1, 2, \dots, 5$ are worldvolume coordinates and $i, j, k, l = 6, 7, 8, 9$ are transverse ones. We parameterize the space transverse to the branes by

spherical coordinates,

$$dx^i dx^i = dr^2 + r^2 d\Omega_3^2. \quad (3.2)$$

To take the limit (2.2) one must send $r \rightarrow 0$ at the same rate as g_s . Defining $r = g_s \exp \sigma$ we have in this limit

$$\begin{aligned} ds^2 &= dx_\mu dx^\mu + N\alpha'(d\sigma^2 + d\Omega_3^2), \\ \Phi &= -\sigma, \end{aligned} \quad (3.3)$$

and we suppress the B -field (3.1). String propagation in this geometry corresponds to an “exact conformal field theory” [4]:

$$\mathbb{R}^{5,1} \times \mathbb{R}_\phi \times SU(2)_N. \quad (3.4)$$

$\mathbb{R}^{5,1}$ is the worldvolume of the fivebranes. \mathbb{R}_ϕ is the real line labeled by $\phi = \sqrt{N\alpha'}\sigma$. The dilaton goes like (3.3):

$$\Phi = -\frac{Q}{2}\phi; \quad Q = \frac{2}{\sqrt{N\alpha'}}. \quad (3.5)$$

The last factor in (3.4) describes the angular three-sphere in (3.3). The B -field (3.1) is precisely such that the CFT on the three-sphere, whose radius is

$$R_{\text{sphere}} = \sqrt{N\alpha'}, \quad (3.6)$$

is described by a level N WZW model. We see that the number of fivebranes N determines the slope of the linear dilaton, Q , and the level of $SU(2)$ current algebra. More precisely, since (3.4) is a background for the superstring, the worldsheet theory contains, in addition to the bosonic coordinates, ten free fermions: ψ^μ , $\mu = 0, 1, 2, \dots, 5$, the superpartners of x^μ ; ψ^i , $i = 3, +, -$, the superpartners of the $SU(2)$ currents J^i ; and ψ^ϕ , the superpartner of ϕ . The total level N of the $SU(2)$ current algebra receives a contribution of $N - 2$ from the worldsheet bosons, and 2 from the fermions ψ^i , which transform in the adjoint of the total $SU(2)$ current algebra. The total central charge of the worldsheet theory (3.4) is

$$\left(6 + \frac{1}{2} \times 6\right) + \left(1 + \frac{6}{N} + \frac{1}{2}\right) + \left(\frac{3(N-2)}{N} + 3 \times \frac{1}{2}\right) = 15, \quad (3.7)$$

which is the correct value for the superstring.

The background (3.4) is thus expected to be holographically dual to the LST on the fivebranes. We next discuss some features of this duality. First note that while the string coupling (3.5) vanishes far from the fivebranes (*i.e.* as $\phi \rightarrow \infty$), it diverges as one approaches the branes ($\phi \rightarrow -\infty$, or $r \rightarrow 0$ in (3.1)). The $NS5$ -branes have the remarkable property that quantum effects near the branes cannot be turned off no matter how small the string coupling is far from the branes [4]. This makes it clear that LST is not a free theory⁷, as argued above, but it raises the question whether one can analyze the physics of the string background (3.4), (3.5) perturbatively. We will return to this question below.

As is familiar from the AdS/CFT correspondence, on-shell observables in the “bulk” theory – string theory on (3.4) – correspond to off-shell observables in the “boundary” theory – the LST corresponding to N $NS5$ -branes. More precisely, off-shell observables in LST correspond to *non-normalizable* observables in string theory on (3.4), whose wavefunctions are supported near the “boundary” at $\phi \rightarrow \infty$. This can be understood as follows (in analogy with the *AdS* case).

Consider (say) a scalar field Ψ on the manifold (3.4), corresponding to one of the modes of the string. As $\phi \rightarrow \infty$, the field behaves as (assuming for simplicity a profile constant on the angular S^3):

$$\Psi(\phi, x^\mu) \sim \sum_k C_k e^{\lambda_k \phi} e^{i k_\mu x^\mu} \quad (3.8)$$

where

$$\lambda_k^2 = k_\mu k^\mu + C. \quad (3.9)$$

C is a constant which depends on the mass of the scalar field. Choosing the positive root of (3.9), we see that the mode (3.8) is non-normalizable and thus the coefficients C_k do not fluctuate – they are not integrated over in the process of integrating over all field configurations in the path integral [21]. Thus, we can think of the C_k as fixed sources. The string partition sum with the fixed boundary conditions (3.8) as $\phi \rightarrow \infty$, $Z_{\text{bulk}}(C_k)$, can be

⁷For $N \geq 2$ fivebranes. Note that for $N = 1$, the bosonic $SU(2)$ current algebra has formally a negative level, $N - 2 = -1$, and the construction breaks down. This is usually taken to mean that a single fivebrane does not have a throat region (3.4) associated with it, and the dynamics on it becomes trivial in the limit (2.5).

interpreted as the generating functional of off-shell Green functions in the six dimensional LST via:

$$Z_{\text{bulk}}(C_k) = \langle \exp \left(- \sum_k C_k \Theta(k) \right) \rangle_{LST}, \quad (3.10)$$

where $\Theta(k_\mu)$ is the off-shell observable which couples to the source C_k . Qualitatively, (3.10) is natural because modes that are non-normalizable in the “near-horizon” geometry (3.4) are nothing but bulk modes in the full geometry (3.1); they are supported at finite r . Thus, they are not part of the LST but rather are fixed background sources (in the limit (2.2)), which couple to the brane modes via couplings like (3.10).

Similarly, *normalizable modes* in the geometry (3.4) correspond to *states* in LST, since in the full geometry (3.1) they correspond to modes localized on the fivebranes (*i.e.* at $r \rightarrow 0$). To illustrate all this, we next give an example each of off-shell observables and states in LST, as described in the holographically dual picture.

3.1 Example 1: Chiral operators in LST

As discussed above, the low energy limit of IIB LST is a $U(N)$ gauge theory with $(1,1)$ supersymmetry. This theory contains four scalar fields in the adjoint of $SU(N)$, X^i , $i = 6, 7, 8, 9$, which parameterize the locations of the N fivebranes in $(6, 7, 8, 9)$. The gauge invariant off-shell operators

$$\text{Tr} X^{i_1} X^{i_2} \cdots X^{i_n}; \quad n = 2, 3, 4, \dots, N, \quad (3.11)$$

where we only take the completely symmetric and traceless combination in (i_1, \dots, i_n) , are lowest components of short multiplets of supersymmetry. Writing the $SO(4)$ symmetry in (2.1) as

$$SO(4) \simeq SU(2)_L \times SU(2)_R, \quad (3.12)$$

the operators (3.11) transform in the spin $(\frac{n}{2}, \frac{n}{2})$ representations. In string theory on (3.4) these chiral operators are described as follows. The $SU(2)_L \times SU(2)_R$ symmetry on (3.12) corresponds to the left and right moving $SU(2)$ symmetries in the $SU(2)_N$ WZW model in (3.4). Physical primaries of this symmetry are $V_{j; m, \bar{m}}$ with the same spin ($2j = 0, 1, 2, \dots, N-2$) under both $SU(2)$ ’s. (m, \bar{m}) are the eigenvalues of (J_3, \bar{J}_3) .

The lowest lying observables have the form (in the -1 picture)

$$\xi_{\alpha\beta} \psi^\alpha \bar{\psi}^\beta e^{\beta\phi} e^{ik_\mu x^\mu} V_j, \quad (3.13)$$

where $\alpha, \beta = 0, 1, 2, \dots, 9$ and $\xi_{\alpha\beta}$ is a polarization tensor satisfying the usual physical state conditions. One can show that (3.11) correspond to⁸

$$\text{Tr} X^{i_1} X^{i_2} \dots X^{i_n} \leftrightarrow (\psi \bar{\psi} V_j)_{j+1} e^{\frac{2j}{N\alpha'} \phi}, \quad j+1 = \frac{n}{2} \quad (3.14)$$

On the right-hand side of (3.14), ψ stands for the three fermions associated with the $SU(2)$ WZW and the brackets mean that ψ , which has spin 1 under $SU(2)_L$, is coupled with V_j into a spin $j+1$ combination (and similarly for the right movers). Thus, the non-normalizable operators (3.14) transform under $SU(2)_L \times SU(2)_R$ as

$$(j+1, j+1); \quad 2j = 0, 1, 2, \dots, N-2, \quad (3.15)$$

in exact agreement with what was found for (3.11) above. Applying the spacetime supercharges gives the other members of the supermultiplets. Thus, the sets of short representations of supersymmetry in LST and in string theory on (3.4) agree.

3.2 Example 2: Normalizable states

A large set of normalizable states is obtained by considering vertex operators of the form

$$V(\phi) \sim e^{(-\frac{Q}{2} + i\lambda)\phi} \quad (3.16)$$

on \mathbb{R}_ϕ . Recall that the vertex operators are related to the wavefunctions (3.8) by a factor of g_s , which here is a function of ϕ (3.5). Therefore, (3.16) actually corresponds to a wavefunction

$$\Psi(\phi) \sim e^{i\lambda\phi}, \quad (3.17)$$

which is (δ -function) normalizable, and thus gives rise to states in LST. Since λ is arbitrary, there is in fact a continuum of such states. To compute their masses, consider the states (3.13) as an example. The mass shell condition reads:

$$k_\mu k^\mu - \beta(\beta + Q) = 0. \quad (3.18)$$

⁸We set k_μ to zero for simplicity.

Plugging in $\beta = -\frac{Q}{2} + i\lambda$, we find

$$M^2 = \frac{1}{N\alpha'} + \lambda^2. \quad (3.19)$$

Thus, we find a continuum above the gap m_s/\sqrt{N} . The gap is given by a natural scale in LST; looking back at (2.4), we see that it is the 't Hooft coupling of the low energy super Yang Mills theory (for IIB fivebranes).

3.3 The strong coupling problem

As we have seen before, the background (3.4) has the property that the string coupling depends on ϕ ; it goes to zero as $\phi \rightarrow \infty$ and diverges as $\phi \rightarrow -\infty$. In this subsection we would like to discuss the physical origin of this behavior and its implications. The strong coupling region $\phi \rightarrow -\infty$ corresponds to the vicinity of the brane ($r \rightarrow 0$). This is the low energy region in the theory on the branes [19].

The low energy behavior of LST is different for IIA and IIB fivebranes. In the IIB case, the low energy limit is a six dimensional $U(N)$ gauge theory, which is weakly coupled in the IR. Thus, in the limit $\phi \rightarrow -\infty$ of the near-horizon geometry, which should be dual to the infrared limit on the brane [3], string theory on (3.4) should reproduce the weakly coupled gauge theory on the branes. Since one does not expect to find two different weakly coupled description of the same physics, the “bulk” description should either be strongly coupled, or exhibit large curvatures (or both). Since in our case the curvature of (3.4) is small, it is natural to find that the string coupling is growing in the infrared region.

In the IIA case the infrared limit of LST is somewhat different. As discussed earlier, one finds in this case a non-trivial superconformal field theory with chiral $(2, 0)$ supersymmetry, the $(2, 0)$ theory. Thus, it is not obvious that one should run into any strong coupling problems in the dual description.

To see what is going on, recall that type IIA string theory can be thought of as an eleven dimensional theory, M-theory, compactified on a circle of radius R_{11} , which is related to the eleven dimensional Planck scale l_{11} , and the string scale m_s and coupling g_s via

$$m_s R_{11} = \ell_{11}^3 m_s^3 = g_s. \quad (3.20)$$

The eleven dimensional theory contains membranes and fivebranes (the $M2$ and $M5$ -branes), which preserve half of the supersymmetry; their tensions

are (up to numerical constants) $1/l_{11}^3$ and $1/l_{11}^6$, respectively. The IIA $NS5$ -branes are M_5 -branes located at points on the circle. Thus, to study them using holography we should construct the background around N coincident $M5$ -branes. Taking the limit (2.2), which corresponds to $R_{11}, l_{11} \rightarrow 0$ with m_s fixed, one finds the eleven dimensional metric

$$ds^2 = H^{-\frac{1}{3}} [dx_\mu dx^\mu + H(dx_{11}^2 + dr^2 + r^2 d\Omega_3^2)], \quad (3.21)$$

where

$$H = \sum_{n=-\infty}^{\infty} \frac{N l_{11}^3}{[r^2 + (x_{11} - 2\pi n R_{11})^2]^{\frac{3}{2}}}. \quad (3.22)$$

x_{11} is a coordinate on the circle; it is periodic with period $2\pi R_{11}$. In the limit $r \rightarrow \infty$, the background (3.21) goes over to (3.4). The radius of the x_{11} circle goes to zero and one finds the linear dilaton behavior discussed above. As $r, x_{11} \rightarrow 0$ only one term in the sum over n in (3.22) (say $n = 0$) contributes, and the metric reduces to the near-horizon background of N coincident $M5$ -branes in eleven dimensions. This background, $AdS_7 \times S^4$, is known to be dual to the $(2, 0)$ superconformal field theory via AdS/CFT [3]. If N is large, it can be studied using eleven dimensional supergravity; otherwise one needs the full M-theory, which is not understood for these backgrounds.

Thus, we see that the growth of the coupling and associated breakdown of string perturbation theory as $\phi \rightarrow -\infty$ in the background (3.4) have slightly different origins in the IIA and IIB cases. However, regardless of the origin of this problem, one can ask what is the dual description of LST good for in view of its existence? We have already seen two examples of applications of the formalism. Since off-shell observables correspond to non-normalizable wavefunctions supported in the region $\phi \rightarrow \infty$, we can classify the observables of LST by analyzing such wavefunctions; since the coupling is small at large ϕ , perturbative string theory is suitable for this. Also, any normalizable states that are supported in the weakly coupled asymptotic region, like those described in section 3.2, can be studied using the formalism.

Correlation functions of the observables discussed above are in general difficult to analyze. Since the string coupling goes to zero as $\phi \rightarrow \infty$, disturbances on the boundary have to propagate to finite ϕ in order to interact. Thus, to compute correlation functions in LST one needs information about the strong coupling region. E.g. for IIA fivebranes, one has to understand

M-theory in the background (3.21), (3.22) which seems difficult⁹.

There are actually some situations in which the strong coupling problem can be avoided. In the next section we describe an example of such a situation, which is in fact of independent interest, the high energy density thermodynamics of LST.

4 High energy thermodynamics of LST

At very high energy density one expects the thermodynamics of fivebranes to be dominated by black brane states. Thus, in this section we will analyze the thermodynamics of near-extremal fivebranes and deduce from it the entropy-energy relation. We will find that the density of states has the Hagedorn behavior

$$\rho(E) \sim E^\alpha e^{\beta_H E} \left[1 + O\left(\frac{1}{E}\right) \right]. \quad (4.1)$$

One of our main purposes is to compute β_H and α . This section is based on [23]. For some additional recent work on LST thermodynamics, see [24, 25, 26].

4.1 Thermodynamics of near-extremal fivebranes

The supergravity solution for N coincident near-extremal $NS5$ -branes in the string frame is [27]:

$$ds^2 = - \left(1 - \frac{r_0^2}{r^2} \right) dt^2 + \left(1 + \frac{N\alpha'}{r^2} \right) \left(\frac{dr^2}{1 - \frac{r_0^2}{r^2}} + r^2 d\Omega_3^2 \right) + dy_5^2, \quad (4.2)$$

$$e^{2\Phi} = g_s^2 \left(1 + \frac{N\alpha'}{r^2} \right). \quad (4.3)$$

$r = r_0$ is the location of the horizon, dy_5^2 denotes the flat metric along the fivebranes, and $d\Omega_3^2$ is the metric on a unit three-sphere, as before. The solution also involves a non-zero NS $B_{\mu\nu}$ field which we suppress. The configuration (4.2), (4.3) has energy per unit volume

$$\frac{E}{V_5} = \frac{1}{(2\pi)^5 \alpha'^3} \left(\frac{N}{g_s^2} + \mu \right), \quad (4.4)$$

⁹For large N and energies much lower than m_s one can use classical eleven dimensional supergravity to compute correlation functions. See [22] for details.

where

$$\mu = \frac{r_0^2}{g_s^2 \alpha'}. \quad (4.5)$$

The first term in (4.4) is the tension of extremal $NS5$ -branes and can be ignored for the thermodynamic considerations below – it is a ground state energy. μ measures the energy density above extremality (in string units) and g_s is the asymptotic string coupling, which goes to zero in the decoupling limit.

The near-horizon geometry is obtained by sending $r_0, g_s \rightarrow 0$, keeping the energy density μ fixed. Changing coordinates to $r = r_0 \cosh \sigma$ and Wick rotating $t \rightarrow it$ to study the thermodynamics, one finds

$$ds^2 = \tanh^2 \sigma dt^2 + N \alpha' d\sigma^2 + N \alpha' d\Omega_3^2 + dy_5^2, \quad (4.6)$$

$$e^{2\Phi} = \frac{N}{\mu \cosh^2 \sigma}. \quad (4.7)$$

This background corresponds to the worldsheet CFT

$$H_3^+ / U(1) \times SU(2)_N \times \mathbb{R}^5, \quad (4.8)$$

where

$$H_3^+ = \frac{SL(2, C)_N}{SU(2)_N} \quad (4.9)$$

is the Euclidean AdS_3 CFT which plays an important role in the AdS-CFT correspondence; the coset $H_3^+ / U(1)$, parametrized by (σ, t) in (4.6), is a semi-infinite cigar [28]. The background (4.8) describes the high energy density thermodynamics of fivebranes; it should be compared to (3.4), which is dual to the zero temperature theory.

The absence of a conical singularity at the tip ($\sigma = 0$ in (4.6)) requires the circumference of the cigar to be

$$\beta_H = 2\pi\sqrt{N\alpha'}. \quad (4.10)$$

Thus, Euclidean time lives on a circle of radius $\sqrt{N\alpha'}$, and the temperature of the system is $T_H = 1/\beta_H$. In particular, the temperature is independent of the energy density μ , which determines the value of the string coupling at the tip of the cigar (4.7).

The fact that the temperature is independent of the energy means that the entropy is proportional to the energy (since $\beta = \frac{\partial S}{\partial E}$). Therefore, the free energy is expected to vanish¹⁰,

$$-\beta\mathcal{F} = S - \beta E = 0. \quad (4.11)$$

In general in string theory the free energy is related to the string partition sum via

$$-\beta\mathcal{F} \equiv \log Z(\beta) = Z_{\text{string}}, \quad (4.12)$$

where Z_{string} is the single string partition sum, given by a sum over connected Riemann surfaces [30]. The string path integral should be performed over geometries in which Euclidean time is compactified on a circle of radius $R = \beta/2\pi$ (asymptotically). As mentioned above, for high energies one expects the thermodynamics to be dominated by the black brane geometry (4.2), (4.6) and thus the free energy is proportional to the partition sum of string theory in the background (4.8).

The string partition sum Z_{string} can be expanded as follows:

$$Z_{\text{string}} = e^{-2\Phi_0} Z_0 + Z_1 + e^{2\Phi_0} Z_2 + \dots, \quad (4.13)$$

where $\exp(\Phi_0)$ is the effective string coupling in the geometry (4.6) and Z_h the genus h partition sum in the background (4.8). Although the string coupling varies along the cigar (see (4.7)), it is bounded from above by its value at the tip,

$$e^{2\Phi_0} = \frac{N}{\mu}. \quad (4.14)$$

Therefore, it is natural to associate (4.14) with the effective coupling in (4.13). We see that the string coupling expansion in the background (4.8) provides an asymptotic expansion of the free energy in powers of $1/\mu$.

The leading term in the free energy (4.12), (4.13) goes like

$$-\beta\mathcal{F} = \frac{\mu}{N} Z_0 \quad (4.15)$$

and corresponds to a free energy that goes like the energy (Z_0 is proportional to the volume of the fivebrane). This term is expected to vanish (see (4.11)),

¹⁰See [29] for a related discussion in the low energy gravity approximation.

and therefore we conclude that the spherical partition sum in the background (4.8) should vanish. The fact that this is indeed the case follows from the results of [31]; we will not discuss it further here (see [23]).

To compute $1/\mu$ corrections to the free energy we have to examine string loop effects in the background (4.8). We next turn to the one loop correction Z_1 (see (4.13)).

4.2 The leading $1/\mu$ correction to classical thermodynamics

As discussed above, one expects the entropy-energy relation to take the form (4.1)

$$S(E) = \beta_H E + \alpha \log \frac{E}{\Lambda} + O\left(\frac{1}{E}\right), \quad (4.16)$$

where Λ is a dimensionful constant (a UV cutoff) which we will not keep track of below. Consider the canonical partition sum

$$Z(\beta) = \int_0^\infty dE \rho(E) e^{-\beta E}. \quad (4.17)$$

Near the Hagedorn temperature one might expect $Z(\beta)$ to be dominated by the contributions of high energy states;¹¹ if this is the case, one can replace $\rho(E)$ by (4.1) and find,

$$Z(\beta) \simeq \int dE E^\alpha e^{(\beta_H - \beta)E} \simeq (\beta - \beta_H)^{-\alpha-1}. \quad (4.18)$$

The free energy (4.12) is thus given by

$$\beta \mathcal{F} \simeq (\alpha + 1) \log(\beta - \beta_H). \quad (4.19)$$

The energy computed in the canonical ensemble is

$$E = \frac{\partial(\beta \mathcal{F})}{\partial \beta} \simeq \frac{\alpha + 1}{\beta - \beta_H}; \quad (4.20)$$

thus the free energy (4.19) can be written as

$$-\beta \mathcal{F} \simeq (\alpha + 1) \log E. \quad (4.21)$$

¹¹We will see that this assumption is valid slightly *above* the Hagedorn temperature, but is *not* valid slightly below it.

Comparing to the expansion (4.12) – (4.14) we see that the leading term in the free energy arises from the torus (one loop) diagram in the background (4.8), since it scales as μ^0 , like Z_1 in (4.13).

The torus partition sum in the background (4.8) is in fact divergent, since it is proportional to the infinite volume of the cigar, associated with the region far from the tip, $\phi \rightarrow \infty$. As is standard in other closely related contexts, we will regulate this divergence by requiring that

$$\phi \leq \phi_{UV}. \quad (4.22)$$

In the fivebrane theory, this can be thought of as introducing a UV cutoff. This makes the partition sum finite, but the bulk of the amplitude still comes from the region far from the tip of the cigar. For the purpose of computing this “bulk contribution” one can replace the cigar by a long cylinder with ϕ bounded on one side by the UV cutoff (4.22) and on the other by the location of the tip of the cigar. Combining (3.5) and (4.14) we find that

$$\frac{1}{Q} \log \frac{\mu}{N} \leq \phi \leq \phi_{UV}. \quad (4.23)$$

Thus, the length of the cut-off cylinder is

$$L_\phi = \phi_{UV} - \frac{1}{Q} \log \frac{\mu}{N} = -\frac{1}{Q} \log E + \text{const.} \quad (4.24)$$

Since we are only interested in the energy dependence, we suppress in (4.24) a large energy independent contribution. Any contributions to the torus partition sum from the region near the tip of the cigar can also be lumped into this constant. Note the minus sign in front of $\log E$ in (4.24). The length L_ϕ is of course positive; the minus sign simply means that L_ϕ decreases as E grows.

To recapitulate, for the purpose of calculating the bulk contribution to the torus partition sum, we can replace the background (4.8) by

$$\mathbb{R}_\phi \times S^1 \times SU(2)_N \times \mathbb{R}_5. \quad (4.25)$$

The linear dilaton direction is regulated as in (4.23). The circumference of the S^1 is β_H (4.10).

The background (4.25) is easy to analyze since it is very similar to that describing flat space at finite temperature (see *e.g.* [32, 33, 34]). The bosonic fields on the worldsheet are seven free fields, one of which (Euclidean time) is

compact, and a level $N-2$ $SU(2)$ WZW model. The worldsheet fermions are free and decoupled from the bosons; their partition sum, and in particular the sum over spin structures, is the same as in the flat space analysis, which we briefly review next.

Collecting all the contributions to the thermal torus partition sum in the background (4.25) we find,¹²

$$\begin{aligned} Z_1 &= \frac{\beta V_5 L_\phi}{4} \int_F \frac{d^2\tau}{\tau_2} \left(\frac{1}{4\pi^2 \alpha' \tau_2} \right)^{7/2} \frac{1}{|\eta(\tau)|^{10}} Z_{N-2}(\tau) \times \\ &\quad \sum_{n,m \in \mathbb{Z}} \sum_{\mu,\nu=1}^4 \delta_\mu U_\mu(n,m) \delta_\nu U_\nu(n,m) \left(\frac{\vartheta_\mu(0,\tau)}{\eta(\tau)} \right)^4 \left(\frac{\vartheta_\nu(0,\bar{\tau})}{\eta(\bar{\tau})} \right)^4 e^{-S_\beta(n,m)}. \end{aligned} \quad (4.26)$$

The modular integral runs over the standard fundamental domain F . Z_{N-2} is the partition sum of level $N-2$ $SU(2)$ WZW¹³ (see for example [35]),

$$Z_{N-2}(\tau) = \sum_{m=0}^{N-2} \chi_m^{(N-2)}(q) \chi_m^{(N-2)}(\bar{q}) = \sum_{m=0}^{N-2} |\chi_m^{(N-2)}(q)|^2, \quad (4.27)$$

where $q = \exp(2\pi i\tau)$ and

$$\chi_m^{(N-2)}(q) = \frac{q^{\frac{(m+1)^2}{4N}}}{\eta(q)^3} \sum_{n \in \mathbb{Z}} [1 + m + 2nN] q^{n(1+m+Nn)}. \quad (4.28)$$

We note for future reference that Z_{N-2} is real and positive.

μ, ν denote the spin structure for left and right moving worldsheet fermions, respectively. $\delta_\mu = (\pm, -, +, -)$ are signs coming from the usual GSO projections for IIA and IIB superstrings at zero temperature; n, m are winding numbers of Euclidean time around the two non-contractible cycles of the torus. The soliton factor $S_\beta(n, m)$ is given by

$$S_\beta(n, m) = \frac{\beta^2}{4\pi\alpha' \tau_2} (m^2 + n^2 |\tau|^2 - 2\tau_1 mn). \quad (4.29)$$

$U_\mu(n, m)$ are additional signs that are associated with finite temperature. Their role is to implement the standard thermal boundary conditions, that

¹²We follow the conventions of [34], which should be consulted for additional details. We also drop the subscript H on β_H , and will reinstate it later.

¹³We choose the A series modular invariant; the D and E series modular invariants can also be studied and correspond to other vacua of LST [18].

spacetime bosons (fermions) are (anti-)periodic around the Euclidean time direction. One can show [34] that this requirement together with modular invariance leads to:

$$\begin{aligned} U_1(n, m) &= \frac{1}{2} (-1 + (-1)^n + (-1)^m + (-1)^{n+m}) \\ U_2(n, m) &= \frac{1}{2} (1 - (-1)^n + (-1)^m + (-1)^{n+m}) \\ U_3(n, m) &= \frac{1}{2} (1 + (-1)^n + (-1)^m - (-1)^{n+m}) \\ U_4(n, m) &= \frac{1}{2} (1 + (-1)^n - (-1)^m + (-1)^{n+m}). \end{aligned} \quad (4.30)$$

The terms with $\mu = 1$ in (4.26) vanish because of the presence of fermionic zero modes for the $(+, +)$ spin structure, or equivalently since $\vartheta_1(0, \tau) = 0$.

The torus partition sum (4.26) can be rewritten in a way that makes it manifest that the coefficient of $\beta V_5 L_\phi / 4$ is positive,

$$\begin{aligned} Z_1 &= \frac{\beta V_5 L_\phi}{4} \int_F \frac{d^2\tau}{\tau_2} \left(\frac{1}{4\pi^2 \alpha' \tau_2} \right)^{7/2} \frac{1}{|\eta(\tau)|^{18}} Z_{N-2}(\tau) \times \\ &\quad \sum_{n, m \in \mathbb{Z}} \left| \sum_{\mu=2}^4 U_\mu(n, m) \delta_\mu \vartheta_\mu^4(0, \tau) \right|^2 e^{-S_\beta(n, m)}. \end{aligned} \quad (4.31)$$

It is not difficult to check that the integral (4.31) is convergent at $\tau_2 \rightarrow \infty$, the only region where a divergence could occur.

To exhibit the interpretation of (4.31) as a sum over the free energies of physical string modes one can proceed as follows [30, 32, 33]. Using the modular invariance of the integrand and the covariance of (n, m) , one can extend the integral from the fundamental domain to the strip

$$S : \quad -\frac{1}{2} \leq \tau \leq \frac{1}{2}; \quad \tau_2 \geq 0, \quad (4.32)$$

while restricting to configurations with $n = 0$ in (4.31). This leads to

$$\begin{aligned} Z_1 &= \frac{\beta V_5 L_\phi}{4} \int_S \frac{d^2\tau}{\tau_2} \left(\frac{1}{4\pi^2 \alpha' \tau_2} \right)^{7/2} \frac{1}{|\eta(\tau)|^{18}} Z_{N-2}(\tau) \times \\ &\quad \sum_{m=-\infty}^{\infty} \left| \sum_{\mu=2}^4 U_\mu(0, m) \delta_\mu \vartheta_\mu^4(0, \tau) \right|^2 e^{-S_\beta(0, m)}. \end{aligned} \quad (4.33)$$

The integral over τ_1 projects on physical states (*i.e.* those with $L_0 = \bar{L}_0$), while τ_2 plays the role of a Schwinger parameter. Because of the Jacobi identity $\vartheta_2^4(0, \tau) - \vartheta_3^4(0, \tau) + \vartheta_4^4(0, \tau) = 0$, and the fact that $U_2(0, m) = (-)^m$, $U_3(0, m) = U_4(0, m) = 1$, the sum over m in (4.33) can be restricted to odd integers. It is not difficult to check in this representation too that the integral over τ_2 is convergent.

We are now ready to determine the parameter α in (4.16), (4.21). Using the relation (4.12) between the free energy \mathcal{F} and the string partition sum, as well as (4.21), we see that Z_1 should be proportional to $\log E$. This is indeed the case in (4.33) since the length L_ϕ goes like $-\log E$ (see (4.24)). Combining these relations we find that

$$\begin{aligned} \alpha + 1 &= -\frac{\beta V_5}{4Q} \int_S \frac{d^2\tau}{\tau_2} \left(\frac{1}{4\pi^2 \alpha' \tau_2} \right)^{7/2} \frac{1}{|\eta(\tau)|^{18}} Z_{N-2}(\tau) \times \\ &\quad \sum_{m=-\infty}^{\infty} \left| \sum_{\mu=2}^4 U_\mu(0, m) \delta_\mu \vartheta_\mu^4(0, \tau) \right|^2 e^{-S_\beta(0, m)}. \end{aligned} \quad (4.34)$$

We see that $\alpha + 1$ is negative.¹⁴ Physically, it is clear that it is counting the free energy of the perturbative string modes which live in the vicinity of the black brane. An interesting point which was mentioned in [36, 37] is that α is an extensive quantity – it is proportional to the volume of the fivebrane V_5 , in contrast, say, to the one particle free energy in critical string theory, where the analogous quantity is of order one.

The integral (4.34) appears in general to be rather formidable and we do not know whether it can be performed exactly. In the remainder of this section we will compute it in the limit $N \rightarrow \infty$, where the computation simplifies.

For large N the partition sum corresponding to the three-sphere, $Z_{N-2}(\tau)$, simplifies significantly. Indeed, for $N \gg 1$ (4.27) can be approximated as

$$Z_{N-2}(\tau) = \frac{1}{|\eta(q)|^6} \sum_{p=0}^{\infty} |q|^{\frac{(p+1)^2}{2N}} (p+1)^2. \quad (4.35)$$

¹⁴Of course, since the r.h.s. of (4.34) is proportional to V_5 which is assumed to be very large, we can neglect the $+1$ on the left-hand side.

Returning to the evaluation of α , (4.34), we have

$$\begin{aligned} \alpha + 1 &= -\frac{\beta V_5}{4Q} \left(\frac{1}{4\pi^2 \alpha'} \right)^{7/2} \int_S \frac{d^2 \tau}{\tau_2^{9/2}} \left| \frac{1}{\eta(\tau)} \right|^{24} \times \\ &\quad \sum_{m \in 2\mathbb{Z}+1} \sum_{p=0}^{\infty} e^{-\frac{(p+1)^2 \tau_2}{2N}} (p+1)^2 e^{-\frac{\beta^2 m^2}{4\pi \alpha' \tau_2}} |\vartheta_2^4 + \vartheta_3^4 - \vartheta_4^4|^2 (0, \tau). \end{aligned} \quad (4.36)$$

At this point it is useful to recall that the inverse temperature β in (4.36) is in fact the Hagedorn temperature of LST, (4.10). In the large N limit, $\beta_H \sim \sqrt{N}$ becomes large (or, equivalently, the Hagedorn temperature is small in string units) and the exponential term in (4.36) suppresses the amplitude, unless τ_2 is large as well (of order N). Therefore, the τ integral in (4.36) is dominated by the large τ_2 region, which corresponds to the free energy of the supergravity modes. To compute the integral we recall the asymptotic forms of the ϑ and η functions at large τ_2 (see *e.g.* [38])

$$\begin{aligned} \vartheta_2(0, \tau) &= \sum_{n=-\infty}^{\infty} q^{\frac{1}{2}(n-\frac{1}{2})^2} = 2q^{\frac{1}{8}}(1 + q + \dots) \\ \vartheta_3(0, \tau) &= \sum_{n=-\infty}^{\infty} q^{\frac{1}{2}n^2} = 1 + 2q^{\frac{1}{2}} + \dots \\ \vartheta_4(0, \tau) &= \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{1}{2}n^2} = 1 - 2q^{\frac{1}{2}} + \dots \\ \eta(\tau) &= q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n) = q^{\frac{1}{24}} + \dots. \end{aligned} \quad (4.37)$$

Plugging in (4.36) and using the definition of the modified Bessel function

$$K_{\nu}(z) = \frac{1}{2} \left(\frac{2}{z} \right)^{\nu} \int_0^{\infty} t^{\nu-1} e^{-\frac{z^2}{4t} - t} dt, \quad (4.38)$$

we find

$$\begin{aligned} \alpha + 1 &= -\frac{8V_5}{\pi^6 (N\alpha')^{5/2}} \sum_{k,p=0}^{\infty} \left(\frac{2\pi(2k+1)^2}{(p+1)^2} \right)^{-7/4} (p+1)^2 \times \\ &\quad K_{-\frac{7}{2}}(\sqrt{2\pi}(p+1)(2k+1)) \simeq -4.08 \cdot 10^{-4} V_5 (N\alpha')^{-5/2} \equiv -a_1 V_5. \end{aligned} \quad (4.39)$$

Note that, as expected, α is negative. Of course, as is clear from (4.36), we can write $\alpha+1$ as $-a_1 V_5$ with a_1 a positive constant for all N , but in general a_1 receives contributions from massive string modes and is thus given by a complicated modular integral. The large N behavior of a_1 is simpler and is given by (4.39). It should be emphasized that, as mentioned above, the large N result (4.39) comes entirely from the thermodynamics of the supergravity modes in the near-extremal fivebrane background (4.6), (4.7), and thus could have been obtained by a supergravity calculation.

The fact that α goes like $N^{-5/2}$ for large N was found in a different way in [36], by analyzing the deformation of the classical solution (4.6) at one string loop. The analysis described here determines the coefficient of $N^{-5/2}$, and in particular its sign, which is important for the thermodynamics.

In the discussion above, the fivebrane was assumed to be effectively non-compact. It is interesting to study the thermodynamics of fivebranes wrapped around compact manifolds, and in particular the dependence of α on the size and shape of the manifold. As an example of the sort of dependence one can expect, consider compactifying the fivebrane on $(S^1)^5$ where all five circles have the same radius R . It is sufficient to consider the case $R \geq \sqrt{\alpha'}$ since smaller radii give rise to the same physics due to T-duality.

As is standard in string theory, the effect of this is to replace the contribution of the non-compact zero modes on R^5 by the momentum and winding sum on $(S^1)^5$:

$$\frac{V_5}{(4\pi^2\alpha'\tau_2)^{5/2}} \longrightarrow \left(\sum_{l,p \in Z} q^{\frac{\alpha'}{4}(\frac{l}{R} + \frac{pR}{\alpha'})^2} \bar{q}^{\frac{\alpha'}{4}(\frac{l}{R} - \frac{pR}{\alpha'})^2} \right)^5. \quad (4.40)$$

Consider for simplicity the limit $N \rightarrow \infty$ discussed above. As mentioned after eq. (4.36), since the Hagedorn temperature is very low, the modular integral is dominated in this case by $\tau_2 \sim N$. If the radius R is much larger than $\sqrt{N\alpha'}$, the sum over momenta on the r.h.s. of (4.40) can be approximated by an integral and gives the same contribution as in the non-compact case (namely the l.h.s. of (4.40)). For $R \sim \sqrt{N\alpha'}$ one has to include a few low lying momentum modes – this is a transition region. For $\sqrt{\alpha'} < R \ll \sqrt{N\alpha'}$ one can neglect all contributions of momentum (and winding) modes, just like one is neglecting the contributions of oscillator

states. Thus, we get in this case

$$\begin{aligned} \alpha + 1 = & -\frac{\beta}{2Q} \left(\frac{1}{4\pi^2 \alpha'} \right) \int_0^\infty \frac{d\tau_2}{\tau_2^2} \cdot 1024 \sum_{k,p=0}^{\infty} e^{-\frac{\beta^2(2k+1)^2}{4\pi\alpha'\tau_2} - \frac{(p+1)^2\tau_2}{2N}} = \\ & -\frac{256}{\pi} \sum_{k,p=0}^{\infty} \left(\frac{2\pi(2k+1)^2}{(p+1)^2} \right)^{-1/2} (p+1)^2 K_{-1}(\sqrt{2\pi}(p+1)(2k+1)) \simeq -3.693. \end{aligned} \quad (4.41)$$

Interestingly, we find that for small fivebranes α is independent of the number of fivebranes N in the $N \rightarrow \infty$ limit. Note also that in this case it is important to keep the $+1$ on the l.h.s. of (4.41), since α is of order one.

To summarize, the power α that appears in the high energy density of states (4.1) is negative, and exhibits an interesting dependence on the size of the spatial manifold that the fivebranes are wrapping. For manifolds of size much larger than the characteristic scale of LST, $\sqrt{N\alpha'}$, α is proportional to the volume of the manifold, while for sizes much smaller than this characteristic scale, it saturates at a finite value, which is independent of N (for large N), (4.41). If the density of states (4.1) is due to strings confined to the fivebranes, then these strings belong to a new universality class, with typical configurations not exceeding the size $\sqrt{N\alpha'}$. It would be interesting to understand this universality class better (see also [36]).

4.3 Comments on the near-Hagedorn thermodynamics of LST

The main result of the previous subsections is that the temperature-energy relation has the form (4.20), with α given by (4.36) or for large N by (4.39), (4.41). Since it is negative, the temperature is above the Hagedorn temperature, and the specific heat is negative. This raises two immediate questions:

- (1) What is the thermodynamics for temperatures slightly below the Hagedorn temperature?
- (2) What is the nature of the instability, reflected by the negative specific heat, above the Hagedorn temperature?

Consider first the behavior well below the Hagedorn temperature, $\beta \gg \beta_H$. In this regime, the thermodynamics is expected to reduce to that corresponding to the extreme IR limit of LST, which is the $(2, 0)$ six dimensional SCFT for type IIA LST, or six dimensional $(1, 1)$ SYM for IIB. From the

point of view of the holographic description, this regime corresponds to the strong coupling region of the near-horizon geometry of the fivebranes, and thus should not be well described by the perturbative theory on the cigar (4.6).

What happens as the temperature approaches T_H from below? One might expect that due to the Hagedorn growth in the density of states (4.1), the high energy part of the spectrum dominates as $\beta \rightarrow \beta_H$, and the partition sum becomes better and better approximated by (4.18). What actually happens depends on the value of α , as we discuss next.

Consider first the case of large V_5 ($R \gg \sqrt{N\alpha'}$ in the discussion at the end of section 4.2). In this case, $|\alpha|$ is large, and the contribution to the partition sum of the high energy part of the spectrum, (4.18), goes rapidly to zero as $\beta \rightarrow \beta_H$. The integral over E is dominated by states with moderate energies, whose contribution to the partition sum is analytic at β_H . It is clear that the mean energy remains finite as we approach the Hagedorn temperature from below, and that thermodynamic fluctuations are suppressed (by a factor of the volume V_5). Since the Hagedorn temperature is reached at a finite energy, it corresponds to a phase transition.

As V_5 decreases, α decreases as well, until it reaches the value (4.41). The fluctuations in energy in the canonical ensemble increase with decreasing α . To see that, consider the case $R \ll \sqrt{N\alpha'}$ in the discussion at the end of section 4.2. Since $-5 < \alpha < -4$ in that case, the expectation values $\langle E^n \rangle$ with $n \geq 4$ in the canonical ensemble diverge as

$$\langle E^n \rangle \sim (\beta - \beta_H)^{-\alpha-n-1}. \quad (4.42)$$

In such situations, one is instructed to pass to the microcanonical ensemble, in which the energy is fixed and the temperature is defined by

$$\beta = \frac{\partial \log \rho}{\partial E} = \beta_H + \frac{\alpha}{E} + \dots \quad (4.43)$$

where on the r.h.s. we included the first two terms in a perturbative expansion in $1/E$. The perturbative evaluation of β in (4.43) gives a temperature *above* the Hagedorn temperature. This of course does not imply that LST cannot be defined at temperatures below T_H ; instead, it means that to study the theory at such temperatures one must compute $S(E)$ to all orders in $1/E$, include non-perturbative corrections, and solve the equation (4.43) to find the energy E corresponding to a particular $\beta > \beta_H$. From the form of the leading terms in $S(E)$ it is clear that the solution of this equation will correspond to finite E . We are led again to the conclusion that the Hagedorn

temperature is reached at a finite energy and thus is associated with a phase transition.

Since the study of the non-extremal fivebrane geometry in the previous sections is perturbative in $1/E$, it is not useful for studying the regime $\beta > \beta_H$. Nevertheless, it seems clear that the specific heat is positive there (this is certainly the case for the infrared theory on the fivebranes). Furthermore, since the energy – temperature relation is such that the Hagedorn temperature is reached at a finite energy, we are led to the second question raised in the beginning of this section: what is the nature of the high temperature phase of LST?

The perturbative analysis of the near-extremal fivebrane, which is valid for β slightly below β_H , predicts that the thermodynamics is unstable. Usually, in such situations the instability is associated with a negative mode in the Euclidean path integral (a tachyon). Examples include the instability of flat space at finite temperature in Einstein gravity [39], and the thermal tachyon that appears above the Hagedorn transition in critical string theory. The one loop instability found above leads one to believe that a similar negative mode should appear in LST above the Hagedorn temperature.

In [23] it was shown that there is a natural candidate for this, a mode that lives near the tip of the cigar and is classically massless. It is likely that one loop corrections give a tachyonic correction to the mass of this state above the Hagedorn temperature, but this has not been proven and we will not discuss the detailed properties of this state here.

5 Weakly coupled LST

In the previous section we saw that the high energy thermodynamics of LST can be analyzed reliably using the holographically dual description, since at large energy density the strongly coupled region on \mathbb{R}_ϕ is eliminated, and the coupling never exceeds (4.14), a value that can be made arbitrarily small by increasing the energy density. In this section we will describe another situation where something similar happens at zero temperature, by studying the theory away from the origin of its moduli space of vacua. This section is based on [40].

Recall that the theory of N fivebranes contains four massless scalars in the adjoint of $U(N)$, X^i , $i = 6, 7, 8, 9$, parameterizing motions in $(6, 7, 8, 9)$. IIA fivebranes have one more scalar X^{11} , which is compact, but we will not discuss it here. The moduli space of vacua of LST is \mathbb{R}^{4N}/S_N for IIB and

$(\mathbb{R}^4 \times S^1)^N / S_N$ for IIA. The origin corresponds to coincident fivebranes; other points are labeled by relative separations of the fivebranes.

The four scalars X^i can be parametrized by two complex $N \times N$ matrices,

$$\begin{aligned} A &\equiv X^8 + iX^9, \\ B &\equiv X^6 + iX^7. \end{aligned} \tag{5.1}$$

Consider a point on the moduli space where

$$\begin{aligned} \langle A \rangle &= 0, \\ \langle B \rangle &= r_0 \text{diag}(1, e^{\frac{2\pi i}{N}}, e^{\frac{4\pi i}{N}}, \dots, e^{\frac{2\pi i(N-1)}{N}}). \end{aligned} \tag{5.2}$$

This corresponds to fivebranes symmetrically distributed around a circle of radius r_0 in the $(6, 7)$ plane. The gauge invariant characterization of this vacuum is

$$\langle \text{Tr } B^N \rangle = r_0^N \tag{5.3}$$

with all other v.e.v.'s of the operators (3.11) set to zero. Since for a single fivebrane the worldvolume dynamics is trivial, in order to get a non-trivial result in the limit (2.2), we have to tune $r_0 \rightarrow 0$ as we take the limit. E.g., in the IIB case the masses of D-strings stretched between $NS5$ -branes

$$M_W \sim \frac{r_0 m_s^2}{g_s} \tag{5.4}$$

must be kept finite in the limit. This leads one to consider the double scaling limit

$$g_s \rightarrow 0; r_0 m_s \rightarrow 0 \tag{5.5}$$

with M_W/m_s (5.4) held fixed.

Distributing the branes on a circle as in (5.2) breaks the $SO(4)$ R -symmetry

$$SO(4) \rightarrow SO(2) \times \mathbb{Z}_N. \tag{5.6}$$

We will next show that this also eliminates the strong coupling singularity at $\phi \rightarrow -\infty$ discussed above.

The first thing we have to understand is how to describe the vacuum (5.3) in the holographically dual theory. In section 3.1 we found the vertex operators corresponding to the gauge invariant operators (3.11). It is not difficult to see that

$$\text{Tr } B^N \leftrightarrow \psi^+ \bar{\psi}^+ V_{\frac{N}{2}-1; \frac{N}{2}-1, \frac{N}{2}-1} \exp \left[\frac{2}{\sqrt{N\alpha'}} \left(\frac{N}{2} - 1 \right) \phi \right]. \quad (5.7)$$

Adding the vertex operator (5.7) to the worldsheet action is equivalent, via the prescription (3.10), to adding the operator $\text{Tr } B^N$ to the action of LST. In order to turn on a v.e.v. of $\text{Tr } B^N$ instead, as in (5.3), we have to use the same vertex operator but replace the charge β in (3.13) by

$$\beta \rightarrow -Q - \beta. \quad (5.8)$$

Thus, to describe the vacuum (5.3) we must study the worldsheet Lagrangian

$$\mathcal{L} = \mathcal{L}_0 + \lambda G_{-\frac{1}{2}} \bar{G}_{-\frac{1}{2}} \psi^+ \bar{\psi}^+ V_{\frac{N}{2}-1; \frac{N}{2}-1, \frac{N}{2}-1} e^{-\sqrt{\frac{N}{\alpha'}} \phi} + \text{c.c.} \quad (5.9)$$

where we explicitly wrote the worldsheet supercharges which are needed to turn a $(-1, -1)$ picture vertex operator to a $(0, 0)$ picture one (the appropriate picture for a term in the worldsheet Lagrangian). λ is a coupling related to r_0 . \mathcal{L}_0 is the free Lagrangian describing string propagation on (3.4). Since the coupling λ breaks explicitly the $SU(2)_L \times SU(2)_R$ symmetry, it is convenient to analyze its effect by rewriting the background (3.4) as

$$\mathbb{R}^{5,1} \times \mathbb{R}_\phi \times \left(S^1 \times \frac{SU(2)}{U(1)} \right) / \mathbb{Z}_N \quad (5.10)$$

where $SU(2)/U(1)$ is an $N = 2$ minimal model, and S^1 a circle of radius $\sqrt{N\alpha'}$. Denoting the coordinate along the circle by Y , one can show that the interaction in (5.9) can be written as

$$\delta\mathcal{L} = \lambda G_{-\frac{1}{2}} \bar{G}_{-\frac{1}{2}} e^{-\frac{2}{\alpha'} Q(\phi+iY)} + \text{c.c.} \quad (5.11)$$

This interaction is familiar in CFT as the $N = 2$ Liouville interaction. Thus, we find that to describe the vacuum (5.3), we must replace the infinite cylinder $\mathbb{R}_\phi \times S^1$ in (5.10) by the $N = 2$ Liouville model. Note that:

- (1) The fact that the interaction (5.9), (5.11) preserves $N = 2$ superconformal invariance is related to the fact that spacetime supersymmetry remains unbroken along the moduli space of LST.

(2) The interaction (5.11) grows as $\phi \rightarrow -\infty$. One can show that it resolves the strong coupling singularity discussed in section 3. We will see this directly momentarily.

To study $N = 2$ Liouville theory, it is convenient to use a dual description of this background. It was argued in [40] that $N = 2$ Liouville is equivalent via strong-weak coupling duality on the worldsheet to CFT on the cigar, $H_3^+/U(1)$, which was discussed in section 2. The parameter N which enters the definition of $N = 2$ Liouville (5.11) via Q is mapped under the duality to the level of the underlying $SL(2)$ current algebra.

I will not describe the duality or the evidence for it here¹⁵, but rather will use it to conclude that the vacuum (5.2), (5.3) is dual to

$$\mathbb{R}^{5,1} \times \left(\frac{SL(2)}{U(1)} \times \frac{SU(2)}{U(1)} \right) / \mathbb{Z}_N . \quad (5.12)$$

Note that the unbroken R -symmetry $SO(2) \times \mathbb{Z}_N$ of the vacuum (5.3) is manifest in the description (5.12). The $SO(2)$ symmetry corresponding to rotations in the $(8, 9)$ plane is realized as the $U(1)$ translation symmetry around the cigar. The rotation symmetry in the $(6, 7)$ plane, which is broken to \mathbb{Z}_N by the v.e.v. of B , corresponds to winding number around the cigar. This quantum number is not conserved, since winding can slip off the tip of the cigar. The \mathbb{Z}_N orbifold in (5.12) leads to a \mathbb{Z}_N remnant of it (since it allows fractional windings $\in \mathbb{Z}/N$).

The radius of the circle on which the fivebranes lie, r_0 in (5.2), is related to the value of the string coupling at the tip of the cigar, g_{cigar} . The precise relation can be determined by noting that D-branes stretched between fivebranes, whose mass is given by (5.4), correspond in (5.12) to D-branes at the tip of the cigar, whose mass is m_s/g_{cigar} . This implies that

$$g_{\text{cigar}} \simeq \frac{m_s}{M_W} . \quad (5.13)$$

Thus, the theory is weakly coupled when $M_W \gg m_s$; as M_W decreases, we recover the original strongly coupled theory described holographically by (3.4). As mentioned above, the behavior (5.13) is very reasonable: as $M_W/m_s \rightarrow \infty$ the fivebranes become infinitely separated and decouple (recall that the dynamics on a single fivebrane is trivial).

The weakly coupled nature of the theory (5.12) for $M_W \gg m_s$ allows one to determine the spectrum in a wide range of energies $0 < E \ll M_W$,

¹⁵See [41, 42] for more detailed discussions.

and to compute various off-shell correlation functions of the observables discussed in section 3. Interactions can be turned on gradually by increasing g_{cigar} (5.13). For energies $E \gg M_W$ one expects the weak coupling expansion to break down. Physically, the reason for that is that in this regime the symmetry breaking in (5.2) can be neglected, and the physics is that of coincident fivebranes. All this is very similar to critical string theory, where the string coupling expansion is associated with a large hierarchy of energy scales, m_s/m_p . For $E \sim m_p$ the string coupling expansion breaks down.

Two and three point functions as well as the spectrum of weakly coupled LST were analyzed in [40]. We next illustrate the resulting structure by discussing an example.

Consider the operator $\text{Tr } B^N(x)$. The dual vertex operator (5.7) can be written in terms of the background (5.12) as

$$\text{Tr } B^N(x) \leftrightarrow e^{-\varphi-\bar{\varphi}} e^{ik_\mu x^\mu} V_{j;m,m} \quad (5.14)$$

with $m = N/2$. $\varphi, \bar{\varphi}$ are the standard bosonized superconformal ghosts needed for the -1 picture, $V_{j;m,\bar{m}}$ is a Virasoro primary on the cigar carrying p units of momentum and w units of winding, with

$$m = \frac{1}{2}(p + wN) ; \quad \bar{m} = -\frac{1}{2}(p - wN) . \quad (5.15)$$

In the case (5.14), $p = 0$ while $w = 1$ (*i.e.* $m = \bar{m} = N/2$). The worldsheet scaling dimension of $V_{j;m,m}$ is

$$\Delta = \bar{\Delta} = \frac{m^2 - j(j+1)}{N} \quad (5.16)$$

Requiring that (5.14) be physical gives rise to the mass-shell condition

$$\alpha' k_\mu k^\mu = \frac{4}{N}(j - m + 1)(j + m). \quad (5.17)$$

To compute the two point function of $\text{Tr } B^N(k_\mu)$ we use the correspondence (3.10):

$$\langle \text{Tr } B^N(k_\mu) \text{Tr } \bar{B}^N(-k_\mu) \rangle = \langle e^{-\varphi-\bar{\varphi}} e^{ik_\mu x^\mu} V_{j;m,m} e^{-\varphi-\bar{\varphi}} e^{-ik_\mu x^\mu} V_{j;-m,-m} \rangle. \quad (5.18)$$

The only non-trivial part of the correlator on the r.h.s. is $\langle VV \rangle$. It was computed in [43]:

$$\langle V_{j;m,\bar{m}} V_{j;-m,-\bar{m}} \rangle = N[\nu(N)]^{2j+1} \frac{\Gamma(1 - \frac{2j+1}{N}) \Gamma(-2j-1) \Gamma(j-m+1) \Gamma(1+j+\bar{m})}{\Gamma(\frac{2j+1}{N}) \Gamma(2j+2) \Gamma(-j-m) \Gamma(\bar{m}-j)} . \quad (5.19)$$

where

$$\nu(N) \equiv \frac{1}{\pi} \frac{\Gamma(1 + \frac{1}{N})}{\Gamma(1 - \frac{1}{N})} . \quad (5.20)$$

The two point function (5.19) has a series of poles; these can be interpreted as contributions of on-shell states in weakly coupled LST, which are created from the vacuum by the operator (5.14). The masses of these states can be computed by using the relation (5.17) between j and $M^2 = -k_\mu k^\mu$. The locations of the poles are given by

$$|m| = j + n; \quad n = 1, 2, 3, \dots \quad (5.21)$$

These values of m and j belong to the principal discrete series representations of $SL(2)$. The corresponding states can be thought of as bound states that live near the tip of the cigar [44]. Such bound states are to be expected since winding modes around the cigar feel an effective attractive potential towards the tip – their energy decreases as they approach the tip and shrink.

For the particular case (5.14), $m = \bar{m} = N/2$, and the masses of these states are given by

$$\begin{aligned} \frac{\alpha'}{2} M_n^2 &= \frac{2}{N} (n - 1)(N - n) , \\ N + 1 &> 2n > 1 . \end{aligned} \quad (5.22)$$

The second line in (5.22) comes from a unitarity constraint on j which must be imposed, $-1/2 < j < (N - 1)/2$. Note that all the masses squared in (5.22) are non-negative; For $n = 1$ one finds massless states, which correspond to the eigenvalues of the scalar matrix B .

A few comments are in order here:

- (1) By analyzing the behavior of the two point function (5.18), (5.19) one can check that the residues of the poles corresponding to the states (5.22) are positive, in agreement with the unitarity of the theory.
- (2) In addition to the discrete spectrum given by (5.22), one also has the continuum discussed in section 3 (3.19). One can show that the continuum starts right above the heaviest state (5.22). Thus the spectrum of states that can be created from the vacuum by the operator (5.14) is a finite discrete set, followed by a continuum (similar to the spectrum of bound states and scattering states in quantum mechanics).

- (3) It is interesting that the low lying spectrum of states associated with N $NS5$ -branes is independent of M_W , or equivalently the radius of the circle on which the fivebranes are placed. This should be contrasted with D-branes, for which masses of open strings stretched between different branes depend on the separation. When the distance between D-branes goes to infinity, states associated with strings stretched between different branes go to infinite mass and decouple. For $NS5$ -branes, the masses of low lying states remain finite, and the decoupling is due to the vanishing of the effective coupling (5.13).
- (4) In addition to the poles (5.21), which correspond to principal discrete series states near the tip of the cigar, the amplitude (5.19) has poles at $0 \leq 2j + 1 \in \mathbb{Z}$, $0 < 2j + 1 \in N\mathbb{Z}$. These poles have a different interpretation than (5.21). They are associated with “bulk scattering processes” which can occur anywhere in the infinite throat corresponding to either the $N = 2$ Liouville (5.11), or $SL(2)/U(1)$ (5.12) description. This is discussed further in [42].
- (5) One can repeat the above discussion for other observables as well. The resulting picture is similar; one always finds a finite set of discrete states which live near the tip of the cigar, followed by a continuum of states which propagate in the semi-infinite throat [40].
- (6) Since there is a Hagedorn growth in the number of observables (coming from oscillator states on (5.12)), one finds a Hagedorn density of states in LST. But the exponent β_H (4.1) does not grow like \sqrt{N} as expected from (4.10). Instead one gets $\beta_H \sim 1/m_s$. This is not particularly surprising since (4.10) is the expected behavior for high energies $E \gg M_W$, whereas the present analysis is only valid in the intermediate regime $m_s \ll E \ll M_W$.
- (7) Three point functions of the off-shell observables discussed above can be computed as well using the results of [43]. One finds a similar analytic structure to that exhibited by the two point functions. There are poles associated with external legs going on-shell; their locations correspond again to the spectrum (5.22). The residues of these poles describe the scattering amplitudes of the physical states; they seem to have sensible physical properties. See [40] for details.

6 Other aspects of LST

In this section we would like to briefly list some additional topics in Little String Theory, which were not covered in detail in the lectures due to lack of time.

6.1 Singular Calabi-Yau manifolds and lower dimensional vacua of LST

The theory of N $NS5$ -branes discussed in sections 2 – 5 is related to string dynamics on an ALE space $\mathbb{C}^2/\mathbb{Z}_N$, which can be described as the manifold

$$z_1^N + z_2^2 + z_3^2 = \mu \quad (6.1)$$

in \mathbb{C}^3 . For $\mu = 0$, (6.1) corresponds to a cone; non-zero μ smoothes out the tip of the cone. String propagation on $\mathbb{R}^{5,1} \times \mathbb{C}^2/\mathbb{Z}_N$ is dual [45, 46] to a vacuum with coincident fivebranes. The blowing up parameter μ is related by duality to the distance between the fivebranes. From the perspective of the geometry (6.1), LST describes the dynamics of the modes localized at the singularity, which can be decoupled from the rest of the theory.

This picture can be naturally generalized to a large class of vacua of LST in $d < 6$ dimensions [47]. Consider, for example, string propagation on

$$\mathbb{R}^{3,1} \times \mathcal{M} , \quad (6.2)$$

where \mathcal{M} is a Calabi-Yau manifold with an isolated singularity, which looks locally like

$$F(z_1, z_2, z_3, z_4) = 0 . \quad (6.3)$$

Here F is a quasi-homogeneous polynomial,

$$F(\lambda^{r_1} z_1, \lambda^{r_2} z_2, \lambda^{r_3} z_3, \lambda^{r_4} z_4) = \lambda F(z_1, z_2, z_3, z_4) \quad (6.4)$$

for some set of charges r_1, r_2, r_3, r_4 . Viewed as a hypersurface in \mathbb{C}^4 , (6.3) describes the vicinity of the singular point $z_1 = z_2 = z_3 = z_4 = 0$.

In analogy to the six dimensional situation (6.1), string theory in the background (6.3) is expected to contain modes localized near the singularity; these modes can be decoupled from the bulk in the same way as in the six dimensional case.

The decoupled dynamics at the singularity (6.3) is holographically dual to string theory in the background

$$\mathbb{R}^{3,1} \times \mathbb{R}_\phi \times (S^1 \times LG(F)) / \Gamma , \quad (6.5)$$

where $LG(F)$ is a Landau-Ginsburg model with the superpotential given by the quasi-homogeneous polynomial $F(z_1, \dots, z_4)$ defining the singularity (6.3). Γ is a discrete group whose origin is the chiral GSO projection in the vacuum (6.5). As before, \mathbb{R}_ϕ is a linear dilaton direction, with the slope Q determined such that the total central charge of (6.5) is fifteen, as appropriate for a critical superstring vacuum. One can show that this implies that

$$\frac{1}{2}Q^2 = \sum_{a=1}^4 r_a - 1 . \quad (6.6)$$

Vacua of the form (6.5) preserve eight supercharges and give rise to $N = 2$ supersymmetric theories in four dimensions.

A simple example is

$$F = z_1^2 + z_2^2 + z_3^3 + z_4^2 , \quad (6.7)$$

which corresponds to the conifold. In this case, (6.5) reduces to

$$\mathbb{R}^{3,1} \times \mathbb{R}_\phi \times S^1 , \quad (6.8)$$

which is the background holographically dual to string theory on the conifold. Smoothing out the singularity as in (6.1) corresponds to replacing the factor $\mathbb{R}_\phi \times S^1$ in (6.8) by the cigar $SL(2)/U(1)$ (or equivalently $N = 2$ Liouville).

In the same way that the ALE space (6.1) is dual to parallel fivebranes, the background (6.7), (6.8) arises from two orthogonal $NS5$ -branes intersecting along 3 + 1 dimensional Minkowski spacetime.

More generally, if

$$F(z_1, \dots, z_4) = H(z_1, z_2) + z_3^2 + z_4^2 , \quad (6.9)$$

the background (6.3) can be thought of as arising from an $NS5$ -brane wrapped around the surface $H(z_1, z_2) = 0$ [48]. An interesting class of examples corresponds to $H(z_1, z_2)$ describing an ADE singularity (*e.g.* $H = z_1^n + z_2^2$ for A_{n-1}), in which case the fivebrane wraps a Seiberg-Witten curve at the corresponding Argyres-Douglas point. For type IIA fivebranes, at low energies

the system approaches an interacting four dimensional $N = 2$ SCFT. In the description (6.5), this SCFT corresponds to the strong coupling region in the background (6.5), where the theory is eleven dimensional, and is difficult to study in detail (beyond the supergravity approximation). In [47] it was shown that certain properties of chiral operators which can be studied at weak coupling (such as the R-charges), agree with known results.

The construction described in this subsection can be generalized to other dimensions and more complicated models in four dimensions. For some work in this direction, see [49, 50, 51, 52, 53, 54]. Other vacua of LST in six dimensions with less than maximal supersymmetry were discussed in [55, 56, 57].

6.2 D-branes in the vicinity of $NS5$ -branes

D -branes stretched between $NS5$ -branes in the weak coupling limit $g_s \rightarrow 0$ have been seen in recent years to be very useful for studying the dynamics of a wide class of gauge theories, which are realized as the low energy theories on such branes [2]. In particular, $D4$ -branes stretched between parallel adjacent fivebranes realize $N = 2$ SYM and are very useful for embedding Seiberg-Witten theory in string theory [58]. $D4$ -branes stretched between orthogonal fivebranes which share $3 + 1$ dimensions, give rise to $N = 1$ SYM and are very useful for studying Seiberg duality in string theory [59].

We have seen above that nearby fivebranes can be described by throat geometries which involve the cigar $SL(2)/U(1)$. Adjacent parallel fivebranes are described by (5.12), while orthogonal fivebranes intersecting on $3 + 1$ dimensional Minkowski spacetime correspond to $\mathbb{R}^{3,1} \times SL(2)/U(1)$. D -branes stretched between the fivebranes correspond in this description to D -branes localized on the cigar and extended in some or all of the non-compact directions. For example, a $D4$ -brane stretched between parallel fivebranes corresponds in the geometry (5.12) to a $D3$ -brane in $\mathbb{R}^{5,1}$, which is localized on the cigar, and is in one of the familiar boundary states in the $N = 2$ minimal model $SU(2)/U(1)$. One can show that different boundary states in the minimal model correspond to D -branes stretched between different pairs of $NS5$ -branes.

Thus, one is led to study D -branes localized on the cigar. It is clear that such D -branes will live near the tip of the cigar, since this is where the string coupling is largest, and thus the energy of the D -branes is smallest. The physics of D -branes living near the tip of the cigar is at present not

completely understood, but progress has been made on two closely related problems. In [60], D-branes in Liouville theory have been constructed which can be thought of as being localized in the Liouville direction. It was found that such D-branes are labeled by two integers, and exhibit very interesting properties, such as having a finite number of Virasoro primaries in the open string sector. This construction was generalized in [61] to Euclidean AdS_3 , where similar D-branes were found.

Both Liouville theory and AdS_3 are known to share many properties with the cigar and $N = 2$ Liouville CFT's. Thus, it is reasonable to expect that localized D-branes exist on the cigar and $N = 2$ Liouville backgrounds as well. It would be interesting to construct them and use their properties to study gauge dynamics.

Another class of objects that figures in many brane constructions is D-branes ending on $NS5$ -branes. In some cases such branes can be studied by analyzing them in the throat region of the fivebranes. For example, consider a D-brane that ends on a stack of N fivebranes. Assuming that the brane extends into the throat of the fivebranes,¹⁶ one can study the behavior of the brane inside the throat region described by (5.10), or for separated fivebranes by (5.12). This analysis was carried out in [62], where many properties of such D-branes that were previously deduced by using spacetime considerations, were verified by using the technology of LST.

6.3 Low dimensional toy models of LST

From the modern perspective, the matrix model description of two dimensional string theory (see *e.g.* [63]) for a review) provides an early example of holography in string theory. Since the “bulk” theory involves in this case a linear dilaton direction, the situation looks like a low dimensional toy model of LST.

Unlike LST, which is difficult to formulate directly (except for the DLCQ construction of [7]), here there is an alternative definition of the theory, which is moreover exactly solvable to all orders in the string coupling expansion (or equivalently, the $1/N$ expansion in the matrix model). This is especially interesting since, like LST, two dimensional string theory is expected to exhibit a Hagedorn growth in the density of states, as in (4.1), with the parameters

¹⁶This is a non-trivial assumption; it is believed that in some cases D-branes that end on fivebranes do not extend into the throat. The simplest example is D-branes ending on a single fivebrane, which does not have a throat region.

β_H and α known from thermodynamic considerations [64]. Therefore, two dimensional string theory is an interesting toy model of the dynamics of LST in higher dimensions.

In [64], the matrix model description was used to study some properties of the Euclidean black hole solution of two dimensional string theory. In particular, some steps were taken towards developing a description of the states that give rise to the Hagedorn entropy (4.1) directly in the matrix model.

7 Some open problems in LST

While a lot has been achieved, many interesting questions regarding Little String Theory await resolution. Some examples of open problems are:

- (1) We have seen in section 4 that LST has at high energies a Hagedorn spectrum of states (4.1). This was established by a thermodynamic analysis; it would be very interesting to exhibit the density of state (4.1) by an explicit counting of states. In the background (3.4) corresponding to coincident fivebranes (and the lower dimensional analogs (6.5)), this is complicated by the fact that string theory in the linear dilaton background is not weakly coupled. The weakly coupled theory described in section 5 does have a Hagedorn spectrum of perturbative states, but the Hagedorn coefficient β_H is smaller than that of the full theory (4.10). As explained in section 5, this is not surprising – most of the states contributing to (4.1) are expected to be non-perturbative. As a first step to counting non-perturbative states in the background (5.12), it would be interesting to enumerate states corresponding to collections of D-branes living near the tip of the cigar, which were briefly discussed in section 6.
- (2) It would be interesting to understand the dynamics of D-branes stretched between $NS5$ -branes, which correspond to D-branes localized near the tip of the cigar in the dual geometry (5.12). This might be useful for application to gauge dynamics, as well as for providing a direct formulation of LST, independent of holography.
- (3) Most of the work on LST concerned spacetime supersymmetric vacua. In the absence of spacetime SUSY one expects to find infrared instabilities such as tachyons, and the system might decay to a more stable

vacuum. It would be interesting to understand the physics associated with supersymmetry breaking and vacuum instabilities in LST. First steps in that direction were recently taken in [65].

Acknowledgements:

I thank O. Aharony, M. Berkooz, S. Elitzur, A. Giveon, V. Kazakov, I. Kostov, O. Pelc, E. Rabinovici, D. Sahakyan, G. Sarkissian, and N. Seiberg for collaboration on the topics presented here. This work is supported in part by DOE grant #DE-FG02-90ER40560. I thank the Weizmann Institute of Science for hospitality during the preparation of these notes.

References

- [1] L. J. Dixon, J. A. Harvey, C. Vafa and E. Witten, “Strings On Orbifolds,” *Nucl. Phys. B* **261**, 678 (1985).
- [2] A. Giveon and D. Kutasov, “Brane dynamics and gauge theory,” *Rev. Mod. Phys.* **71**, 983 (1999) [hep-th/9802067].
- [3] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity,” *Phys. Rept.* **323**, 183 (2000) [hep-th/9905111].
- [4] C. G. Callan, J. A. Harvey and A. Strominger, “Supersymmetric string solitons,” hep-th/9112030.
- [5] P. S. Aspinwall, “Enhanced gauge symmetries and K3 surfaces,” *Phys. Lett. B* **357**, 329 (1995) [hep-th/9507012].
- [6] A. Losev, G. W. Moore and S. L. Shatashvili, “M & m’s,” *Nucl. Phys. B* **522**, 105 (1998) [hep-th/9707250].
- [7] O. Aharony, M. Berkooz, S. Kachru, N. Seiberg and E. Silverstein, “Matrix description of interacting theories in six dimensions,” *Adv. Theor. Math. Phys.* **1**, 148 (1998) [hep-th/9707079].
- [8] E. Witten, “On the conformal field theory of the Higgs branch,” *JHEP* **9707**, 003 (1997) [hep-th/9707093].
- [9] O. Aharony, M. Berkooz and N. Seiberg, “Light-cone description of (2,0) superconformal theories in six dimensions,” *Adv. Theor. Math. Phys.* **2**, 119 (1998) [hep-th/9712117].
- [10] O. Aharony, “A brief review of little string theories”, *Class. Quant. Grav.* **17** (2000) 929, [hep-th/9911147].
- [11] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, “M theory as a matrix model: A conjecture,” *Phys. Rev. D* **55**, 5112 (1997) [hep-th/9610043].
- [12] M. Berkooz, M. Rozali and N. Seiberg, “On transverse fivebranes in M(atrix) theory on T^{**5} ,” *Phys. Lett. B* **408**, 105 (1997) [hep-th/9704089].

- [13] N. Seiberg, “New theories in six dimensions and matrix description of M-theory on T^{*5} and $T^{*5}/Z(2)$,” *Phys. Lett. B* **408**, 98 (1997) [hep-th/9705221].
- [14] I. Antoniadis, S. Dimopoulos and A. Giveon, “Little string theory at a TeV,” *JHEP* **0105**, 055 (2001) [hep-th/0103033].
- [15] J. Polchinski, “String theory”, vol. 2, Cambridge University Press, 1998.
- [16] E. Witten, “Some comments on string dynamics,” hep-th/9507121.
- [17] A. Strominger, “Open p-branes,” *Phys. Lett. B* **383**, 44 (1996) [hep-th/9512059].
- [18] O. Aharony, M. Berkooz, D. Kutasov and N. Seiberg, “Linear dilatons, NS5-branes and holography,” *JHEP* **9810**, 004 (1998) [hep-th/9808149].
- [19] N. Itzhaki, J. M. Maldacena, J. Sonnenschein and S. Yankielowicz, “Supergravity and the large N limit of theories with sixteen supercharges,” *Phys. Rev. D* **58**, 046004 (1998) [hep-th/9802042].
- [20] H. J. Boonstra, K. Skenderis and P. K. Townsend, “The domain wall/QFT correspondence,” *JHEP* **9901**, 003 (1999) [hep-th/9807137].
- [21] N. Seiberg and S. H. Shenker, “A Note on background (in)dependence,” *Phys. Rev. D* **45**, 4581 (1992) [hep-th/9201017].
- [22] S. Minwalla and N. Seiberg, “Comments on the IIA NS5-brane,” *JHEP* **9906**, 007 (1999) [hep-th/9904142].
- [23] D. Kutasov and D. A. Sahakyan, “Comments on the thermodynamics of little string theory,” *JHEP* **0102**, 021 (2001) [hep-th/0012258].
- [24] M. Rangamani, “Little string thermodynamics,” *JHEP* **0106**, 042 (2001) [hep-th/0104125].
- [25] K. Narayan and M. Rangamani, “Hot little string correlators: A view from supergravity,” *JHEP* **0108**, 054 (2001) [hep-th/0107111].
- [26] A. Buchel, “On the thermodynamic instability of LST,” hep-th/0107102.

- [27] G. Horowitz and A. Strominger, “Black Strings and p -Branes”, *Nucl. Phys.* **B360** (1991) 197; J. Maldacena and A. Strominger, “Semiclassical decay of near extremal fivebranes”, *JHEP* **9712** (1997) 008, hep-th/9710014.
- [28] E. Witten, “On String Theory and Black Holes”, *Phys. Rev.* **D44** (1991) 314.
- [29] G. Gibbons and M. Perry, “The Physics of 2-d Stringy Spacetimes”, *Int. J. Mod. Phys.* **D1** (1992) 335, hep-th/9204090; C. R. Nappi and A. Pasquinucci, “Thermodynamics of Two-Dimensional Black-Holes”, *Mod. Phys. Lett.* **A7** (1992) 3337, gr-qc/9208002.
- [30] J. Polchinski, “Evaluation of the One Loop String Path Integral”, *Commun. Math. Phys.* **104** (1986) 37.
- [31] D. Kutasov and N. Seiberg, “Noncritical Superstrings,” *Phys. Lett. B* **251**, 67 (1990); D. Kutasov, “Some properties of (non)critical strings,” hep-th/9110041.
- [32] B. McClain and B. D. B. Roth, “Modular Invariance for Interacting Bosonic String at Finite Temperature”, *Commun. Math. Phys.* **111** (1987) 539.
- [33] K. H. O’Brien and C.-I. Tan, “Modular Invariance of the thermo-partition function and global phase structure of the heterotic string”, *Phys. Rev.* **D36** (1987) 1184.
- [34] J. J. Atick and E. Witten, “The Hagedorn Transition and the Number of Degrees of Freedom of String Theory”, *Nucl. Phys.* **B310** (1988) 291.
- [35] P. Di Francesco, P. Mathieu, D. Senechal, “Conformal Field Theory”, Springer-Verlag, 1997.
- [36] M. Berkooz and M. Rozali, “Near Hagedorn dynamics of NS fivebranes, or a new universality class of coiled strings,” *JHEP* **0005**, 040 (2000) [hep-th/0005047].
- [37] T. Harmark and N. A. Obers, “Hagedorn behaviour of little string theory from string corrections to NS5-branes,” *Phys. Lett. B* **485**, 285 (2000) [hep-th/0005021].

- [38] J. Polchinski, “String theory”, vol. 1, Cambridge University Press, 1998.
- [39] D. J. Gross, M. J. Perry and L. G. Yaffe, “Instability of Flat Space at Finite Temperature”, *Phys. Rev. D* **25** (1982) 330.
- [40] A. Giveon and D. Kutasov, “Little string theory in a double scaling limit,” *JHEP* **9910**, 034 (1999) [hep-th/9909110]; “Comments on double scaled little string theory,” *JHEP* **0001**, 023 (2000) [hep-th/9911039].
- [41] K. Hori and A. Kapustin, “Duality of the fermionic 2d black hole and $N = 2$ Liouville theory as mirror symmetry,” *JHEP* **0108**, 045 (2001) hep-th/0104202.
- [42] A. Giveon and D. Kutasov, “Notes on $AdS(3)$,” hep-th/0106004.
- [43] J. Teschner, “On structure constants and fusion rules in the $SL(2, C)/SU(2)$ WZNW model”, *Nucl. Phys. B* **546** (1999) 390, hep-th/9712256; “The Mini-Superspace Limit of the $SL(2, C)/SU(2)$ -WZNW Model”, *Nucl. Phys. B* **546** (1999) 369, hep-th/9712258; “Operator product expansion and factorization in the H_3^+ -WZNW model”, *Nucl. Phys. B* **571** (2000) 555, hep-th/9906215.
- [44] R. Dijkgraaf, H. Verlinde and E. Verlinde, “String propagation in a black hole geometry,” *Nucl. Phys. B* **371**, 269 (1992).
- [45] H. Ooguri and C. Vafa, “Two-Dimensional Black Hole and Singularities of CY Manifolds,” *Nucl. Phys. B* **463**, 55 (1996) hep-th/9511164.
- [46] D. Kutasov, “Orbifolds and Solitons,” *Phys. Lett. B* **383**, 48 (1996) hep-th/9512145.
- [47] A. Giveon, D. Kutasov and O. Pelc, “Holography for non-critical superstrings,” *JHEP* **9910**, 035 (1999) hep-th/9907178.
- [48] A. Klemm, W. Lerche, P. Mayr, C. Vafa and N. P. Warner, “Self-Dual Strings and $N=2$ Supersymmetric Field Theory,” *Nucl. Phys. B* **477**, 746 (1996) [hep-th/9604034].
- [49] O. Pelc, “Holography, singularities on orbifolds and 4D $N = 2$ SQCD,” *JHEP* **0003**, 012 (2000) [hep-th/0001054].

- [50] T. Eguchi and Y. Sugawara, “Modular invariance in superstring on Calabi-Yau n-fold with A-D-E singularity,” *Nucl. Phys. B* **577**, 3 (2000) [hep-th/0002100].
- [51] S. Mizoguchi, “Modular invariant critical superstrings on four-dimensional Minkowski space x two-dimensional black hole,” *JHEP* **0004**, 014 (2000) [hep-th/0003053]; “Noncompact Gepner models for type II strings on a conifold and an ALE instanton,” hep-th/0009240.
- [52] S. Yamaguchi, “Gepner-like description of a string theory on a non-compact singular Calabi-Yau manifold,” *Nucl. Phys. B* **594**, 190 (2001) [hep-th/0007069].
- [53] M. Naka and M. Nozaki, “Singular Calabi-Yau manifolds and ADE classification of CFTs,” *Nucl. Phys. B* **599**, 334 (2001) [hep-th/0010002].
- [54] K. Ohta and T. Yokono, “Linear dilaton background and fully localized intersecting five-branes,” *Phys. Rev. D* **63**, 105011 (2001) [hep-th/0012030].
- [55] D. E. Diaconescu and J. Gomis, “Neveu-Schwarz five-branes at orbifold singularities and holography,” *Nucl. Phys. B* **548**, 258 (1999) [hep-th/9810132].
- [56] D. Kutasov, F. Larsen and R. G. Leigh, “String theory in magnetic monopole backgrounds,” *Nucl. Phys. B* **550**, 183 (1999) [hep-th/9812027].
- [57] M. Gremm and A. Kapustin, “Heterotic little string theories and holography,” *JHEP* **9911**, 018 (1999) [hep-th/9907210].
- [58] E. Witten, “Solutions of four-dimensional field theories via M-theory,” *Nucl. Phys. B* **500**, 3 (1997) [hep-th/9703166].
- [59] S. Elitzur, A. Giveon and D. Kutasov, “Branes and $N = 1$ duality in string theory,” *Phys. Lett. B* **400**, 269 (1997) [hep-th/9702014].
- [60] A. B. Zamolodchikov and A. B. Zamolodchikov, “Liouville field theory on a pseudosphere,” hep-th/0101152.
- [61] A. Giveon, D. Kutasov and A. Schwimmer, “Comments on D-branes in $AdS(3)$,” *Nucl. Phys. B* **615**, 133 (2001) [hep-th/0106005].

- [62] S. Elitzur, A. Giveon, D. Kutasov, E. Rabinovici and G. Sarkissian, “D-branes in the background of NS fivebranes,” JHEP **0008**, 046 (2000) [hep-th/0005052].
- [63] P. Ginsparg and G. W. Moore, “Lectures On 2-D Gravity And 2-D String Theory,” hep-th/9304011.
- [64] V. Kazakov, I. K. Kostov and D. Kutasov, “A matrix model for the two-dimensional black hole,” hep-th/0101011.
- [65] J. A. Harvey, D. Kutasov, E. J. Martinec and G. Moore, “Localized tachyons and RG flows,” hep-th/0111154.

