

Quantum Science and Technology

**OPEN ACCESS****PAPER**

Conditional quantum thermometry—enhancing precision by measuring less

RECEIVED
15 October 2023**REVISED**
8 July 2024**ACCEPTED FOR PUBLICATION**
23 July 2024**PUBLISHED**
2 August 2024**Akira Sone**^{1,*} **Diogo O Soares-Pinto**² **and Sebastian Deffner**³ ¹ Department of Physics, University of Massachusetts, Boston, MA 02125, United States of America² Instituto de Física de São Carlos, Universidade de São Paulo, CP 369, 13560-970 São Carlos, São Paulo, Brazil³ Department of Physics, University of Maryland, Baltimore County, Baltimore, MD 21250, United States of America

* Author to whom any correspondence should be addressed.

E-mail: akirasone628@gmail.com**Keywords:** conditional thermal state, pointer basis, quantum Fisher information

Original Content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.

**Abstract**

Taking accurate measurements of the temperature of quantum systems is a challenging task. The mathematical peculiarities of quantum information make it virtually impossible to measure with infinite precision. In the present paper, we introduce a generalized thermal state, which is conditioned on the pointer states of the available measurement apparatus. We show that this conditional thermal state outperforms the Gibbs state in quantum thermometry. The origin for the enhanced precision can be sought in its asymmetry quantified by the Wigner–Yanase–Dyson skew information. This additional resource is further clarified in a fully resource-theoretic analysis, and we show that there is a Gibbs-preserving map to convert a target state into the conditional thermal state. We relate the quantum J-divergence between the conditional thermal state and the same target state to quantum heat.

1. Introduction

Quantum metrology is a task to utilize the peculiar properties of quantum systems, such as quantum coherence and entanglement, to achieve parameter estimation at precision beyond the classical limit [1–4]. Hence, the quest to identify quantum states that are uniquely suited for metrological tasks with limited resources [5, 6] is of crucial practical relevance. In the following, we address this issue and impose the plausible and realistic condition that only the observable *eigenstates* of the measurement apparatus, aka pointer states [7–11] are available.

One of the most prominent metrological tasks is quantum thermometry of stationary states. In fact, quantum thermometry with Gibbs states has been well studied [12–20]. In this simplest scenario it is easy to see that the optimal measurement for estimating the inverse temperature β is the Hamiltonian. However, when the system size is very large and supports quantum correlations, even energy measurements are a challenging task [14, 15]. Therefore, it is desirable to find *better* states whose corresponding optimal measurements are experimentally implementable, and which ideally even outperform Gibbs states in the low-temperature limit [21–23].

In this paper, we discuss a quantum state, that indeed fulfills the aforementioned ‘wishlist’. The *conditional thermal state* (CTS) is constructed as Gibbsian-distributed quantum state of the pointer basis corresponding to the available measurement apparatus. The CTS originally appeared in the one-time measurement approach to quantum work [24–29] and correspondingly tighter maximum work theorems. In the following, we demonstrate that the CTS outperforms the Gibbs state in quantum thermometry. To elucidate its properties further we then show that it can be understood as a non-passive state with useful resources [30–34]. To this end, we first relate its asymmetry, which is quantified by the Wigner–Yanase–Dyson (WYD) skew information [35–40], to its QFI. Then, focusing on a system undergoing unitary evolution, we discuss the state convertibility between the exact final state and the CTS constructed by the pointer states given by the evolved post-measurement state. Finally, we demonstrate that their symmetric

divergence, also known as quantum J-divergence, can be interpreted as quantum heat [41]. Our results demonstrate the usefulness of the CTS as a resource state from both fundamental and practical perspectives.

2. Conditional thermal state

We begin by establishing notions and notations. For the ease of the presentation, and to avoid clutter in the formulas, we work in units for which the Boltzmann constant k_B and the reduced Planck constant \hbar are $k_B = \hbar = 1$.

The concept of the CTS is illustrated in figure 1. In a d -dimensional Hilbert space \mathcal{H} , given a Hamiltonian H and the pointer states $\{|\psi_k\rangle\}_{k=1}^d$ of an implementable measurement M , the CTS is defined as

$$\rho_\beta \equiv \sum_{k=1}^d \frac{e^{-\beta \langle \psi_k | H | \psi_k \rangle}}{Z_\beta} |\psi_k\rangle\langle \psi_k|. \quad (1)$$

Here, Z_β is the normalization factor

$$Z_\beta \equiv \sum_{k=1}^d e^{-\beta \langle \psi_k | H | \psi_k \rangle}, \quad (2)$$

which can be interpreted as a generalized partition function [24–28]. The CTS is defined as the thermal state conditioned on the choice of the measurement implementable in the laboratory. The CTS maximizes the von Neumann entropy [42] under the constraint that the ensemble average of the Hamiltonian is fixed (see appendix A). Note that when $\{|\psi_k\rangle\}_{k=1}^d$ are the pointer states of H , ρ_β becomes the Gibbs state $\rho_\beta^{\text{eq}} \equiv e^{-\beta H}/Z_\beta^{\text{eq}}$, where $Z_\beta^{\text{eq}} \equiv \text{tr} \{e^{-\beta H}\}$ is the standard canonical partition function. Therefore, the CTS can be regarded as a generalized thermal state.

For later convenience, we also define the following conditional thermal separable state in a composite Hilbert space $\mathcal{H}_1 \otimes \mathcal{H}_2$

$$\tilde{\rho}_\beta \equiv \sum_{k=1}^d \frac{e^{-\beta \langle \psi_k | H | \psi_k \rangle}}{Z_\beta} |\psi_k\rangle\langle \psi_k|_1 \otimes |\phi_k\rangle\langle \phi_k|_2, \quad (3)$$

where $\{|\phi_k\rangle\}_{k=1}^d$ are the arbitrary orthogonal eigenstates of \mathcal{H}_2 . From these definitions, we also have $\rho_\beta = \text{tr}_2 \{\tilde{\rho}_\beta\}$, which will become important in our discussion of the relation between the quantum Fisher information (QFI) for quantum thermometry and the asymmetry measure of ρ_β .

3. Quantum thermometry

3.1. Quantum Fisher information

An important figure of merit in quantum metrology is the QFI. Given a quantum state ρ_θ parameterized by a certain parameter $\theta \in \mathbb{R}$, the variance of an unbiased estimator $(\delta\theta)^2$ is bounded by the quantum Cramér–Rao bound (QCRB), $(\delta\theta)^2 \geq 1/\mathcal{I}(\rho_\theta; \theta)$, where $\mathcal{I}(\rho_\theta; \theta)$ is the QFI defined by $\mathcal{I}(\rho_\theta; \theta) \equiv -2 \lim_{\epsilon \rightarrow 0} \partial_\epsilon^2 \mathcal{F}(\rho_\theta, \rho_{\theta+\epsilon})$ [43–48]. The definition stems from the relation between the Bures distance and the symmetric logarithmic derivative. Thus, the QFI can be obtained by optimizing the classical Fisher information over all possible measurements, as detailed in [45]. Here, $\mathcal{F}(\rho, \sigma) \equiv \|\sqrt{\rho}\sqrt{\sigma}\|_1^2$ is the quantum fidelity between two quantum states ρ and σ [49], and $\|A\|_1 \equiv \text{tr} \{\sqrt{A^\dagger A}\}$ is the trace norm. We denote $\partial^m/\partial x^m$ simply by ∂_x^m .

3.2. Quantum thermometry with the CTS


For our present purposes we now analyze the precision, with which the inverse temperature β can be estimated from the CTS. This will also allow us to relate the QFI of the CTS to the WYD skew information $I_\alpha(\rho_\beta, H)$ contained by ρ_β with respect to the Hamiltonian H , which quantifies the asymmetry of ρ_β .

The QFI $\mathcal{I}(\rho_\beta; \beta)$ of the CTS ρ_β is given by (see appendix B for the proof.)

$$\mathcal{I}(\rho_\beta; \beta) = \partial_\beta^2 \ln Z_\beta, \quad (4)$$

and the optimal measurement achieving the quantum Cramér–Rao bound is

$$G = \sum_{k=1}^d \langle \psi_k | H | \psi_k \rangle |\psi_k\rangle\langle \psi_k|. \quad (5)$$



To compare the sensitivity of the CTS ρ_β and the Gibbs state ρ_β^{eq} , we define the QFI difference as $\Delta\mathcal{I}_\beta \equiv \mathcal{I}(\rho_\beta; \beta) - \mathcal{I}(\rho_\beta^{\text{eq}}; \beta)$. We have

$$\Delta\mathcal{I}_\beta = -\partial_\beta^2 S(\rho_\beta || \rho_\beta^{\text{eq}}) = \partial_\beta^2 \left(\ln \frac{Z_\beta}{Z_\beta^{\text{eq}}} \right), \quad (6)$$

where $S(\rho_\beta || \rho_\beta^{\text{eq}}) \equiv \text{tr} \{ \rho_\beta \ln \rho_\beta \} - \text{tr} \{ \rho_\beta \ln \rho_\beta^{\text{eq}} \}$ denotes the quantum relative entropy [50] of ρ_β with respect to ρ_β^{eq} . The relative entropy measures the distinguishability of ρ_β and ρ_β^{eq} . Therefore, the condition that the CTS outperforms the Gibbs state in quantum thermometry at a certain temperature β_0 is

$$\partial_\beta^2 S(\rho_\beta || \rho_\beta^{\text{eq}}) \Big|_{\beta=\beta_0} < 0. \quad (7)$$

Therefore, the curvature of the quantum relative entropy $S(\rho_\beta || \rho_\beta^{\text{eq}})$ with respect to β can be regarded as the criteria quantifying the performance of the CTS for quantum thermometry.

4. Single-qubit example

As a pedagogical example, we show the single-qubit case. Let σ_x, σ_y and σ_z be the Pauli matrices. The Hamiltonian is $H = \omega \sigma_z$, so that the eigenstates of H are $|0\rangle = (1 \ 0)^T$ and $|1\rangle = (0 \ 1)^T$. Therefore, $Z_\beta^{\text{eq}} = \text{tr} \{ e^{-\beta H} \} = 2 \cosh(\beta\omega)$. Considering the pointer states $|\psi_0(\theta)\rangle = e^{i\frac{\theta}{2}\sigma_x}|0\rangle$ and $|\psi_1(\theta)\rangle = e^{i\frac{\theta}{2}\sigma_x}|1\rangle$ with $\theta \in \mathbb{R}$, the normalization factor of the CTS becomes $Z_\beta(\theta) = 2 \cosh(\beta\omega \cos(\theta))$.

Therefore, we obtain from equation (6),

$$\Delta\mathcal{I}_\beta(\theta) = \omega^2 \left(-1 + \frac{\cos^2(\theta)}{\cosh^2(\beta\omega \cos(\theta))} + \tanh^2(\beta\omega) \right). \quad (8)$$

In the low-temperature limit ($\beta\omega \gg 1$), we have

$$\Delta\mathcal{I}_\beta(\theta) \simeq \left(\frac{\omega \cos(\theta)}{\cosh(\beta\omega \cos(\theta))} \right)^2 \geq 0 \quad \forall \theta \in \mathbb{R}. \quad (9)$$

This means that the CTS can outperform the Gibbs state for any choice of pointer states in the low-temperature limit. As an example, choose $\theta = \pi/4$ and $\omega = 1$, which is illustrated in figure 2.

5. Asymmetry

The natural question arises, where exactly this enhanced performance of the CTS in thermometry originates. To this end, we now analyze the relation between the QFI of ρ_β and the asymmetry measure, which is quantified by the WYD skew information $I_\alpha(\rho_\beta, H)$ contained by ρ_β with respect to the Hamiltonian H .

The asymmetry of quantum state is related to its quantum coherence, which is regarded as a resource of breaking the symmetry of a group. This is characterized by the *resource theory of asymmetry* (RTA) [30–34], whose development has also contributed to exploring quantum thermodynamic resources [51–56]. In particular, when we consider time translations $\{\exp(-iHt), |, t \in \mathbb{R}\}$, a state ρ is symmetric if and only if $[\rho, H] = 0$, and asymmetric if and only if $[\rho, H] \neq 0$. Here, the symmetric state is considered the free state,




Figure 2. Relation between ΔI_β and β : We choose $\omega = 1$ and $\theta = \pi/4$. In the low-temperature limit, we have $I(\rho_\beta; \beta) \geq I(\rho_\beta^{\text{eq}}; \beta)$, meaning that the CTS outperforms the Gibbs state in thermometry.

while the asymmetric state is regarded as the resource state. Covariant operations [57], which are the relevant set of free operations, cannot generate an asymmetric state from a symmetric one or convert one asymmetric state into another. An asymmetry measure must satisfy two conditions: (i) it vanishes if and only if the state is symmetric, and (ii) it does not increase under all covariant operations. One quantity satisfying these conditions is the skew information [35–40].

For a quantum state ρ , the WYD skew information is defined as $I_\alpha(\rho, H) \equiv \text{tr}\{\rho H^2\} - \text{tr}\{\rho^\alpha H^{1-\alpha} H\}$ with $0 < \alpha < 1$. With $\text{Var}_\rho\{H\} \equiv \text{tr}\{\rho H^2\} - (\text{tr}\{\rho H\})^2$ the variance of H with respect to ρ , we can define the general variance (covariance) $Q_\alpha(\rho, H)$ as $Q_\alpha(\rho, H) \equiv \text{Var}_\rho\{H\} - I_\alpha(\rho, H)$ [58–60].

Hence, the QFI of ρ_β can be alternatively written as the covariance of the Hamiltonian $H_{12} \equiv H \otimes \mathbb{1}_2$ with respect to the conditional thermal separable state $\tilde{\rho}_\beta$ (3). By using the fact that $\text{Var}_{\tilde{\rho}_\beta}\{H_{12}\} = \text{Var}_{\rho_\beta}\{H\}$ and the asymmetry monotone of WYD skew information $I_\alpha(\tilde{\rho}_\beta, H_{12}) \geq I_\alpha(\rho_\beta, H)$ [35–37], we obtain (see appendix C for the proof.)

$$I(\rho_\beta; \beta) = Q_\alpha(\tilde{\rho}_\beta, H_{12}) \leq Q_\alpha(\rho_\beta, H). \quad (10)$$

Note that when the chosen basis comprises the pointer states of the Hamiltonian, i.e. $\rho_\beta = \rho_\beta^{\text{eq}}$, we have $I_\alpha(\rho_\beta^{\text{eq}}, H) = 0$ because of $[\rho_\beta^{\text{eq}}, H] = 0$. In this case, the upper bound becomes $\text{Var}_{\rho_\beta}\{H\}$, which is exactly the QFI of ρ_β for estimating β , and equation (10) saturates.

These results demonstrate that the asymmetry of the CTS quantifies the maximum ultimate precision limit of the quantum thermometry. In our scenario, since the asymmetry resource contains the quantum coherence over the eigenstates of the Hamiltonian [35], from equation (10), the quantum coherence of the conditional thermal separable state $\tilde{\rho}_\beta$ can be engineered by choosing appropriate pointer states maximizing the QFI of ρ_β .

6. State preparation of the CTS

In the preceding section we have shown that the inherent asymmetry of the CTS can be exploited as a resource. Hence, we now continue with a more formal analysis of the resource-theoretic properties of the CTS arising from the state distinguishability with the Gibbs state. To this end, we consider a quantum system, which is initially prepared in a Gibbs state, $\rho_\beta^{\text{eq}}(0)$, and evolves under a time-dependent Hamiltonian H_t from $t = 0$ to $t = \tau$ with a corresponding unitary U_τ . Our main interest is the state convertibility of the exact final state into the CTS by considering the information-theoretic properties of the averaged excess work.

Let us write $\rho(\tau)$ as the exact final state $\rho(\tau) = U_\tau \rho_\beta^{\text{eq}}(0) U_\tau$. In this case, the exact averaged work $\langle W \rangle$ is given by $\langle W \rangle = \text{tr}\{\rho(\tau) H_\tau\} - \text{tr}\{\rho_\beta^{\text{eq}}(0) H_0\}$. When ΔF^{eq} is the equilibrium free energy difference $\Delta F^{\text{eq}} \equiv -\beta^{-1} \ln(Z_\beta^{\text{eq}}(\tau)/Z_\beta^{\text{eq}}(0))$, the excess work is given by [61, 62]

$$\langle W_{\text{ex}} \rangle \equiv \langle W \rangle - \Delta F^{\text{eq}} = \beta^{-1} S(\rho(\tau) || \rho_\beta^{\text{eq}}(\tau)). \quad (11)$$

We initially measure the system with $H_0 \equiv \sum_{k=1}^d E_k |E_k\rangle\langle E_k|$, where $\{|E_k\rangle\}_{k=1}^d$ are the eigenstates of H_0 . The post-measurement state of the system after the evolution is $U_\tau |E_k\rangle$. Thus, the CTS constructed from the

post-measurement state for the final Hamiltonian $H_\tau \equiv \sum_{k=1}^d E'_k |E'_k\rangle\langle E'_k|$ is given by [24]

$$\rho_\beta(\tau) = \sum_{k=1}^d \frac{e^{-\beta\langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)} U_\tau |E_k\rangle\langle E_k| U_\tau^\dagger, \quad (12)$$

where $Z_\beta(\tau)$ is again the conditional partition function.

In [24] it was shown that the excess work is lower bounded by

$$\langle W_{\text{ex}} \rangle \geq \beta^{-1} S(\rho_\beta(\tau) || \rho_\beta^{\text{eq}}(\tau)). \quad (13)$$

Therefore, from equations (11) and (13), we have

$$S(\rho(\tau) || \rho_\beta^{\text{eq}}(\tau)) \geq S(\rho_\beta(\tau) || \rho_\beta^{\text{eq}}(\tau)). \quad (14)$$

From [63–65], equation (14) becomes tight if and only if there exists a sufficiently large n_0 and there exists a Gibbs-preserving map $\mathcal{E}_n : \mathcal{H}^{\otimes n} \rightarrow \mathcal{H}^{\otimes n}$ for $n \geq n_0$ such that

$$\mathcal{E}_n(\rho_\beta^{\text{eq}}(\tau)^{\otimes n}) = \rho_\beta^{\text{eq}}(\tau)^{\otimes n}, \quad \mathcal{E}_n(\rho(\tau)^{\otimes n}) = \Xi_n, \quad (15)$$

where for any $\epsilon > 0$, the quantum state Ξ_n satisfies

$$T_n \equiv \frac{1}{2} \|\rho_\beta(\tau)^{\otimes n} - \Xi_n\|_1 < \epsilon. \quad (16)$$

This means that the CTS is not just a mathematical object but can be prepared, allowing for an arbitrarily small error via the Gibbs-preserving map with an *asymptotic* number of copies of the state.

Also, from the quantum relative entropy of the exact final state $\rho(\tau)$ with respect to the CTS $\rho_\beta(\tau)$ $S(\rho(\tau) || \rho_\beta(\tau))$, we obtain (see appendix D for the proof.)

$$S(\rho(\tau) || \rho_\beta(\tau)) + S(\rho_\beta(\tau) || \rho_\beta^{\text{eq}}(\tau)) = S(\rho(\tau) || \rho_\beta^{\text{eq}}(\tau)), \quad (17)$$

which we call *thermodynamic triangle equality*. Therefore, when $S(\rho(\tau) || \rho_\beta^{\text{eq}}(\tau)) = S(\rho_\beta(\tau) || \rho_\beta^{\text{eq}}(\tau))$, we must have $\rho(\tau) = \rho_\beta(\tau)$ (i.e. $U_\tau^\dagger H_\tau U_\tau = H_0$).

To analyze the scaling of the required number of copies n given an arbitrarily small error ϵ , we employ the inequality $S(\rho_\beta(\tau)^{\otimes n} || \Xi_n) \leq (T_n + \lambda_n) \ln(1 + T_n / \lambda_n)$ [66], where λ_n denotes the minimum non-zero eigenvalue of Ξ_n . Since $d = \text{rank}(\rho_\beta(\tau))$, we can write $\lambda_n \sim 1/d^n$ [66]. Therefore, when $T_n < \epsilon \ll 1$, we have

$$\begin{aligned} S(\rho_\beta(\tau)^{\otimes n} || \Xi_n) &\lesssim \left(\epsilon + \frac{1}{d^n} \right) \ln(1 + \epsilon d^n) \\ &= \epsilon \left(1 + \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k(k+1)} \epsilon^k d^{kn} \right) \\ &= \epsilon \left(1 + \frac{d^n}{2} \epsilon + O(\epsilon^2 d^{2n}) \right). \end{aligned} \quad (18)$$

To guarantee $\lim_{n \rightarrow 0} S(\rho_\beta(\tau)^{\otimes n} || \Xi_n) = 0$, the simplest condition is $\epsilon d^n \sim 1$. Therefore, the required number of copies of $\rho_\beta(\tau)$ can be approximately given by

$$n \sim \log_d \left(\frac{1}{\epsilon} \right). \quad (19)$$

Here, we note that this state preparation protocol inherently depends on an unknown parameter. Therefore, in the context of thermometry, this protocol remains limited. We need to further explore state preparation protocols that are independent of such parameters.

7. Quantum heat from the CTS

Furthermore, the symmetric divergence between $\rho(\tau)$ and $\rho_\beta(\tau)$, i.e. the quantum J-divergence defined as [67],

$$J(\rho(\tau), \rho_\beta(\tau)) \equiv S(\rho(\tau) \parallel \rho_\beta(\tau)) + S(\rho_\beta(\tau) \parallel \rho(\tau)), \quad (20)$$

is related to the concept of quantum heat [41]. This becomes obvious when the quantum J-divergence is written as (see appendix E for the proof.)

$$J(\rho(\tau), \rho_\beta(\tau)) = \beta(\langle W \rangle - \mathcal{W}_0(\rho_\beta(\tau)) - \Delta E(\rho_\beta(\tau))). \quad (21)$$

Here, the internal energy change of $\rho_\beta(\tau)$ is $\Delta E(\rho_\beta(\tau)) \equiv \text{tr}\{\rho_\beta(\tau)(H_\tau - H_0)\}$, and the quantum ergotropy of $\rho_\beta(\tau)$ with respect to H_0 is $\mathcal{W}_0(\rho_\beta(\tau)) \equiv \text{tr}\{\rho_\beta(\tau)H_0\} - \text{tr}\{\Gamma\rho_\beta(\tau)\Gamma^\dagger H_0\}$, where $\Gamma \equiv \sum_{k=1}^d |E_k\rangle\langle E_k|U_\tau^\dagger = U_\tau^\dagger$ is the ergotropic transformation [68]. From the first law of thermodynamics, $\beta^{-1}J(\rho(\tau), \rho_\beta(\tau))$ can be regarded as a heat, particularly the *quantum heat*, which has been discussed in the literature as the heat induced by the measurement [41]. In essence, the CTS $\rho_\beta(\tau)$ is conditioned on the first energy measurement outcome; therefore, its relation to the quantum heat is consistent in this context. Also, note that when U_τ is an adiabatic passage, then $\rho(\tau) = \rho_\beta(\tau) = \rho_\beta^{\text{eq}}(\tau)$, so that $\beta^{-1}J(\rho(\tau), \rho_\beta(\tau)) = 0$, which is consistent with zero heat exchange in the adiabatic process.

8. Conclusion

In conclusion, we have introduced a conditional thermal state, which is a thermal state conditioned on the pointer states, and demonstrated that this conditional thermal state can outperform the Gibbs state in the quantum thermometry as a useful resource state. We also have explored its resource-theoretic properties in terms of the asymmetry, state convertibility and its relation to the quantum heat. As an application, these results could help experimentalists to achieve a better sensitivity in quantum thermometry under the constraints in the implementable measurements. From the fundamental point of view, these results provide insightful perspective on the implications of the conditional thermal state in the thermodynamic protocols in the quantum systems in a resource-theoretic approach. Finally, the present analysis also provides an additional *a posteriori* motivation and justification for the one-time energy measurement approach to quantum work [24–29]. Future work will involve further exploration of concrete state preparation protocols for generating the conditional thermal state.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgments

We would like to thank to Ryuji Takagi for helpful discussions. A S gratefully acknowledges startup funding supported by the University of Massachusetts, Boston. D O S P acknowledges the Brazilian funding agencies CNPq (Grant No. 307028/2019-4), FAPESP (Grant No. 2017/03727-0), and the Brazilian National Institute of Science and Technology of Quantum Information (INCT-IQ) Grant No. 465469/2014-0. S D acknowledges support from the John Templeton Foundation under Grant No. 62422.

Appendix A. Derivation of conditional thermal state

Consider a density matrix ρ with the following spectral decomposition

$$\rho = \sum_{k=1}^d p_k |\psi_k\rangle\langle\psi_k|, \quad (A1)$$

where $\{|\psi_k\rangle\}_{k=1}^d$ are the pointer states of a given measurement M . Let H be the Hamiltonian of the system. Then, we want to maximize the von Neumann entropy

$$S(\rho) \equiv -\text{tr}\{\rho \ln \rho\} \quad (A2)$$

under the conditions

$$\text{tr}\{\rho\} = 1 \quad \text{and} \quad E(\rho) \equiv \text{tr}\{\rho H\} = \text{Const.} \quad (\text{A3})$$

By using the optimization method of Lagrange multipliers with constraints, we have:

$$\delta(S(\rho) - \gamma E(\rho) - \alpha) = - \sum_{k=1}^d \delta p_k (\ln p_k + \beta \langle \psi_k | H | \psi_k \rangle + \alpha + 1) = 0. \quad (\text{A4})$$

For any δp_k , equation (A4) has to be valid so that each term has to be independently 0. Therefore, we must have $p_k \propto e^{-\beta \langle \psi_k | H | \psi_k \rangle}$. From the requirement of $\sum_{k=1}^d p_k = 1$, we obtain the CTS ρ_β .

Appendix B. Detailed derivations of equations (4)–(7)

From the definition of QFI, we have

$$\mathcal{I}(\rho_\theta; \theta) = -2 \lim_{\epsilon \rightarrow 0} \partial_\epsilon^2 \mathcal{F}(\rho_\theta, \rho_{\theta+\epsilon}) \quad (\text{B1})$$

where

$$\begin{aligned} \mathcal{F}(\rho_\beta, \rho_{\beta+\epsilon}) &\equiv \|\sqrt{\rho_\beta} \sqrt{\rho_{\beta+\epsilon}}\|_1^2 \\ &= \left(\text{tr} \left\{ \sqrt{\rho_\beta^{1/2} \rho_{\beta+\epsilon} \rho_\beta^{1/2}} \right\} \right)^2 \\ &= \frac{1}{Z_\beta Z_{\beta+\epsilon}} \left(\sum_{k=1}^d e^{-\left(\beta + \frac{\epsilon}{2}\right) \langle \psi_k | H | \psi_k \rangle} \right)^2 \\ &= \frac{Z_{\beta+\epsilon/2}^2}{Z_\beta Z_{\beta+\epsilon}}. \end{aligned} \quad (\text{B2})$$

Before calculating the quantum Fisher information, let us show the following fact:

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \partial_\epsilon Z_{\beta+\epsilon} &= - \sum_{k=1}^d e^{-\beta \langle \psi_k | H | \psi_k \rangle} \langle \psi_k | H | \psi_k \rangle = \partial_\beta Z_\beta \\ \lim_{\epsilon \rightarrow 0} \partial_\epsilon^2 Z_{\beta+\epsilon} &= \sum_{k=1}^d e^{-\beta \langle \psi_k | H | \psi_k \rangle} \langle \psi_k | H | \psi_k \rangle^2 = \partial_\beta^2 Z_\beta \\ \lim_{\epsilon \rightarrow 0} \partial_\epsilon Z_{\beta+\epsilon/2} &= - \frac{1}{2} \sum_{k=1}^d e^{-\beta \langle \psi_k | H | \psi_k \rangle} \langle \psi_k | H | \psi_k \rangle = \frac{1}{2} \partial_\beta Z_\beta \\ \lim_{\epsilon \rightarrow 0} \partial_\epsilon^2 Z_{\beta+\epsilon/2} &= \frac{1}{4} \sum_{k=1}^d e^{-\beta \langle \psi_k | H | \psi_k \rangle} \langle \psi_k | H | \psi_k \rangle^2 = \frac{1}{4} \partial_\beta^2 Z_\beta. \end{aligned} \quad (\text{B3})$$

For two functions $f(x)$ and $g(x)$, where $g(x) \neq 0$, we have:

$$\partial_x^2 \left(\frac{f^2}{g} \right) = 2 \left(\partial_x^2 f \right) \frac{f}{g} + \frac{2}{g} (\partial_x f)^2 - \frac{4f}{g^2} (\partial_x f) (\partial_x g) - \left(\partial_x^2 g \right) \frac{f^2}{g^2} + \frac{2f^2}{g^3} (\partial_x g)^2.$$

Therefore, if we define $x = \epsilon$, $f = Z_{\beta+\epsilon/2}$ and $g = Z_{\beta+\epsilon}$, we can obtain:

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \partial_\epsilon^2 F(\rho_\beta, \rho_{\beta+\epsilon}) &= \lim_{\epsilon \rightarrow 0} \partial_\epsilon^2 \left(\frac{Z_{\beta+\epsilon/2}^2}{Z_\beta Z_{\beta+\epsilon}} \right) \\ &= \frac{1}{2} \frac{\partial_\beta^2 Z_\beta}{Z_\beta} + \frac{1}{2} \left(\frac{\partial_\beta Z_\beta}{Z_\beta} \right)^2 - 2 \left(\frac{\partial_\beta Z_\beta}{Z_\beta} \right)^2 - \frac{\partial_\beta^2 Z_\beta}{Z_\beta} + 2 \left(\frac{\partial_\beta Z_\beta}{Z_\beta} \right)^2 \\ &= \frac{1}{2} \left(\frac{\partial_\beta Z_\beta}{Z_\beta} \right)^2 - \frac{1}{2} \frac{\partial_\beta^2 Z_\beta}{Z_\beta} \\ &= \frac{1}{2} \left((\partial_\beta \ln Z_\beta)^2 - Z_\beta^{-1} \partial_\beta^2 Z_\beta \right). \end{aligned} \quad (\text{B4})$$

Therefore, from equation (B1), we can obtain

$$\mathcal{I}(\rho_\beta; \beta) = Z_\beta^{-1} \partial_\beta^2 Z_\beta - (\partial_\beta \ln Z_\beta)^2 = \partial_\beta^2 \ln Z_\beta. \quad (\text{B5})$$

For the optimal measurement, we need to prove that the uncertainty of the unbiased estimator with the measurement

$$G = \sum_{k=1}^d \langle \psi_k | H | \psi_k \rangle | \psi_k \rangle \langle \psi_k | \quad (\text{B6})$$

achieves the quantum Cramér–Rao bound. The uncertainty is given by

$$(\delta\beta)_G^2 \equiv \frac{\text{Var}_{\rho_\beta}\{G\}}{|\partial_\beta \text{tr}\{\rho_\beta G\}|^2}. \quad (\text{B7})$$

Here, we have

$$\text{tr}\{\rho_\beta G\} = \sum_{k=1}^d \frac{e^{-\beta \langle \psi_k | H | \psi_k \rangle}}{Z_\beta} \langle \psi_k | H | \psi_k \rangle = \text{tr}\{\rho_\beta H\} = -\partial_\beta \ln Z_\beta \quad (\text{B8})$$

and

$$\begin{aligned} \text{Var}_{\rho_\beta}\{G\} &\equiv \text{tr}\{\rho_\beta H^2\} - (\text{tr}\{\rho_\beta H\})^2 \\ &= \sum_{k=1}^d \frac{e^{-\beta \langle \psi_k | H | \psi_k \rangle}}{Z_\beta} \langle \psi_k | H | \psi_k \rangle^2 - (\partial_\beta \ln Z_\beta)^2 \\ &= Z_\beta^{-1} \partial_\beta^2 Z_\beta - (\partial_\beta \ln Z_\beta)^2 \\ &= \partial_\beta^2 \ln Z_\beta. \end{aligned} \quad (\text{B9})$$

From equations (B5) and (B7), we can obtain

$$(\delta\beta)_G^2 = \frac{1}{\partial_\beta^2 \ln Z_\beta} = \frac{1}{\mathcal{I}(\rho_\beta; \beta)}, \quad (\text{B10})$$

which is the quantum Cramér–Rao bound; therefore, equation (B6) is the optimal measurement for the quantum thermometry.

From equation (B5), we can write

$$\Delta\mathcal{I}_\beta \equiv \mathcal{I}(\rho_\beta; \beta) - \mathcal{I}\left(\rho_\beta^{\text{eq}}; \beta\right) = \partial_\beta^2 \ln \frac{Z_\beta}{Z_\beta^{\text{eq}}}. \quad (\text{B11})$$

Since the von Neumann entropy $S(\rho_\beta) \equiv -\text{tr}\{\rho_\beta \ln \rho_\beta\}$ is

$$S(\rho_\beta) = \beta \text{tr}\{\rho_\beta H\} + \ln Z_\beta, \quad (\text{B12})$$

the quantum relative entropy $S(\rho_\beta || \rho_\beta^{\text{eq}})$ becomes

$$S\left(\rho_\beta || \rho_\beta^{\text{eq}}\right) = -S(\rho_\beta) - \text{tr}\left\{\rho_\beta \ln \rho_\beta^{\text{eq}}\right\} = -\ln \frac{Z_\beta}{Z_\beta^{\text{eq}}}. \quad (\text{B13})$$

Therefore, we can write

$$\Delta\mathcal{I}_\beta = -\partial_\beta^2 S\left(\rho_\beta || \rho_\beta^{\text{eq}}\right). \quad (\text{B14})$$

When the CTS outperforms the Gibbs state for the quantum thermometry at $\beta = \beta_0$, i.e.

$$\Delta\mathcal{I}_{\beta_0} > 0, \quad (\text{B15})$$

we must have

$$\partial_\beta^2 S\left(\rho_\beta || \rho_\beta^{\text{eq}}\right) \Big|_{\beta=\beta_0} < 0. \quad (\text{B16})$$

Appendix C. Proof of equation (10)

First, let us prove $\mathcal{I}(\rho_\beta; \beta) = Q_\alpha(\tilde{\rho}_\beta, H_{12})$, where

$$H_{12} \equiv H \otimes \mathbb{1}_2. \quad (\text{C1})$$

Since

$$\tilde{\rho}_\beta \equiv \sum_{k=1}^d \frac{e^{-\beta \langle \psi_k | H | \psi_k \rangle}}{Z_\beta} |\psi_k\rangle\langle\psi_k|_1 \otimes |\phi_k\rangle\langle\phi_k|_2 \quad (\text{C2})$$

and

$$\rho_\beta = \text{tr}_2 \{ \tilde{\rho}_\beta \}, \quad (\text{C3})$$

we have

$$\text{tr} \{ \tilde{\rho}_\beta H_{12} \} = \text{tr} \{ \rho_\beta H \} = -\partial_\beta \ln Z_\beta \quad (\text{C4})$$

and

$$\text{tr} \{ \tilde{\rho}_\beta H_{12}^2 \} = \text{tr} \{ \rho_\beta H^2 \}, \quad (\text{C5})$$

so that

$$\text{Var}_{\tilde{\rho}_\beta} \{ H_{12} \} = \text{Var}_{\rho_\beta} \{ H \}. \quad (\text{C6})$$

Next, let us compute the WYD skew information $I_\alpha(\tilde{\rho}_\beta, H)$ contained by $\tilde{\rho}_\beta$ with respect to H_{12} . By using the fact that for a pure state $|\varphi\rangle$ we have $|\varphi\rangle\langle\varphi|^\alpha = |\varphi\rangle\langle\varphi|$ for any α , we can obtain

$$\begin{aligned} I_\alpha(\tilde{\rho}_\beta, H_{12}) &= \text{tr} \{ \tilde{\rho}_\beta H_{12}^2 \} - \text{tr} \{ \tilde{\rho}_\beta^\alpha H_{12} \tilde{\rho}_\beta^{1-\alpha} H_{12} \} \\ &= \text{tr} \{ \tilde{\rho}_\beta H_{12}^2 \} - \sum_{k=1}^d \frac{e^{-\beta \langle \psi_k | H | \psi_k \rangle}}{Z_\beta} \langle \psi_k | H | \psi_k \rangle^2 \\ &= \text{tr} \{ \tilde{\rho}_\beta H_{12}^2 \} - Z_\beta^{-1} \partial_\beta^2 Z_\beta. \end{aligned} \quad (\text{C7})$$

Therefore, we can obtain

$$\begin{aligned} Q_\alpha(\tilde{\rho}_\beta, H_{12}) &\equiv \text{Var}_{\tilde{\rho}_\beta} \{ H \} - I_\alpha(\tilde{\rho}_\beta, H_{12}) \\ &= Z_\beta^{-1} \partial_\beta^2 Z_\beta - (\partial_\beta \ln Z_\beta)^2 \\ &= \partial_\beta^2 \ln Z_\beta. \end{aligned} \quad (\text{C8})$$

From equation (B5), we can finally write

$$\mathcal{I}(\rho_\beta; \beta) = Q_\alpha(\tilde{\rho}_\beta, H_{12}). \quad (\text{C9})$$

Next, let us prove $I(\rho_\beta; \beta) \leq Q_\alpha(\rho_\beta, H)$. The WYD skew information has asymmetry monotone [36]

$$I_\alpha(\tilde{\rho}_\beta, H_{12}) \geq I_\alpha(\rho_\beta, H). \quad (\text{C10})$$

From equation (C6), we can obtain

$$Q_\alpha(\tilde{\rho}_\beta, H_{12}) \leq Q_\alpha(\rho_\beta, H) = \text{Var}_{\rho_\beta} \{ H \} - I_\alpha(\rho_\beta, H). \quad (\text{C11})$$

Because of equation (C9), we can obtain

$$\mathcal{I}(\rho_\beta; \beta) \leq Q_\alpha(\rho_\beta, H). \quad (\text{C12})$$

Appendix D. Proof of equation (17)

The quantum relative entropy $S(\rho(\tau) \parallel \rho_\beta(\tau))$ can be written as

$$\begin{aligned} S(\rho(\tau) \parallel \rho_\beta(\tau)) &= -S(\rho(\tau)) - \text{tr}\{\rho(\tau) \ln \rho_\beta(\tau)\} \\ &= -S(\rho(\tau)) + \beta \sum_{k=1}^d \langle E_k | U_\tau^\dagger \rho(\tau) U_\tau | E_k \rangle \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle + \ln Z_\beta(\tau). \end{aligned} \quad (\text{D1})$$

Because

$$\rho(\tau) = U_\tau \rho_\beta^{\text{eq}}(0) U_\tau^\dagger = \sum_{k=1}^d \frac{e^{-\beta E_k}}{Z_\beta^{\text{eq}}(0)} U_\tau |E_k\rangle\langle E_k| U_\tau^\dagger, \quad (\text{D2})$$

we have

$$\sum_{k=1}^d \langle E_k | U_\tau^\dagger \rho(\tau) U_\tau | E_k \rangle \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle = \text{tr} \left\{ U_\tau \rho_\beta^{\text{eq}}(0) U_\tau^\dagger H_\tau \right\} = \text{tr} \{ \rho(\tau) H_\tau \}. \quad (\text{D3})$$

Therefore,

$$S(\rho(\tau) \parallel \rho_\beta(\tau)) = -S(\rho(\tau)) + \beta \text{tr} \{ \rho(\tau) H_\tau \} + \ln Z_\beta(\tau). \quad (\text{D4})$$

From the unitary invariance of the von Neumann entropy, we can write

$$\begin{aligned} S(\rho(\tau) \parallel \rho_\beta(\tau)) &= -S(\rho_\beta^{\text{eq}}(0)) + \ln Z_\beta(\tau) + \beta \text{tr} \{ \rho(\tau) H_\tau \} \\ &= \beta \left(\text{tr} \{ \rho(\tau) H_\tau \} - \text{tr} \left\{ \rho_\beta^{\text{eq}}(0) H_0 \right\} \right) + \ln \frac{Z_\beta(\tau)}{Z_\beta^{\text{eq}}(0)} \\ &= \beta \langle W \rangle + \ln \frac{Z_\beta(\tau)}{Z_\beta^{\text{eq}}(0)}, \end{aligned} \quad (\text{D5})$$

where

$$\langle W \rangle \equiv \text{tr} \{ \rho(\tau) H_\tau \} - \text{tr} \left\{ \rho_\beta^{\text{eq}}(0) H_0 \right\} \quad (\text{D6})$$

is the exact averaged work. Because

$$\ln \frac{Z_\beta(\tau)}{Z_\beta^{\text{eq}}(0)} = \ln \frac{Z_\beta(\tau)}{Z_\beta^{\text{eq}}(\tau)} + \ln \frac{Z_\beta^{\text{eq}}(\tau)}{Z_\beta^{\text{eq}}(0)}, \quad (\text{D7})$$

from [24], we have

$$\ln \frac{Z_\beta(\tau)}{Z_\beta^{\text{eq}}(0)} = -S(\rho_\beta(\tau) \parallel \rho_\beta^{\text{eq}}(\tau)) - \beta \Delta F^{\text{eq}}. \quad (\text{D8})$$

Therefore, we have

$$S(\rho(\tau) \parallel \rho_\beta(\tau)) + S(\rho_\beta(\tau) \parallel \rho_\beta^{\text{eq}}(\tau)) = \beta (\langle W \rangle - \Delta F^{\text{eq}}). \quad (\text{D9})$$

Because the exact excess work is [62]

$$\langle W_{\text{ex}} \rangle = \langle W \rangle - \Delta F^{\text{eq}} = \beta^{-1} S(\rho(\tau) \parallel \rho_\beta^{\text{eq}}(\tau)), \quad (\text{D10})$$

we can obtain the thermodynamic triangle equality

$$S(\rho(\tau) \parallel \rho_\beta(\tau)) + S(\rho_\beta(\tau) \parallel \rho_\beta^{\text{eq}}(\tau)) = S(\rho(\tau) \parallel \rho_\beta^{\text{eq}}(\tau)). \quad (\text{D11})$$

Appendix E. Proof of equation (21)

The quantum J-divergence of $\rho(\tau)$ and $\rho_\beta(\tau)$ is defined as

$$J(\rho(\tau), \rho_\beta(\tau)) \equiv S(\rho(\tau) || \rho_\beta(\tau)) + S(\rho_\beta(\tau) || \rho(\tau)). \quad (\text{E1})$$

In equations (D9) and (D11), we have already proved the thermodynamic triangle equality. Next, let us consider the quantum relative entropy $S(\rho_\beta(\tau) || \rho(\tau))$, which is given by

$$S(\rho_\beta(\tau) || \rho(\tau)) = -S(\rho_\beta(\tau)) - \text{tr}\{\rho_\beta(\tau) \ln \rho(\tau)\}. \quad (\text{E2})$$

The von Neumann entropy of $\rho_\beta(\tau)$ is

$$S(\rho_\beta(\tau)) = \beta \text{tr}\{\rho_\beta(\tau) H_\tau\} + \ln Z_\beta(\tau). \quad (\text{E3})$$

Also

$$\begin{aligned} \text{tr}\{\rho_\beta(\tau) \ln \rho(\tau)\} &= -\beta \sum_{k=1}^d E_k \langle E_k | U_\tau^\dagger \rho_\beta(\tau) U_\tau | E_k \rangle - \ln Z_\beta^{\text{eq}}(0) \\ &= -\beta \sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)} - \ln Z_\beta^{\text{eq}}(0). \end{aligned} \quad (\text{E4})$$

Therefore,

$$S(\rho_\beta(\tau) || \rho(\tau)) = \beta \left(\sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)} - \text{tr}\{\rho_\beta(\tau) H_\tau\} \right) - \ln \frac{Z_\beta(\tau)}{Z_\beta^{\text{eq}}(0)}. \quad (\text{E5})$$

From equation (D8), we can obtain

$$S(\rho_\beta(\tau) || \rho(\tau)) - S(\rho_\beta(\tau) || \rho_\beta^{\text{eq}}(\tau)) - \beta \Delta F_\beta^{\text{eq}} = \beta \left(\sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)} - \text{tr}\{\rho_\beta(\tau) H_\tau\} \right). \quad (\text{E6})$$

By using $\text{tr}\{\rho_\beta(\tau) H_0\}$, we can write

$$\begin{aligned} &\sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)} - \text{tr}\{\rho_\beta(\tau) H_\tau\} \\ &= \sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)} - \text{tr}\{\rho_\beta(\tau) H_0\} - \text{tr}\{\rho_\beta(\tau) (H_\tau - H_0)\}. \end{aligned} \quad (\text{E7})$$

Let Γ be the ergotropic transformation

$$\Gamma \equiv \sum_{k=1}^d |E_k\rangle\langle E_k| U_\tau^\dagger = U_\tau^\dagger. \quad (\text{E8})$$

Then, we have

$$\text{tr}\{\Gamma \rho_\beta(\tau) \Gamma^\dagger H_0\} = \sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)}. \quad (\text{E9})$$

The ergotropy of $\rho_\beta(\tau)$ with respect to the Hamiltonian H_0 is defined as [68]

$$\mathcal{W}_0(\rho_\beta(\tau)) \equiv \text{tr}\{\rho_\beta(\tau) H_0\} - \text{tr}\{\Gamma \rho_\beta(\tau) \Gamma^\dagger H_0\} = \text{tr}\{\rho_\beta(\tau) H_0\} - \sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)}. \quad (\text{E10})$$

Therefore, by defining

$$\Delta E(\rho_\beta(\tau)) \equiv \text{tr}\{\rho_\beta(\tau) (H_\tau - H_0)\}, \quad (\text{E11})$$

we can write

$$\sum_{k=1}^d E_k \frac{e^{-\beta \langle E_k | U_\tau^\dagger H_\tau U_\tau | E_k \rangle}}{Z_\beta(\tau)} - \text{tr}\{\rho_\beta(\tau) H_\tau\} = -\mathcal{W}_0(\rho_\beta(\tau)) - \Delta E(\rho_\beta(\tau)). \quad (\text{E12})$$

Therefore,

$$S(\rho_\beta(\tau) || \rho(\tau)) - S(\rho_\beta(\tau) || \rho_\beta^{\text{eq}}(\tau)) = \beta(\Delta F^{\text{eq}} - \mathcal{W}_0(\rho_\beta(\tau)) - \Delta E(\rho_\beta(\tau))). \quad (\text{E13})$$

From equations (E1), (D9) and (E13), we can obtain

$$J(\rho(\tau), \rho_\beta(\tau)) = \beta(\langle W \rangle - \mathcal{W}_0(\rho_\beta(\tau)) - \Delta E(\rho_\beta(\tau))). \quad (\text{E14})$$

ORCID iDs

Akira Sone  <https://orcid.org/0000-0003-3539-6140>

Diogo O Soares-Pinto  <https://orcid.org/0000-0002-4293-6144>

Sebastian Deffner  <https://orcid.org/0000-0003-0504-6932>

References

- [1] Giovannetti V, Lloyd S and Maccone L 2004 Quantum-enhanced measurements: beating the standard quantum limit *Science* **306** 1330
- [2] Giovannetti V, Lloyd S and Maccone L 2006 Quantum metrology *Phys. Rev. Lett.* **96** 010401
- [3] Giovannetti V, Lloyd S and Maccone L 2011 Advances in quantum metrology *Nat. Photon.* **5** 222
- [4] Degen C L, Reinhard F and Cappellaro P 2017 Quantum sensing *Rev. Mod. Phys.* **89** 035002
- [5] Rubio J, Knott P and Dunningham J 2018 Non-asymptotic analysis of quantum metrology protocols beyond the Cramér–Rao bound *J. Phys. Commun.* **2** 015027
- [6] Rubio J and Dunningham J 2019 Quantum metrology in the presence of limited data *New J. Phys.* **21** 043037
- [7] Zurek W H 1981 Pointer basis of quantum apparatus: into what mixture does the wave packet collapse? *Phys. Rev. D* **24** 1516
- [8] Schlosshauer M A 2007 *Decoherence: and the Quantum-to-Classical Transition* (Springer Science & Business Media)
- [9] Brasil C A and de Castro L A 2015 Understanding the pointer states *Eur. J. Phys.* **36** 065024
- [10] Touil A, Yan B, Girolami D, Deffner S and Zurek W H 2022 Eavesdropping on the decohering environment: quantum darwinism, amplification and the origin of objective classical reality *Phys. Rev. Lett.* **128** 010401
- [11] Zurek W H 2003 Decoherence, einselection and the quantum origins of the classical *Rev. Mod. Phys.* **75** 715
- [12] Correa L A, Mehboudi M, Adesso G and Sanpera A 2015 Individual quantum probes for optimal thermometry *Phys. Rev. Lett.* **114** 220405
- [13] Pasquale A D, Rossini D, Fazio R and Giovannetti V 2016 Local quantum thermal susceptibility *Nat. Commun.* **7** 12782
- [14] Sone A, Zhuang Q and Cappellaro P 2018 Quantifying precision loss in local quantum thermometry via diagonal discord *Phys. Rev. A* **98** 012115
- [15] Sone A, Zhuang Q, Li C, Liu Y-X and Cappellaro P 2019 Nonclassical correlations for quantum metrology in thermal equilibrium *Phys. Rev. A* **99** 052318
- [16] Brunelli M, Olivares S, Paternostro M and Paris M G 2012 Qubit-assisted thermometry of a quantum harmonic oscillator *Phys. Rev. A* **86** 012125
- [17] Mancino L, Sbroscia M, Gianani I, Roccia E and Barbieri M 2017 Quantum simulation of single-qubit thermometry using linear optics *Phys. Rev. Lett.* **118** 130502
- [18] Campbell S, Genoni M G and Deffner S 2018 Precision thermometry and the quantum speed limit *Quantum Sci. Technol.* **3** 025002
- [19] Mehboudi M, Moreno-Cardoner M, De Chiara G and Sanpera A 2015 Thermometry precision in strongly correlated ultracold lattice gases *New J. Phys.* **17** 055020
- [20] Pasquale A D and Stace T M 2018 Quantum thermometry *Thermodynamics in the Quantum Regime* (Springer) pp 503–27
- [21] Mehboudi M, Sanpera A and Correa L A 2019 Thermometry in the quantum regime: recent theoretical progress *J. Phys. A: Math. Theor.* **52** 303001
- [22] Narasimhachar V and Gour G 2015 Low-temperature thermodynamics with quantum coherence *Nat. Commun.* **6** 7689
- [23] Correa L A, Perarnau-Llobet M, Hovhannyan K V, Hernández-Santana S, Mehboudi M and Sanpera A 2017 Low-temperature thermometry can be enhanced by strong coupling *Phys. Rev. A* **96** 062103
- [24] Deffner S, Paz J P and Zurek W H 2016 Quantum work and the thermodynamic cost of quantum measurements *Phys. Rev. E* **94** 010103(R)
- [25] Sone A, Liu Y-X and Cappellaro P 2020 Quantum Jarzynski equality in open quantum systems from the one-time measurement scheme *Phys. Rev. Lett.* **125** 060602
- [26] Sone A and Deffner S 2021 Jarzynski equality for stochastic conditional work *J. Stat. Phys.* **183** 11
- [27] Beyer K, Luoma K and Strunz W T 2020 Work as an external quantum observable and an operational quantum work fluctuation theorem *Phys. Rev. Res.* **2** 033508
- [28] Sone A, Soares-Pinto D O and Deffner S 2023 Exchange fluctuation theorems for strongly interacting quantum pumps *AVS Quantum Sci.* **5** 032001
- [29] Sone A, Yamamoto N, Holdsworth T and Narang P 2023 Jarzynski-like equality of nonequilibrium information production based on quantum cross-entropy *Phys. Rev. Res.* **5** 023039
- [30] Chitambar E and Gour G 2019 Quantum resource theories *Rev. Mod. Phys.* **91** 025001
- [31] Marvian I and Spekkens R W 2014 Asymmetry properties of pure quantum states *Phys. Rev. A* **90** 014102

[32] Gour G and Spekkens R W 2008 The resource theory of quantum reference frames: manipulations and monotones *New J. Phys.* **10** 033023

[33] Marvian I and Spekkens R W 2013 The theory of manipulations of pure state asymmetry: I. Basic tools, equivalence classes and single copy transformations *New J. Phys.* **15** 033001

[34] Gour G, Marvian I and Spekkens R W 2009 Measuring the quality of a quantum reference frame: the relative entropy of frameness *Phys. Rev. A* **80** 012307

[35] Marvian I and Spekkens R W 2014 Extending Noether's theorem by quantifying the asymmetry of quantum states *Nat. Commun.* **5** 3821

[36] Takagi R 2019 Skew informations from an operational view via resource theory of asymmetry *Sci. Rep.* **9** 14562

[37] Li W 2019 Monotonicity of skew information and its applications in quantum resource theory *Quantum Inf. Process.* **18** 1

[38] Yamaguchi K and Tajima H 2023 Smooth metric adjusted skew information rates *Quantum* **7** 1012

[39] Marvian I, Spekkens R W and Zanardi P 2016 Quantum speed limits, coherence and asymmetry *Phys. Rev. A* **93** 052331

[40] Ahmadi M, Jennings D and Rudolph T 2013 The Wigner-Araki-Yanase theorem and the quantum resource theory of asymmetry *New J. Phys.* **15** 013057

[41] Elouard C, Herrera-Martí D A, Clusel M and Auffèves A 2017 The role of quantum measurement in stochastic thermodynamics *npj Quantum Inf.* **3** 9

[42] Jaynes E T 1957 Information theory and statistical mechanics *Phys. Rev.* **106** 620

[43] Liu J, Yuan H, Lu X-M and Wang X 2020 Quantum Fisher information matrix and multiparameter estimation *J. Phys. A: Math. Theor.* **53** 023001

[44] Fiderer L J, Fraïssé J M and Braun D 2019 Maximal quantum Fisher information for mixed states *Phys. Rev. Lett.* **123** 250502

[45] Jacobs K 2014 *Quantum Measurement Theory and its Applications* (Cambridge University Press)

[46] Sone A, Cerezo M, Beckey J L and Coles P J 2021 Generalized measure of quantum Fisher information *Phys. Rev. A* **104** 062602

[47] Cerezo M, Sone A, Beckey J L and Coles P J 2021 Sub-quantum Fisher information *Quantum Sci. Technol.* **6** 035008

[48] Beckey J L, Cerezo M, Sone A and Coles P J 2022 Variational quantum algorithm for estimating the quantum Fisher information *Phys. Rev. Res.* **4** 013083

[49] Jozsa R 1994 Fidelity for mixed quantum states *J. Mod. Opt.* **41** 2315

[50] Nielsen M A and Chuang I L 2010 *Quantum Computation and Quantum Information: 10th Anniversary edn* (Cambridge University Press)

[51] Gour G, Müller M P, Narasimhachar V, Spekkens R W and Halpern N Y 2015 The resource theory of informational nonequilibrium in thermodynamics *Phys. Rep.* **583** 1

[52] Lostaglio M 2019 An introductory review of the resource theory approach to thermodynamics *Rep. Prog. Phys.* **82** 114001

[53] Lostaglio M, Korzekwa K, Jennings D and Rudolph T 2015 Quantum coherence, time-translation symmetry and thermodynamics *Phys. Rev. X* **5** 021001

[54] Lostaglio M, Jennings D and Rudolph T 2015 Description of quantum coherence in thermodynamic processes requires constraints beyond free energy *Nat. Commun.* **6** 6383

[55] Marvian I 2022 Operational interpretation of quantum Fisher information in quantum thermodynamics *Phys. Rev. Lett.* **129** 190502

[56] Marvian I 2020 Coherence distillation machines are impossible in quantum thermodynamics *Nat. Commun.* **11** 25

[57] Bartlett S D and Wiseman H M 2003 Entanglement constrained by superselection rules *Phys. Rev. Lett.* **91** 097903

[58] Petz D 2002 Covariance and Fisher information in quantum mechanics *J. Phys. A: Math. Gen.* **35** 929

[59] Manzano G, Parrondo J M and Landi G T 2022 Non-Abelian quantum transport and thermosqueezing effects *PRX Quantum* **3** 010304

[60] Pires D P, Céleri L C and Soares-Pinto D O 2015 Geometric lower bound for a quantum coherence measure *Phys. Rev. A* **91** 042330

[61] Deffner S and Lutz E 2010 Generalized Clausius inequality for nonequilibrium quantum processes *Phys. Rev. Lett.* **105** 170402

[62] Landi G T and Paternostro M 2021 Irreversible entropy production: from classical to quantum *Rev. Mod. Phys.* **93** 035008

[63] Shiraishi N and Sagawa T 2021 Quantum thermodynamics of correlated-catalytic state conversion at small scale *Phys. Rev. Lett.* **126** 150502

[64] Shiraishi N and Sagawa T 2022 Erratum: Quantum thermodynamics of correlated-catalytic state conversion at small scale [Phys. Rev. Lett. 126, 150502 (2021)] *Phys. Rev. Lett.* **128** 089901

[65] Takagi R and Shiraishi N 2022 Correlation in catalysts enables arbitrary manipulation of quantum coherence *Phys. Rev. Lett.* **128** 240501

[66] Audenaert K M and Eisert J 2005 Continuity bounds on the quantum relative entropy *J. Math. Phys.* **46** 102104

[67] Audenaert K M 2013 On the asymmetry of the relative entropy *J. Math. Phys.* **54** 073506

[68] Allahverdyan A E, Balian R and Nieuwenhuizen T M 2004 Maximal work extraction from finite quantum systems *Europhys. Lett.* **67** 565