

THE SABINA TERAHERTZ/INFRARED BEAMLINE AT SPARC-LAB FACILITY

S. Macis¹, M. Bellaveglia, M. C. Guidi, E. Chiadroni, F. Dipace¹, A. Doria², A. Ghigo,
L. Giannessi³, A. Giribono², A. Petralia², L. Sabbatini, C. Vaccarezza, S. Lupi¹

INFN - Laboratori Nazionali di Frascati, Frascati (Rome), Italy

V. Petrillo, INFN - Sezione di Milano, Segrate (Milan), Italy

¹also at Department of Physics, Sapienza University, Rome, Italy

²also at ENEA, C.R. Frascati, Frascati (Rome), Italy

³also at Elettra-Sincrotrone Trieste, Trieste, Italy

Abstract

Following the EU Terahertz (THz) Road Map [1], high-intensity, ps-long, THz/Infrared (IR) radiation is going to become a fundamental spectroscopy tool for probing and control quantum materials ranging from graphene [2], and Topological Insulators, to strongly correlated oxides [3-6] and novel superconductors [7, 8]. In the framework of the SABINA project a novel THz/IR beamline based on an APPLE-X undulator emission will be developed at the SPARC-Lab facility at LNF-INFN. Light will be propagated from the SPARC-Lab to a new user lab facility nearly 25 m far away from the SPARC laboratory. This beamline will cover a broad spectral region from 3 THz to 30 THz, showing ps- pulses and energy of hundreds of μJ with variable polarization from linear to circular. The corresponding electric fields up to 10 MV/cm, are able to induce non-linear phenomena in many quantum systems. The beamline, open to user experiments, will be equipped with a 5 T magnetic cryostat, and will be synchronized with a fs laser for THz/IR pump, VIS/UV probe experiment.

INTRODUCTION

The spectacular advancements observed in the last decade leads to an increasing interest to terahertz (THz) technology in the efforts to harness the power of the thermal radiation in the region from ~ 50 GHz to 10 THz ($1.5 - 350 \text{ cm}^{-1}$, $7 \text{ mm} - 15 \mu\text{m}$ wavelength, $0.2 - 45 \text{ meV}$) [1, 9-11]. The THz region of the electromagnetic spectrum is a frontier area for research in Physics, Chemistry, Biology, Materials Science and Biomedicine. Indeed, small molecules rotate at THz frequencies; biologically-important collective modes of proteins, DNA and RNA vibrate at THz frequencies; frustrated rotations and collective modes cause polar liquids such as water to absorb at THz frequencies; electrons in semiconductors and their nanostructures resonate at THz frequencies; electrons in highly-excited atomic Rydberg states orbit at THz frequencies; superconducting energy gaps fall at THz frequencies; gaseous and solid-state plasmas also oscillate at THz frequencies so that this radiation can be used to study and control an extraordinary vast number of fundamental systems and phenomena [1, 11-15]. The roadmap for the development of THz technologies [1, 16, 17] considers applications in the fields of outdoor and indoor communications, security, drug detection, biometrics, food quality control,



agriculture, medicine, semiconductors, air pollution, etc. Their exploitation and realization demands high-power compact THz sources, more sensitive detectors, and more functional integrated THz systems. A further important use of THz radiation concerns the excitation and control of quantum materials. These systems which show quantum macroscopic phenomena at room temperature, are characterized by excitations resonating in the THz and Infrared (IR) frequencies [18-20].

Radiation sources of high quality in the THz region of the e.m. spectrum have been scarce [21], but this “THz gap”, after continuous research efforts, has been filled by a wide range of new technologies ranging from accelerated relativistic electrons [22, 23], to high-power femtosecond laser-based sources [24, 25] and Quantum Cascade Lasers [26]. Thus, THz radiation is now available in both CW and pulsed form, down to single-cycles, with peak powers up to tens of MW [27] and several THz facilities are worldwide distributed for fundamental experiments, users’ applications and industrial R&D.

However, a real bridge between THz and IR radiation is not complete in particular for what concerns high-intense ps pulsed beams. Previous discussed THz sources indeed cover a broad spectral range up to 10 THz, while IR sources, often based on the Difference Frequency Generation mechanism, reach, 20 THz at low-frequency. In this paper, we will discuss a new THz/IR source based on the Self Amplified Spontaneous Emission (SASE) mechanism from a relativistic electronic beam from the Free-Electron Facility SPARC at LNF-INFN of Frascati, Italy. This source will be extremely competitive in the international scenario and will be open for national and international users.

DISCUSSION

The Sabina beamline (Source of Advanced Beam Imaging for Novel Applications) is based on an Apple-X undulator emission of quasi-monochromatic light, with sub-picosecond photon pulses of high intensity (hundreds of μJ each), in the frequency range between 3 THz ($100 \mu\text{m}$) to 30 THz ($10 \mu\text{m}$). Polarization can be changed from linear to elliptical to circular. The intensity per pulse corresponds to associate electric fields up to 10 MV/cm, which are useful to perform non-linear and pump-probe spectroscopy measures.



- [1] S. S. Dhillon *et al.*, “The 2017 terahertz science and technology roadmap”, *J. Phys. D: Appl. Phys.*, vol. 50, no. 4, p. 043001, Jan. 2017. doi:10.1088/1361-6463/50/4/043001
- [2] E. Tamburri *et al.*, “Shungite Carbon as Unexpected Natural Source of Few-Layer Graphene Platelets in a Low Oxidation State”, *Inorg. Chem.*, vol. 57, no. 14, p. 8487-8498, Jul. 2018. doi:10.1021/acs.inorgchem.8b01164
- [3] F. Giorgianni, J. Sakai and S. Lupi, “Overcoming the thermal regime for the electric-field driven Mott transition in vanadium sesquioxide”, *Nat. Commun.*, vol. 10, p. 1159, Mar. 2019. doi:10.1038/s41467-019-09137-6
- [4] S. Macis *et al.*, “Angular Dependence of Copper Surface Damage Induced by an Intense Coherent THz Radiation Beam”, *Condens. Matter.*, vol. 5, p. 16, Mar. 2020. doi:10.3390/condmat5010016
- [5] S. Macis *et al.*, “MoO₃ films grown on polycrystalline Cu: Morphological, structural, and electronic properties”, *J. Vac. Sci. Technol.*, vol. 37, p. 021513, Jan. 2019. doi:10.1116/1.5078794

- [6] A. D'Elia *et al.*, "Interplay among work function, electronic structure and stoichiometry in nanostructured VOx films", *Phys. Chem. Chem. Phys.*, vol. 22, p. 6282-6290, Feb. 2020. doi:10.1039/D0CP000216J
- [7] F. Giorgianni *et al.*, "Strong nonlinear terahertz response induced by Dirac surface states in Bi₂Se₃ topological insulator", *Nat. Commun.*, vol. 7, p. 11421, Jan. 2017. doi:10.1088/1361-6463/50/4/043001
- [8] P. Di Pietro *et al.*, "Terahertz Tuning of Dirac Plasmons in Bi₂Se₃ Topological Insulator", *Phys. Rev. Lett.*, vol. 124, p. 226403, Jun. 2020. doi:10.1103/PhysRevLett.124.226403
- [9] Y. Kawano, "Terahertz waves: a tool for condensed matter, the life sciences and astronomy", *Contemp. Phys.*, vol. 54, p. 143-165, Sep. 2013. doi:10.1080/00107514.2013.817194
- [10] M. Tonouchi, "Cutting-edge terahertz technology", *Nat. Photonics*, vol. 1, p. 97-105, Feb. 2007. doi:10.1038/nphoton.2007.3
- [11] D. M. Mittelman, "Perspective: Terahertz science and technology", *J. Appl. Phys.*, vol. 122, p. 230901, Sep. 2013. doi:10.1063/1.5007683
- [12] "Terahertz optics taking off", *Nat. Photonics*, vol. 7, p. 665, Sep. 2013. doi:10.1038/nphoton.2013.239
- [13] H. B. Wang, P. H. Wu, and T. Yamashita, "Responses of Intrinsic Josephson Junctions in High Tc Superconductors", *Phys. Rev. Lett.*, vol. 87, p. 107002, Aug. 2001. doi:10.1103/PhysRevLett.87.107002
- [14] P. Innocenzi *et al.*, "Application of Terahertz spectroscopy to time-dependent chemical-physical phenomena", *J. Phys. Chem. A*, vol. 113, pp. 9418-9423, Aug. 2009. doi:10.1021/jp902502z
- [15] O. Limaj *et al.*, "Superconductivity-Induced Transparency in Terahertz Metamaterials", *ACS Phot.*, vol. 1, no. 7, pp. 570-575, Jun. 2014. doi:10.1021/ph500104k
- [16] X. C. Zhang, A. Shkurinov, and Y. Zhang, "Extreme terahertz science", *Nat. Photonics*, vol. 11, pp. 16-18, Jan. 2017. doi:10.1038/nphoton.2016.249
- [17] J. Ma *et al.*, "Security and eavesdropping in terahertz wireless links", *Nature*, vol. 563, pp. 89-96, Oct. 2018. doi:10.1038/s41586-018-0609-x
- [18] V. Galstyan *et al.*, "A novel approach for green synthesis of WO₃ nanomaterials and their highly selective chemical sensing properties", *J. Mater. Chem. A*, vol. 8, pp. 20373-20385, Aug. 2020. doi:10.1039/D0TA06418A
- [19] M. Mitrano *et al.*, "Possible light-induced superconductivity in K₃C₆₀ at high temperature", *Nature*, vol. 530, no. 7591, pp. 461-464, Feb. 2016. doi:10.1038/nature16522
- [20] P. Di Pietro *et al.*, "Observation of Dirac plasmons in a topological insulator", *Nat. Nanotech.*, vol. 8, pp. 556-560, Jul. 2013. doi:10.1038/nnano.2013.134
- [21] G. P. Gallerano and S. G. Biedron, "Overview of Terahertz Radiation Sources", in *Proc. 26th Int. Free Electron Laser Conf. & 11th FEL Users Workshop (FEL'04)*, Trieste, Italy, Aug.-Sep. 2004, paper FRBIS02, pp. 216-221.
- [22] E. Chiadroni *et al.*, "The SPARC linear accelerator based terahertz source", *Appl. Phys. Lett.*, vol. 102, no. 9, p. 094101, Feb. 2013. doi:10.1063/1.4794014
- [23] A. Perucchi, S. Di Mitri, G. Penco, E. Allaria, and S. Lupi, "The TeraFERMI terahertz source at the seeded FERMI free-electron-laser facility", *Rev. Sci. Instrum.*, vol. 84, no. 2, p. 022702, Feb. 2013. doi:10.1063/1.4790428
- [24] J. A. Fülöp, S. Tzortzakis, and T. Kampfrath, "Laser-Driven Strong-Field Terahertz Sources", *Adv. Opt. Mater.*, vol. 8, no. 3, p. 1900681, Dec. 2019. doi:10.1002/adom.201900681
- [25] B. S. Williams, "Terahertz quantum-cascade lasers", *Nat. Photonics*, vol. 1, pp. 517-525, Sep. 2007. doi:10.1038/nphoton.2007.166
- [26] M. S. Vitiello and A. Tredicucci, "Physics and technology of Terahertz quantum cascade lasers", *Adv. Phys.: X*, vol. 6, no. 1, p. 1893809, Jan. 2021. doi:10.1080/23746149.2021.1893809
- [27] A. Irizawa *et al.*, "Spatially Resolved Spectral Imaging by A THz-FEL", *Condens. Matter*, vol. 5, no. 2, p. 38, Jun. 2020. doi:10.3390/condmat5020038