

# New search for a sterile neutrino at MicroBooNE with BNB and NuMI beams

FERMILAB-POSTER-24-0113-V


Sergey Martynenko, Brookhaven National Laboratory, smartynen@bnl.gov

Xiangpan Ji, Nankai University, jixp@mail.nankai.edu.cn


On behalf of the MicroBooNE Collaboration



## MicroBooNE Experiment



- A 85 ton active-mass liquid-argon TPC
- Two neutrino beams: BNB (on-axis) and NuMI (off-axis)
- On-axis Booster Neutrino Beam (BNB) at a baseline of  $\sim 470$  m with mean neutrino energy at 800 MeV
- Off-axis Neutrino from the Main Injector (NuMI) beam at a baseline of  $\sim 680$  m with neutrinos up to a few GeV



- Beam intrinsic muon neutrinos ( $\nu_\mu$ ) and electron neutrinos ( $\nu_e$ ): significant difference in the  $\nu_\mu/\nu_e$  ratio in BNB and NuMI [1,2]

## Sterile Neutrino Search

- The existence of a light eV-scale sterile neutrino has been postulated to explain several experimental anomalies.
- 3 active + 1 sterile neutrino framework (3+1 oscillation)

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \delta_{\alpha\beta} + (-1)^{\delta_{\alpha\beta}} \cdot \sin^2 2\theta_{\alpha\beta} \cdot \sin^2 \left( \frac{\Delta m_{41}^2 L}{4E} \right)$$

$$\begin{aligned} \sin^2 2\theta_{ee} &= \sin^2 2\theta_{14} \\ \sin^2 2\theta_{\mu\mu} &= 4 \cos^2 \theta_{14} \sin^2 \theta_{24} (1 - \cos^2 \theta_{14} \sin^2 \theta_{24}) \\ \sin^2 2\theta_{\mu e} &= \sin^2 2\theta_{14} \sin^2 \theta_{24} \\ \sin^2 2\theta_{e s} &= \sin^2 2\theta_{14} \cos^2 \theta_{24} \cos^2 \theta_{34} \\ \sin^2 2\theta_{\mu s} &= \cos^4 \theta_{14} \sin^2 2\theta_{24} \cos^2 \theta_{34} \end{aligned}$$


This analysis involves three independent oscillation parameters ( $\Delta m_{14}^2$ ,  $\sin^2 \theta_{14}$ ,  $\sin^2 \theta_{24}$ ), fixing  $\theta_{34}$  at zero [3].

## Cancellation between $\nu_e$ disappearance and appearance oscillations


- Cancellation between  $\nu_e$  disappearance and  $\nu_\mu$  to  $\nu_e$  appearance oscillations leads to a degeneracy in the oscillation parameters. Expected degeneracy to be at  $\sin^2 \theta_{24} \sim 0.005$  for the BNB.
- The degeneracy can be mitigated by using both BNB and NuMI because of different beam intrinsic  $\nu_e/\nu_\mu$  ratios.

$$N_{\nu_e} = N_{\text{intrinsic } \nu_e} \cdot P_{\nu_e \rightarrow \nu_e} + N_{\text{intrinsic } \nu_\mu} \cdot P_{\nu_\mu \rightarrow \nu_e} \quad \text{where } R_{\nu_\mu/\nu_e} \text{ beam intrinsic } \nu_\mu \text{ and } \nu_e$$

$$= N_{\text{intrinsic } \nu_e} \cdot [1 + (R_{\nu_\mu/\nu_e} \cdot \sin^2 \theta_{24} - 1) \cdot \sin^2 2\theta_{14} \cdot \sin^2 \Delta_{41}] \text{ ratio}$$





## Results of 3+1 oscillation analysis



- The best-fit value from BNB data on the 7 channels is:

$$(\Delta m_{14}^2, \sin^2 \theta_{14}, \sin^2 \theta_{24}) = (1.295 \text{ eV}^2, 0.936, 0), \text{ with } \chi^2/NDF = 86.62/179$$

- The BNB result is consistent with the 3-neutrino hypothesis at  $0.80\sigma$ . Exclusion limits are placed on the sterile neutrino parameter space
- The addition of NuMI alongside BNB both increases the overall sensitivity and breaks the degeneracy considerably



[1] Abratenko, P., et al. "First constraints on light sterile neutrino oscillations from combined appearance and disappearance searches with the MicroBooNE detector." *Physical review letters* 130.1 (2023)

[2] Supporting Public Notes: MICROBOONE-NOTE 1129-PUB and MICROBOONE-NOTE1132-PUB

[3] Aartsen, M. G., et al. "Searching for eV-scale sterile neutrinos with eight years of atmospheric neutrinos at the IceCube Neutrino Telescope." *Physical Review D* 102.5 (2020)

\*All the data or MC samples correspond to BNB run1-3 period with  $6.4 \times 10^{20}$  POT, NuMI run1-3 period with  $1.054 \times 10^{21}$  POT